

345a Pregnancy Induced Hypertension

Kansas Risk Factor Summary

Risk Factor Code	Kansas Risk Factor Title	High Risk	Auto-Assigned	Based on MD Diagnosis	Category and Priority					USDA Revised Date
					PG	BF	PP	I	C	
345a	Pregnancy Induced Hypertension		X	X	1					05/19

Kansas Risk Factor Definition

Presence of pregnancy induced hypertension diagnosed by a physician as self-reported by client; or as reported or documented by a physician, or someone working under physician's orders. The term "pregnancy-induced hypertension" includes gestational hypertension, preeclampsia and eclampsia.

- Gestational Hypertension – Onset of hypertension during pregnancy, usually after 20 weeks gestation, and without proteinuria. It usually resolves after delivery but does increase the risk of developing chronic hypertension.
- Preeclampsia – Onset of hypertension during pregnancy, typically with proteinuria, and usually after 20 weeks gestation.
- Eclampsia – Presence of new-onset grand mal seizures in a woman with preeclampsia.

KWIC will auto-assign if "Yes" is marked for the KWIC Health Interview question "Did you develop high blood pressure or pre-high blood pressure ($\geq 120/80$) during this pregnancy including pregnancy induced hypertension?"

[USDA Justification](#)

Note: USDA Justification is provided because it explains nicely why the risk is important. However, you must use the Kansas risk factor names and definitions which may differ slightly from the USDA document.