

Table 4A
2008 Pediatric Nutrition Surveillance (1)
Kansas
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years (2)

Comparison of Racial and Ethnic Distribution by Clinic								
Clinic	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	Multiple Races %	All Other / Unknown %
Allen	363	87.3	4.1	3.9	0.6	0.3	3.9	0.0
Anderson	222	88.3	1.8	6.3	0.0	1.4	2.3	0.0
Atchison	482	82.6	9.5	2.1	0.6	0.0	5.2	0.0
Barber-Kio	8	*	*	*	*	*	*	*
Barber-Med	105	91.4	3.8	3.8	0.0	1.0	0.0	0.0
Barton	1,014	59.0	2.7	36.4	0.1	0.2	1.7	0.0
Bourbon	499	86.4	5.4	4.0	0.4	0.2	3.6	0.0
Brown	414	73.9	2.2	5.3	15.5	0.2	2.9	0.0
Butler	1,023	88.0	2.7	6.4	0.4	0.0	2.5	0.0
Chase	52	*	*	*	*	*	*	*
Chautauqua	133	86.5	0.8	9.8	1.5	0.0	1.5	0.0
Cherokee	773	87.8	1.8	4.7	1.2	1.2	3.4	0.0
Cheyenne	1	*	*	*	*	*	*	*
Clark	31	*	*	*	*	*	*	*
Clay	258	94.6	0.8	1.9	1.6	0.8	0.4	0.0
Cloud	309	90.0	1.3	8.1	0.3	0.3	0.0	0.0
Coffey	99	*	*	*	*	*	*	*
Comanche	22	*	*	*	*	*	*	*
Cowley-Ark	802	64.7	2.9	29.9	1.0	0.1	1.4	0.0

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 4A
2008 Pediatric Nutrition Surveillance (1)
Kansas
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years (2)

Comparison of Racial and Ethnic Distribution by Clinic								
Clinic	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	Multiple Races %	All Other / Unknown %
Cowley-Win	618	80.4	6.0	8.1	0.5	1.8	3.2	0.0
Crawford-G	4	*	*	*	*	*	*	*
Crawford-P	1,340	79.0	4.6	11.6	0.6	1.0	3.3	0.0
Decatur	43	*	*	*	*	*	*	*
DickinsonA	382	87.4	1.3	8.9	0.3	0.0	2.1	0.0
DickinsonH	68	*	*	*	*	*	*	*
Doniphan	146	84.9	6.2	5.5	1.4	0.0	2.1	0.0
Douglas	1,630	52.2	12.1	14.8	8.6	5.1	7.2	0.0
Edwards	104	51.0	0.0	49.0	0.0	0.0	0.0	0.0
Elk	79	*	*	*	*	*	*	*
Ellis	490	76.5	2.7	18.2	0.8	0.2	1.6	0.0
Ellsworth	140	88.6	1.4	7.9	0.0	0.0	2.1	0.0
Finney	2,754	22.7	1.3	72.9	0.2	2.1	0.7	0.0
Ford	2,421	16.5	1.0	81.9	0.0	0.4	0.2	0.0
Franklin	649	85.4	2.6	7.1	0.5	0.0	4.5	0.0
Geary	1,087	43.5	23.9	22.2	0.4	2.6	7.5	0.0
Gove	53	*	*	*	*	*	*	*
Graham	66	*	*	*	*	*	*	*
Grant	314	19.7	0.0	79.3	0.6	0.3	0.0	0.0

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 4A
2008 Pediatric Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Comparison of Racial and Ethnic Distribution by Clinic								
Clinic	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	Multiple Races %	All Other / Unknown %
Gray	250	61.6	1.6	36.0	0.0	0.0	0.8	0.0
Greeley	34	*	*	*	*	*	*	*
Greenwood	208	91.3	1.4	3.8	0.5	0.0	2.9	0.0
Hamilton	96	*	*	*	*	*	*	*
Harper	213	80.8	0.0	17.8	0.9	0.0	0.5	0.0
Harvey	1,059	59.8	3.1	32.5	1.0	0.3	3.3	0.0
Haskell	121	53.7	0.8	45.5	0.0	0.0	0.0	0.0
Hodgeman	51	*	*	*	*	*	*	*
Jackson	357	71.4	3.9	8.1	11.8	0.0	4.8	0.0
Jefferson	261	93.1	3.1	2.3	0.4	0.0	1.1	0.0
Jewell	57	*	*	*	*	*	*	*
Johnson-M	3,274	34.9	15.9	43.3	0.4	2.5	3.0	0.0
Johnson-O	3,825	37.2	12.5	43.8	0.3	2.5	3.6	0.0
Kearny	127	39.4	0.8	59.8	0.0	0.0	0.0	0.0
Kingman	219	92.2	1.8	4.6	0.5	0.9	0.0	0.0
Kiowa	53	*	*	*	*	*	*	*
Labette	649	76.9	10.8	6.8	1.2	0.3	4.0	0.0
Lane	64	*	*	*	*	*	*	*
Leavenwort	1,176	55.2	21.0	14.1	0.4	0.9	8.4	0.0

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 4A
2008 Pediatric Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Comparison of Racial and Ethnic Distribution by Clinic								
Clinic	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	Multiple Races %	All Other / Unknown %
Lincoln	92	*	*	*	*	*	*	*
Linn	260	94.2	1.2	3.5	0.0	0.0	1.2	0.0
Logan	63	*	*	*	*	*	*	*
Lyon	1,318	47.9	4.6	45.9	0.1	0.5	1.1	0.0
Marion	204	86.8	0.0	9.3	0.5	1.5	2.0	0.0
Marshall	232	86.6	1.7	4.7	1.7	0.0	5.2	0.0
Mc Pherson	547	82.3	3.1	10.4	1.3	0.7	2.2	0.0
Meade	169	56.2	0.6	42.6	0.0	0.0	0.6	0.0
Miami	492	82.9	5.1	8.5	0.2	0.4	2.8	0.0
Mitchell	143	93.7	0.7	4.2	0.0	0.0	1.4	0.0
Montgome-C	601	63.9	14.5	12.8	4.2	0.5	4.2	0.0
Montgome-I	669	76.1	8.4	9.6	1.3	0.7	3.9	0.0
Morris	129	84.5	1.6	10.1	0.8	0.0	3.1	0.0
Nemaha	103	91.3	1.9	4.9	0.0	0.0	1.9	0.0
Neosho-Ch	526	83.3	3.0	12.0	0.6	0.4	0.8	0.0
Neosho-Er	47	*	*	*	*	*	*	*
Ness	48	*	*	*	*	*	*	*
Norton	123	87.0	2.4	8.1	0.8	0.8	0.8	0.0
Osage	324	94.1	1.9	1.9	0.3	0.0	1.9	0.0

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 4A
2008 Pediatric Nutrition Surveillance (1)
Kansas
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years (2)

Comparison of Racial and Ethnic Distribution by Clinic								
Clinic	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	Multiple Races %	All Other / Unknown %
Osborne	96	*	*	*	*	*	*	*
Ottawa	94	*	*	*	*	*	*	*
Pawnee	205	74.6	3.4	19.0	0.5	0.0	2.4	0.0
Phillips	148	79.1	2.7	14.2	0.7	0.7	2.7	0.0
Pottawat-S	98	*	*	*	*	*	*	*
Pottawat-W	94	*	*	*	*	*	*	*
Pratt	273	81.7	0.7	15.4	1.1	0.0	1.1	0.0
Rawlins	54	*	*	*	*	*	*	*
Reno	1,720	73.4	4.9	17.8	0.6	0.3	3.0	0.0
Republic	138	87.0	2.9	9.4	0.0	0.0	0.7	0.0
Rice	276	73.9	1.1	23.9	0.0	0.0	1.1	0.0
Riley-Ft	1,794	68.2	11.3	14.7	0.8	2.0	3.0	0.0
Riley-Manh	1,299	64.4	12.8	12.6	0.5	4.2	5.5	0.0
Rooks	172	91.3	2.3	4.7	0.6	0.0	1.2	0.0
Rush	53	*	*	*	*	*	*	*
Russell	169	85.2	8.3	5.3	0.6	0.0	0.6	0.0
Saline	1,726	59.0	5.4	27.3	0.2	1.9	6.2	0.0
Scott	192	55.7	2.1	41.7	0.0	0.5	0.0	0.0
Sedgwick-C	4,315	32.6	13.8	42.1	1.1	5.3	5.1	0.0

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 4A
2008 Pediatric Nutrition Surveillance (1)
Kansas
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years (2)

Comparison of Racial and Ethnic Distribution by Clinic								
Clinic	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	Multiple Races %	All Other / Unknown %
Sedgwick-M	7,335	24.5	24.5	42.7	1.8	2.1	4.4	0.0
Sedgwick-S	4,560	57.3	10.5	23.6	1.6	1.9	5.0	0.0
Seneca	52	*	*	*	*	*	*	*
Seward	1,807	11.8	1.9	82.7	2.3	0.8	0.4	0.0
ShawneeM&I	290	21.0	11.7	59.3	1.4	0.3	6.2	0.0
ShawneeMa	4,966	43.1	17.9	31.6	1.4	0.3	5.7	0.0
Sheridan	41	*	*	*	*	*	*	*
Sherman	296	61.5	0.7	33.4	1.0	0.0	3.4	0.0
Smith	93	*	*	*	*	*	*	*
Stafford	174	62.6	0.6	36.2	0.0	0.0	0.6	0.0
Stanton	95	*	*	*	*	*	*	*
Stevens	322	37.3	0.0	61.5	0.0	0.3	0.9	0.0
Sumner	576	85.9	2.1	9.4	0.5	0.2	1.9	0.0
Thomas	214	76.6	1.9	18.7	0.5	0.5	1.9	0.0
Trego	74	*	*	*	*	*	*	*
Wabaunsee	88	*	*	*	*	*	*	*
Wallace	3	*	*	*	*	*	*	*
Washington	113	79.6	4.4	12.4	0.0	0.9	2.7	0.0
Wichita	83	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 4A
2008 Pediatric Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Comparison of Racial and Ethnic Distribution by Clinic								
Clinic	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	Multiple Races %	All Other / Unknown %
Wilson	347	90.5	2.9	5.5	0.0	0.3	0.9	0.0
Woodson	27	*	*	*	*	*	*	*
Wyandotte	7,900	19.7	28.6	47.8	0.2	2.2	1.5	0.0
Unknown Clinics	2	*	*	*	*	*	*	*
Kansas	79,556	49.1	11.6	33.2	1.1	1.6	3.3	0.0
Nation (Prior Year)	7,995,600	31.6	19.4	40.9	1.0	2.5	1.8	2.7

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 5A
2008 Pediatric Nutrition Surveillance (1)
Kansas
Comparison of Age Distribution by Clinic
Children Aged < 5 Years (2)

Comparison of Age Distribution by Clinic				
Clinic	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
Allen	363	37.7	24.2	38.0
Anderson	222	28.4	26.6	45.0
Atchison	482	28.0	28.2	43.8
Barber-Kio	8	*	*	*
Barber-Med	105	38.1	26.7	35.2
Barton	1,014	31.9	24.5	43.7
Bourbon	499	34.9	23.8	41.3
Brown	414	28.7	24.2	47.1
Butler	1,023	34.9	21.6	43.5
Chase	52	*	*	*
Chautauqua	133	32.3	16.5	51.1
Cherokee	773	33.6	22.5	43.9
Cheyenne	1	*	*	*
Clark	31	*	*	*
Clay	258	30.2	22.5	47.3
Cloud	309	27.2	28.5	44.3
Coffey	99	*	*	*
Comanche	22	*	*	*

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Records rejected if date of birth and date of visit unknown.
* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 5A
2008 Pediatric Nutrition Surveillance (1)
Kansas
Comparison of Age Distribution by Clinic
Children Aged < 5 Years (2)

Comparison of Age Distribution by Clinic				
Clinic	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
Cowley-Ark	802	31.4	22.7	45.9
Cowley-Win	618	29.3	23.3	47.4
Crawford-G	4	*	*	*
Crawford-P	1,340	34.0	22.9	43.1
Decatur	43	*	*	*
DickinsonA	382	33.0	19.9	47.1
DickinsonH	68	*	*	*
Doniphan	146	34.2	24.0	41.8
Douglas	1,630	36.1	24.0	39.9
Edwards	104	21.2	26.0	52.9
Elk	79	*	*	*
Ellis	490	35.1	20.6	44.3
Ellsworth	140	25.0	24.3	50.7
Finney	2,754	30.4	22.5	47.1
Ford	2,421	28.9	23.6	47.5
Franklin	649	32.2	23.0	44.8
Geary	1,087	36.7	24.5	38.8
Gove	53	*	*	*

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Records rejected if date of birth and date of visit unknown.
* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 5A
2008 Pediatric Nutrition Surveillance (1)
Kansas
Comparison of Age Distribution by Clinic
Children Aged < 5 Years (2)

Comparison of Age Distribution by Clinic				
Clinic	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
Graham	66	*	*	*
Grant	314	26.1	25.2	48.7
Gray	250	23.6	25.2	51.2
Greeley	34	*	*	*
Greenwood	208	23.6	21.2	55.3
Hamilton	96	*	*	*
Harper	213	27.2	24.9	47.9
Harvey	1,059	29.6	23.7	46.7
Haskell	121	22.3	21.5	56.2
Hodgeman	51	*	*	*
Jackson	357	26.3	24.4	49.3
Jefferson	261	28.4	27.6	44.1
Jewell	57	*	*	*
Johnson-M	3,274	33.2	24.7	42.1
Johnson-O	3,825	32.7	23.6	43.7
Kearny	127	24.4	20.5	55.1
Kingman	219	28.3	21.9	49.8
Kiowa	53	*	*	*

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Records rejected if date of birth and date of visit unknown.
* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 5A
2008 Pediatric Nutrition Surveillance (1)
Kansas
Comparison of Age Distribution by Clinic
Children Aged < 5 Years (2)

Comparison of Age Distribution by Clinic				
Clinic	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
Labette	649	31.7	28.0	40.2
Lane	64	*	*	*
Leavenwort	1,176	35.7	23.5	40.8
Lincoln	92	*	*	*
Linn	260	26.9	31.9	41.2
Logan	63	*	*	*
Lyon	1,318	31.6	22.8	45.7
Marion	204	27.9	26.0	46.1
Marshall	232	32.8	19.8	47.4
Mc Pherson	547	32.7	24.1	43.1
Meade	169	33.1	18.9	47.9
Miami	492	34.8	23.8	41.5
Mitchell	143	32.9	24.5	42.7
Montgome-C	601	28.8	24.0	47.3
Montgome-I	669	32.1	23.8	44.1
Morris	129	27.1	27.9	45.0
Nemaha	103	29.1	25.2	45.6
Neosho-Ch	526	34.2	20.9	44.9

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Records rejected if date of birth and date of visit unknown.
* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 5A
2008 Pediatric Nutrition Surveillance (1)
Kansas
Comparison of Age Distribution by Clinic
Children Aged < 5 Years (2)

Comparison of Age Distribution by Clinic				
Clinic	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
Neosho-Er	47	*	*	*
Ness	48	*	*	*
Norton	123	26.0	22.0	52.0
Osage	324	34.3	21.6	44.1
Osborne	96	*	*	*
Ottawa	94	*	*	*
Pawnee	205	28.3	24.4	47.3
Phillips	148	24.3	30.4	45.3
Pottawat-S	98	*	*	*
Pottawat-W	94	*	*	*
Pratt	273	28.9	27.8	43.2
Rawlins	54	*	*	*
Reno	1,720	34.1	23.2	42.7
Republic	138	34.8	19.6	45.7
Rice	276	28.6	23.2	48.2
Riley-Ft	1,794	31.5	26.8	41.6
Riley-Manh	1,299	36.2	22.5	41.3
Rooks	172	24.4	27.3	48.3

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Records rejected if date of birth and date of visit unknown.
* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 5A
2008 Pediatric Nutrition Surveillance (1)
Kansas
Comparison of Age Distribution by Clinic
Children Aged < 5 Years (2)

Comparison of Age Distribution by Clinic				
Clinic	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
Rush	53	*	*	*
Russell	169	29.6	29.0	41.4
Saline	1,726	34.1	24.0	41.9
Scott	192	21.9	27.6	50.5
Sedgwick-C	4,315	30.4	25.1	44.5
Sedgwick-M	7,335	34.5	22.6	43.0
Sedgwick-S	4,560	31.4	24.7	43.8
Seneca	52	*	*	*
Seward	1,807	33.1	22.6	44.3
ShawneeM&I	290	55.5	11.7	32.8
ShawneeMa	4,966	30.5	23.5	46.0
Sheridan	41	*	*	*
Sherman	296	29.7	22.6	47.6
Smith	93	*	*	*
Stafford	174	24.7	23.0	52.3
Stanton	95	*	*	*
Stevens	322	23.6	26.1	50.3
Sumner	576	32.5	24.7	42.9

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Records rejected if date of birth and date of visit unknown.
* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 5A
2008 Pediatric Nutrition Surveillance (1)
Kansas
Comparison of Age Distribution by Clinic
Children Aged < 5 Years (2)

Comparison of Age Distribution by Clinic				
Clinic	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
Thomas	214	28.0	24.3	47.7
Trego	74	*	*	*
Wabaunsee	88	*	*	*
Wallace	3	*	*	*
Washington	113	30.1	21.2	48.7
Wichita	83	*	*	*
Wilson	347	28.0	25.4	46.7
Woodson	27	*	*	*
Wyandotte	7,900	31.3	23.8	44.9
Unknown Clinics	2	*	*	*
Kansas	79,556	31.9	23.8	44.4
Nation (Prior Year)	7,995,600	37.6	22.1	40.3

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Records rejected if date of birth and date of visit unknown.
* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2008 Pediatric Nutrition Surveillance (1)
Kansas
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Clinic (3,4)												
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Obese (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Allen	110	7.3 (18)	6.4 (21)	356	9.3 (71)	1.1 (5)	17.1 (74)	132	20.5 (50)	13.6 (34)	222	6.3 (13)
Anderson	50	* *	* *	211	6.6 (40)	3.3 (50)	11.8 (33)	96	* *	* *	148	15.5 (56)
Atchison	105	5.7 (5)	5.7 (15)	467	4.7 (16)	4.7 (64)	12.8 (47)	199	17.6 (33)	14.1 (41)	319	11.9 (44)
Barber-Kio	1	* *	* *	8	* *	* *	* *	5	* *	* *	5	* *
Barber-Med	32	* *	* *	105	10.5 (76)	1.9 (21)	13.3 (54)	33	* *	* *	57	* *
Barton	256	8.2 (25)	5.5 (12)	975	7.0 (45)	3.2 (47)	9.1 (12)	430	14.7 (12)	9.1 (9)	635	11.7 (41)
Bourbon	123	7.3 (18)	5.7 (15)	485	8.5 (67)	1.2 (8)	15.7 (70)	206	18.4 (39)	17.0 (51)	318	12.3 (45)
Brown	86	* *	* *	386	3.1 (7)	3.1 (45)	11.7 (32)	187	16.0 (19)	11.8 (19)	257	13.2 (50)
Butler	288	10.8 (36)	5.9 (18)	995	10.6 (77)	3.6 (54)	11.0 (27)	435	13.6 (9)	9.9 (12)	634	20.5 (64)
Chase	13	* *	* *	51	* *	* *	* *	26	* *	* *	29	* *
Chautauqua	28	* *	* *	128	7.0 (45)	2.3 (29)	21.1 (78)	66	* *	* *	91	* *

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2008 Pediatric Nutrition Surveillance (1)
Kansas
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Clinic (3,4)												
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Obese (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Cherokee	200	7.5 (20)	5.5 (12)	750	6.5 (37)	4.3 (60)	12.3 (41)	324	18.2 (37)	16.0 (48)	497	7.8 (21)
Cheyenne	0	* *	* *	1	* *	* *	* *	1	* *	* *	1	* *
Clark	9	* *	* *	27	* *	* *	* *	10	* *	* *	19	* *
Clay	60	* *	* *	253	9.1 (70)	4.3 (60)	20.2 (77)	111	13.5 (8)	21.6 (55)	166	16.3 (59)
Cloud	67	* *	* *	304	7.9 (59)	3.0 (41)	10.9 (26)	138	18.1 (36)	10.9 (14)	214	11.7 (41)
Coffey	25	* *	* *	96	* *	* *	* *	47	* *	* *	62	* *
Comanche	5	* *	* *	22	* *	* *	* *	8	* *	* *	17	* *
Cowley-Ark	205	6.3 (12)	8.3 (36)	776	5.8 (30)	1.8 (20)	13.8 (60)	355	15.8 (16)	12.7 (23)	526	5.1 (8)
Cowley-Win	146	8.2 (25)	6.8 (25)	605	6.6 (40)	1.3 (9)	12.2 (40)	287	19.2 (44)	15.0 (45)	406	11.6 (39)
Crawford-G	1	* *	* *	4	* *	* *	* *	2	* *	* *	3	* *
Crawford-P	336	9.5 (31)	5.4 (10)	1,291	6.8 (43)	1.6 (14)	14.1 (62)	568	16.0 (19)	12.9 (26)	831	8.2 (23)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2008 Pediatric Nutrition Surveillance (1)
Kansas
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Clinic (3,4)												
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Obese (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Decatur	7	* *	* *	41	* *	* *	* *	21	* *	* *	32	* *
DickinsonA	85	* *	* *	372	4.8 (17)	5.4 (67)	11.0 (27)	178	7.3 (1)	12.9 (26)	257	3.1 (4)
DickinsonH	11	* *	* *	64	* *	* *	* *	29	* *	* *	46	* *
Doniphan	38	* *	* *	131	15.3 (80)	6.1 (71)	22.9 (79)	57	* *	* *	85	* *
Douglas	438	6.2 (10)	6.8 (25)	1,587	8.3 (64)	1.4 (11)	14.2 (63)	625	18.7 (40)	13.3 (31)	1,058	12.6 (48)
Edwards	21	* *	* *	104	4.8 (17)	1.9 (21)	12.5 (44)	52	* *	* *	71	* *
Elk	14	* *	* *	78	* *	* *	* *	39	* *	* *	62	* *
Ellis	130	10.0 (35)	1.5 (1)	478	7.3 (52)	1.7 (17)	9.2 (13)	215	21.4 (53)	9.8 (11)	293	3.4 (5)
Ellsworth	30	* *	* *	135	11.1 (79)	3.0 (41)	13.3 (54)	67	* *	* *	94	* *
Finney	647	6.0 (8)	7.0 (30)	2,654	5.4 (24)	6.4 (72)	11.6 (31)	1,242	9.4 (4)	7.7 (6)	1,821	25.1 (66)
Ford	591	5.8 (6)	6.3 (19)	2,294	7.6 (54)	1.9 (21)	19.4 (76)	1,107	18.2 (37)	18.1 (54)	1,535	6.5 (14)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2008 Pediatric Nutrition Surveillance (1)
Kansas
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Clinic (3,4)												
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Obese (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- (Rank)	% <95th (Rank)	% ≥95th (Rank)	Number
Franklin	159	6.9 (16)	6.3 (19)	629	7.8 (57)	2.2 (27)	12.4 (42)	273	17.6 (33)	11.0 (16)	404	5.9 (11)
Geary	332	6.0 (8)	4.8 (4)	1,040	8.5 (67)	3.2 (47)	10.3 (19)	411	17.3 (31)	10.9 (14)	656	14.9 (54)
Gove	10	* *	* *	53	* *	* *	* *	26	* *	* *	42	* *
Graham	11	* *	* *	63	* *	* *	* *	31	* *	* *	42	* *
Grant	67	* *	* *	301	5.3 (22)	1.7 (17)	15.3 (69)	150	15.3 (13)	17.3 (52)	217	31.8 (67)
Gray	39	* *	* *	242	1.7 (1)	11.6 (78)	3.3 (1)	125	10.4 (5)	3.2 (2)	176	5.7 (10)
Greeley	10	* *	* *	32	* *	* *	* *	13	* *	* *	14	* *
Greenwood	41	* *	* *	203	7.4 (53)	2.5 (30)	11.8 (33)	107	16.8 (25)	14.0 (40)	147	6.8 (17)
Hamilton	13	* *	* *	91	* *	* *	* *	44	* *	* *	71	* *
Harper	38	* *	* *	202	5.9 (31)	3.0 (41)	15.8 (71)	101	18.8 (41)	13.9 (37)	135	8.9 (26)
Harvey	224	9.8 (33)	5.4 (10)	1,012	7.2 (49)	2.1 (25)	13.0 (50)	489	20.0 (49)	14.1 (41)	714	9.2 (27)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2008 Pediatric Nutrition Surveillance (1)
Kansas
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Clinic (3,4)												
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Obese (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Haskell	24	* *	* *	119	5.0 (20)	0.8 (3)	16.8 (73)	69	* *	* *	75	* *
Hodgeman	12	* *	* *	51	* *	* *	* *	26	* *	* *	38	* *
Jackson	70	* *	* *	348	4.9 (19)	1.1 (5)	14.9 (67)	171	22.2 (55)	17.5 (53)	208	8.2 (23)
Jefferson	64	* *	* *	256	5.9 (31)	5.9 (69)	12.9 (48)	118	22.0 (54)	6.8 (5)	159	18.2 (61)
Jewell	15	* *	* *	54	* *	* *	* *	24	* *	* *	34	* *
Johnson-M	771	9.9 (34)	6.9 (28)	3,105	6.0 (34)	2.2 (27)	12.7 (46)	1,347	16.5 (24)	13.1 (30)	2,053	6.8 (17)
Johnson-O	923	6.5 (14)	6.9 (28)	3,679	5.3 (22)	3.0 (41)	11.9 (35)	1,636	15.9 (18)	12.4 (22)	2,454	7.0 (20)
Kearny	22	* *	* *	123	9.8 (74)	5.7 (68)	11.4 (30)	71	* *	* *	93	* *
Kingman	43	* *	* *	209	4.3 (13)	9.1 (75)	8.1 (9)	104	8.7 (2)	5.8 (3)	139	6.5 (14)
Kiowa	10	* *	* *	52	* *	* *	* *	27	* *	* *	34	* *
Labette	163	5.5 (3)	9.2 (38)	634	6.0 (34)	2.5 (30)	12.6 (45)	261	13.0 (7)	13.0 (28)	425	8.0 (22)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2008 Pediatric Nutrition Surveillance (1)
Kansas
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Clinic (3,4)												
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under-weight (7)	Obese (7)		≥ 2 Yrs Over-weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Lane	13	* *	* *	61	* *	* *	* *	35	* *	* *	49	* *
Leavenwort	308	9.7 (32)	4.2 (3)	1,120	7.9 (59)	2.9 (37)	9.8 (16)	470	19.4 (46)	13.0 (28)	711	15.9 (58)
Lincoln	22	* *	* *	89	* *	* *	* *	39	* *	* *	63	* *
Linn	53	* *	* *	249	9.6 (72)	0.8 (3)	12.9 (48)	106	23.6 (56)	12.3 (21)	174	10.3 (32)
Logan	11	* *	* *	61	* *	* *	* *	35	* *	* *	38	* *
Lyon	324	6.2 (10)	6.8 (25)	1,293	5.0 (20)	2.6 (33)	12.4 (42)	588	17.0 (28)	16.3 (49)	855	5.3 (9)
Marion	43	* *	* *	203	5.9 (31)	0.5 (1)	10.3 (19)	87	* *	* *	146	11.6 (39)
Marshall	55	* *	* *	211	2.8 (3)	17.5 (80)	8.5 (11)	102	8.8 (3)	2.0 (1)	143	8.4 (25)
Mc Pherson	142	4.2 (1)	4.9 (6)	540	6.5 (37)	3.1 (45)	6.1 (4)	240	13.8 (10)	5.8 (3)	346	12.4 (47)
Meade	38	* *	* *	158	4.4 (15)	4.4 (62)	10.8 (24)	78	* *	* *	106	6.6 (16)
Miami	140	8.6 (28)	5.7 (15)	474	9.7 (73)	1.1 (5)	13.1 (52)	202	17.3 (31)	13.9 (37)	299	19.4 (62)

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2008 Pediatric Nutrition Surveillance (1)
Kansas
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Clinic (3,4)												
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Obese (7)		≥ 2 Yrs Over- weight (7)			Number	%
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- (Rank)	% <95th (Rank)	% ≥95th (Rank)	Number
Mitchell	40	* *	* *	143	2.8 (3)	3.5 (53)	8.4 (10)	61	* *	* *	93	* *
Montgome-C	141	7.8 (23)	5.0 (7)	583	7.7 (55)	2.7 (36)	12.0 (37)	273	19.0 (43)	12.8 (24)	396	11.4 (38)
Montgome-I	163	11.7 (38)	3.1 (2)	661	7.9 (59)	2.6 (33)	13.5 (58)	286	16.8 (25)	13.6 (34)	454	10.8 (37)
Morris	29	* *	* *	123	5.7 (29)	1.6 (14)	4.9 (3)	58	* *	* *	85	* *
Nemaha	23	* *	* *	100	4.0 (11)	6.0 (70)	16.0 (72)	50	* *	* *	60	* *
Neosho-Ch	112	8.0 (24)	7.1 (31)	514	8.4 (66)	3.3 (50)	13.2 (53)	223	17.0 (28)	10.8 (13)	321	14.3 (52)
Neosho-Er	11	* *	* *	47	* *	* *	* *	21	* *	* *	33	* *
Ness	8	* *	* *	48	* *	* *	* *	24	* *	* *	32	* *
Norton	21	* *	* *	122	9.8 (74)	7.4 (74)	9.8 (16)	60	* *	* *	85	* *
Osage	85	* *	* *	315	6.7 (42)	2.5 (30)	13.7 (59)	139	20.9 (51)	11.5 (18)	205	1.5 (1)
Osborne	20	* *	* *	92	* *	* *	* *	43	* *	* *	67	* *

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2008 Pediatric Nutrition Surveillance (1)
Kansas
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Clinic (3,4)													
Clinic	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Obese (7)		≥ 2 Yrs Over- weight (7)			(8)		
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- (Rank)	% <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Ottawa	19	* *	* *	90	* *	* *	* *	41	* *	* *	* *	63	* *
Pawnee	43	* *	* *	199	5.5 (25)	4.0 (56)	12.1 (38)	91	* *	* *	* *	147	2.0 (2)
Phillips	27	* *	* *	146	6.2 (36)	2.1 (25)	17.1 (74)	67	* *	* *	* *	107	20.6 (65)
Pottawat-S	10	* *	* *	97	* *	* *	* *	59	* *	* *	* *	78	* *
Pottawat-W	25	* *	* *	90	* *	* *	* *	34	* *	* *	* *	60	* *
Pratt	64	* *	* *	269	7.8 (57)	5.2 (66)	11.9 (35)	116	19.8 (48)	13.8 (36)		183	9.3 (28)
Rawlins	9	* *	* *	54	* *	* *	* *	27	* *	* *	* *	38	* *
Reno	478	6.7 (15)	7.9 (35)	1,701	6.5 (37)	4.1 (58)	8.0 (8)	716	15.4 (14)	11.9 (20)		1,079	5.0 (7)
Republic	33	* *	* *	136	11.0 (78)	1.5 (13)	14.0 (61)	63	* *	* *	* *	89	* *
Rice	60	* *	* *	269	7.1 (47)	0.7 (2)	24.5 (80)	134	16.4 (23)	26.1 (56)		186	11.8 (43)
Riley-Ft	422	7.6 (21)	7.3 (32)	1,747	5.5 (25)	2.9 (37)	9.7 (15)	744	12.8 (6)	8.1 (7)		819	3.8 (6)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2008 Pediatric Nutrition Surveillance (1)
Kansas
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Clinic (3,4)												
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under-weight (7)	Obese (7)	≥ 2 Yrs Over-weight (7)					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Riley-Manh	363	6.3 (12)	5.5 (12)	1,261	5.5 (25)	2.9 (37)	9.8 (16)	532	16.9 (27)	11.1 (17)	835	9.7 (30)
Rooks	33	* *	* *	166	4.2 (12)	3.6 (54)	4.2 (2)	83	* *	* *	117	10.3 (32)
Rush	12	* *	* *	51	* *	* *	* *	26	* *	* *	36	* *
Russell	44	* *	* *	153	3.9 (9)	13.1 (79)	7.2 (5)	65	* *	* *	121	2.5 (3)
Saline	473	9.3 (30)	5.1 (8)	1,673	9.0 (69)	1.4 (11)	15.1 (68)	713	18.8 (41)	15.4 (46)	1,111	6.9 (19)
Scott	28	* *	* *	181	2.8 (3)	9.4 (76)	11.0 (27)	92	* *	* *	130	10.0 (31)
Sedgwick-C	1,084	8.3 (27)	7.6 (33)	4,195	6.8 (43)	1.6 (14)	12.1 (38)	1,868	17.8 (35)	16.3 (49)	2,897	16.6 (60)
Sedgwick-M	1,907	11.4 (37)	6.6 (22)	7,152	8.2 (62)	1.7 (17)	13.0 (50)	3,063	17.1 (30)	14.7 (44)	4,703	13.0 (49)
Sedgwick-S	1,130	7.0 (17)	6.6 (22)	4,477	7.7 (55)	1.9 (21)	14.3 (64)	1,983	19.5 (47)	15.9 (47)	3,000	12.3 (45)
Seneca	12	* *	* *	49	* *	* *	* *	22	* *	* *	35	* *
Seward	437	5.5 (3)	4.8 (4)	1,737	3.8 (8)	4.0 (56)	10.5 (21)	772	16.2 (21)	13.9 (37)	1,074	10.7 (36)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2008 Pediatric Nutrition Surveillance (1)
Kansas
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Clinic (3,4)													
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)			
	Low	High		Short Stature (6)	Under- weight (7)	Obese (7)		≥ 2 Yrs Over- weight (7)					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- (Rank)	% <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
ShawneeM&I	131	4.6 (2)	8.4 (37)	282	2.1 (2)	3.2 (47)	9.6 (14)	98	* *	* *	* *	137	20.4 (63)
ShawneeMa	1,089	7.7 (22)	5.1 (8)	4,806	7.1 (47)	2.6 (33)	14.8 (66)	2,225	19.3 (45)	14.6 (43)		3,211	13.3 (51)
Sheridan	7	* *	* *	40	* *	* *	* *	22	* *	* *	* *	27	* *
Sherman	68	* *	* *	288	8.3 (64)	4.2 (59)	7.6 (6)	142	14.1 (11)	8.5 (8)		185	10.3 (32)
Smith	17	* *	* *	90	* *	* *	* *	44	* *	* *	* *	68	* *
Stafford	34	* *	* *	164	4.3 (13)	6.7 (73)	14.6 (65)	91	* *	* *	* *	124	10.5 (35)
Stanton	23	* *	* *	91	* *	* *	* *	43	* *	* *	* *	59	* *
Stevens	35	* *	* *	307	2.9 (6)	1.3 (9)	13.4 (57)	157	21.0 (52)	13.4 (32)		222	15.8 (57)
Sumner	155	5.8 (6)	7.7 (34)	557	7.2 (49)	2.9 (37)	13.3 (54)	234	16.2 (21)	9.4 (10)		365	15.3 (55)
Thomas	46	* *	* *	208	7.2 (49)	4.8 (65)	7.7 (7)	98	* *	* *	* *	144	14.6 (53)
Trego	11	* *	* *	73	* *	* *	* *	32	* *	* *	* *	53	* *

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2008 Pediatric Nutrition Surveillance (1)
Kansas
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Clinic (3,4)												
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Obese (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Wabaunsee	18	* *	* *	86	* *	* *	* *	38	* *	* *	64	* *
Wallace	1	* *	* *	3	* *	* *	* *	2	* *	* *	2	* *
Washington	25	* *	* *	102	3.9 (9)	10.8 (77)	10.8 (24)	54	* *	* *	84	* *
Wichita	13	* *	* *	80	* *	* *	* *	44	* *	* *	61	* *
Wilson	79	* *	* *	341	8.2 (62)	4.4 (62)	10.6 (22)	156	15.4 (14)	13.5 (33)	232	6.0 (12)
Woodson	2	* *	* *	27	* *	* *	* *	15	* *	* *	21	* *
Wyandotte	1,930	9.1 (29)	6.7 (24)	7,598	5.6 (28)	3.3 (50)	10.6 (22)	3,404	15.8 (16)	12.8 (24)	5,324	9.6 (29)
Unknown Clinics	0	* *	* *	1	* *	* *	* *	0	* *	* *	1	* *
Kansas	19,418	8.1	6.4	76,976	6.7	2.8	12.4	34,352	16.7	13.3	51,194	11.4
Nation (Prior Year)	1,953,242	9.1	6.4	7,526,585	6.2	4.5	13.9	3,058,130	16.4	14.9	4,759,588	15.0

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2008 Pediatric Nutrition Surveillance (1)
Kansas

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
(4,5)

Clinic	Breastfeeding (5)		Exclusive Breastfeeding (6)		≤2 Hours TV Viewing/Day (7)		Smoking in Household							
	Number (Rank)	% Ever Breastfed (Rank)	Number (Rank)	% Breastfed At Least 6 Months (Rank)	Number (Rank)	% Breastfed At Least 12 Months (Rank)	Number (Rank)	% (Rank)	Number (Rank)	% (Rank)				
Allen	113	47.8 (37)	34	* *	61	* *	16	* *	34	* *	146	82.2 (24)	363	48.8 (72)
Anderson	49	* *	24	* *	36	* *	12	* *	23	* *	103	78.6 (38)	216	20.8 (41)
Atchison	104	52.9 (34)	52	* *	92	* *	26	* *	51	* *	214	79.4 (36)	466	29.2 (52)
Barber-Kio	1	* *	1	* *	2	* *	0	* *	1	* *	5	* *	8	* *
Barber-Med	35	* *	14	* *	19	* *	14	* *	14	* *	39	* *	105	60.0 (80)
Barton	253	73.5 (11)	108	22.2 (8)	158	13.9 (6)	64	* *	106	4.7 (5)	467	80.9 (30)	983	7.4 (12)
Bourbon	126	64.3 (21)	50	* *	81	* *	33	* *	50	* *	216	71.3 (51)	491	40.7 (64)
Brown	101	43.6 (39)	40	* *	67	* *	19	* *	38	* *	201	87.6 (5)	405	44.9 (69)
Butler	289	68.5 (18)	97	* *	153	9.8 (15)	60	* *	96	* *	467	66.4 (57)	1,002	12.0 (24)
Chase	12	* *	3	* *	6	* *	1	* *	3	* *	27	* *	51	* *
Chautauqua	28	* *	10	* *	14	* *	4	* *	9	* *	70	* *	127	43.3 (68)
Cherokee	196	59.2 (30)	82	* *	111	7.2 (27)	40	* *	82	* *	354	86.4 (9)	757	34.1 (57)
Cheyenne	0	* *	0	* *	0	* *	0	* *	0	* *	1	* *	1	* *

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in

(6) the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) Children ages 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 7A
2008 Pediatric Nutrition Surveillance (1)
Kansas

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
(4,5)

Clinic	Breastfeeding (5)		Exclusive Breastfeeding (6)		≤2 Hours TV Viewing/Day (7)		Smoking in Household							
	Number (Rank)	% Ever Breastfed (Rank)	Number (Rank)	% Breastfed At Least 6 Months (Rank)	Number (Rank)	% Breastfed At Least 12 Months (Rank)	Number (Rank)	% (Rank)						
Clark	8	* *	2	* *	6	* *	1	* *	2	* *	11	* *	26	* *
Clay	61	* *	22	* *	39	* *	13	* *	22	* *	129	82.9 (20)	255	31.8 (54)
Cloud	65	* *	37	* *	51	* *	17	* *	36	* *	145	84.1 (17)	305	3.9 (2)
Coffey	25	* *	8	* *	16	* *	6	* *	8	* *	48	* *	99	* *
Comanche	5	* *	4	* *	6	* *	1	* *	4	* *	8	* *	22	* *
Cowley-Ark	195	45.1 (38)	66	* *	122	7.4 (24)	40	* *	66	* *	383	86.4 (9)	777	7.9 (14)
Cowley-Win	151	61.6 (26)	64	* *	89	* *	34	* *	63	* *	311	77.5 (40)	611	17.8 (36)
Crawford-G	1	* *	1	* *	0	* *	1	* *	1	* *	2	* *	4	* *
Crawford-P	335	58.5 (32)	145	13.1 (21)	203	7.4 (24)	81	* *	143	0.7 (20)	610	76.1 (45)	1,301	41.5 (67)
Decatur	7	* *	6	* *	9	* *	4	* *	5	* *	20	* *	41	* *
DickinsonA	80	* *	43	* *	45	* *	34	* *	42	* *	190	86.8 (8)	368	7.9 (14)
DickinsonH	9	* *	11	* *	11	* *	2	* *	11	* *	31	* *	66	* *
Doniphan	39	* *	13	* *	21	* *	8	* *	13	* *	61	* *	143	55.2 (78)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in

(6) the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) Children ages 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 7A
2008 Pediatric Nutrition Surveillance (1)
Kansas

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
(4,5)

Clinic	Breastfeeding (5)				Exclusive Breastfeeding (6)				≤2 Hours TV Viewing/Day (7)		Smoking in Household			
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% (Rank)	Number	% (Rank)		
Douglas	428	74.8 (10)	202	35.1 (1)	290	22.4 (1)	102	25.5 (1)	197	7.6 (1)	670	77.5 (40)	1,578	6.4 (8)
Edwards	21	* *	11	* *	16	* *	8	* *	11	* *	59	* *	104	24.0 (47)
Elk	14	* *	9	* *	8	* *	5	* *	9	* *	40	* *	79	* *
Ellis	139	82.0 (1)	48	* *	65	* *	33	* *	47	* *	234	82.9 (20)	486	6.4 (8)
Ellsworth	30	* *	14	* *	23	* *	14	* *	14	* *	77	* *	140	12.9 (26)
Finney	656	75.5 (9)	227	22.5 (7)	437	13.3 (10)	110	9.1 (8)	226	5.3 (3)	1,341	75.5 (48)	2,679	17.4 (35)
Ford	573	73.1 (12)	218	22.9 (6)	388	12.4 (12)	94	* *	217	1.8 (14)	1,188	82.8 (22)	2,344	3.7 (1)
Franklin	160	53.1 (33)	54	* *	102	13.7 (7)	33	* *	53	* *	294	79.6 (35)	633	11.1 (22)
Geary	312	66.7 (19)	114	15.8 (16)	185	8.1 (20)	80	* *	113	3.5 (8)	438	76.0 (47)	1,036	40.9 (65)
Gove	10	* *	7	* *	12	* *	6	* *	7	* *	26	* *	53	* *
Graham	12	* *	12	* *	9	* *	4	* *	12	* *	25	* *	44	* *
Grant	68	* *	29	* *	59	* *	13	* *	29	* *	166	81.3 (28)	309	16.5 (33)
Gray	49	* *	34	* *	45	* *	17	* *	32	* *	125	87.2 (7)	248	10.9 (21)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in

(6) the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) Children ages 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 7A
2008 Pediatric Nutrition Surveillance (1)
Kansas

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
(4,5)

Clinic	Breastfeeding (5)		Exclusive Breastfeeding (6)		≤2 Hours TV Viewing/Day (7)		Smoking in Household							
	Number (Rank)	% Ever Breastfed (Rank)	Number (Rank)	% Breastfed At Least 6 Months (Rank)	Number (Rank)	% Breastfed At Least 12 Months (Rank)	Number (Rank)	% (Rank)	Number (Rank)	% (Rank)				
Greeley	10	* *	2	* *	5	* *	1	* *	2	* *	15	* *	34	* *
Greenwood	40	* *	22	* *	32	* *	15	* *	22	* *	121	98.3 (1)	205	21.5 (43)
Hamilton	14	* *	11	* *	17	* *	5	* *	11	* *	51	* *	94	* *
Harper	45	* *	17	* *	32	* *	5	* *	17	* *	108	65.7 (58)	210	30.5 (53)
Harvey	231	76.6 (6)	117	22.2 (8)	164	15.9 (4)	69	* *	115	0.9 (18)	530	88.5 (3)	1,043	5.1 (4)
Haskell	24	* *	5	* *	14	* *	2	* *	5	* *	70	* *	119	14.3 (29)
Hodgeman	9	* *	3	* *	10	* *	2	* *	3	* *	29	* *	48	* *
Jackson	71	* *	35	* *	49	* *	17	* *	35	* *	181	85.1 (12)	356	15.7 (31)
Jefferson	64	* *	25	* *	43	* *	18	* *	25	* *	115	95.7 (2)	242	45.0 (70)
Jewell	16	* *	8	* *	10	* *	3	* *	8	* *	25	* *	54	* *
Johnson-M	825	76.6 (6)	318	28.6 (3)	521	17.5 (2)	196	13.3 (5)	316	2.8 (9)	1,453	82.0 (25)	3,171	22.1 (44)
Johnson-O	928	77.9 (4)	406	28.3 (4)	589	16.1 (3)	226	17.3 (4)	402	2.7 (10)	1,752	84.5 (15)	3,717	17.3 (34)
Kearny	21	* *	11	* *	13	* *	7	* *	11	* *	74	* *	125	24.0 (47)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in

(6) the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) Children ages 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 7A
2008 Pediatric Nutrition Surveillance (1)
Kansas

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
(4,5)

Clinic	Breastfeeding (5)		Exclusive Breastfeeding (6)		≤2 Hours TV Viewing/Day (7)		Smoking in Household							
	Number (Rank)	% Ever Breastfed (Rank)	Number (Rank)	% Breastfed At Least 6 Months (Rank)	Number (Rank)	% Breastfed At Least 12 Months (Rank)	Number (Rank)	% (Rank)	Number (Rank)	% (Rank)				
Kingman	49	* *	24	* *	35	* *	14	* *	24	* *	109	81.7 (26)	214	37.9 (61)
Kiowa	10	* *	7	* *	10	* *	6	* *	7	* *	29	* *	52	* *
Labette	163	50.9 (36)	67	* *	129	7.0 (28)	40	* *	67	* *	282	84.4 (16)	643	32.0 (55)
Lane	12	* *	4	* *	8	* *	4	* *	4	* *	39	* *	60	* *
Leavenworth	317	62.1 (25)	106	15.1 (18)	176	7.4 (24)	67	* *	106	0.9 (18)	497	67.4 (53)	1,118	52.0 (75)
Lincoln	23	* *	9	* *	14	* *	5	* *	8	* *	42	* *	90	* *
Linn	55	* *	26	* *	51	* *	11	* *	26	* *	113	84.1 (17)	252	18.7 (37)
Logan	11	* *	5	* *	5	* *	2	* *	5	* *	38	* *	62	* *
Lyon	315	71.7 (14)	129	18.6 (13)	208	8.7 (19)	66	* *	127	2.4 (12)	641	80.8 (31)	1,297	10.2 (19)
Marion	43	* *	15	* *	35	* *	11	* *	15	* *	99	* *	202	5.9 (6)
Marshall	60	* *	21	* *	32	* *	10	* *	21	* *	112	65.2 (59)	228	53.1 (77)
Mc Pherson	137	72.3 (13)	54	* *	85	* *	36	* *	52	* *	251	76.1 (45)	539	6.1 (7)
Meade	43	* *	18	* *	21	* *	9	* *	17	* *	84	* *	164	20.1 (40)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in

(6) the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) Children ages 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 7A
2008 Pediatric Nutrition Surveillance (1)
Kansas

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
(4,5)

Clinic	Breastfeeding (5)				Exclusive Breastfeeding (6)				≤2 Hours TV Viewing/Day (7)		Smoking in Household			
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% (Rank)	Number	% (Rank)		
Miami	136	77.2 (5)	48	* *	75	* *	31	* *	47	* *	214	85.0 (13)	477	22.9 (45)
Mitchell	40	* *	11	* *	26	* *	11	* *	11	* *	64	* *	141	5.0 (3)
Montgome-C	143	62.9 (24)	48	* *	96	* *	33	* *	48	* *	300	78.7 (37)	593	36.6 (60)
Montgome-I	165	60.6 (29)	70	* *	122	9.0 (18)	36	* *	67	* *	309	80.3 (33)	666	34.7 (58)
Morris	29	* *	12	* *	26	* *	8	* *	12	* *	61	* *	128	7.8 (13)
Nemaha	27	* *	13	* *	17	* *	5	* *	12	* *	49	* *	103	49.5 (74)
Neosho-Ch	140	52.9 (34)	49	* *	72	* *	28	* *	47	* *	241	79.7 (34)	520	19.2 (38)
Neosho-Er	11	* *	4	* *	8	* *	4	* *	4	* *	22	* *	47	* *
Ness	8	* *	4	* *	7	* *	1	* *	4	* *	25	* *	47	* *
Norton	23	* *	17	* *	23	* *	7	* *	17	* *	66	* *	123	8.1 (16)
Osage	87	* *	26	* *	39	* *	18	* *	26	* *	145	86.2 (11)	318	21.1 (42)
Osborne	20	* *	11	* *	17	* *	6	* *	11	* *	46	* *	94	* *
Ottawa	18	* *	15	* *	20	* *	9	* *	14	* *	44	* *	92	* *

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in

(6) the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) Children ages 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 7A
2008 Pediatric Nutrition Surveillance (1)
Kansas

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
(4,5)

Clinic	Breastfeeding (5)				Exclusive Breastfeeding (6)				≤2 Hours TV Viewing/Day (7)		Smoking in Household			
	Number (Rank)	% Ever Breastfed (Rank)	Number Breastfed At Least 6 Months (Rank)	% Breastfed At Least 12 Months (Rank)	Number (Rank)	% Breastfed At Least 3 Months (Rank)	Number Breastfed At Least 6 Months (Rank)	% (Rank)	Number (Rank)	% (Rank)				
Pawnee	43	* *	15	* *	42	* *	13	* *	15	* *	101	73.3 (50)	205	12.7 (25)
Phillips	29	* *	15	* *	33	* *	7	* *	15	* *	68	* *	147	35.4 (59)
Pottawat-S	10	* *	9	* *	15	* *	4	* *	8	* *	64	* *	98	* *
Pottawat-W	25	* *	6	* *	16	* *	5	* *	5	* *	38	* *	94	* *
Pratt	66	* *	22	* *	49	* *	14	* *	21	* *	125	83.2 (19)	267	39.3 (62)
Rawlins	9	* *	5	* *	10	* *	2	* *	4	* *	32	* *	54	* *
Reno	464	58.6 (31)	163	20.9 (12)	268	13.1 (11)	92	* *	161	3.7 (7)	761	87.8 (4)	1,682	16.4 (32)
Republic	35	* *	16	* *	16	* *	8	* *	16	* *	67	* *	138	52.2 (76)
Rice	58	* *	24	* *	42	* *	12	* *	24	* *	135	87.4 (6)	268	23.5 (46)
Riley-Ft	378	76.5 (8)	188	16.5 (15)	306	7.8 (22)	118	18.6 (2)	185	3.8 (6)	758	67.2 (55)	1,699	20.0 (39)
Riley-Manh	340	78.5 (3)	126	31.7 (2)	197	11.2 (13)	66	* *	121	5.8 (2)	560	77.9 (39)	1,246	8.3 (17)
Rooks	34	* *	19	* *	33	* *	7	* *	19	* *	90	* *	170	40.0 (63)
Rush	11	* *	5	* *	9	* *	0	* *	5	* *	29	* *	52	* *

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in

(6) the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) Children ages 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 7A
2008 Pediatric Nutrition Surveillance (1)
Kansas

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
(4,5)

Clinic	Breastfeeding (5)			Exclusive Breastfeeding (6)			≤2 Hours TV Viewing/Day (7)		Smoking in Household					
	Number (Rank)	% Ever Breastfed (Rank)	Number Breastfed At Least 6 Months (Rank)	Number Breastfed At Least 12 Months (Rank)	Number Breastfed At Least 3 Months (Rank)	Number Breastfed At Least 6 Months (Rank)	Number (Rank)	% (Rank)	Number (Rank)	% (Rank)				
Russell	44	* *	26	* *	37	* *	15	* *	26	* *	79	* *	169	56.2 (79)
Saline	470	60.9 (28)	170	14.7 (19)	279	7.9 (21)	93	* *	169	1.2 (16)	752	82.7 (23)	1,680	6.6 (10)
Scott	31	* *	17	* *	38	* *	9	* *	16	* *	101	76.2 (44)	186	33.9 (56)
Sedgwick-C	1,046	71.3 (15)	395	15.7 (17)	758	9.8 (15)	238	7.6 (9)	391	1.8 (14)	2,011	66.8 (56)	4,192	9.1 (18)
Sedgwick-M	1,880	64.3 (21)	800	17.0 (14)	1,118	9.3 (17)	469	10.7 (6)	791	2.7 (10)	3,325	70.2 (52)	7,198	14.2 (28)
Sedgwick-S	1,103	61.3 (27)	462	14.7 (19)	728	7.8 (22)	271	10.3 (7)	454	1.1 (17)	2,132	84.9 (14)	4,482	7.3 (11)
Seneca	13	* *	5	* *	9	* *	4	* *	5	* *	23	* *	52	* *
Seward	439	71.1 (16)	173	24.3 (5)	265	13.6 (8)	91	* *	166	0.6 (21)	823	64.3 (60)	1,740	15.5 (30)
ShawneeM&I	133	81.2 (2)	29	* *	24	* *	21	* *	28	* *	99	* *	281	5.7 (5)
ShawneeMa	1,095	65.1 (20)	499	22.2 (8)	793	14.5 (5)	246	17.9 (3)	495	5.1 (4)	2,372	73.7 (49)	4,826	10.7 (20)
Sheridan	7	* *	6	* *	9	* *	2	* *	6	* *	22	* *	39	* *
Sherman	64	* *	26	* *	40	* *	21	* *	26	* *	145	76.6 (43)	288	28.1 (51)
Smith	17	* *	9	* *	17	* *	5	* *	9	* *	49	* *	93	* *

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in

(6) the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) Children ages 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 7A
2008 Pediatric Nutrition Surveillance (1)
Kansas

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
(4,5)

Clinic	Breastfeeding (5)		Exclusive Breastfeeding (6)		≤2 Hours TV Viewing/Day (7)		Smoking in Household							
	Number (Rank)	% Ever Breastfed (Rank)	Number (Rank)	% Breastfed At Least 6 Months (Rank)	Number (Rank)	% Breastfed At Least 12 Months (Rank)	Number (Rank)	% (Rank)	Number (Rank)	% (Rank)				
Stafford	30	* *	20	* *	28	* *	8	* *	18	* *	96	* *	168	11.3 (23)
Stanton	21	* *	7	* *	10	* *	4	* *	7	* *	48	* *	92	* *
Stevens	58	* *	39	* *	57	* *	19	* *	39	* *	169	81.7 (26)	316	13.0 (27)
Sumner	155	63.2 (23)	58	* *	101	9.9 (14)	38	* *	57	* *	253	77.5 (40)	556	49.3 (73)
Thomas	43	* *	17	* *	32	* *	10	* *	17	* *	102	80.4 (32)	207	25.1 (49)
Trego	13	* *	11	* *	14	* *	8	* *	11	* *	34	* *	73	* *
Wabaunsee	18	* *	11	* *	16	* *	4	* *	11	* *	38	* *	83	* *
Wallace	1	* *	0	* *	0	* *	1	* *	0	* *	2	* *	3	* *
Washington	25	* *	7	* *	13	* *	3	* *	7	* *	59	* *	109	41.3 (66)
Wichita	13	* *	8	* *	12	* *	6	* *	8	* *	48	* *	82	* *
Wilson	76	* *	31	* *	56	* *	15	* *	30	* *	170	81.2 (29)	342	47.7 (71)
Woodson	2	* *	4	* *	6	* *	2	* *	4	* *	16	* *	27	* *
Wyandotte	1,900	70.9 (17)	726	21.6 (11)	1,268	13.6 (8)	345	6.1 (10)	718	2.4 (12)	3,684	67.3 (54)	7,664	26.3 (50)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in

(6) the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) Children ages 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 7A
2008 Pediatric Nutrition Surveillance (1)
Kansas

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
(4,5)

Clinic	Breastfeeding (5)				Exclusive Breastfeeding (6)				≤2 Hours TV Viewing/Day (7)		Smoking in Household			
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% (Rank)	Number	% (Rank)		
Unknown Clinics	0	* *	0	* *	1	* *	0	* *	0	* *	1	* *		
Kansas	19,342	68.0	7,878	20.6	12,678	12.2	4,390	14.0	7,769	2.9	36,868	76.8	77,483	18.6
Nation (Prior Year)	1,747,004	59.8	874,434	25.4	979,105	17.5	127,047	10.2	182,064	5.2	687,613	75.0	3,047,308	11.3

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in

(6) the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) Children ages 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 6A
2006 - 2008 Pediatric Nutrition Surveillance (1)
Kansas
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Clinic (3,4)												
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under-weight (7)	Obese (7)	≥ 2 Yrs Over-weight (7)					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Allen	193	7.3 (22)	4.7 (12)	812	7.1 (77)	2.2 (32)	16.5(101)	346	22.8 (88)	14.5 (62)	572	6.5 (26)
Anderson	117	8.5 (35)	5.1 (20)	520	6.0 (51)	3.5 (75)	12.3 (61)	244	15.2 (32)	15.2 (71)	358	11.7 (65)
Atchison	285	8.4 (34)	5.3 (24)	1,233	6.6 (64)	5.7 (99)	11.3 (43)	522	13.6 (15)	14.2 (60)	795	14.3 (79)
Barber-Kio	4	* *	* *	32	* *	* *	* *	20	* *	* *	24	* *
Barber-Med	76	* *	* *	256	9.4(104)	0.8 (2)	13.3 (75)	89	* *	* *	150	12.7 (69)
Barton	604	9.1 (42)	5.1 (20)	2,528	5.9 (49)	3.7 (82)	9.1 (20)	1,136	14.3 (20)	11.1 (31)	1,684	7.4 (31)
Bourbon	235	7.7 (25)	4.7 (12)	1,005	7.6 (88)	1.1 (5)	16.0 (98)	444	15.8 (36)	17.6 (81)	691	8.1 (42)
Brown	145	9.0 (41)	7.6 (49)	967	5.0 (29)	3.5 (75)	12.2 (58)	476	14.7 (26)	9.9 (23)	685	13.6 (75)
Butler	639	9.7 (48)	5.8 (26)	2,602	9.9(108)	3.6 (80)	9.8 (26)	1,116	13.8 (18)	9.3 (18)	1,696	21.2 (98)
Chase	28	* *	* *	155	1.3 (1)	3.2 (67)	19.4(112)	79	* *	* *	106	10.4 (56)
Chautauqua	53	* *	* *	339	5.0 (29)	3.2 (67)	19.2(111)	173	14.5 (24)	19.7 (86)	244	7.4 (31)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2006 - 2008 Pediatric Nutrition Surveillance (1)
Kansas
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Clinic (3,4)												
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under-weight (7)	Obese (7)	≥ 2 Yrs Over-weight (7)					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Cherokee	702	8.0 (28)	5.0 (18)	2,396	6.8 (70)	3.4 (73)	11.8 (53)	834	16.9 (48)	15.1 (70)	1,275	7.1 (30)
Cheyenne	4	* *	* *	40	* *	* *	* *	22	* *	* *	32	* *
Clark	14	* *	* *	66	* *	* *	* *	28	* *	* *	49	* *
Clay	144	7.6 (24)	4.2 (8)	635	9.6(106)	2.5 (43)	18.6(108)	283	16.6 (44)	16.6 (76)	414	11.1 (61)
Cloud	162	9.9 (49)	1.9 (1)	748	6.8 (70)	3.2 (67)	11.0 (38)	335	17.9 (61)	12.5 (43)	488	13.1 (73)
Coffey	48	* *	* *	289	3.5 (11)	1.7 (12)	16.3 (99)	149	18.8 (71)	24.2 (92)	206	4.4 (11)
Comanche	13	* *	* *	74	* *	* *	* *	38	* *	* *	60	* *
Cowley-Ark	424	8.5 (35)	8.0 (52)	1,886	5.5 (40)	1.9 (20)	13.1 (74)	865	15.8 (36)	10.6 (27)	1,301	6.4 (25)
Cowley-Win	339	8.0 (28)	7.1 (42)	1,610	6.8 (70)	1.2 (7)	11.9 (55)	736	17.4 (55)	14.5 (62)	1,084	11.5 (63)
Crawford-G	1	* *	* *	43	* *	* *	* *	26	* *	* *	41	* *
Crawford-P	916	9.6 (47)	4.9 (15)	3,647	7.8 (92)	1.9 (20)	14.1 (83)	1,504	18.5 (69)	14.0 (58)	2,255	6.2 (24)

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2006 - 2008 Pediatric Nutrition Surveillance (1)
Kansas
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Clinic (3,4)												
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Obese (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Decatur	20	* *	* *	100	5.0 (29)	4.0 (88)	8.0 (13)	54	* *	* *	71	* *
DickinsonA	218	6.4 (9)	8.3 (54)	971	4.9 (27)	4.0 (88)	8.8 (16)	439	10.5 (9)	9.6 (21)	666	3.3 (6)
DickinsonH	31	* *	* *	182	3.8 (16)	2.7 (51)	4.9 (3)	93	* *	* *	135	5.9 (20)
Doniphan	87	* *	* *	338	15.4(114)	5.9(102)	24.0(114)	156	25.6 (93)	39.7 (93)	240	5.8 (19)
Douglas	1,048	7.8 (26)	7.3 (45)	4,159	7.5 (86)	1.5 (9)	12.6 (66)	1,656	18.0 (63)	12.9 (48)	2,715	15.0 (84)
Edwards	44	* *	* *	262	8.0 (94)	2.3 (34)	14.1 (83)	131	13.7 (16)	12.2 (40)	188	3.2 (5)
Elk	34	* *	* *	217	9.7(107)	1.8 (14)	12.4 (63)	99	* *	* *	164	9.8 (52)
Ellis	315	6.3 (8)	4.4 (10)	1,241	5.4 (36)	1.8 (14)	11.7 (51)	543	20.4 (84)	13.3 (52)	787	3.3 (6)
Ellsworth	70	* *	* *	334	10.8(112)	3.0 (57)	11.4 (45)	169	17.2 (53)	9.5 (19)	233	7.7 (38)
Finney	1,625	6.5 (12)	7.0 (39)	7,357	5.5 (40)	4.2 (92)	13.0 (71)	3,401	10.9 (10)	9.5 (19)	4,937	21.9 (99)
Ford	1,365	5.8 (5)	5.8 (26)	5,954	5.9 (49)	2.1 (27)	17.5(106)	2,862	18.3 (66)	18.9 (85)	3,960	8.4 (44)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2006 - 2008 Pediatric Nutrition Surveillance (1)
Kansas
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Clinic (3,4)												
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Obese (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Franklin	367	9.3 (43)	4.9 (15)	1,553	8.1 (97)	2.3 (34)	11.4 (45)	679	17.4 (55)	10.5 (25)	1,031	6.0 (22)
Ft. Riley	187	4.3 (2)	13.9 (62)	1,163	5.6 (44)	3.3 (72)	11.9 (55)	483	14.9 (27)	9.7 (22)	801	3.5 (9)
Geary	749	6.9 (16)	6.9 (36)	2,578	7.6 (88)	2.8 (53)	9.0 (18)	1,020	18.9 (72)	10.5 (25)	1,619	15.4 (86)
Gove	20	* *	* *	130	6.2 (55)	3.8 (84)	13.8 (78)	59	* *	* *	98	* *
Graham	32	* *	* *	156	10.3(109)	3.8 (84)	14.1 (83)	71	* *	* *	105	28.6(103)
Grant	139	9.4 (44)	5.0 (18)	821	7.6 (88)	2.1 (27)	15.6 (96)	442	13.3 (14)	17.0 (80)	637	27.3(102)
Gray	96	* *	* *	596	2.2 (4)	8.9(110)	5.5 (5)	315	8.9 (6)	6.0 (7)	419	4.5 (13)
Greeley	16	* *	* *	100	2.0 (3)	7.0(105)	11.0 (38)	50	* *	* *	66	* *
Greenwood	102	10.8 (56)	3.9 (6)	509	5.1 (32)	4.9 (96)	11.6 (49)	240	17.9 (61)	12.5 (43)	359	6.1 (23)
Hamilton	38	* *	* *	229	7.0 (75)	3.9 (87)	12.2 (58)	108	7.4 (4)	14.8 (67)	175	26.9(101)
Harper	79	* *	* *	497	6.8 (70)	3.8 (84)	14.5 (90)	248	16.1 (42)	10.9 (30)	346	9.0 (48)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2006 - 2008 Pediatric Nutrition Surveillance (1)
Kansas
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Clinic (3,4)												
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Obese (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Harvey	528	8.9 (40)	6.1 (28)	2,554	6.7 (66)	2.5 (43)	13.5 (77)	1,228	19.6 (78)	13.8 (54)	1,739	9.9 (53)
Haskell	41	* *	* *	331	4.5 (22)	3.0 (57)	13.0 (71)	198	15.2 (32)	10.6 (27)	261	18.0 (93)
Hodgeman	27	* *	* *	125	3.2 (9)	0.8 (2)	14.4 (89)	58	* *	* *	93	* *
Jackson	156	8.3 (33)	9.0 (58)	918	4.7 (24)	1.6 (11)	12.9 (70)	444	19.8 (80)	14.9 (68)	591	11.5 (63)
Jefferson	145	6.2 (7)	11.0 (61)	627	6.2 (55)	3.5 (75)	15.5 (95)	286	22.7 (87)	11.9 (36)	420	15.7 (87)
Jewell	34	* *	* *	128	7.0 (75)	2.3 (34)	11.7 (51)	58	* *	* *	82	* *
Johnson-M	1,877	8.7 (39)	6.9 (36)	7,759	5.8 (47)	2.1 (27)	12.5 (65)	3,201	16.4 (43)	12.9 (48)	5,019	7.5 (33)
Johnson-O	2,145	6.4 (9)	7.7 (51)	9,024	5.7 (46)	2.6 (47)	11.6 (49)	3,865	17.1 (50)	12.3 (41)	5,850	6.9 (29)
Kearny	38	* *	* *	267	7.9 (93)	7.5(106)	10.9 (37)	155	9.0 (7)	7.7 (13)	209	35.4(104)
Kingman	90	* *	* *	525	3.4 (10)	9.0(111)	6.5 (7)	262	8.8 (5)	3.8 (4)	360	11.1 (61)
Kiowa	24	* *	* *	172	4.1 (18)	3.5 (75)	12.2 (58)	87	* *	* *	122	0.8 (1)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2006 - 2008 Pediatric Nutrition Surveillance (1)
Kansas
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Clinic (3,4)												
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Obese (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Labette	331	6.9 (16)	7.6 (49)	1,574	5.3 (34)	2.4 (40)	10.5 (31)	721	14.4 (22)	10.8 (29)	1,145	7.6 (35)
Lane	24	* *	* *	140	2.9 (7)	5.7 (99)	9.3 (22)	85	* *	* *	114	8.8 (46)
Leavenwort	744	9.5 (46)	3.5 (3)	2,867	7.4 (83)	3.2 (67)	12.4 (63)	1,175	19.7 (79)	13.9 (55)	1,808	14.7 (82)
Lincoln	42	* *	* *	218	7.3 (80)	7.8(108)	8.7 (15)	111	17.1 (50)	3.6 (3)	169	11.8 (68)
Linn	136	8.1 (31)	5.1 (20)	625	9.1(102)	1.8 (14)	12.6 (66)	273	20.9 (85)	12.8 (47)	445	9.7 (51)
Logan	17	* *	* *	130	6.2 (55)	6.2(104)	8.5 (14)	75	* *	* *	95	* *
Lyon	859	7.2 (21)	7.2 (43)	3,549	4.5 (22)	2.5 (43)	13.9 (81)	1,556	18.4 (67)	18.0 (83)	2,319	4.9 (14)
Marion	103	5.8 (5)	2.9 (2)	541	6.1 (53)	2.2 (32)	11.3 (43)	247	19.0 (74)	12.6 (46)	380	13.7 (76)
Marshall	110	13.6 (63)	4.5 (11)	557	2.9 (7)	13.3(114)	7.4 (10)	279	9.0 (7)	4.3 (5)	381	7.6 (35)
Mc Pherson	313	6.4 (9)	7.0 (39)	1,434	7.2 (78)	2.3 (34)	7.1 (8)	645	15.0 (28)	6.7 (11)	951	13.0 (71)
Meade	86	* *	* *	421	3.6 (12)	3.6 (80)	10.0 (28)	213	16.0 (40)	6.1 (8)	308	5.5 (17)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2006 - 2008 Pediatric Nutrition Surveillance (1)
Kansas
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Clinic (3,4)												
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Obese (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- (Rank)	% <95th (Rank)	% ≥95th (Rank)	Number
Miami	311	6.8 (15)	5.1 (20)	1,263	10.4(111)	1.7 (12)	14.0 (82)	561	17.3 (54)	13.0 (50)	868	21.1 (97)
Mitchell	78	* *	* *	369	4.1 (18)	3.0 (57)	10.8 (36)	176	15.3 (35)	15.9 (73)	253	7.5 (33)
Montgome-C	268	8.2 (32)	3.7 (5)	1,370	6.3 (59)	2.6 (47)	11.4 (45)	665	18.5 (69)	12.5 (43)	995	8.9 (47)
Montgome-I	335	10.1 (52)	3.6 (4)	1,496	6.6 (64)	2.9 (55)	14.1 (83)	686	16.6 (44)	15.0 (69)	1,074	8.2 (43)
Morris	60	* *	* *	286	6.3 (59)	3.1 (64)	9.4 (23)	148	21.6 (86)	13.5 (53)	191	9.4 (49)
Nemaha	52	* *	* *	259	4.2 (21)	3.1 (64)	15.4 (93)	125	16.0 (40)	21.6 (90)	152	7.9 (40)
Neosho-Ch	270	11.1 (58)	7.4 (47)	1,303	7.4 (83)	4.6 (95)	11.8 (53)	556	14.4 (22)	10.1 (24)	802	15.7 (87)
Neosho-Er	21	* *	* *	146	10.3(109)	2.1 (27)	16.4(100)	74	* *	* *	103	16.5 (90)
Ness	13	* *	* *	111	3.6 (12)	4.5 (93)	3.6 (1)	67	* *	* *	83	* *
Norton	65	* *	* *	328	7.3 (80)	6.1(103)	7.3 (9)	152	15.1 (29)	6.6 (9)	229	1.3 (2)
Osage	164	5.5 (4)	6.7 (35)	791	5.3 (34)	2.0 (25)	15.4 (93)	374	18.4 (67)	14.7 (66)	538	3.0 (4)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2006 - 2008 Pediatric Nutrition Surveillance (1)
Kansas
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Clinic (3,4)												
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Obese (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Osborne	41	* *	* *	206	5.8 (47)	2.4 (40)	17.0(104)	93	* *	* *	147	17.7 (92)
Ottawa	50	* *	* *	242	5.4 (36)	2.5 (43)	18.6(108)	112	23.2 (90)	18.8 (84)	172	14.5 (81)
Pawnee	111	11.7 (60)	6.3 (30)	508	5.1 (32)	5.1 (97)	9.4 (23)	244	11.1 (11)	11.9 (36)	352	2.6 (3)
Phillips	73	* *	* *	356	6.7 (66)	1.1 (5)	16.9(103)	171	22.8 (88)	19.9 (88)	251	19.9 (96)
Pottawat-S	33	* *	* *	232	3.9 (17)	2.6 (47)	3.9 (2)	139	11.5 (13)	2.9 (1)	180	6.7 (28)
Pottawat-W	45	* *	* *	227	9.3(103)	1.3 (8)	11.0 (38)	105	14.3 (20)	14.3 (61)	170	5.9 (20)
Pratt	175	13.1 (61)	9.1 (59)	711	8.0 (94)	4.5 (93)	12.0 (57)	304	14.5 (24)	14.1 (59)	454	10.6 (58)
Rawlins	19	* *	* *	114	2.6 (6)	1.8 (14)	11.4 (45)	52	* *	* *	84	* *
Reno	1,071	6.7 (14)	8.6 (56)	4,395	6.3 (59)	3.7 (82)	8.8 (16)	1,833	15.2 (32)	12.0 (38)	2,803	6.6 (27)
Republic	54	* *	* *	337	8.0 (94)	1.5 (9)	14.2 (87)	167	19.8 (80)	16.8 (79)	244	14.8 (83)
Rice	157	10.2 (53)	8.3 (54)	735	6.7 (66)	0.7 (1)	22.7(113)	366	19.9 (82)	23.8 (91)	525	7.6 (35)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2006 - 2008 Pediatric Nutrition Surveillance (1)
Kansas
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Clinic (3,4)												
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under-weight (7)	Obese (7)	≥ 2 Yrs Over-weight (7)					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Riley-Ft	896	7.1 (20)	8.6 (56)	3,394	6.2 (55)	2.6 (47)	9.7 (25)	1,447	13.7 (16)	7.9 (14)	1,561	4.4 (11)
Riley-Manh	852	7.9 (27)	6.1 (28)	3,326	5.5 (40)	2.9 (55)	10.7 (35)	1,377	15.9 (38)	11.7 (35)	2,166	10.7 (59)
Rooks	85	* *	* *	458	4.8 (25)	3.5 (75)	5.2 (4)	240	17.1 (50)	5.8 (6)	299	8.4 (44)
Rush	29	* *	* *	140	12.1(113)	2.1 (27)	10.0 (28)	67	* *	* *	103	4.9 (14)
Russell	100	11.0 (57)	4.0 (7)	367	6.5 (63)	11.4(113)	7.9 (12)	160	6.3 (3)	3.1 (2)	301	3.3 (6)
Saline	1,042	9.9 (49)	5.6 (25)	4,135	8.6(100)	2.3 (34)	13.8 (78)	1,787	18.2 (65)	13.9 (55)	2,760	5.7 (18)
Scott	79	* *	* *	449	3.6 (12)	10.0(112)	9.8 (26)	226	3.1 (1)	7.1 (12)	313	12.8 (70)
Sedgwick-C	2,399	8.0 (28)	6.9 (36)	10,639	6.7 (66)	1.9 (20)	13.3 (75)	4,841	19.0 (74)	16.6 (76)	7,414	15.1 (85)
Sedgwick-E	733	11.5 (59)	6.3 (30)	3,111	8.9(101)	1.0 (4)	16.7(102)	1,250	19.0 (74)	16.5 (75)	1,957	10.4 (56)
Sedgwick-M	4,164	10.7 (55)	6.6 (33)	15,994	8.5 (98)	1.8 (14)	13.8 (78)	6,722	18.0 (63)	16.3 (74)	10,325	11.7 (65)
Sedgwick-S	2,406	7.3 (22)	7.4 (47)	10,716	7.6 (88)	1.9 (20)	14.9 (92)	4,769	18.9 (72)	16.7 (78)	7,320	14.0 (78)

- (1) Reporting period is January 1 through December 31.
- (2) Analyses based on one record per child.
- (3) Excludes records with unknown data or errors.
- (4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
- (5) Infants born during the reporting period included in the analysis.
- (6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
- (7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.
- (8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
- * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2006 - 2008 Pediatric Nutrition Surveillance (1)
Kansas
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Clinic (3,4)												
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Obese (7)		≥ 2 Yrs Over- weight (7)			Number	%
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Seneca	15	* *	* *	111	5.4 (36)	1.8 (14)	18.9(110)	56	* *	* *	90	* *
Seward	151	6.6 (13)	13.9 (62)	1,073	2.5 (5)	3.1 (64)	10.6 (34)	457	11.4 (12)	11.6 (33)	738	11.0 (60)
Seward	862	5.1 (3)	6.3 (30)	3,335	3.6 (12)	5.6 (98)	10.2 (30)	1,528	15.1 (29)	12.4 (42)	1,990	13.1 (73)
ShawneeM&I	404	6.9 (16)	7.2 (43)	955	5.4 (36)	4.1 (91)	11.2 (41)	266	20.3 (83)	15.8 (72)	364	13.7 (76)
ShawneeMa	2,525	8.6 (38)	4.9 (15)	12,627	6.4 (62)	2.7 (51)	14.2 (87)	5,886	17.7 (58)	14.6 (65)	8,603	13.0 (71)
Sheridan	21	* *	* *	105	1.9 (2)	7.6(107)	5.7 (6)	51	* *	* *	70	* *
Sherman	171	13.5 (62)	4.7 (12)	741	9.4(104)	2.8 (53)	9.2 (21)	327	14.1 (19)	11.3 (32)	476	7.8 (39)
Smith	53	* *	* *	230	6.1 (53)	3.0 (57)	12.6 (66)	112	23.2 (90)	11.6 (33)	166	5.4 (16)
Stafford	74	* *	* *	428	5.6 (44)	5.8(101)	14.5 (90)	223	17.0 (49)	19.7 (86)	320	10.0 (55)
Stanton	40	* *	* *	251	7.2 (78)	2.4 (40)	11.2 (41)	135	17.8 (59)	8.9 (16)	194	24.7(100)
Stevens	100	10.0 (51)	10.0 (60)	611	4.9 (27)	2.0 (25)	15.9 (97)	302	23.5 (92)	13.9 (55)	411	17.5 (91)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2006 - 2008 Pediatric Nutrition Surveillance (1)
Kansas
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Clinic (3,4)												
Clinic	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)	
	Low	High		Short Stature (6)	Under-weight (7)	Obese (7)		≥ 2 Yrs Over-weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Sumner	314	7.0 (19)	7.3 (45)	1,444	7.4 (83)	3.0 (57)	12.3 (61)	637	17.4 (55)	8.5 (15)	950	19.7 (95)
Thomas	113	3.5 (1)	8.0 (52)	537	4.8 (25)	3.4 (73)	7.8 (11)	239	15.9 (38)	9.2 (17)	373	9.9 (53)
Trego	31	* *	* *	177	8.5 (98)	4.0 (88)	9.0 (18)	80	* *	* *	114	15.8 (89)
Wabaunsee	46	* *	* *	219	4.1 (18)	2.3 (34)	18.3(107)	101	17.8 (59)	17.8 (82)	154	18.2 (94)
Wallace	4	* *	* *	29	* *	* *	* *	17	* *	* *	25	* *
Washington	42	* *	* *	267	6.0 (51)	8.6(109)	10.5 (31)	137	5.1 (2)	6.6 (9)	215	14.4 (80)
Wichita	30	* *	* *	259	7.3 (80)	1.9 (20)	17.4(105)	128	19.5 (77)	20.3 (89)	188	3.7 (10)
Wilson	213	9.4 (44)	4.2 (8)	897	6.9 (74)	3.0 (57)	12.6 (66)	397	16.6 (44)	13.1 (51)	585	8.0 (41)
Woodson	20	* *	* *	97	* *	* *	* *	35	* *	* *	67	* *
Wyandotte	4,802	8.5 (35)	7.0 (39)	19,583	5.5 (40)	3.2 (67)	10.5 (31)	8,382	15.1 (29)	12.1 (39)	13,283	9.6 (50)
Unknown Clinics	486	10.3 (54)	6.6 (33)	2,203	7.5 (86)	3.0 (57)	13.0 (71)	675	16.7 (47)	14.5 (62)	1,201	11.7 (65)

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2006 - 2008 Pediatric Nutrition Surveillance (1)
Kansas
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Clinic (3,4)												
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Obese (7)	≥ 2 Yrs Over- weight (7)					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- (Rank)	% <95th (Rank)	% ≥95th (Rank)	Number
Kansas	46,466	8.2	6.5	200,509	6.6	2.8	12.6	88,232	16.7	13.5	132,589	11.1

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2006 - 2008 Pediatric Nutrition Surveillance (1)
Kansas

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
 (4,5)

Clinic	Breastfeeding (5)				Exclusive Breastfeeding (6)				≤2 Hours TV Viewing/Day (7)		Smoking in Household			
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% (Rank)	Number	% (Rank)		
Allen	319	49.2 (77)	203	12.3 (69)	205	5.9 (65)	194	17.0 (29)	195	2.1 (66)	378	80.4 (53)	674	43.5 (96)
Anderson	202	50.0 (76)	124	18.5 (48)	130	15.4 (16)	116	14.7 (41)	119	5.0 (36)	253	81.4 (49)	456	17.3 (46)
Atchison	475	48.8 (79)	322	29.8 (16)	306	29.7 (4)	278	14.0 (44)	304	8.2 (16)	567	78.3 (68)	1,124	35.6 (84)
Barber-Kio	9	* *	8	* *	6	* *	7	* *	8	* *	20	* *	26	* *
Barber-Med	135	71.1 (26)	72	* *	63	* *	74	* *	67	* *	97	* *	241	46.1 (101)
Barton	1,012	70.5 (31)	680	22.2 (27)	629	13.5 (26)	593	16.0 (36)	622	5.3 (35)	1,195	80.5 (52)	2,223	10.9 (22)
Bourbon	387	55.0 (69)	264	14.0 (63)	252	8.7 (50)	240	17.5 (27)	252	5.6 (33)	477	75.9 (74)	828	38.4 (88)
Brown	346	51.2 (75)	270	38.5 (2)	237	31.6 (2)	228	14.0 (44)	241	7.1 (23)	506	82.8 (41)	882	48.3 (103)
Butler	1,037	68.9 (35)	712	19.0 (45)	621	9.2 (49)	630	12.5 (55)	656	4.3 (47)	1,188	66.4 (90)	2,390	13.1 (26)
Chase	54	* *	34	* *	36	* *	30	* *	32	* *	82	* *	140	37.1 (86)
Chautauqua	98	* *	93	* *	76	* *	66	* *	81	* *	175	92.0 (4)	281	50.9 (107)
Cherokee	1,162	54.0 (70)	807	13.3 (66)	519	6.9 (58)	748	13.0 (52)	723	4.4 (43)	909	84.3 (32)	2,304	32.7 (82)
Cheyenne	8	* *	12	* *	11	* *	8	* *	10	* *	22	* *	34	* *

- (1) Reporting period is January 1 through December 31.
- (2) TV Viewing and Smoking in Household data are not reported by all contributors.
- (3) Analyses based on one record per child.
- (4) Excludes records with unknown data or errors.
- (5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
- (6) the Breastfed At Least 6 Months analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
- (7) Children ages 2 years and older included in the analyses.
- * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 7A
2006 - 2008 Pediatric Nutrition Surveillance (1)
Kansas

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
(4,5)

Clinic	Breastfeeding (5)		Exclusive Breastfeeding (6)		≤2 Hours TV Viewing/Day (7)		Smoking in Household							
	Number (Rank)	% Ever Breastfed (Rank)	Number (Rank)	% Breastfed At Least 6 Months (Rank)	Number (Rank)	% Breastfed At Least 3 Months (Rank)	Number (Rank)	% (Rank)						
Clark	26	* *	14	* *	16	* *	15	* *	14	* *	30	* *	57	* *
Clay	267	61.0 (50)	163	31.9 (10)	151	18.5 (11)	157	26.1 (8)	153	8.5 (13)	275	82.9 (40)	573	27.1 (70)
Cloud	291	60.1 (55)	215	13.0 (67)	188	6.4 (63)	197	15.7 (37)	197	3.6 (57)	318	87.1 (13)	659	6.4 (7)
Coffey	77	* *	58	* *	59	* *	51	* *	56	* *	154	90.9 (6)	247	29.1 (74)
Comanche	18	* *	14	* *	16	* *	13	* *	14	* *	26	* *	68	* *
Cowley-Ark	726	44.4 (81)	490	14.9 (60)	426	6.6 (61)	447	10.3 (61)	461	2.8 (64)	856	85.2 (26)	1,575	9.6 (18)
Cowley-Win	588	58.7 (59)	431	14.6 (61)	401	7.5 (55)	350	14.0 (44)	396	3.3 (60)	769	78.4 (66)	1,416	16.1 (42)
Crawford-G	2	* *	4	* *	7	* *	2	* *	2	* *	27	* *	56	* *
Crawford-P	1,561	57.6 (63)	1,076	15.7 (58)	825	6.5 (62)	931	13.2 (50)	986	4.8 (39)	1,631	79.7 (56)	3,364	39.1 (89)
Decatur	33	* *	26	* *	26	* *	22	* *	23	* *	57	* *	97	* *
DickinsonA	378	72.5 (16)	252	19.4 (44)	215	11.2 (37)	219	20.5 (16)	227	4.8 (39)	463	90.7 (7)	848	6.8 (12)
DickinsonH	52	* *	51	* *	52	* *	44	* *	46	* *	97	* *	167	6.0 (5)
Doniphan	157	47.1 (80)	82	* *	80	* *	85	* *	77	* *	165	78.8 (63)	333	57.7 (109)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in

(6) the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) Children ages 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 7A
2006 - 2008 Pediatric Nutrition Surveillance (1)
Kansas

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
(4,5)

Clinic	Breastfeeding (5)				Exclusive Breastfeeding (6)				≤2 Hours TV Viewing/Day (7)		Smoking in Household			
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% (Rank)	Number	% (Rank)		
Douglas	1,725	73.8 (12)	1,230	30.3 (14)	1,039	18.1 (12)	1,043	26.2 (7)	1,112	11.7 (6)	1,761	78.4 (66)	3,734	8.7 (16)
Edwards	100	76.0 (9)	77	* *	66	* *	68	* *	65	* *	142	76.8 (72)	237	28.7 (73)
Elk	84	* *	56	* *	57	* *	58	* *	56	* *	105	83.8 (34)	192	30.2 (75)
Ellis	511	73.4 (13)	315	21.9 (28)	286	10.8 (39)	264	18.6 (24)	284	3.2 (62)	581	86.6 (19)	1,146	6.5 (8)
Ellsworth	123	69.1 (34)	85	* *	74	* *	70	* *	74	* *	185	81.6 (46)	300	17.7 (47)
Finney	2,729	72.4 (18)	1,901	24.3 (23)	1,720	12.2 (33)	1,580	14.0 (44)	1,693	5.6 (33)	3,670	72.3 (81)	6,638	19.5 (49)
Ford	2,202	71.5 (22)	1,515	23.4 (25)	1,452	11.7 (35)	1,255	8.7 (65)	1,355	3.9 (52)	3,066	85.8 (24)	5,366	5.0 (3)
Franklin	657	51.9 (73)	411	16.5 (54)	408	7.6 (54)	362	12.2 (56)	382	3.7 (55)	709	77.3 (69)	1,427	15.3 (37)
Ft. Riley	454	72.0 (19)	342	25.4 (20)	243	16.9 (13)	229	21.4 (15)	230	14.3 (4)	483	61.7 (94)	1,812	26.8 (69)
Geary	1,148	63.0 (42)	703	16.1 (56)	638	7.8 (53)	655	14.4 (42)	650	3.8 (53)	1,084	79.2 (58)	2,359	41.4 (95)
Gove	49	* *	39	* *	35	* *	38	* *	39	* *	63	* *	108	39.8 (91)
Graham	60	* *	39	* *	39	* *	36	* *	37	* *	28	* *	47	* *
Grant	247	71.3 (23)	193	24.9 (22)	198	12.6 (30)	159	11.3 (59)	180	4.4 (43)	474	78.7 (64)	699	18.5 (48)

- (1) Reporting period is January 1 through December 31.
- (2) TV Viewing and Smoking in Household data are not reported by all contributors.
- (3) Analyses based on one record per child.
- (4) Excludes records with unknown data or errors.
- (5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
- (6) the Breastfed At Least 6 Months analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
- (7) Children ages 2 years and older included in the analyses.
- * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 7A
2006 - 2008 Pediatric Nutrition Surveillance (1)
Kansas

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
(4,5)

Clinic	Breastfeeding (5)		Exclusive Breastfeeding (6)		≤2 Hours TV Viewing/Day (7)		Smoking in Household							
	Number (Rank)	% Ever Breastfed (Rank)	Number (Rank)	% Breastfed At Least 6 Months (Rank)	Number (Rank)	% Breastfed At Least 12 Months (Rank)	Number (Rank)	% (Rank)	Number (Rank)	% (Rank)				
Gray	213	81.2 (1)	165	38.2 (3)	152	20.4 (7)	141	24.1 (11)	151	8.6 (12)	298	87.9 (11)	523	14.0 (29)
Greeley	33	* *	22	* *	18	* *	19	* *	20	* *	55	* *	97	* *
Greenwood	165	55.8 (67)	129	18.6 (47)	116	10.3 (44)	110	17.3 (28)	120	9.2 (11)	249	98.8 (1)	411	21.2 (55)
Hamilton	70	* *	54	* *	57	* *	47	* *	52	* *	121	76.0 (73)	184	14.1 (30)
Harper	207	58.9 (58)	121	37.2 (4)	120	30.0 (3)	112	18.8 (23)	118	15.3 (3)	183	67.2 (88)	406	24.6 (64)
Harvey	972	74.5 (11)	728	26.8 (17)	642	14.3 (19)	591	20.1 (17)	632	7.4 (20)	1,219	86.9 (17)	2,244	4.9 (2)
Haskell	71	* *	58	* *	77	* *	49	* *	55	* *	212	81.6 (46)	289	20.1 (52)
Hodgeman	38	* *	30	* *	31	* *	27	* *	30	* *	64	* *	104	14.4 (32)
Jackson	320	71.3 (23)	233	36.5 (5)	217	23.5 (5)	190	31.1 (2)	211	18.5 (1)	443	83.7 (35)	792	19.7 (50)
Jefferson	254	53.5 (71)	156	19.9 (42)	162	9.9 (45)	142	26.8 (6)	143	11.2 (7)	298	88.6 (10)	554	40.8 (93)
Jewell	54	* *	31	* *	26	* *	31	* *	31	* *	60	* *	109	3.7 (1)
Johnson-M	3,260	76.6 (7)	2,252	31.0 (13)	1,881	18.6 (10)	1,984	15.7 (37)	2,056	5.8 (32)	3,232	84.2 (33)	6,919	23.2 (59)
Johnson-O	3,612	78.4 (4)	2,496	30.3 (14)	2,187	16.7 (14)	2,132	18.5 (25)	2,250	6.4 (27)	4,037	85.3 (25)	7,877	16.5 (43)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in

(6) the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) Children ages 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 7A
2006 - 2008 Pediatric Nutrition Surveillance (1)
Kansas

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
(4,5)

Clinic	Breastfeeding (5)				Exclusive Breastfeeding (6)				≤2 Hours TV Viewing/Day (7)		Smoking in Household			
	Number (Rank)	% Ever Breastfed (Rank)	Number At Least 6 Months (Rank)	% Breastfed At Least 12 Months (Rank)	Number At Least 3 Months (Rank)	% Breastfed At Least 6 Months (Rank)	Number (Rank)	% (Rank)	Number (Rank)	% (Rank)				
Kearny	69	* *	61	* *	57	* *	51	* *	57	* *	165	66.7 (89)	210	23.8 (61)
Kingman	196	71.9 (21)	141	31.2 (12)	121	22.3 (6)	107	21.5 (14)	122	7.4 (20)	206	79.6 (57)	442	32.6 (81)
Kiowa	58	* *	46	* *	48	* *	32	* *	40	* *	83	* *	173	40.5 (92)
Labette	565	49.2 (77)	410	14.6 (61)	426	6.8 (60)	366	13.1 (51)	385	4.2 (48)	776	85.2 (26)	1,344	31.6 (79)
Lane	42	* *	30	* *	30	* *	24	* *	26	* *	73	* *	115	27.8 (72)
Leavenworth	1,271	58.5 (60)	714	16.8 (53)	623	7.4 (56)	732	13.0 (52)	661	3.8 (53)	1,235	68.8 (85)	2,689	50.8 (106)
Lincoln	82	* *	46	* *	43	* *	46	* *	44	* *	104	37.5 (95)	196	45.9 (100)
Linn	230	59.1 (57)	174	25.9 (18)	159	13.8 (21)	155	27.7 (4)	161	9.9 (9)	282	84.4 (30)	539	25.0 (66)
Logan	36	* *	28	* *	32	* *	22	* *	25	* *	83	* *	106	30.2 (75)
Lyon	1,379	70.8 (29)	961	20.3 (39)	832	10.5 (41)	810	11.5 (58)	870	4.7 (42)	1,651	84.9 (29)	3,138	15.7 (39)
Marion	204	80.4 (3)	151	33.8 (7)	144	15.3 (17)	119	32.8 (1)	134	8.2 (16)	278	94.2 (2)	502	6.0 (5)
Marshall	199	57.8 (61)	143	20.3 (39)	123	13.8 (21)	112	17.0 (29)	129	6.2 (29)	312	66.0 (91)	546	49.5 (104)
Mc Pherson	560	72.9 (14)	372	21.8 (29)	343	11.4 (36)	321	16.2 (35)	337	4.2 (48)	682	80.8 (50)	1,313	8.1 (14)

- (1) Reporting period is January 1 through December 31.
- (2) TV Viewing and Smoking in Household data are not reported by all contributors.
- (3) Analyses based on one record per child.
- (4) Excludes records with unknown data or errors.
- (5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
- (6) the Breastfed At Least 6 Months analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
- (7) Children ages 2 years and older included in the analyses.
- * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 7A
2006 - 2008 Pediatric Nutrition Surveillance (1)
Kansas

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
(4,5)

Clinic	Breastfeeding (5)				Exclusive Breastfeeding (6)				≤2 Hours TV Viewing/Day (7)		Smoking in Household			
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% (Rank)	Number	% (Rank)		
Meade	142	68.3 (36)	107	42.1 (1)	108	35.2 (1)	79	* *	97	* *	230	86.5 (20)	391	20.5 (53)
Miami	505	63.4 (41)	316	18.0 (50)	277	8.3 (51)	274	17.9 (26)	290	4.1 (50)	598	86.3 (21)	1,207	27.7 (71)
Mitchell	118	61.0 (50)	84	* *	89	* *	76	* *	75	* *	181	89.5 (9)	322	8.7 (16)
Montgome-C	485	56.7 (65)	351	13.4 (65)	373	6.4 (63)	302	12.9 (54)	329	4.9 (37)	731	80.2 (54)	1,221	31.2 (77)
Montgome-I	534	56.2 (66)	407	12.5 (68)	386	5.7 (66)	356	11.0 (60)	382	2.9 (63)	747	83.4 (38)	1,276	34.5 (83)
Morris	103	70.9 (27)	78	* *	67	* *	66	* *	71	* *	155	93.5 (3)	251	14.7 (33)
Nemaha	101	60.4 (53)	77	* *	62	* *	57	* *	70	* *	128	84.4 (30)	233	43.8 (98)
Neosho-Ch	554	51.4 (74)	375	13.6 (64)	319	4.4 (68)	329	12.2 (56)	338	6.2 (29)	508	83.5 (37)	1,092	24.0 (62)
Neosho-Er	42	* *	34	* *	30	* *	29	* *	33	* *	64	* *	122	23.0 (58)
Ness	27	* *	18	* *	21	* *	17	* *	17	* *	69	* *	92	* *
Norton	113	62.8 (43)	91	* *	84	* *	75	* *	82	* *	167	82.0 (44)	287	15.7 (39)
Osage	298	57.0 (64)	190	22.6 (26)	195	13.8 (21)	160	16.9 (31)	176	8.5 (13)	396	87.1 (13)	744	24.9 (65)
Osborne	78	* *	61	* *	53	* *	47	* *	55	* *	103	82.5 (43)	179	41.3 (94)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in

(6) the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) Children ages 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 7A
2006 - 2008 Pediatric Nutrition Surveillance (1)
Kansas

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
(4,5)

Clinic	Breastfeeding (5)				Exclusive Breastfeeding (6)				≤2 Hours TV Viewing/Day (7)		Smoking in Household			
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% (Rank)	Number	% (Rank)		
Ottawa	90	* *	67	* *	54	* *	58	* *	61	* *	123	86.2 (22)	211	14.2 (31)
Pawnee	188	66.0 (38)	144	20.8 (34)	129	12.4 (31)	136	16.9 (31)	137	4.4 (43)	261	78.9 (62)	451	22.4 (57)
Phillips	134	76.1 (8)	99	* *	95	* *	98	* *	97	* *	179	79.9 (55)	312	32.1 (80)
Pottawat-S	60	* *	54	* *	52	* *	41	* *	50	* *	146	90.4 (8)	182	6.6 (9)
Pottawat-W	70	* *	50	* *	48	* *	44	* *	47	* *	109	80.7 (51)	191	12.0 (23)
Pratt	302	70.2 (32)	183	19.7 (43)	170	7.1 (57)	165	26.1 (8)	170	6.5 (26)	278	82.0 (44)	512	25.2 (67)
Rawlins	35	* *	29	* *	34	* *	26	* *	28	* *	61	* *	89	* *
Reno	1,805	61.8 (47)	1,214	18.8 (46)	1,056	9.8 (47)	1,066	19.4 (20)	1,103	4.1 (50)	1,935	87.1 (13)	3,938	10.8 (21)
Republic	115	71.3 (23)	82	* *	65	* *	71	* *	74	* *	171	65.5 (93)	297	45.8 (99)
Rice	261	61.3 (48)	169	18.3 (49)	166	9.6 (48)	152	19.7 (18)	160	7.5 (19)	384	87.2 (12)	663	26.4 (68)
Riley-Ft	1,506	72.9 (14)	982	20.9 (33)	888	9.9 (45)	952	25.8 (10)	950	4.8 (39)	1,521	71.3 (82)	2,625	21.2 (55)
Riley-Manh	1,433	77.6 (6)	984	32.6 (8)	816	16.7 (14)	815	27.5 (5)	863	9.6 (10)	1,436	79.0 (61)	3,113	8.4 (15)
Rooks	141	61.0 (50)	95	* *	76	* *	84	* *	91	* *	224	77.2 (71)	381	38.1 (87)

(1) Reporting period is January 1 through December 31.
(2) TV Viewing and Smoking in Household data are not reported by all contributors.
(3) Analyses based on one record per child.
(4) Excludes records with unknown data or errors.
(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
(6) the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
(7) Children ages 2 years and older included in the analyses.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 7A
2006 - 2008 Pediatric Nutrition Surveillance (1)
Kansas

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
(4,5)

Clinic	Breastfeeding (5)				Exclusive Breastfeeding (6)				≤2 Hours TV Viewing/Day (7)		Smoking in Household			
	Number (Rank)	% Ever Breastfed (Rank)	Number Breastfed At Least 6 Months (Rank)	% Breastfed At Least 12 Months (Rank)	Number Breastfed At Least 3 Months (Rank)	% Breastfed At Least 6 Months (Rank)	Number (Rank)	% (Rank)	Number (Rank)	% (Rank)				
Rush	53	* *	33	* *	32	* *	29	* *	31	* *	72	* *	145	17.2 (45)
Russell	154	55.2 (68)	116	16.4 (55)	98	* *	98	* *	107	0.9 (67)	179	74.3 (78)	348	50.9 (107)
Saline	1,725	60.2 (54)	1,095	17.6 (51)	1,044	6.9 (58)	972	15.5 (39)	991	3.3 (60)	1,875	83.7 (35)	3,798	6.6 (9)
Scott	173	53.2 (72)	141	20.6 (36)	134	11.9 (34)	116	21.6 (13)	130	8.5 (13)	244	81.6 (46)	414	21.0 (54)
Sedgwick-C	3,877	65.8 (39)	2,687	20.7 (35)	2,702	10.4 (43)	2,274	13.8 (48)	2,492	3.7 (55)	5,181	68.6 (86)	9,156	13.1 (26)
Sedgwick-E	742	64.3 (40)	640	20.3 (39)	248	13.7 (24)	581	8.6 (66)	512	7.4 (20)	1,259	82.8 (41)	2,527	6.7 (11)
Sedgwick-M	7,209	62.7 (44)	4,585	21.4 (31)	4,099	11.0 (38)	4,027	16.5 (34)	4,247	6.0 (31)	7,295	74.7 (75)	14,719	12.7 (25)
Sedgwick-S	4,130	61.1 (49)	2,847	17.4 (52)	2,679	7.9 (52)	2,526	14.8 (40)	2,666	3.6 (57)	5,107	83.4 (38)	9,205	9.9 (20)
Seneca	37	* *	25	* *	30	* *	22	* *	23	* *	61	* *	109	39.4 (90)
Seward	411	70.6 (30)	289	35.6 (6)	241	14.1 (20)	150	13.3 (49)	158	16.5 (2)	459	69.1 (84)	1,597	9.6 (18)
Seward	1,314	70.9 (27)	866	25.1 (21)	821	13.2 (27)	786	8.8 (64)	833	2.3 (65)	1,676	67.4 (87)	2,376	15.9 (41)
ShawneeM&I	560	72.0 (19)	323	31.3 (11)	179	20.1 (8)	254	19.7 (18)	268	11.2 (7)	211	74.4 (77)	957	6.9 (13)
ShawneeMa	4,595	62.0 (46)	3,255	21.5 (30)	3,015	13.2 (27)	2,867	19.4 (20)	3,030	6.4 (27)	6,067	74.1 (79)	11,231	13.4 (28)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in

(6) the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) Children ages 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 7A
2006 - 2008 Pediatric Nutrition Surveillance (1)
Kansas

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
(4,5)

Clinic	Breastfeeding (5)			Exclusive Breastfeeding (6)			≤2 Hours TV Viewing/Day (7)		Smoking in Household					
	Number (Rank)	% Ever Breastfed (Rank)	Number % Breastfed At Least 6 Months (Rank)	Number % Breastfed At Least 12 Months (Rank)	Number % Breastfed At Least 3 Months (Rank)	Number % Breastfed At Least 6 Months (Rank)	Number (Rank)	% (Rank)	Number (Rank)	% (Rank)				
Sheridan	37	* *	26	* *	29	* *	24	* *	25	* *	51	* *	92	* *
Sherman	302	69.2 (33)	210	20.5 (37)	169	12.4 (31)	187	19.3 (22)	189	6.9 (24)	340	85.0 (28)	673	16.6 (44)
Smith	77	* *	69	* *	50	* *	63	* *	62	* *	119	73.1 (80)	196	14.8 (34)
Stafford	129	80.6 (2)	108	21.3 (32)	105	10.5 (41)	81	* *	94	* *	236	86.0 (23)	371	12.4 (24)
Stanton	68	* *	55	* *	55	* *	42	* *	47	* *	149	74.5 (76)	210	14.8 (34)
Stevens	250	74.8 (10)	176	32.4 (9)	176	19.9 (9)	170	29.4 (3)	173	8.1 (18)	327	78.6 (65)	439	15.3 (37)
Stevens	41	* *	19	* *	8	* *	13	* *	14	* *	0	* *	111	19.8 (51)
Sumner	556	57.7 (62)	385	25.7 (19)	298	14.4 (18)	344	24.1 (11)	373	12.1 (5)	484	77.3 (69)	909	46.9 (102)
Thomas	193	72.5 (16)	156	20.5 (37)	126	12.7 (29)	126	16.7 (33)	142	4.9 (37)	256	86.7 (18)	474	14.8 (34)
Trego	64	* *	49	* *	45	* *	42	* *	46	* *	76	* *	134	50.0 (105)
Wabaunsee	79	* *	58	* *	49	* *	50	* *	52	* *	101	91.1 (5)	180	5.0 (3)
Wallace	5	* *	6	* *	6	* *	4	* *	3	* *	19	* *	32	* *
Washington	106	78.3 (5)	66	* *	60	* *	59	* *	60	* *	148	70.9 (83)	258	36.8 (85)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in

(6) the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) Children ages 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 7A
2006 - 2008 Pediatric Nutrition Surveillance (1)
Kansas

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
(4,5)

Clinic	Breastfeeding (5)				Exclusive Breastfeeding (6)				≤2 Hours TV Viewing/Day (7)		Smoking in Household			
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% (Rank)	Number	% (Rank)		
Wichita	84	* *	66	* *	57	* *	59	* *	58	* *	138	87.0 (16)	230	31.3 (78)
Wilson	353	59.5 (56)	251	15.1 (59)	215	4.7 (67)	212	14.2 (43)	231	3.5 (59)	430	79.1 (59)	795	43.5 (96)
Woodson	38	* *	32	* *	31	* *	34	* *	30	* *	43	* *	92	* *
Wyandotte	7,818	68.3 (36)	5,379	24.2 (24)	4,736	13.7 (24)	4,550	9.6 (62)	4,829	4.4 (43)	9,002	65.7 (92)	17,260	24.3 (63)
Unknown Clinics	851	62.4 (45)	658	15.8 (57)	338	10.7 (40)	467	9.2 (63)	497	6.6 (25)	642	79.1 (59)	2,623	23.4 (60)
Kansas	78,940	66.0	54,147	22.6	48,063	12.3	46,838	16.0	49,238	5.7	92,641	77.7	178,993	18.8

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in

(6) the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) Children ages 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

