

302 Gestational Diabetes

Definition/Cut-off Value

Gestational diabetes mellitus (GDM) is defined as any degree of glucose/carbohydrate intolerance with onset or first recognition during pregnancy (1, 2).

Presence of condition diagnosed, documented, or reported by a physician or someone working under a physician's orders, or as self reported by applicant/participant/caregiver. See Clarification for more information about self-reporting a diagnosis.

Participant Category and Priority Level

Category	Priority
Pregnant Women	I

Justification

The definition of GDM applies regardless of whether insulin or only diet modification is used for treatment, or whether the condition persists after pregnancy. Included in this classification are women who may have had undiagnosed diabetes prior to pregnancy but who are first diagnosed during pregnancy (1, 2). Pregnant women requiring the use of exogenous steroids, tocolytics, or other medications, or who have medical conditions that alter glucose tolerance, may develop GDM (2). GDM represents nearly 90% of all pregnancies complicated by diabetes (1). The criteria for the diagnosis of GDM (3) are shown in Table 1 (see Clarification).

Pregnancy is an insulin-resistant and diabetogenic state (2). Deterioration of glucose tolerance occurs normally during pregnancy, particularly in the 3rd trimester (1, 2). Untreated or poorly treated GDM results in a higher risk of morbidity and mortality for both the mother and the fetus (2).

Established risk factors for GDM are advanced maternal age, obesity, and family history of diabetes (4). Risk assessment for GDM should be undertaken at the first prenatal visit. Women with clinical characteristics consistent with a high risk for GDM (e.g., those with marked obesity, personal history of GDM or delivery of a previous large-for-gestation-age infant, glycosuria, polycystic ovary syndrome, or a strong family history of diabetes) should undergo glucose testing as soon as possible (5). Unquestionably, there are also ethnic differences in the prevalence of GDM. In the U.S., Native Americans, Asians, Hispanics, and African American women are at a higher risk for GDM than non-Hispanic White women. Besides obesity, there is a suggestion that physical inactivity, diets high in saturated fat and smoking are associated with increasing risk for GDM or recurrent GDM (4).

Infants of women with GDM are at an increased risk of developing obesity, impaired glucose tolerance or diabetes as children or young adults (4). GDM is associated with a higher incidence of maternal and fetal complications. Maternal complications include polycythemia, respiratory distress syndrome, and increased rate of stillbirth (6). Although rarely seen in GDM, congenital anomalies, neural tube defects, cardiac abnormalities and/or caudal regression may occur if a woman has GDM in the early first trimester (6, 7).

Since GDM is a risk factor for subsequent type 2 diabetes after delivery, lifestyle modifications aimed at reducing weight and increasing physical activity are recommended (8). The National Diabetes Education

Program (NDEP) is currently promoting a GDM Prevention Initiative, targeting both providers and women with a GDM history (9). Key messages are illustrated in Table 2 (see Clarification).

Medical Nutrition Therapy (MNT) is the primary treatment for the management of GDM (7). MNT for GDM primarily involves a carbohydrate-controlled meal plan that promotes optimal nutrition for maternal and fetal health with adequate energy for appropriate gestational weight gain, achievement and maintenance of normoglycemia, and absence of ketosis (7, 8). Breastfeeding should be strongly encouraged as it is associated with maternal weight loss and reduced insulin resistance for both mother and offspring (10). WIC nutrition services can reinforce and support the medical and diet therapies (such as MNT) that participants with GDM receive from their health care providers.

References

1. American Diabetes Association: Diagnosis and classification of diabetes mellitus. *Diabetes Care*. Jan 2008; 31 Suppl 1:S55-60.
2. Franz MJ, Biastre SA, Slocum J. Diabetes in the life cycle and research. In: *Gestational Diabetes – A core curriculum for diabetes education*, American Association of Diabetes Educators. 5th Ed. 2003.
3. American Diabetes Association. Gestational diabetes mellitus (position statement). *Diabetes Care*. 2003; 26 Suppl. 1:S103-105.
4. Ferrara, A. Increasing prevalence of gestational diabetes mellitus: a public health perspective. *Proceedings of the fifth international workshop – conference on Gestational Diabetes Mellitus*. *Diabetes Care*. Jul 2007; 30 Suppl. 2:S141-46.
5. American Diabetes Association. Standards of medical care in diabetes (position statement). *Diabetes Care*. Jan 2007; 30 Suppl. 2:S4-41.
6. Thomas AM, Gutierrez YM. American Dietetic Association guide to gestational diabetes mellitus in postpartum considerations. Eds. American Dietetic Association; 2005:101-113.
7. Brian SR, Nickless N, Thung SF, Inzucchio SE. Gestational diabetes update: screening, medical management and follow-up. *Practical Diabetology*. Mar 2007; 10-18.
8. American Diabetes Association. Diagnosis and classification of diabetes mellitus. *Diabetes Care*. 2008 Jan; 31 Suppl. 1:S55-60.
9. Ratner, RE. Prevention of type 2 diabetes in women with previous gestational diabetes. *Proceedings of the fifth international workshop – conference on gestational diabetes mellitus*. *Diabetes Care*. Jul 2007; 30 Suppl. 2:S242-245.
10. Evert AG, Vande Hei K. Gestational diabetes education and diabetes prevention strategies. *Diabetes Spectrum*. 2006; 19(3):135-139.

Clarification

Self-reporting of a diagnosis by a medical professional should not be confused with self-diagnosis, where a person simply claims to have or to have had a medical condition without any reference to professional diagnosis. A self-reported medical diagnosis (“My doctor says that I have/my son or daughter has...”) should prompt the CPA to validate the presence of the condition by asking more pointed questions related to that diagnosis.

Women at high risk for GDM who have tested negative at the initial screening, and women at average risk for GDM should be tested by a licensed medical provider, between 24 and 28 weeks of gestation. Women of average risk should be tested at 24-28 weeks of gestation. Testing should follow one of two approaches:

1. One-step approach: perform a diagnostic 100-g OGTT (Oral Glucose Tolerance Test).
2. Two-step approach:
 - A screening test (glucose challenge test) that measures plasma or serum glucose is done 1 hour after a 50-g oral glucose load without regard for time of day or time of last meal. If a plasma or serum glucose level meets or exceeds the threshold (≥ 130 mg/dl [7.2 mmol/L] or ≥ 140 mg/dl [7.8 mmol/L], respectively), an OGTT is performed (3).
 - A diagnosis of GDM is made with a 100-g oral glucose load after an overnight fast. Using a 3-hour test, if two or more plasma or serum glucose levels meet or exceed the threshold, a diagnosis of GDM is made. Alternatively, the diagnosis can be made using a 75-g oral glucose load. The glucose threshold values for both tests are listed in Table 1 (10). The 75-g glucose load test is not as well validated as the 100-g OGTT.

With either the 75-g OGTT or the 100-g OGTT, it is recommended that the test be performed after an overnight fast of at least 8 hours but no longer than 14 hours. For 3 days prior to the test the woman should consume an unrestricted diet (≥ 150 g carbohydrate per day) and maintain unrestricted physical activity. Women need to remain seated and not smoke during the test. (1, 2).

Table 1. Diagnosis of Gestational Diabetes Mellitus with a 100-g or 75-g Oral Glucose Load

Time (h)	100-g Oral Glucose Load	75-g Oral Glucose Load
Fasting	95 mg/dL (5.3 mmol/L)	95 mg/dL (5.3 mmol/L)
1	180 mg/dL (10.0 mmol/L)	180 mg/dL (10.0 mmol/L)
2	155 mg/dL (8.6 mmol/L)	155 mg/dL (8.6 mmol/L)
3	140 mg/dL (7.8 mmol/L)	

Two or more of the venous plasma concentrations must be met or exceeded for a positive diagnosis.
Source: American Diabetes Association (3).

Table 2. Gestational Diabetes Mellitus (GDM) Prevention Initiative from the National Diabetes Education Program

- GDM imparts lifelong risk for diabetes, mostly type 2.
- Modest weight loss and physical activity can delay or prevent type 2 diabetes.
- Offspring can lower risk of diabetes by eating healthy foods, being active, and not becoming overweight.

Conservative recommendations to patients include:

- Let health care practitioners know of any history of GDM.



- Get glucose testing at 6 to 12 weeks postpartum, then every 1-2 years.
- Reach pre-pregnancy weight 6 to 12 months postpartum.
- If still overweight, lose at least 5 to 7% of weight slowly, over time, and keep it off.

Adapted from the National Diabetes Education Program (9).

