

Table 3B
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Racial and Ethnic Distribution by County

Comparison of Racial and Ethnic Distribution ⁽²⁾								
County	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaska Native	Asian/Pacific Islander	Multiple Races	All Other/Unknown
		%	%	%	%	%	%	%
Allen	84	*	*	*	*	*	*	*
Anderson	56	*	*	*	*	*	*	*
Atchison	140	88.6	9.3	1.4	0.0	0.0	0.7	0.0
Barber	56	*	*	*	*	*	*	*
Barton	255	68.6	2.4	28.2	0.0	0.8	0.0	0.0
Bourbon	127	93.7	1.6	2.4	0.8	0.8	0.8	0.0
Brown	91	*	*	*	*	*	*	*
Butler	285	89.8	1.4	7.0	0.7	0.0	1.1	0.0
Chase	9	*	*	*	*	*	*	*
Chautauqua	19	*	*	*	*	*	*	*
Cherokee	183	90.7	1.6	1.1	1.6	2.7	2.2	0.0
Cheyenne	10	*	*	*	*	*	*	*
Clark	2	*	*	*	*	*	*	*
Clay	51	*	*	*	*	*	*	*
Cloud	93	*	*	*	*	*	*	*
Coffey	33	*	*	*	*	*	*	*
Comanche	11	*	*	*	*	*	*	*
Cowley	363	75.5	2.2	19.0	0.8	1.7	0.8	0.0
Crawford	339	83.2	4.1	9.4	1.5	0.6	1.2	0.0
Decatur	14	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3B
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Racial and Ethnic Distribution by County

Comparison of Racial and Ethnic Distribution ⁽²⁾								
County	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Dickinson	155	90.3	0.0	6.5	1.3	0.6	1.3	0.0
Doniphan	53	*	*	*	*	*	*	*
Douglas	530	67.5	9.4	10.2	7.5	3.0	2.3	0.0
Edwards	25	*	*	*	*	*	*	*
Elk	24	*	*	*	*	*	*	*
Ellis	158	82.9	1.3	12.0	1.3	0.6	1.9	0.0
Ellsworth	37	*	*	*	*	*	*	*
Finney	703	27.9	1.6	65.6	0.0	4.6	0.4	0.0
Ford	697	21.8	2.3	75.0	0.3	0.6	0.0	0.0
Franklin	185	94.1	0.0	2.2	2.2	0.5	1.1	0.0
Geary	421	58.0	20.0	15.2	1.0	3.1	2.9	0.0
Gove	10	*	*	*	*	*	*	*
Graham	20	*	*	*	*	*	*	*
Grant	66	*	*	*	*	*	*	*
Gray	46	*	*	*	*	*	*	*
Greeley	12	*	*	*	*	*	*	*
Greenwood	47	*	*	*	*	*	*	*
Hamilton	27	*	*	*	*	*	*	*
Harper	50	*	*	*	*	*	*	*
Harvey	283	70.0	3.2	21.9	1.4	1.4	2.1	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3B
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Racial and Ethnic Distribution by County

Comparison of Racial and Ethnic Distribution ⁽²⁾								
County	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Haskell	17	*	*	*	*	*	*	*
Hodgeman	11	*	*	*	*	*	*	*
Jackson	88	*	*	*	*	*	*	*
Jefferson	58	*	*	*	*	*	*	*
Jewell	7	*	*	*	*	*	*	*
Johnson	1,990	47.8	13.0	34.0	0.5	3.3	1.5	0.0
Kearny	21	*	*	*	*	*	*	*
Kingman	49	*	*	*	*	*	*	*
Kiowa	8	*	*	*	*	*	*	*
Labette	176	89.2	3.4	4.0	1.7	0.0	1.7	0.0
Lane	11	*	*	*	*	*	*	*
Leavenworth	364	69.2	13.2	12.4	0.8	1.6	2.7	0.0
Lincoln	20	*	*	*	*	*	*	*
Linn	71	*	*	*	*	*	*	*
Logan	12	*	*	*	*	*	*	*
Lyon	294	59.2	5.1	33.0	0.3	1.4	1.0	0.0
Marion	125	82.4	1.6	13.6	0.8	1.6	0.0	0.0
Marshall	61	*	*	*	*	*	*	*
Mcperson	52	*	*	*	*	*	*	*
Meade	38	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3B
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Racial and Ethnic Distribution by County

Comparison of Racial and Ethnic Distribution ⁽²⁾								
County	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Miami	151	90.1	2.6	4.0	0.7	0.7	2.0	0.0
Mitchell	36	*	*	*	*	*	*	*
Montgomery	345	75.7	11.0	9.3	1.7	1.2	1.2	0.0
Morris	39	*	*	*	*	*	*	*
Nemaha	38	*	*	*	*	*	*	*
Neosho	166	92.2	1.2	4.2	1.2	0.0	1.2	0.0
Ness	22	*	*	*	*	*	*	*
Norton	38	*	*	*	*	*	*	*
Osage	85	*	*	*	*	*	*	*
Osborne	26	*	*	*	*	*	*	*
Ottawa	29	*	*	*	*	*	*	*
Pawnee	58	*	*	*	*	*	*	*
Phillips	41	*	*	*	*	*	*	*
Pottawatom	36	*	*	*	*	*	*	*
Pratt	82	*	*	*	*	*	*	*
Rawlins	7	*	*	*	*	*	*	*
Reno	453	77.5	2.2	16.6	0.7	0.9	2.2	0.0
Republic	27	*	*	*	*	*	*	*
Rice	89	*	*	*	*	*	*	*
Riley	1,010	70.2	11.1	13.5	0.3	4.2	0.8	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3B
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Racial and Ethnic Distribution by County

Comparison of Racial and Ethnic Distribution ⁽²⁾								
County	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Rooks	50	*	*	*	*	*	*	*
Rush	10	*	*	*	*	*	*	*
Russell	44	*	*	*	*	*	*	*
Saline	546	72.5	3.7	20.3	0.0	1.6	1.8	0.0
Scott	36	*	*	*	*	*	*	*
Sedgwick	4,551	45.9	18.1	29.4	1.3	3.4	1.9	0.0
Seward	428	15.0	2.1	82.0	0.5	0.5	0.0	0.0
Shawnee	1,504	54.3	15.2	24.7	2.8	1.0	2.0	0.0
Sheridan	8	*	*	*	*	*	*	*
Sherman	72	*	*	*	*	*	*	*
Smith	15	*	*	*	*	*	*	*
Stafford	33	*	*	*	*	*	*	*
Stanton	16	*	*	*	*	*	*	*
Stevens	70	*	*	*	*	*	*	*
Sumner	164	87.8	1.2	7.9	1.2	0.6	1.2	0.0
Thomas	50	*	*	*	*	*	*	*
Trego	14	*	*	*	*	*	*	*
Wabaunsee	17	*	*	*	*	*	*	*
Washington	22	*	*	*	*	*	*	*
Wichita	18	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3B
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Racial and Ethnic Distribution by County

Comparison of Racial and Ethnic Distribution ⁽²⁾								
County	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Wilson	75	*	*	*	*	*	*	*
Woodson	18	*	*	*	*	*	*	*
Wyandotte	2,100	26.8	29.4	38.3	0.5	4.0	1.1	0.0
Unknown Counties	20	*	*	*	*	*	*	*
Kansas	21,902	57.3	11.2	26.7	1.2	2.3	1.4	0.0
Nation (Prior Year)	1,304,320	41.6	24.1	28.3	1.1	3.2	1.2	0.5

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4B
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Age Distribution by County

Comparison of Age Distribution ⁽²⁾								
County	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Allen	84	*	*	*	*	*	*	*
Anderson	56	*	*	*	*	*	*	*
Atchison	140	0.0	7.9	12.9	61.4	17.1	0.7	0.0
Barber	56	*	*	*	*	*	*	*
Barton	255	0.0	7.5	12.5	60.8	16.1	3.1	0.0
Bourbon	127	0.8	7.9	15.7	59.1	15.7	0.8	0.0
Brown	91	*	*	*	*	*	*	*
Butler	285	0.0	5.6	14.7	63.9	15.1	0.4	0.4
Chase	9	*	*	*	*	*	*	*
Chautauqua	19	*	*	*	*	*	*	*
Cherokee	183	0.5	10.9	22.4	54.6	11.5	0.0	0.0
Cheyenne	10	*	*	*	*	*	*	*
Clark	2	*	*	*	*	*	*	*
Clay	51	*	*	*	*	*	*	*
Cloud	93	*	*	*	*	*	*	*
Coffey	33	*	*	*	*	*	*	*
Comanche	11	*	*	*	*	*	*	*
Cowley	363	0.6	6.6	17.1	62.5	12.9	0.3	0.0
Crawford	339	0.0	4.1	14.2	69.3	12.4	0.0	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4B
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Age Distribution by County

Comparison of Age Distribution ⁽²⁾								
County	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Decatur	14	*	*	*	*	*	*	*
Dickinson	155	0.0	2.6	14.2	67.7	15.5	0.0	0.0
Doniphan	53	*	*	*	*	*	*	*
Douglas	530	0.0	4.2	12.5	64.3	17.9	1.1	0.0
Edwards	25	*	*	*	*	*	*	*
Elk	24	*	*	*	*	*	*	*
Ellis	158	1.3	6.3	17.7	58.2	15.8	0.0	0.6
Ellsworth	37	*	*	*	*	*	*	*
Finney	703	0.4	6.7	14.1	57.6	18.9	2.3	0.0
Ford	697	0.3	6.0	12.1	53.8	24.8	3.0	0.0
Franklin	185	0.5	5.4	14.1	64.9	14.6	0.5	0.0
Geary	421	0.2	4.0	12.8	70.3	11.4	1.0	0.2
Gove	10	*	*	*	*	*	*	*
Graham	20	*	*	*	*	*	*	*
Grant	66	*	*	*	*	*	*	*
Gray	46	*	*	*	*	*	*	*
Greeley	12	*	*	*	*	*	*	*
Greenwood	47	*	*	*	*	*	*	*
Hamilton	27	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4B
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Age Distribution by County

Comparison of Age Distribution ⁽²⁾								
County	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Harper	50	*	*	*	*	*	*	*
Harvey	283	0.0	4.6	13.4	64.0	17.3	0.7	0.0
Haskell	17	*	*	*	*	*	*	*
Hodgeman	11	*	*	*	*	*	*	*
Jackson	88	*	*	*	*	*	*	*
Jefferson	58	*	*	*	*	*	*	*
Jewell	7	*	*	*	*	*	*	*
Johnson	1,990	0.1	3.0	10.0	60.9	23.8	2.2	0.1
Kearny	21	*	*	*	*	*	*	*
Kingman	49	*	*	*	*	*	*	*
Kiowa	8	*	*	*	*	*	*	*
Labette	176	0.0	6.8	20.5	59.1	13.1	0.6	0.0
Lane	11	*	*	*	*	*	*	*
Leavenworth	364	0.3	5.5	10.2	69.8	13.7	0.5	0.0
Lincoln	20	*	*	*	*	*	*	*
Linn	71	*	*	*	*	*	*	*
Logan	12	*	*	*	*	*	*	*
Lyon	294	1.0	5.8	15.6	60.9	15.3	1.4	0.0
Marion	125	0.8	3.2	12.8	62.4	18.4	2.4	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4B
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Age Distribution by County

Comparison of Age Distribution ⁽²⁾								
County	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Marshall	61	*	*	*	*	*	*	*
Mcpherson	52	*	*	*	*	*	*	*
Meade	38	*	*	*	*	*	*	*
Miami	151	0.7	4.0	18.5	61.6	13.9	1.3	0.0
Mitchell	36	*	*	*	*	*	*	*
Montgomery	345	0.6	7.0	17.7	60.3	12.2	2.3	0.0
Morris	39	*	*	*	*	*	*	*
Nemaha	38	*	*	*	*	*	*	*
Neosho	166	0.0	5.4	15.1	65.1	13.3	1.2	0.0
Ness	22	*	*	*	*	*	*	*
Norton	38	*	*	*	*	*	*	*
Osage	85	*	*	*	*	*	*	*
Osborne	26	*	*	*	*	*	*	*
Ottawa	29	*	*	*	*	*	*	*
Pawnee	58	*	*	*	*	*	*	*
Phillips	41	*	*	*	*	*	*	*
Pottawatom	36	*	*	*	*	*	*	*
Pratt	82	*	*	*	*	*	*	*
Rawlins	7	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4B
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Age Distribution by County

Comparison of Age Distribution ⁽²⁾								
County	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Reno	453	0.2	7.7	14.3	63.1	14.1	0.4	0.0
Republic	27	*	*	*	*	*	*	*
Rice	89	*	*	*	*	*	*	*
Riley	1,010	0.1	1.9	11.2	73.8	12.9	0.2	0.0
Rooks	50	*	*	*	*	*	*	*
Rush	10	*	*	*	*	*	*	*
Russell	44	*	*	*	*	*	*	*
Saline	546	0.0	5.5	16.5	61.0	15.9	0.9	0.2
Scott	36	*	*	*	*	*	*	*
Sedgwick	4,551	0.3	6.6	13.6	60.2	18.0	1.2	0.0
Seward	428	0.2	7.7	11.0	58.2	21.7	1.2	0.0
Shawnee	1,504	0.3	6.3	14.0	60.3	18.0	1.1	0.1
Sheridan	8	*	*	*	*	*	*	*
Sherman	72	*	*	*	*	*	*	*
Smith	15	*	*	*	*	*	*	*
Stafford	33	*	*	*	*	*	*	*
Stanton	16	*	*	*	*	*	*	*
Stevens	70	*	*	*	*	*	*	*
Sumner	164	0.0	8.5	13.4	59.8	17.1	1.2	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4B
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Age Distribution by County

Comparison of Age Distribution ⁽²⁾								
County	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Thomas	50	*	*	*	*	*	*	*
Trego	14	*	*	*	*	*	*	*
Wabaunsee	17	*	*	*	*	*	*	*
Washington	22	*	*	*	*	*	*	*
Wichita	18	*	*	*	*	*	*	*
Wilson	75	*	*	*	*	*	*	*
Woodson	18	*	*	*	*	*	*	*
Wyandotte	2,100	0.5	9.0	13.3	55.5	20.1	1.7	0.0
Unknown Counties	20	*	*	*	*	*	*	*
Kansas	21,902	0.3	5.9	13.5	61.3	17.7	1.3	0.0
Nation (Prior Year)	1,304,320	0.3	5.8	12.3	59.4	20.4	1.5	0.3

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5B
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Health Indicators by County

Comparison of Maternal Health Indicators ^(2, 3)														
County	Prepregnancy BMI ⁽⁴⁾			Weight Gain ⁽⁵⁾			Anemia (Low Hb/Hct) ⁽⁶⁾				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Allen	82	* *	* *	72	* *	* *	9	* *	23	* *	75	* *	75	* *
Anderson	54	* *	* *	51	* *	* *	5	* *	11	* *	53	* *	53	* *
Atchison	138	4.3 (15)	48.6 (3)	119	12.6 (1)	63.0 (31)	18	* *	24	* *	105	4.8 (6)	106	11.3(14)
Barber	56	* *	* *	47	* *	* *	6	* *	12	* *	47	* *	47	* *
Barton	244	5.7 (26)	57.8 (25)	213	23.9 (22)	51.2 (16)	23	* *	63	* *	222	8.6 (25)	222	14.0(26)
Bourbon	125	0.8 (2)	56.8 (21)	108	29.6 (31)	48.1 (9)	9	* *	20	* *	108	8.3 (24)	109	15.6(29)
Brown	90	* *	* *	74	* *	* *	11	* *	18	* *	75	* *	75	* *
Butler	280	6.1 (27)	53.9 (11)	253	21.3 (16)	57.3 (29)	41	* *	44	* *	260	4.6 (5)	260	11.2(13)
Chase	9	* *	* *	7	* *	* *	0	* *	0	* *	8	* *	8	* *
Chautauqua	18	* *	* *	15	* *	* *	2	* *	4	* *	16	* *	16	* *
Cherokee	170	7.6 (31)	48.8 (4)	152	21.7 (17)	53.3 (25)	24	* *	35	* *	163	8.0 (21)	162	11.7(19)
Cheyenne	9	* *	* *	6	* *	* *	3	* *	1	* *	7	* *	7	* *
Clark	2	* *	* *	2	* *	* *	1	* *	0	* *	2	* *	2	* *
Clay	48	* *	* *	41	* *	* *	8	* *	14	* *	45	* *	45	* *
Cloud	90	* *	* *	81	* *	* *	9	* *	23	* *	83	* *	84	* *
Coffey	31	* *	* *	23	* *	* *	5	* *	5	* *	25	* *	25	* *
Comanche	11	* *	* *	9	* *	* *	1	* *	2	* *	9	* *	9	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5B
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Health Indicators by County

Comparison of Maternal Health Indicators ^(2, 3)														
County	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	%	%	Number	%	%	Number	%	Number	%	Number	%	Number	%
	(Rank)	(Rank)		(Rank)	(Rank)		(Rank)		(Rank)		(Rank)		(Rank)	
Cowley	355	3.4 (9)	54.9 (17)	307	22.1 (18)	50.8 (14)	40	* *	57	* *	315	8.9 (29)	316	9.8(10)
Crawford	337	6.5 (28)	52.2 (8)	283	24.0 (23)	46.3 (8)	45	* *	43	* *	290	10.0 (31)	290	12.4(20)
Decatur	14	* *	* *	14	* *	* *	4	* *	4	* *	14	* *	14	* *
Dickinson	149	0.7 (1)	65.1 (31)	124	25.8 (26)	44.4 (5)	19	* *	27	* *	131	6.1 (11)	131	13.7(24)
Doniphan	52	* *	* *	44	* *	* *	10	* *	10	* *	46	* *	46	* *
Douglas	513	2.9 (6)	48.3 (1)	423	19.4 (10)	51.3 (18)	62	* *	130	32.3 (6)	439	7.5 (18)	439	9.6(9)
Edwards	25	* *	* *	21	* *	* *	6	* *	3	* *	21	* *	21	* *
Elk	24	* *	* *	20	* *	* *	3	* *	5	* *	19	* *	19	* *
Ellis	155	6.5 (28)	51.0 (6)	135	15.6 (3)	49.6 (11)	28	* *	27	* *	139	2.9 (1)	139	6.5(2)
Ellsworth	35	* *	* *	31	* *	* *	7	* *	3	* *	33	* *	33	* *
Finney	690	4.9 (21)	53.2 (10)	614	25.7 (25)	43.2 (2)	92	* *	95	* *	634	8.7 (26)	634	7.6(4)
Ford	687	2.2 (3)	59.1 (27)	595	28.4 (29)	44.4 (5)	84	* *	91	* *	596	6.5 (13)	595	8.6(5)
Franklin	172	5.2 (24)	48.3 (1)	148	18.2 (5)	50.0 (12)	20	* *	48	* *	139	5.0 (7)	139	10.1(12)
Geary	405	4.2 (14)	56.8 (21)	317	20.2 (11)	52.7 (23)	61	* *	64	* *	332	3.9 (2)	331	11.5(16)
Gove	8	* *	* *	6	* *	* *	1	* *	5	* *	8	* *	8	* *
Graham	20	* *	* *	16	* *	* *	1	* *	3	* *	16	* *	16	* *
Grant	64	* *	* *	59	* *	* *	7	* *	17	* *	62	* *	62	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5B
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Health Indicators by County

Comparison of Maternal Health Indicators ^(2, 3)														
County	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Gray	42	* *	* *	33	* *	* *	6	* *	13	* *	37	* *	37	* *
Greeley	12	* *	* *	9	* *	* *	0	* *	1	* *	9	* *	9	* *
Greenwood	45	* *	* *	37	* *	* *	5	* *	9	* *	36	* *	36	* *
Hamilton	27	* *	* *	26	* *	* *	10	* *	7	* *	25	* *	25	* *
Harper	47	* *	* *	43	* *	* *	4	* *	4	* *	44	* *	44	* *
Harvey	276	4.7 (18)	59.4 (30)	244	18.9 (8)	51.2 (16)	38	* *	39	* *	251	4.4 (4)	250	11.6(17)
Haskell	17	* *	* *	16	* *	* *	0	* *	1	* *	16	* *	16	* *
Hodgeman	10	* *	* *	8	* *	* *	1	* *	1	* *	8	* *	8	* *
Jackson	84	* *	* *	77	* *	* *	9	* *	17	* *	78	* *	79	* *
Jefferson	56	* *	* *	48	* *	* *	10	* *	13	* *	50	* *	50	* *
Jewell	7	* *	* *	7	* *	* *	1	* *	2	* *	7	* *	7	* *
Johnson	1,916	3.2 (7)	54.8 (16)	1,651	18.7 (6)	51.0 (15)	281	22.4 (1)	462	17.5 (1)	1,701	7.8 (20)	1,695	6.8(3)
Kearny	21	* *	* *	21	* *	* *	2	* *	1	* *	21	* *	21	* *
Kingman	49	* *	* *	42	* *	* *	2	* *	9	* *	43	* *	43	* *
Kiowa	7	* *	* *	7	* *	* *	2	* *	2	* *	8	* *	8	* *
Labette	172	4.7 (18)	59.3 (28)	154	26.0 (27)	50.0 (12)	35	* *	39	* *	154	6.5 (13)	154	14.3(27)
Lane	10	* *	* *	9	* *	* *	2	* *	3	* *	10	* *	10	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5B
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Health Indicators by County

Comparison of Maternal Health Indicators ^(2, 3)														
County	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Leavenworth	348	4.0 (11)	56.0 (19)	292	20.5 (13)	56.2 (28)	51	* *	92	* *	304	5.6 (10)	303	13.2(22)
Lincoln	19	* *	* *	17	* *	* *	6	* *	3	* *	17	* *	17	* *
Linn	71	* *	* *	63	* *	* *	11	* *	9	* *	63	* *	64	* *
Logan	11	* *	* *	9	* *	* *	2	* *	1	* *	9	* *	9	* *
Lyon	288	2.4 (4)	56.6 (20)	249	25.3 (24)	44.2 (4)	34	* *	57	* *	252	6.7 (16)	252	13.5(23)
Marion	122	3.3 (8)	58.2 (26)	105	29.5 (30)	43.8 (3)	14	* *	27	* *	109	9.2 (30)	110	9.1(7)
Marshall	59	* *	* *	46	* *	* *	8	* *	11	* *	50	* *	51	* *
Mcperson	51	* *	* *	49	* *	* *	5	* *	12	* *	49	* *	48	* *
Meade	37	* *	* *	34	* *	* *	1	* *	6	* *	33	* *	33	* *
Miami	147	3.4 (9)	55.1 (18)	125	13.6 (2)	58.4 (30)	30	* *	34	* *	132	6.8 (17)	132	11.4(15)
Mitchell	35	* *	* *	30	* *	* *	2	* *	9	* *	32	* *	31	* *
Montgomery	333	4.8 (20)	49.8 (5)	291	23.7 (21)	49.1 (10)	71	* *	70	* *	296	8.8 (28)	296	13.9(25)
Morris	38	* *	* *	34	* *	* *	5	* *	5	* *	35	* *	35	* *
Nemaha	33	* *	* *	30	* *	* *	10	* *	6	* *	34	* *	34	* *
Neosho	164	5.5 (25)	54.3 (13)	147	23.1 (20)	51.7 (21)	26	* *	27	* *	148	6.1 (11)	147	18.4(31)
Ness	21	* *	* *	17	* *	* *	3	* *	3	* *	17	* *	17	* *
Norton	38	* *	* *	35	* *	* *	8	* *	13	* *	35	* *	35	* *

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one county's rate to other counties. Rank 1 = best rate.
- (4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
- (5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
- (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5B
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Health Indicators by County

Comparison of Maternal Health Indicators ^(2, 3)														
County	Prepregnancy BMI ⁽⁴⁾			Weight Gain ⁽⁵⁾			Anemia (Low Hb/Hct) ⁽⁶⁾				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Osage	82	* *	* *	73	* *	* *	11	* *	16	* *	74	* *	74	* *
Osborne	25	* *	* *	22	* *	* *	2	* *	5	* *	23	* *	23	* *
Ottawa	27	* *	* *	24	* *	* *	5	* *	2	* *	29	* *	29	* *
Pawnee	56	* *	* *	44	* *	* *	4	* *	8	* *	46	* *	45	* *
Phillips	39	* *	* *	28	* *	* *	5	* *	6	* *	34	* *	34	* *
Pottawatom	36	* *	* *	33	* *	* *	4	* *	11	* *	33	* *	33	* *
Pratt	82	* *	* *	71	* *	* *	11	* *	14	* *	72	* *	72	* *
Rawlins	7	* *	* *	6	* *	* *	1	* *	2	* *	6	* *	6	* *
Reno	441	4.1 (12)	54.6 (14)	390	19.2 (9)	53.8 (26)	62	* *	76	* *	396	4.3 (3)	397	14.6(28)
Republic	27	* *	* *	22	* *	* *	5	* *	4	* *	22	* *	22	* *
Rice	87	* *	* *	76	* *	* *	12	* *	16	* *	78	* *	78	* *
Riley	960	4.4 (16)	52.8 (9)	765	18.7 (6)	52.7 (23)	152	37.5 (4)	249	30.1 (5)	791	5.1 (8)	792	12.5(21)
Rooks	50	* *	* *	42	* *	* *	7	* *	8	* *	40	* *	40	* *
Rush	10	* *	* *	10	* *	* *	1	* *	3	* *	10	* *	10	* *
Russell	43	* *	* *	39	* *	* *	7	* *	4	* *	40	* *	40	* *
Saline	538	5.0 (23)	57.4 (24)	467	16.5 (4)	55.9 (27)	96	* *	77	* *	471	8.7 (26)	470	9.1(7)
Scott	36	* *	* *	28	* *	* *	2	* *	5	* *	28	* *	28	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5B
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Health Indicators by County

Comparison of Maternal Health Indicators ^(2, 3)														
County	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight	Over- weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Sedgwick	4,426	4.1 (12)	54.2 (12)	3,947	20.6 (14)	51.5 (19)	834	29.4 (2)	986	23.5 (2)	4,070	6.5 (13)	4,063	11.6(17)
Seward	423	2.4 (4)	59.3 (28)	374	23.0 (19)	44.9 (7)	63	* *	69	* *	380	8.2 (23)	378	6.3(1)
Shawnee	1,464	4.9 (21)	54.6 (14)	1,246	20.4 (12)	52.0 (22)	135	49.6 (5)	239	27.6 (4)	1,282	8.0 (21)	1,282	8.8(6)
Sheridan	8	* *	* *	6	* *	* *	0	* *	1	* *	7	* *	7	* *
Sherman	69	* *	* *	57	* *	* *	7	* *	14	* *	60	* *	60	* *
Smith	15	* *	* *	12	* *	* *	2	* *	1	* *	12	* *	12	* *
Stafford	32	* *	* *	28	* *	* *	4	* *	5	* *	28	* *	29	* *
Stanton	15	* *	* *	14	* *	* *	1	* *	8	* *	15	* *	14	* *
Stevens	69	* *	* *	60	* *	* *	15	* *	17	* *	60	* *	60	* *
Sumner	161	7.5 (30)	51.6 (7)	144	26.4 (28)	40.3 (1)	23	* *	43	* *	146	7.5 (18)	147	17.7(30)
Thomas	50	* *	* *	46	* *	* *	5	* *	8	* *	47	* *	47	* *
Trego	14	* *	* *	10	* *	* *	2	* *	0	* *	10	* *	10	* *
Wabaunsee	16	* *	* *	15	* *	* *	4	* *	8	* *	17	* *	17	* *
Washington	21	* *	* *	19	* *	* *	2	* *	8	* *	20	* *	20	* *
Wichita	16	* *	* *	15	* *	* *	2	* *	0	* *	17	* *	17	* *
Wilson	74	* *	* *	69	* *	* *	13	* *	10	* *	69	* *	69	* *
Woodson	18	* *	* *	16	* *	* *	6	* *	8	* *	16	* *	16	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5B
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Health Indicators by County

Comparison of Maternal Health Indicators ^(2, 3)														
County	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Wyandotte	2,037	4.5 (17)	57.1 (23)	1,785	20.8 (15)	51.5 (19)	310	31.0 (3)	390	27.4 (3)	1,811	5.4 (9)	1,809	9.8(10)
Unknown Counties	18	* *	* *	17	* *	* *	1	* *	2	* *	19	* *	19	* *
Kansas	21,267	4.1	55.0	18,455	21.1	50.7	3,176	29.0	4,262	23.7	18,929	6.8	18,913	10.7
Nation (Prior Year)	1,226,875	4.5	52.9	917,032	21.2	48.2	189,944	33.9	412,633	28.8	389,161	5.0	394,368	6.8

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6B
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Behavioral Indicators by County

Comparison of Maternal Behavioral Indicators ^(2, 3)										
County	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	% (Rank)		Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Allen	67	* *	* *		84	* *	84	* *	67	* *
Anderson	41	* *	* *		56	* *	56	* *	50	* *
Atchison	119	89.1 (1)	2.5 (12)		139	40.3 (7)	140	10.7 (27)	120	58.3 (20)
Barber	47	* *	* *		56	* *	56	* *	44	* *
Barton	225	87.6 (4)	0.9 (3)		254	48.8 (1)	255	21.2 (8)	201	59.2 (17)
Bourbon	99	* *	* *		127	40.2 (8)	127	11.8 (24)	94	* *
Brown	54	* *	* *		91	* *	91	* *	76	* *
Butler	255	74.9 (21)	3.1 (13)		285	33.3 (23)	285	14.0 (16)	224	37.9 (30)
Chase	8	* *	* *		7	* *	9	* *	9	* *
Chautauqua	16	* *	* *		19	* *	19	* *	16	* *
Cherokee	117	73.5 (23)	9.4 (25)		183	35.5 (20)	183	11.5 (25)	148	55.4 (23)
Cheyenne	6	* *	* *		10	* *	10	* *	10	* *
Clark	2	* *	* *		2	* *	2	* *	2	* *
Clay	46	* *	* *		51	* *	51	* *	46	* *
Cloud	85	* *	* *		92	* *	93	* *	82	* *
Coffey	25	* *	* *		33	* *	33	* *	31	* *
Comanche	9	* *	* *		11	* *	11	* *	9	* *
Cowley	297	72.4 (24)	6.4 (23)		360	41.4 (6)	363	13.2 (19)	303	56.1 (22)
Crawford	271	87.5 (6)	4.4 (19)		339	40.1 (9)	339	14.2 (14)	282	59.2 (17)
Decatur	13	* *	* *		14	* *	14	* *	10	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6B
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Behavioral Indicators by County

Comparison of Maternal Behavioral Indicators ^(2, 3)										
County	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Prepregnancy ⁽⁴⁾		During Pregnancy ⁽⁵⁾	
	Number	% (Rank)	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Dickinson	133	88.0 (3)	1.5 (8)		153	37.3 (18)	155	21.9 (6)	125	69.6 (4)
Doniphan	47	* *	* *		53	* *	53	* *	41	* *
Douglas	443	76.1 (17)	5.6 (21)		528	38.6 (14)	530	21.5 (7)	434	64.5 (12)
Edwards	18	* *	* *		24	* *	25	* *	23	* *
Elk	19	* *	* *		24	* *	24	* *	21	* *
Ellis	141	86.5 (8)	0.7 (2)		158	39.9 (10)	158	25.9 (2)	137	65.0 (10)
Ellsworth	33	* *	* *		37	* *	37	* *	30	* *
Finney	603	52.4 (29)	29.2 (29)		694	35.2 (21)	703	12.4 (23)	589	48.6 (29)
Ford	533	72.2 (25)	12.4 (27)		695	44.7 (4)	697	13.9 (17)	591	52.1 (27)
Franklin	76	* *	* *		185	31.4 (25)	185	8.6 (28)	127	63.8 (13)
Geary	317	76.0 (18)	1.6 (10)		421	30.2 (28)	421	22.1 (5)	342	58.5 (19)
Gove	8	* *	* *		10	* *	10	* *	7	* *
Graham	16	* *	* *		20	* *	20	* *	17	* *
Grant	58	* *	* *		65	* *	66	* *	52	* *
Gray	37	* *	* *		46	* *	46	* *	44	* *
Greeley	8	* *	* *		12	* *	12	* *	10	* *
Greenwood	37	* *	* *		47	* *	47	* *	43	* *
Hamilton	24	* *	* *		27	* *	27	* *	25	* *
Harper	44	* *	* *		48	* *	50	* *	43	* *
Harvey	220	73.6 (22)	11.4 (26)		281	48.4 (2)	283	19.4 (10)	235	66.0 (9)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6B
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Behavioral Indicators by County

Comparison of Maternal Behavioral Indicators ^(2, 3)										
County	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Pregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	%	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Haskell	15	* *	* *	* *	16	* *	17	* *	15	* *
Hodgeman	8	* *	* *	* *	11	* *	11	* *	10	* *
Jackson	38	* *	* *	* *	88	* *	88	* *	74	* *
Jefferson	48	* *	* *	* *	58	* *	58	* *	49	* *
Jewell	7	* *	* *	* *	7	* *	7	* *	6	* *
Johnson	1,518	65.5 (27)	4.0 (18)		1,983	34.1 (22)	1,990	13.5 (18)	1,598	60.8 (16)
Kearny	19	* *	* *	* *	21	* *	21	* *	16	* *
Kingman	43	* *	* *	* *	49	* *	49	* *	39	* *
Kiowa	8	* *	* *	* *	8	* *	8	* *	6	* *
Labette	128	86.7 (7)	1.6 (10)		175	30.9 (26)	176	11.4 (26)	146	67.8 (7)
Lane	10	* *	* *	* *	11	* *	11	* *	10	* *
Leavenworth	307	83.7 (11)	1.0 (4)		364	28.0 (30)	364	22.3 (4)	273	68.5 (6)
Lincoln	17	* *	* *	* *	20	* *	20	* *	20	* *
Linn	65	* *	* *	* *	71	* *	71	* *	60	* *
Logan	7	* *	* *	* *	12	* *	12	* *	11	* *
Lyon	253	83.8 (10)	1.2 (5)		294	38.8 (13)	294	17.7 (13)	237	64.6 (11)
Marion	109	84.4 (9)	0.0 (1)		125	38.4 (15)	125	28.0 (1)	106	73.6 (1)
Marshall	48	* *	* *	* *	60	* *	61	* *	48	* *
Mcperson	50	* *	* *	* *	52	* *	52	* *	43	* *
Meade	33	* *	* *	* *	38	* *	38	* *	36	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6B
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Behavioral Indicators by County

Comparison of Maternal Behavioral Indicators ^(2, 3)										
County	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	%	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Miami	131	79.4 (14)	1.5	(8)	151	37.7 (17)	151	18.5 (11)	127	55.1 (24)
Mitchell	32	* *	* *		36	* *	36	* *	31	* *
Montgomery	217	87.6 (4)	1.4	(7)	345	25.2 (31)	345	14.2 (14)	287	69.0 (5)
Morris	32	* *	* *		38	* *	39	* *	32	* *
Nemaha	33	* *	* *		38	* *	38	* *	32	* *
Neosho	133	88.7 (2)	3.8	(17)	165	39.4 (11)	166	12.7 (22)	139	71.9 (2)
Ness	13	* *	* *		22	* *	22	* *	20	* *
Norton	35	* *	* *		38	* *	38	* *	32	* *
Osage	59	* *	* *		85	* *	85	* *	70	* *
Osborne	21	* *	* *		26	* *	26	* *	18	* *
Ottawa	28	* *	* *		29	* *	29	* *	25	* *
Pawnee	45	* *	* *		57	* *	58	* *	49	* *
Phillips	34	* *	* *		41	* *	41	* *	36	* *
Pottawatom	30	* *	* *		36	* *	36	* *	32	* *
Pratt	72	* *	* *		82	* *	82	* *	66	* *
Rawlins	6	* *	* *		7	* *	7	* *	5	* *
Reno	388	83.2 (12)	3.6	(15)	451	39.2 (12)	453	20.8 (9)	382	63.6 (14)
Republic	22	* *	* *		27	* *	27	* *	23	* *
Rice	77	* *	* *		88	* *	89	* *	69	* *
Riley	740	82.7 (13)	1.2	(5)	1,010	33.2 (24)	1,010	22.5 (3)	793	70.9 (3)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6B
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Behavioral Indicators by County

Comparison of Maternal Behavioral Indicators ^(2, 3)										
County	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	%	(Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Rooks	41	* *	* *	* *	49	* *	50	* *	42	* *
Rush	10	* *	* *	* *	10	* *	10	* *	10	* *
Russell	40	* *	* *	* *	43	* *	44	* *	39	* *
Saline	402	76.4 (16)	3.2 (14)		544	37.9 (16)	546	4.6 (31)	478	54.6 (25)
Scott	27	* *	* *	* *	35	* *	36	* *	32	* *
Sedgwick	3,982	75.1 (20)	5.9 (22)		4,538	28.6 (29)	4,551	18.2 (12)	3,758	50.4 (28)
Seward	336	66.4 (26)	13.7 (28)		423	30.7 (27)	428	8.4 (29)	345	67.0 (8)
Shawnee	1,131	75.7 (19)	3.7 (16)		1,497	45.4 (3)	1,504	13.2 (19)	1,244	53.5 (26)
Sheridan	7	* *	* *	* *	8	* *	8	* *	8	* *
Sherman	55	* *	* *	* *	72	* *	72	* *	56	* *
Smith	11	* *	* *	* *	15	* *	15	* *	14	* *
Stafford	29	* *	* *	* *	33	* *	33	* *	26	* *
Stanton	15	* *	* *	* *	16	* *	16	* *	12	* *
Stevens	60	* *	* *	* *	69	* *	70	* *	60	* *
Sumner	147	78.2 (15)	4.8 (20)		164	42.1 (5)	164	12.8 (21)	136	62.5 (15)
Thomas	40	* *	* *	* *	50	* *	50	* *	38	* *
Trego	11	* *	* *	* *	14	* *	14	* *	13	* *
Wabaunsee	15	* *	* *	* *	17	* *	17	* *	14	* *
Washington	19	* *	* *	* *	21	* *	22	* *	18	* *
Wichita	17	* *	* *	* *	18	* *	18	* *	13	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6B
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Behavioral Indicators by County

Comparison of Maternal Behavioral Indicators ^(2, 3)										
County	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Pregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	%	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Wilson	68	* *	* *	* *	74	* *	75	* *	63	* *
Woodson	18	* *	* *	* *	18	* *	18	* *	15	* *
Wyandotte	1,726	64.1 (28)	8.6	(24)	2,082	36.0 (19)	2,100	6.6 (30)	1,778	57.9 (21)
Unknown Counties	17	* *	* *	* *	20	* *	20	* *	17	* *
Kansas	17,720	75.1	6.1		21,806	36.1	21,902	16.2	18,060	58.1
Nation (Prior Year)	527,184	82.2	3.1		1,213,982	34.1	360,369	18.7	372,743	83.5

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7B
2010 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by County

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Allen	79	* *	60	* *	26	* *	66	* *	62	* *
Anderson	56	* *	52	* *	21	* *	50	* *	53	* *
Atchison	118	42.4 (23)	81	* *	26	* *	100	31.0 (23)	82	* *
Barber	55	* *	44	* *	14	* *	44	* *	45	* *
Barton	253	33.6 (10)	223	23.3 (17)	58	* *	201	10.0 (7)	222	5.9 (5)
Bourbon	118	43.2 (24)	93	* *	29	* *	88	* *	96	* *
Brown	85	* *	60	* *	20	* *	71	* *	62	* *
Butler	281	51.6 (30)	253	31.6 (27)	97	* *	223	13.0 (14)	258	9.7 (13)
Chase	9	* *	8	* *	4	* *	9	* *	8	* *
Chautauqua	18	* *	11	* *	5	* *	15	* *	11	* *
Cherokee	180	44.4 (27)	157	29.9 (25)	54	* *	146	38.4 (27)	159	40.9 (27)
Cheyenne	10	* *	7	* *	2	* *	10	* *	7	* *
Clark	2	* *	1	* *	0	* *	2	* *	1	* *
Clay	51	* *	44	* *	14	* *	46	* *	44	* *
Cloud	93	* *	84	* *	23	* *	82	* *	84	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7B
2010 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by County

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Coffey	32	* *	25	* *	10	* *	30	* *	25	* *
Comanche	11	* *	9	* *	5	* *	9	* *	9	* *
Cowley	355	40.8 (19)	278	28.1 (23)	88	* *	297	19.5 (20)	278	18.0 (21)
Crawford	320	49.1 (29)	247	30.8 (26)	99	* *	271	39.1 (29)	248	36.3 (25)
Decatur	14	* *	12	* *	3	* *	10	* *	13	* *
Dickinson	152	44.1 (26)	130	24.6 (19)	46	* *	122	7.4 (3)	131	6.1 (6)
Doniphan	52	* *	37	* *	8	* *	40	* *	37	* *
Douglas	524	34.9 (12)	431	17.2 (8)	117	40.2 (3)	429	10.3 (10)	429	6.3 (8)
Edwards	25	* *	21	* *	9	* *	23	* *	21	* *
Elk	21	* *	16	* *	6	* *	19	* *	16	* *
Ellis	158	41.1 (20)	136	19.1 (11)	48	* *	136	7.4 (3)	138	3.6 (3)
Ellsworth	37	* *	33	* *	12	* *	30	* *	33	* *
Finney	696	9.3 (3)	616	5.2 (3)	40	* *	586	11.8 (13)	616	10.7 (16)
Ford	671	8.0 (2)	538	4.3 (2)	32	* *	581	8.4 (5)	540	8.0 (11)
Franklin	184	40.2 (18)	156	28.8 (24)	34	* *	126	15.1 (17)	157	10.8 (17)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7B
2010 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by County

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Geary	415	35.2 (13)	328	21.6 (14)	77	* *	338	35.2 (25)	328	34.5 (24)
Gove	10	* *	8	* *	1	* *	7	* *	8	* *
Graham	20	* *	16	* *	6	* *	17	* *	16	* *
Grant	66	* *	62	* *	2	* *	52	* *	62	* *
Gray	46	* *	35	* *	2	* *	44	* *	35	* *
Greeley	12	* *	8	* *	1	* *	10	* *	8	* *
Greenwood	39	* *	13	* *	7	* *	39	* *	21	* *
Hamilton	27	* *	26	* *	3	* *	25	* *	26	* *
Harper	47	* *	41	* *	10	* *	40	* *	38	* *
Harvey	280	40.0 (17)	243	21.0 (13)	78	* *	233	5.6 (1)	242	1.7 (2)
Haskell	17	* *	16	* *	2	* *	15	* *	16	* *
Hodgeman	9	* *	7	* *	4	* *	9	* *	7	* *
Jackson	87	* *	76	* *	33	* *	74	* *	77	* *
Jefferson	15	* *	11	* *	3	* *	13	* *	12	* *
Jewell	7	* *	6	* *	2	* *	6	* *	6	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7B
2010 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by County

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Johnson	1,963	23.9 (5)	1,697	10.4 (4)	278	45.3 (1)	1,573	17.9 (19)	1,700	17.4 (19)
Kearny	21	* *	20	* *	4	* *	16	* *	20	* *
Kingman	43	* *	37	* *	13	* *	37	* *	38	* *
Kiowa	8	* *	8	* *	2	* *	6	* *	8	* *
Labette	170	37.1 (14)	150	27.3 (22)	44	* *	143	30.1 (22)	150	34.0 (23)
Lane	5	* *	1	* *	0	* *	5	* *	2	* *
Leavenworth	361	47.6 (28)	301	23.9 (18)	98	* *	271	38.4 (27)	299	38.8 (26)
Lincoln	20	* *	18	* *	10	* *	20	* *	18	* *
Linn	69	* *	63	* *	37	* *	59	* *	63	* *
Logan	11	* *	10	* *	4	* *	11	* *	10	* *
Lyon	292	31.2 (8)	257	18.7 (10)	61	* *	236	10.6 (11)	257	6.2 (7)
Marion	125	33.6 (10)	110	17.3 (9)	31	* *	105	5.7 (2)	110	0.9 (1)
Marshall	60	* *	50	* *	8	* *	48	* *	50	* *
Mcperson	51	* *	45	* *	21	* *	41	* *	45	* *
Meade	35	* *	32	* *	3	* *	36	* *	33	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7B
2010 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by County

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Miami	143	39.9 (16)	121	19.8 (12)	36	* *	119	24.4 (21)	121	17.4 (19)
Mitchell	36	* *	31	* *	12	* *	31	* *	31	* *
Montgomery	326	42.0 (22)	278	26.3 (21)	94	* *	282	37.9 (26)	277	32.9 (22)
Morris	39	* *	35	* *	14	* *	32	* *	34	* *
Nemaha	31	* *	20	* *	5	* *	28	* *	21	* *
Neosho	139	43.9 (25)	74	* *	27	* *	126	34.1 (24)	76	* *
Ness	17	* *	14	* *	3	* *	17	* *	14	* *
Norton	37	* *	34	* *	13	* *	32	* *	34	* *
Osage	76	* *	57	* *	17	* *	62	* *	57	* *
Osborne	26	* *	23	* *	8	* *	18	* *	23	* *
Ottawa	28	* *	12	* *	6	* *	25	* *	23	* *
Pawnee	55	* *	37	* *	7	* *	48	* *	38	* *
Phillips	36	* *	23	* *	8	* *	31	* *	23	* *
Pottawatom	36	* *	32	* *	8	* *	32	* *	33	* *
Pratt	79	* *	67	* *	18	* *	66	* *	66	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7B
2010 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by County

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Rawlins	7	* *	6	* *	1	* *	5	* *	6	* *
Reno	448	41.7 (21)	392	21.9 (15)	139	41.0 (2)	378	13.5 (15)	392	11.2 (18)
Republic	27	* *	21	* *	6	* *	22	* *	21	* *
Rice	85	* *	73	* *	20	* *	66	* *	74	* *
Riley	961	29.1 (6)	664	13.7 (6)	140	39.3 (4)	774	15.4 (18)	666	10.2 (14)
Rooks	48	* *	37	* *	19	* *	40	* *	39	* *
Rush	10	* *	10	* *	7	* *	10	* *	10	* *
Russell	44	* *	39	* *	16	* *	39	* *	39	* *
Saline	543	39.4 (15)	470	25.3 (20)	165	29.7 (7)	478	10.9 (12)	469	7.7 (10)
Scott	34	* *	22	* *	8	* *	30	* *	22	* *
Sedgwick	4,518	29.2 (7)	4,065	16.1 (7)	911	33.3 (5)	3,742	9.2 (6)	4,055	6.6 (9)
Seward	422	5.7 (1)	377	3.2 (1)	17	* *	342	10.2 (9)	382	8.9 (12)
Shawnee	1,386	32.3 (9)	1,079	22.0 (16)	266	25.9 (8)	1,119	10.0 (7)	1,092	4.0 (4)
Sheridan	8	* *	6	* *	2	* *	8	* *	6	* *
Sherman	70	* *	59	* *	20	* *	54	* *	59	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7B
2010 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by County

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Smith	13	* *	10	* *	5	* *	12	* *	10	* *
Stafford	31	* *	27	* *	6	* *	25	* *	27	* *
Stanton	16	* *	15	* *	1	* *	12	* *	15	* *
Stevens	70	* *	60	* *	4	* *	60	* *	60	* *
Sumner	163	54.6 (31)	144	34.0 (28)	63	* *	135	44.4 (30)	144	44.4 (28)
Thomas	50	* *	46	* *	9	* *	38	* *	47	* *
Trego	14	* *	10	* *	4	* *	13	* *	10	* *
Wabaunsee	17	* *	16	* *	6	* *	14	* *	16	* *
Washington	22	* *	19	* *	1	* *	18	* *	19	* *
Wichita	17	* *	16	* *	4	* *	12	* *	16	* *
Wilson	74	* *	69	* *	35	* *	63	* *	69	* *
Woodson	15	* *	9	* *	0	* *	13	* *	10	* *
Wyandotte	2,056	20.6 (4)	1,754	10.9 (5)	271	32.5 (6)	1,761	14.7 (16)	1,807	10.6 (15)
Unknown Counties	19	* *	19	* *	10	* *	17	* *	20	* *
Kansas	21,283	30.7	17,950	17.5	4,227	33.0	17,640	16.5	18,063	13.3

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7B
2010 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by County

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal	Postpartum			
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)		
Nation (Prior Year)	1,040,568	22.9	838,781	13.0	156,709	36.7	650,648	17.1	559,995	16.8

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one county's rate to other counties. Rank 1 = best rate.
- (4) Among women who smoked 3 months prior to pregnancy.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8B
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Infant Health Indicators by County

Comparison of Infant Health Indicators ^(2, 3)										
County	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)		Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
	Number	% (Rank)	% (Rank)	Number						
Allen	69	* * * *	* * * *	73	* *	62	* *	72	* *	
Anderson	43	* * * *	* * * *	51	* *	39	* *	50	* *	
Atchison	115	9.6 (28)	3.5 (3)	122	6.6 (4)	106	7.5 (28)	121	48.8 (31)	
Barber	43	* * * *	* * * *	43	* *	38	* *	43	* *	
Barton	215	7.4 (20)	4.2 (5)	216	5.6 (3)	202	5.9 (27)	217	77.9 (7)	
Bourbon	103	13.6 (30)	1.9 (1)	106	10.4 (27)	92	* *	105	62.9 (23)	
Brown	67	* * * *	* * * *	73	* *	54	* *	73	* *	
Butler	252	7.1 (15)	6.0 (14)	257	10.1 (26)	226	3.5 (10)	256	65.2 (22)	
Chase	8	* * * *	* * * *	6	* *	6	* *	8	* *	
Chautauqua	15	* * * *	* * * *	16	* *	15	* *	16	* *	
Cherokee	160	9.4 (27)	5.6 (11)	163	9.2 (20)	145	2.8 (6)	163	53.4 (30)	
Cheyenne	7	* * * *	* * * *	7	* *	5	* *	7	* *	
Clark	2	* * * *	* * * *	2	* *	2	* *	2	* *	
Clay	45	* * * *	* * * *	45	* *	40	* *	45	* *	

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8B
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Infant Health Indicators by County

Comparison of Infant Health Indicators ^(2, 3)										
County	Birthweight				Preterm ⁽⁶⁾		Full Term LBW ⁽⁷⁾		Ever Breastfed	
	Low ⁽⁴⁾		High ⁽⁵⁾		Number	%	Number	%	Number	%
	Number	% (Rank)	% (Rank)	(Rank)						
Cloud	84	* * * *	* * * *	84	* *	76	* *	83	* *	
Coffey	25	* * * *	* * * *	25	* *	25	* *	25	* *	
Comanche	7	* * * *	* * * *	7	* *	7	* *	7	* *	
Cowley	302	6.6 (13)	7.9 (25)	317	10.7 (29)	267	3.7 (13)	317	56.8 (29)	
Crawford	277	5.8 (7)	3.2 (2)	282	10.6 (28)	247	3.2 (8)	282	62.1 (25)	
Decatur	13	* * * *	* * * *	13	* *	12	* *	13	* *	
Dickinson	129	6.2 (8)	7.8 (24)	127	11.8 (30)	112	0.9 (2)	129	74.4 (13)	
Doniphan	48	* * * *	* * * *	48	* *	39	* *	48	* *	
Douglas	435	8.3 (22)	8.5 (28)	441	7.7 (11)	400	5.3 (24)	438	79.5 (4)	
Edwards	19	* * * *	* * * *	18	* *	15	* *	19	* *	
Elk	20	* * * *	* * * *	21	* *	19	* *	21	* *	
Ellis	130	6.2 (8)	6.2 (17)	136	8.8 (18)	118	3.4 (9)	136	80.1 (3)	
Ellsworth	32	* * * *	* * * *	32	* *	32	* *	32	* *	
Finney	598	7.2 (17)	6.0 (14)	611	9.2 (20)	539	4.5 (18)	613	77.8 (8)	

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8B
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Infant Health Indicators by County

Comparison of Infant Health Indicators ^(2, 3)										
County	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)							
	Number	% (Rank)	%	(Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Ford	564	5.0 (3)	6.0	(14)	580	7.1 (8)	523	2.9 (7)	575	77.4 (9)
Franklin	153	5.2 (4)	5.9	(13)	153	5.2 (2)	145	4.8 (21)	153	60.8 (26)
Geary	326	6.4 (10)	8.3	(27)	327	8.9 (19)	297	2.0 (3)	324	71.0 (15)
Gove	6	* *	* *	* *	6	* *	5	* *	6	* *
Graham	15	* *	* *	* *	15	* *	12	* *	14	* *
Grant	60	* *	* *	* *	60	* *	56	* *	61	* *
Gray	24	* *	* *	* *	36	* *	23	* *	36	* *
Greeley	9	* *	* *	* *	9	* *	7	* *	9	* *
Greenwood	39	* *	* *	* *	39	* *	37	* *	39	* *
Hamilton	25	* *	* *	* *	26	* *	24	* *	26	* *
Harper	45	* *	* *	* *	43	* *	41	* *	45	* *
Harvey	237	8.4 (23)	4.2	(5)	240	6.7 (5)	219	4.6 (19)	241	83.4 (1)
Haskell	15	* *	* *	* *	15	* *	14	* *	15	* *
Hodgeman	9	* *	* *	* *	9	* *	9	* *	9	* *

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one county's rate to other counties. Rank 1 = best rate.
- (4) <2500 g.
- (5) >4000 g.
- (6) <37 weeks gestation.
- (7) 37 or more weeks gestation and <2500 g.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8B
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Infant Health Indicators by County

Comparison of Infant Health Indicators ^(2, 3)										
County	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)							
	Number	% (Rank)	%	(Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Jackson	76	* * * *	* *	* *	80	* *	73	* *	79	* *
Jefferson	49	* * * *	* *	* *	49	* *	46	* *	48	* *
Jewell	6	* * * *	* *	* *	7	* *	5	* *	7	* *
Johnson	1,601	7.1 (15)	6.7 (20)		1,690	7.3 (9)	1,476	3.5 (10)	1,683	81.8 (2)
Kearny	19	* * * *	* *	* *	20	* *	16	* *	20	* *
Kingman	21	* * * *	* *	* *	41	* *	20	* *	41	* *
Kiowa	7	* * * *	* *	* *	8	* *	7	* *	8	* *
Labette	152	7.2 (17)	7.9 (25)		151	9.3 (23)	137	5.1 (22)	152	57.2 (28)
Lane	10	* * * *	* *	* *	10	* *	7	* *	10	* *
Leavenworth	300	5.3 (5)	8.7 (29)		304	9.2 (20)	272	2.6 (5)	302	66.2 (20)
Lincoln	18	* * * *	* *	* *	18	* *	14	* *	18	* *
Linn	58	* * * *	* *	* *	62	* *	50	* *	62	* *
Logan	11	* * * *	* *	* *	11	* *	11	* *	11	* *
Lyon	252	8.7 (25)	6.7 (20)		252	8.3 (14)	231	5.2 (23)	252	77.0 (11)

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one county's rate to other counties. Rank 1 = best rate.
- (4) <2500 g.
- (5) >4000 g.
- (6) <37 weeks gestation.
- (7) 37 or more weeks gestation and <2500 g.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8B
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Infant Health Indicators by County

Comparison of Infant Health Indicators ^(2, 3)									
County	Birthweight			Preterm ⁽⁶⁾		Full Term LBW ⁽⁷⁾		Ever Breastfed	
	Low ⁽⁴⁾	High ⁽⁵⁾		Number	% ^(Rank)	Number	% ^(Rank)	Number	% ^(Rank)
	Number	% ^(Rank)	% ^(Rank)						
Marion	105	4.8 (2)	4.8 (7)	107	7.5 (10)	97	* *	106	77.4 (9)
Marshall	46	* *	* *	46	* *	42	* *	47	* *
Mcperson	45	* *	* *	48	* *	40	* *	47	* *
Meade	32	* *	* *	36	* *	28	* *	36	* *
Miami	132	3.0 (1)	10.6 (31)	132	4.5 (1)	126	0.8 (1)	132	75.0 (12)
Mitchell	28	* *	* *	32	* *	26	* *	31	* *
Montgomery	296	6.8 (14)	3.7 (4)	299	6.7 (5)	276	3.6 (12)	296	62.2 (24)
Morris	30	* *	* *	34	* *	27	* *	35	* *
Nemaha	15	* *	* *	33	* *	13	* *	29	* *
Neosho	130	8.5 (24)	5.4 (9)	144	8.3 (14)	117	4.3 (17)	143	59.4 (27)
Ness	16	* *	* *	18	* *	16	* *	18	* *
Norton	34	* *	* *	34	* *	30	* *	33	* *
Osage	70	* *	* *	76	* *	59	* *	76	* *
Osborne	22	* *	* *	22	* *	20	* *	22	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8B
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Infant Health Indicators by County

Comparison of Infant Health Indicators ^(2, 3)										
County	Birthweight				Preterm ⁽⁶⁾		Full Term LBW ⁽⁷⁾		Ever Breastfed	
	Low ⁽⁴⁾		High ⁽⁵⁾		Number	% ^(Rank)	Number	% ^(Rank)	Number	% ^(Rank)
	Number	% ^(Rank)	% ^(Rank)							
Ottawa	28	* * * *			29	* *	26	* *	29	* *
Pawnee	46	* * * *			45	* *	40	* *	46	* *
Phillips	31	* * * *			32	* *	31	* *	30	* *
Pottawatom	32	* * * *			32	* *	30	* *	32	* *
Pratt	70	* * * *			72	* *	65	* *	72	* *
Rawlins	5	* * * *			5	* *	5	* *	5	* *
Reno	388	9.0 (26)	7.0 (22)		395	8.6 (17)	354	5.4 (25)	394	70.3 (18)
Republic	21	* * * *			22	* *	19	* *	22	* *
Rice	69	* * * *			71	* *	63	* *	72	* *
Riley	792	5.7 (6)	9.3 (30)		809	6.8 (7)	737	2.4 (4)	802	78.7 (6)
Rooks	39	* * * *			38	* *	35	* *	39	* *
Rush	10	* * * *			10	* *	10	* *	10	* *
Russell	37	* * * *			36	* *	30	* *	37	* *
Saline	471	6.4 (10)	5.5 (10)		476	8.0 (13)	432	4.2 (15)	475	65.9 (21)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8B
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Infant Health Indicators by County

Comparison of Infant Health Indicators ^(2, 3)										
County	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)							
	Number	% (Rank)	%	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Scott	26	* * * *	* *		26	* *	24	* *	26	* *
Sedgwick	3,976	9.6 (28)	7.1 (23)		4,005	8.5 (16)	3,627	5.7 (26)	3,993	70.4 (17)
Seward	346	6.4 (10)	6.6 (19)		368	7.9 (12)	313	3.8 (14)	370	78.9 (5)
Shawnee	1,249	7.2 (17)	6.2 (17)		1,262	9.3 (23)	1,128	4.2 (15)	1,254	70.5 (16)
Sheridan	7	* * * *	* *		7	* *	6	* *	7	* *
Sherman	56	* * * *	* *		56	* *	52	* *	56	* *
Smith	12	* * * *	* *		12	* *	11	* *	12	* *
Stafford	27	* * * *	* *		27	* *	25	* *	27	* *
Stanton	14	* * * *	* *		14	* *	14	* *	14	* *
Stevens	39	* * * *	* *		59	* *	38	* *	58	* *
Sumner	141	15.6 (31)	5.0 (8)		145	14.5 (31)	120	9.2 (29)	144	70.1 (19)
Thomas	44	* * * *	* *		44	* *	41	* *	44	* *
Trego	11	* * * *	* *		11	* *	9	* *	11	* *
Wabaunsee	16	* * * *	* *		16	* *	16	* *	16	* *

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one county's rate to other counties. Rank 1 = best rate.
- (4) <2500 g.
- (5) >4000 g.
- (6) <37 weeks gestation.
- (7) 37 or more weeks gestation and <2500 g.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8B
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Infant Health Indicators by County

Comparison of Infant Health Indicators ^(2, 3)										
County	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)							
	Number	% (Rank)	%	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Washington	16	* * * *	* * * *		18	* *	15	* *	18	* *
Wichita	15	* * * *	* * * *		15	* *	14	* *	15	* *
Wilson	71	* * * *	* * * *		71	* *	63	* *	71	* *
Woodson	15	* * * *	* * * *		18	* *	15	* *	18	* *
Wyandotte	1,715	8.2 (21)	5.7 (12)		1,796	9.6 (25)	1,541	4.6 (19)	1,794	71.9 (14)
Unknown Counties	20	* * * *	* * * *		20	* *	20	* *	19	* *
Kansas	18,226	7.6	6.6		18,726	8.4	16,603	4.4	18,670	72.2
Nation (Prior Year)	1,018,145	8.2	6.7		1,017,333	11.1	848,957	3.5	997,136	66.7

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3B
2008 - 2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Racial and Ethnic Distribution by County
Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾								
County	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Allen	287	92.0	3.8	1.4	1.4	0.0	1.4	0.0
Anderson	156	94.9	0.0	3.8	0.0	1.3	0.0	0.0
Atchison	413	88.9	7.5	1.5	0.7	0.0	1.5	0.0
Barber	141	95.0	2.1	1.4	0.0	1.4	0.0	0.0
Barton	791	68.1	2.1	28.4	0.3	0.5	0.5	0.0
Bourbon	379	92.6	2.1	2.6	0.5	0.5	1.6	0.0
Brown	300	76.3	2.3	4.3	14.3	0.0	2.7	0.0
Butler	853	90.3	1.2	6.1	0.5	0.2	1.8	0.0
Chase	40	*	*	*	*	*	*	*
Chautauqua	83	*	*	*	*	*	*	*
Cherokee	614	90.1	1.3	3.3	1.6	2.3	1.5	0.0
Cheyenne	13	*	*	*	*	*	*	*
Clark	10	*	*	*	*	*	*	*
Clay	177	94.4	0.6	2.3	2.3	0.0	0.6	0.0
Cloud	254	92.5	1.6	2.8	1.6	0.8	0.8	0.0
Coffey	82	*	*	*	*	*	*	*
Comanche	28	*	*	*	*	*	*	*
Cowley	1,082	74.9	2.5	19.3	0.8	1.7	0.8	0.0
Crawford	1,122	83.4	3.6	10.2	0.4	1.2	1.2	0.0
Decatur	33	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3B
2008 - 2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Racial and Ethnic Distribution by County
Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾								
County	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Dickinson	394	91.9	0.0	5.6	1.3	0.5	0.8	0.0
Doniphan	133	88.0	2.3	6.8	0.8	0.0	2.3	0.0
Douglas	1,500	65.5	8.7	10.7	8.5	4.5	2.1	0.0
Edwards	77	*	*	*	*	*	*	*
Elk	57	*	*	*	*	*	*	*
Ellis	481	83.8	1.0	13.3	0.8	0.4	0.6	0.0
Ellsworth	98	*	*	*	*	*	*	*
Finney	2,109	29.7	1.0	65.5	0.0	3.3	0.4	0.0
Ford	2,029	21.8	1.5	76.0	0.2	0.4	0.0	0.0
Franklin	538	91.4	0.9	4.5	1.5	0.2	1.5	0.0
Geary	1,154	53.9	22.7	15.3	0.6	3.2	4.3	0.0
Gove	29	*	*	*	*	*	*	*
Graham	51	*	*	*	*	*	*	*
Grant	205	22.4	0.0	77.1	0.0	0.0	0.5	0.0
Gray	148	68.9	0.0	30.4	0.0	0.0	0.7	0.0
Greeley	37	*	*	*	*	*	*	*
Greenwood	140	93.6	0.7	4.3	0.0	0.0	1.4	0.0
Hamilton	76	*	*	*	*	*	*	*
Harper	156	80.1	0.0	17.9	0.6	0.0	1.3	0.0
Harvey	818	70.3	2.4	23.7	1.2	0.7	1.6	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3B
2008 - 2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Racial and Ethnic Distribution by County
Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾								
County	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Haskell	61	*	*	*	*	*	*	*
Hodgeman	39	*	*	*	*	*	*	*
Jackson	233	76.8	0.0	5.6	15.5	0.0	2.1	0.0
Jefferson	191	94.2	0.5	2.1	2.6	0.0	0.5	0.0
Jewell	30	*	*	*	*	*	*	*
Johnson	5,750	46.4	13.0	35.3	0.5	3.2	1.5	0.0
Kearny	61	*	*	*	*	*	*	*
Kingman	165	90.9	0.0	6.7	1.2	1.2	0.0	0.0
Kiowa	33	*	*	*	*	*	*	*
Labette	536	87.9	6.2	3.2	1.5	0.2	1.1	0.0
Lane	37	*	*	*	*	*	*	*
Leavenworth	1,079	69.0	14.6	10.6	0.6	1.7	3.7	0.0
Lincoln	63	*	*	*	*	*	*	*
Linn	191	95.8	0.0	2.1	1.0	0.0	1.0	0.0
Logan	40	*	*	*	*	*	*	*
Lyon	960	57.6	4.8	34.8	0.3	1.6	0.9	0.0
Marion	414	87.4	1.7	9.2	0.7	0.5	0.5	0.0
Marshall	154	90.9	0.0	7.8	0.6	0.6	0.0	0.0
Mcperson	163	92.0	0.0	2.5	3.1	0.0	2.5	0.0
Meade	131	67.2	0.8	31.3	0.0	0.0	0.8	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3B
2008 - 2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Racial and Ethnic Distribution by County
Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾								
County	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Miami	433	91.0	2.8	3.5	0.2	0.9	1.6	0.0
Mitchell	133	92.5	0.0	7.5	0.0	0.0	0.0	0.0
Montgomery	1,019	77.4	8.9	9.3	2.5	0.9	1.0	0.0
Morris	94	*	*	*	*	*	*	*
Nemaha	118	89.0	0.0	8.5	0.0	0.8	1.7	0.0
Neosho	465	91.6	2.4	4.1	0.6	0.2	1.1	0.0
Ness	35	*	*	*	*	*	*	*
Norton	96	*	*	*	*	*	*	*
Osage	269	96.3	0.0	1.9	0.7	0.0	1.1	0.0
Osborne	59	*	*	*	*	*	*	*
Ottawa	79	*	*	*	*	*	*	*
Pawnee	163	78.5	2.5	17.2	0.6	0.0	1.2	0.0
Phillips	110	91.8	0.0	5.5	0.0	1.8	0.9	0.0
Pottawatom	114	86.0	0.0	12.3	0.9	0.0	0.9	0.0
Pratt	235	86.0	0.0	13.6	0.0	0.4	0.0	0.0
Rawlins	29	*	*	*	*	*	*	*
Reno	1,446	79.7	2.2	15.4	0.6	0.7	1.5	0.0
Republic	103	91.3	1.0	5.8	0.0	0.0	1.9	0.0
Rice	234	73.9	2.1	22.2	0.9	0.9	0.0	0.0
Riley	2,789	71.8	11.4	12.1	0.5	3.8	0.5	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3B
2008 - 2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Racial and Ethnic Distribution by County
Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾								
County	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Rooks	136	93.4	1.5	2.2	2.2	0.7	0.0	0.0
Rush	31	*	*	*	*	*	*	*
Russell	126	90.5	5.6	2.4	1.6	0.0	0.0	0.0
Saline	1,570	71.5	4.2	20.0	0.2	2.2	1.9	0.0
Scott	123	61.8	0.8	37.4	0.0	0.0	0.0	0.0
Sedgwick	13,421	45.2	18.2	29.6	1.5	3.7	1.8	0.0
Seward	1,456	15.4	1.6	80.6	1.5	0.7	0.2	0.0
Shawnee	4,365	54.9	15.4	24.9	2.2	0.8	1.7	0.0
Sheridan	25	*	*	*	*	*	*	*
Sherman	202	73.8	0.0	24.3	0.0	1.0	1.0	0.0
Smith	52	*	*	*	*	*	*	*
Stafford	116	68.1	0.9	30.2	0.0	0.9	0.0	0.0
Stanton	53	*	*	*	*	*	*	*
Stevens	225	45.3	0.0	52.9	1.3	0.4	0.0	0.0
Sumner	479	88.9	0.8	6.7	1.5	0.4	1.7	0.0
Thomas	160	87.5	1.9	8.1	0.6	0.6	1.3	0.0
Trego	46	*	*	*	*	*	*	*
Wabaunsee	60	*	*	*	*	*	*	*
Wallace	1	*	*	*	*	*	*	*
Washington	74	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3B
2008 - 2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Racial and Ethnic Distribution by County
Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾								
County	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Wichita	47	*	*	*	*	*	*	*
Wilson	257	96.1	0.8	1.9	0.4	0.0	0.8	0.0
Woodson	28	*	*	*	*	*	*	*
Wyandotte	6,498	26.7	30.1	39.4	0.3	2.8	0.6	0.0
Unknown Counties	23	*	*	*	*	*	*	*
Kansas	65,016	56.9	11.3	27.2	1.2	2.1	1.3	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4B
2008 - 2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Age Distribution by County
Combined 3 Years

Comparison of Age Distribution ⁽²⁾								
County	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Allen	287	0.0	8.4	16.7	62.0	11.5	1.0	0.3
Anderson	156	0.6	5.1	15.4	62.8	16.0	0.0	0.0
Atchison	413	0.0	7.3	15.5	62.2	14.3	0.7	0.0
Barber	141	0.0	5.7	19.9	62.4	11.3	0.7	0.0
Barton	791	0.0	7.5	15.4	59.9	15.7	1.5	0.0
Bourbon	379	0.3	6.6	18.2	61.5	12.9	0.5	0.0
Brown	300	0.3	7.3	13.7	63.7	14.0	1.0	0.0
Butler	853	0.1	6.8	16.9	61.9	13.6	0.5	0.2
Chase	40	*	*	*	*	*	*	*
Chautauqua	83	*	*	*	*	*	*	*
Cherokee	614	0.3	9.3	19.9	57.2	12.4	0.7	0.3
Cheyenne	13	*	*	*	*	*	*	*
Clark	10	*	*	*	*	*	*	*
Clay	177	1.1	8.5	16.4	59.9	13.6	0.6	0.0
Cloud	254	0.4	7.5	16.5	59.8	13.4	2.4	0.0
Coffey	82	*	*	*	*	*	*	*
Comanche	28	*	*	*	*	*	*	*
Cowley	1,082	0.6	7.9	16.4	61.6	13.0	0.5	0.0
Crawford	1,122	0.2	6.8	14.8	65.1	12.7	0.4	0.1

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4B
2008 - 2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Age Distribution by County
Combined 3 Years

Comparison of Age Distribution ⁽²⁾								
County	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Decatur	33	*	*	*	*	*	*	*
Dickinson	394	0.3	5.1	14.0	64.5	16.0	0.3	0.0
Doniphan	133	0.8	4.5	13.5	63.2	18.0	0.0	0.0
Douglas	1,500	0.0	4.1	12.5	63.7	18.4	1.4	0.0
Edwards	77	*	*	*	*	*	*	*
Elk	57	*	*	*	*	*	*	*
Ellis	481	0.4	4.6	13.1	67.6	12.7	1.5	0.2
Ellsworth	98	*	*	*	*	*	*	*
Finney	2,109	0.4	8.7	15.3	55.1	18.6	1.8	0.1
Ford	2,029	0.3	7.0	13.0	54.2	23.5	1.9	0.0
Franklin	538	0.4	5.9	18.0	60.0	14.7	0.7	0.2
Geary	1,154	0.3	4.8	14.9	67.9	11.4	0.6	0.1
Gove	29	*	*	*	*	*	*	*
Graham	51	*	*	*	*	*	*	*
Grant	205	0.0	6.3	13.2	63.4	14.1	2.9	0.0
Gray	148	0.0	5.4	10.1	60.1	23.0	1.4	0.0
Greeley	37	*	*	*	*	*	*	*
Greenwood	140	0.0	6.4	17.1	62.9	13.6	0.0	0.0
Hamilton	76	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4B
2008 - 2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Age Distribution by County
Combined 3 Years

Comparison of Age Distribution ⁽²⁾								
County	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Harper	156	0.0	6.4	10.3	73.1	10.3	0.0	0.0
Harvey	818	0.2	5.9	13.4	63.4	15.8	1.2	0.0
Haskell	61	*	*	*	*	*	*	*
Hodgeman	39	*	*	*	*	*	*	*
Jackson	233	0.0	5.2	18.9	63.5	12.0	0.4	0.0
Jefferson	191	0.0	5.2	18.8	58.6	16.8	0.5	0.0
Jewell	30	*	*	*	*	*	*	*
Johnson	5,750	0.2	3.8	10.9	60.4	23.1	1.6	0.1
Kearny	61	*	*	*	*	*	*	*
Kingman	165	0.0	7.3	15.8	63.6	12.1	1.2	0.0
Kiowa	33	*	*	*	*	*	*	*
Labette	536	0.0	7.5	20.1	61.2	10.4	0.7	0.0
Lane	37	*	*	*	*	*	*	*
Leavenworth	1,079	0.1	6.0	15.6	64.7	12.5	1.1	0.0
Lincoln	63	*	*	*	*	*	*	*
Linn	191	0.0	6.3	15.7	61.8	15.7	0.5	0.0
Logan	40	*	*	*	*	*	*	*
Lyon	960	0.6	6.3	14.2	62.3	14.5	2.2	0.0
Marion	414	0.5	3.9	12.8	64.5	17.4	1.0	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4B
2008 - 2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Age Distribution by County
Combined 3 Years

Comparison of Age Distribution ⁽²⁾								
County	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Marshall	154	0.0	3.2	13.6	64.9	14.3	3.9	0.0
Mcperson	163	0.0	6.1	12.9	68.1	12.3	0.6	0.0
Meade	131	0.0	9.9	9.2	58.0	21.4	1.5	0.0
Miami	433	0.5	5.3	15.7	64.0	13.9	0.7	0.0
Mitchell	133	0.0	1.5	18.0	65.4	15.0	0.0	0.0
Montgomery	1,019	0.3	5.4	17.5	63.8	11.9	1.0	0.2
Morris	94	*	*	*	*	*	*	*
Nemaha	118	0.0	3.4	12.7	65.3	13.6	4.2	0.8
Neosho	465	0.2	6.2	18.3	61.9	12.5	0.9	0.0
Ness	35	*	*	*	*	*	*	*
Norton	96	*	*	*	*	*	*	*
Osage	269	0.0	5.2	20.4	62.1	10.8	1.1	0.4
Osborne	59	*	*	*	*	*	*	*
Ottawa	79	*	*	*	*	*	*	*
Pawnee	163	1.2	6.7	12.3	68.1	11.0	0.6	0.0
Phillips	110	0.0	3.6	17.3	62.7	15.5	0.9	0.0
Pottawatom	114	0.0	5.3	7.0	58.8	25.4	3.5	0.0
Pratt	235	0.4	8.1	14.5	61.3	14.9	0.9	0.0
Rawlins	29	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4B
2008 - 2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Age Distribution by County
Combined 3 Years

Comparison of Age Distribution ⁽²⁾								
County	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Reno	1,446	0.1	6.8	15.6	64.0	12.9	0.6	0.0
Republic	103	0.0	3.9	20.4	56.3	16.5	2.9	0.0
Rice	234	0.4	5.6	13.2	63.7	16.7	0.4	0.0
Riley	2,789	0.0	1.8	12.1	74.0	11.7	0.4	0.0
Rooks	136	0.0	2.2	15.4	62.5	17.6	1.5	0.7
Rush	31	*	*	*	*	*	*	*
Russell	126	0.0	10.3	16.7	62.7	10.3	0.0	0.0
Saline	1,570	0.1	6.7	15.9	62.4	14.0	1.0	0.1
Scott	123	0.8	3.3	13.0	68.3	13.8	0.8	0.0
Sedgwick	13,421	0.3	6.8	13.9	60.7	17.1	1.3	0.0
Seward	1,456	0.2	7.3	13.6	56.6	20.9	1.4	0.0
Shawnee	4,365	0.3	6.6	14.6	60.7	16.7	1.2	0.0
Sheridan	25	*	*	*	*	*	*	*
Sherman	202	0.0	5.4	15.8	59.4	17.3	2.0	0.0
Smith	52	*	*	*	*	*	*	*
Stafford	116	0.0	6.0	13.8	60.3	17.2	2.6	0.0
Stanton	53	*	*	*	*	*	*	*
Stevens	225	0.0	7.1	18.2	57.3	16.4	0.9	0.0
Sumner	479	0.0	6.7	16.7	61.2	14.6	0.8	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4B
2008 - 2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Age Distribution by County
Combined 3 Years

Comparison of Age Distribution ⁽²⁾								
County	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Thomas	160	0.0	6.3	15.0	68.1	10.6	0.0	0.0
Trego	46	*	*	*	*	*	*	*
Wabaunsee	60	*	*	*	*	*	*	*
Wallace	1	*	*	*	*	*	*	*
Washington	74	*	*	*	*	*	*	*
Wichita	47	*	*	*	*	*	*	*
Wilson	257	0.4	8.2	15.6	62.3	13.6	0.0	0.0
Woodson	28	*	*	*	*	*	*	*
Wyandotte	6,498	0.5	8.4	13.5	57.2	19.0	1.4	0.0
Unknown Counties	23	*	*	*	*	*	*	*
Kansas	65,016	0.3	6.2	14.2	61.3	16.7	1.2	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5B
2008 - 2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Health Indicators by County
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)														
County	Prepregnancy BMI ⁽⁴⁾			Weight Gain ⁽⁵⁾			Anemia (Low Hb/Hct) ⁽⁶⁾				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Allen	283	4.9 (40)	56.9 (48)	260	18.1 (15)	55.4 (55)	48	* *	82	* *	254	7.5 (53)	254	15.7(60)
Anderson	153	6.5 (58)	51.0 (8)	141	23.4 (51)	46.8 (11)	22	* *	25	* *	136	2.9 (5)	136	14.7(57)
Atchison	401	5.2 (46)	53.9 (25)	357	16.2 (6)	57.4 (58)	61	* *	83	* *	329	5.5 (27)	329	12.8(51)
Barber	139	2.9 (11)	51.8 (13)	113	30.1 (61)	48.7 (19)	15	* *	34	* *	112	0.9 (2)	112	16.1(61)
Barton	770	4.9 (40)	55.3 (35)	685	20.9 (36)	49.6 (24)	78	* *	161	26.1(21)	658	5.5 (27)	656	11.6(43)
Bourbon	373	2.1 (6)	56.6 (45)	329	23.7 (52)	48.9 (21)	34	* *	72	* *	317	6.3 (43)	318	11.9(47)
Brown	296	1.7 (5)	57.4 (50)	259	23.2 (50)	51.0 (34)	38	* *	66	* *	257	5.1 (21)	257	13.6(54)
Butler	826	5.9 (52)	51.0 (8)	755	19.7 (26)	53.4 (43)	114	43.9 (20)	188	29.3(25)	723	5.4 (25)	723	11.3(42)
Chase	38	* *	* *	31	* *	* *	4	* *	2	* *	34	* *	34	* *
Chautauqua	82	* *	* *	74	* *	* *	8	* *	13	* *	73	* *	73	* *
Cherokee	580	6.6 (60)	51.9 (14)	517	18.0 (13)	54.0 (45)	72	* *	119	25.2(18)	515	5.6 (29)	515	10.3(34)
Cheyenne	12	* *	* *	9	* *	* *	3	* *	2	* *	10	* *	10	* *
Clark	9	* *	* *	8	* *	* *	2	* *	2	* *	9	* *	9	* *
Clay	172	4.7 (39)	55.8 (42)	151	19.2 (20)	60.9 (61)	25	* *	43	* *	150	8.0 (58)	150	10.7(38)
Cloud	251	4.4 (32)	53.4 (20)	226	20.8 (34)	54.0 (45)	27	* *	66	* *	216	6.9 (49)	221	10.0(30)
Coffey	78	* *	* *	66	* *	* *	8	* *	21	* *	61	* *	61	* *
Comanche	28	* *	* *	25	* *	* *	6	* *	7	* *	23	* *	23	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5B
2008 - 2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Health Indicators by County
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)														
County	Prepregnancy BMI ⁽⁴⁾			Weight Gain ⁽⁵⁾			Anemia (Low Hb/Hct) ⁽⁶⁾				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
							Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)
Cowley	1,066	4.5 (33)	54.7 (33)	952	20.7 (33)	52.4 (40)	132	32.6 (16)	160	22.5(12)	917	5.9 (36)	919	9.9(28)
Crawford	1,102	6.3 (57)	54.3 (29)	954	20.6 (29)	52.3 (38)	166	19.3 (2)	169	18.9 (6)	939	7.2 (52)	938	10.8(40)
Decatur	33	* *	* *	32	* *	* *	5	* *	8	* *	30	* *	30	* *
Dickinson	381	2.9 (11)	57.2 (49)	334	21.9 (40)	50.0 (26)	49	* *	84	* *	332	6.3 (43)	328	11.6(43)
Doniphan	130	0.8 (1)	62.3 (63)	110	13.6 (2)	65.5 (62)	23	* *	25	* *	107	5.6 (29)	107	12.1(48)
Douglas	1,463	3.8 (24)	48.4 (4)	1,253	17.6 (11)	54.7 (49)	199	26.6 (7)	384	26.8(22)	1,218	6.0 (37)	1,217	7.8(17)
Edwards	77	* *	* *	67	* *	* *	13	* *	14	* *	61	* *	61	* *
Elk	56	* *	* *	50	* *	* *	4	* *	11	* *	47	* *	47	* *
Ellis	462	3.9 (26)	51.1 (10)	400	18.3 (16)	49.0 (22)	103	23.3 (4)	111	17.1 (3)	397	2.5 (4)	397	6.3(9)
Ellsworth	96	* *	* *	91	* *	* *	14	* *	22	* *	84	* *	84	* *
Finney	2,072	6.5 (58)	50.9 (7)	1,882	24.5 (55)	45.2 (4)	290	29.7 (12)	297	21.9(11)	1,849	7.6 (56)	1,847	6.2(7)
Ford	1,999	2.2 (7)	57.5 (52)	1,777	23.7 (52)	46.7 (10)	254	21.3 (3)	230	13.5 (1)	1,729	5.3 (24)	1,727	7.6(15)
Franklin	516	3.7 (22)	51.6 (11)	453	16.8 (8)	53.0 (41)	52	* *	136	14.0 (2)	386	4.1 (10)	386	9.8(27)
Geary	1,119	4.6 (37)	53.4 (20)	900	18.0 (13)	53.7 (44)	171	39.2 (18)	197	34.0(26)	902	4.1 (10)	901	10.7(38)
Gove	24	* *	* *	21	* *	* *	3	* *	9	* *	26	* *	26	* *
Graham	51	* *	* *	41	* *	* *	7	* *	6	* *	42	* *	42	* *
Grant	198	2.5 (9)	67.2 (66)	183	23.0 (48)	48.1 (16)	24	* *	59	* *	181	6.1 (38)	179	5.0(2)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5B
2008 - 2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Health Indicators by County
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)														
County	Prepregnancy BMI ⁽⁴⁾			Weight Gain ⁽⁵⁾			Anemia (Low Hb/Hct) ⁽⁶⁾				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Gray	144	3.5 (18)	54.2 (28)	121	23.1 (49)	52.1 (37)	15	* *	39	* *	118	7.6 (56)	118	11.0(41)
Greeley	35	* *	* *	31	* *	* *	3	* *	5	* *	33	* *	33	* *
Greenwood	135	2.2 (7)	60.0 (58)	123	22.0 (41)	55.3 (53)	13	* *	21	* *	120	5.0 (19)	121	18.2(62)
Hamilton	73	* *	* *	65	* *	* *	15	* *	22	* *	66	* *	66	* *
Harper	150	6.0 (54)	52.7 (17)	130	18.5 (17)	48.5 (18)	16	* *	21	* *	125	5.6 (29)	124	8.1(18)
Harvey	803	4.5 (33)	54.5 (31)	713	22.6 (45)	48.0 (15)	108	36.1 (17)	122	21.3 (9)	689	4.6 (16)	686	10.3(34)
Haskell	61	* *	* *	55	* *	* *	8	* *	16	* *	52	* *	52	* *
Hodgeman	37	* *	* *	35	* *	* *	6	* *	7	* *	33	* *	33	* *
Jackson	224	3.6 (20)	55.4 (37)	206	19.9 (27)	56.8 (57)	32	* *	48	* *	203	8.4 (60)	204	14.7(57)
Jefferson	188	3.7 (22)	58.0 (53)	169	16.0 (4)	58.0 (59)	31	* *	51	* *	165	6.7 (46)	165	9.1(23)
Jewell	30	* *	* *	30	* *	* *	3	* *	5	* *	29	* *	29	* *
Johnson	5,547	4.2 (29)	54.0 (27)	4,944	18.8 (19)	50.9 (33)	902	24.2 (5)	1,454	17.4 (4)	4,815	6.8 (48)	4,806	6.2(7)
Kearny	61	* *	* *	57	* *	* *	8	* *	12	* *	57	* *	57	* *
Kingman	165	3.6 (20)	46.1 (1)	144	22.2 (42)	50.0 (26)	16	* *	27	* *	135	6.7 (46)	135	14.1(55)
Kiowa	31	* *	* *	26	* *	* *	5	* *	7	* *	29	* *	29	* *
Labette	527	5.9 (52)	56.2 (44)	482	24.7 (57)	50.6 (30)	92	* *	118	24.6(15)	465	5.2 (23)	465	12.3(49)
Lane	34	* *	* *	30	* *	* *	6	* *	6	* *	30	* *	30	* *

- (1) Reporting period is January 1 through December 31.
 - (2) Excludes records with unknown data or errors.
 - (3) Rank compares one county's rate to other counties. Rank 1 = best rate.
 - (4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
 - (5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
 - (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5B
2008 - 2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Health Indicators by County
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)														
County	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight	Over- weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
							Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)
Leavenworth	1,046	4.2 (29)	55.4 (37)	911	19.5 (23)	55.3 (53)	167	26.9 (8)	266	21.8(10)	860	5.6 (29)	859	9.7(26)
Lincoln	60	* *	* *	56	* *	* *	11	* *	15	* *	52	* *	52	* *
Linn	189	5.8 (51)	62.4 (64)	173	28.9 (60)	46.2 (8)	24	* *	29	* *	169	4.7 (17)	170	5.3(3)
Logan	39	* *	* *	32	* *	* *	6	* *	6	* *	31	* *	31	* *
Lyon	935	3.3 (17)	59.5 (56)	828	21.6 (39)	48.7 (19)	127	11.0 (1)	172	20.3 (8)	800	5.4 (25)	800	12.5(50)
Marion	408	5.4 (47)	54.7 (33)	368	22.3 (44)	45.4 (6)	62	* *	92	* *	357	5.0 (19)	359	8.4(20)
Marshall	149	7.4 (64)	48.3 (3)	124	18.5 (17)	48.4 (17)	19	* *	31	* *	123	2.4 (3)	124	8.9(22)
Mcperson	159	5.7 (50)	53.5 (22)	144	20.8 (34)	50.0 (26)	21	* *	35	* *	137	5.1 (21)	136	9.6(25)
Meade	130	3.1 (14)	61.5 (62)	115	22.6 (45)	45.2 (4)	12	* *	24	* *	109	8.3 (59)	109	9.2(24)
Miami	419	6.0 (54)	55.6 (39)	375	16.0 (4)	55.2 (51)	84	* *	120	25.8(19)	367	7.1 (51)	367	10.4(36)
Mitchell	130	3.1 (14)	58.5 (54)	122	32.0 (62)	45.9 (7)	14	* *	34	* *	120	9.2 (61)	119	14.3(56)
Montgomery	998	6.1 (56)	53.6 (24)	883	23.7 (52)	47.2 (12)	184	32.1 (15)	251	25.1(17)	852	5.6 (29)	852	10.1(32)
Morris	92	* *	* *	82	* *	* *	13	* *	19	* *	81	* *	81	* *
Nemaha	106	0.9 (2)	55.7 (40)	99	* *	* *	27	* *	32	* *	105	2.9 (5)	105	5.7(5)
Neosho	451	7.1 (62)	52.3 (15)	413	20.6 (29)	49.9 (25)	71	* *	71	* *	398	5.8 (35)	397	13.4(53)
Ness	33	* *	* *	28	* *	* *	3	* *	11	* *	26	* *	26	* *
Norton	96	* *	* *	86	* *	* *	14	* *	27	* *	81	* *	81	* *

- (1) Reporting period is January 1 through December 31.
 - (2) Excludes records with unknown data or errors.
 - (3) Rank compares one county's rate to other counties. Rank 1 = best rate.
 - (4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
 - (5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
 - (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5B
2008 - 2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Health Indicators by County
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)														
County	Prepregnancy BMI ⁽⁴⁾			Weight Gain ⁽⁵⁾			Anemia (Low Hb/Hct) ⁽⁶⁾				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Osage	264	4.9 (40)	52.3 (15)	230	16.5 (7)	55.2 (51)	33	* *	56	* *	228	4.8 (18)	228	11.8(46)
Osborne	58	* *	* *	50	* *	* *	5	* *	13	* *	50	* *	50	* *
Ottawa	75	* *	* *	67	* *	* *	11	* *	13	* *	70	* *	70	* *
Pawnee	159	5.0 (45)	56.6 (45)	132	28.0 (59)	46.2 (8)	13	* *	31	* *	134	7.5 (53)	133	7.5(14)
Phillips	106	3.8 (24)	60.4 (60)	89	* *	* *	8	* *	20	* *	90	* *	90	* *
Pottawatom	112	5.4 (47)	54.5 (31)	102	20.6 (29)	45.1 (3)	12	* *	46	* *	99	* *	99	* *
Pratt	230	3.5 (18)	53.5 (22)	196	20.9 (36)	49.0 (22)	29	* *	52	* *	189	0.5 (1)	189	7.4(11)
Rawlins	28	* *	* *	25	* *	* *	3	* *	9	* *	25	* *	25	* *
Reno	1,423	4.6 (37)	53.9 (25)	1,278	20.6 (29)	50.7 (31)	182	27.5 (10)	242	23.1(13)	1,210	3.4 (8)	1,211	13.1(52)
Republic	101	6.9 (61)	57.4 (50)	87	* *	* *	15	* *	20	* *	86	* *	86	* *
Rice	228	3.9 (26)	60.1 (59)	198	19.2 (20)	50.5 (29)	28	* *	49	* *	199	6.5 (45)	198	7.6(15)
Riley	2,674	4.5 (33)	50.8 (6)	2,177	17.5 (10)	54.2 (47)	418	28.5 (11)	632	27.4(23)	2,122	4.3 (13)	2,122	10.0(30)
Rooks	136	8.1 (66)	46.3 (2)	122	24.6 (56)	39.3 (1)	20	* *	21	* *	115	6.1 (38)	115	10.4(36)
Rush	31	* *	* *	30	* *	* *	4	* *	8	* *	29	* *	29	* *
Russell	125	3.2 (16)	50.4 (5)	113	25.7 (58)	47.8 (14)	18	* *	18	* *	108	3.7 (9)	108	4.6(1)
Saline	1,543	4.9 (40)	55.3 (35)	1,397	17.0 (9)	54.3 (48)	294	24.8 (6)	308	20.1 (7)	1,366	6.2 (42)	1,369	7.4(11)
Scott	122	4.1 (28)	59.0 (55)	107	19.6 (24)	55.1 (50)	15	* *	19	* *	103	2.9 (5)	103	5.8(6)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5B
2008 - 2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Health Indicators by County
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)														
County	Prepregnancy BMI ⁽⁴⁾			Weight Gain ⁽⁵⁾			Anemia (Low Hb/Hct) ⁽⁶⁾				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Sedgwick	13,120	4.5 (33)	53.3 (19)	11,998	20.2 (28)	51.5 (35)	2,570	31.5 (14)	3,130	24.7 (16)	11,695	5.6 (29)	11,690	9.9 (28)
Seward	1,432	2.5 (9)	56.7 (47)	1,290	22.2 (42)	47.7 (13)	238	27.3 (9)	214	24.3 (14)	1,251	7.5 (53)	1,251	6.5 (10)
Shawnee	4,260	4.9 (40)	54.3 (29)	3,732	19.4 (22)	52.3 (38)	426	42.7 (19)	708	29.1 (24)	3,648	6.9 (49)	3,653	7.4 (11)
Sheridan	25	* *	* *	21	* *	* *	5	* *	4	* *	22	* *	22	* *
Sherman	198	3.0 (13)	59.6 (57)	171	21.1 (38)	56.7 (56)	18	* *	36	* *	165	4.2 (12)	165	8.5 (21)
Smith	52	* *	* *	43	* *	* *	9	* *	13	* *	43	* *	43	* *
Stafford	114	0.9 (2)	65.8 (65)	99	* *	* *	11	* *	16	* *	93	* *	94	* *
Stanton	52	* *	* *	49	* *	* *	5	* *	21	* *	50	* *	49	* *
Stevens	221	0.9 (2)	61.1 (61)	191	15.7 (3)	50.8 (32)	37	* *	60	* *	183	14.2 (62)	183	5.5 (4)
Sumner	466	5.6 (49)	51.7 (12)	426	22.8 (47)	43.9 (2)	68	* *	123	17.9 (5)	407	6.1 (38)	408	15.0 (59)
Thomas	155	7.1 (62)	52.9 (18)	139	11.5 (1)	59.7 (60)	21	* *	29	* *	138	4.3 (13)	138	11.6 (43)
Trego	44	* *	* *	37	* *	* *	5	* *	5	* *	38	* *	38	* *
Wabaunsee	58	* *	* *	57	* *	* *	7	* *	24	* *	56	* *	56	* *
Wallace	1	* *	* *	1	* *	* *	0	* *	1	* *	0	* *	0	* *
Washington	71	* *	* *	64	* *	* *	5	* *	21	* *	63	* *	62	* *
Wichita	45	* *	* *	41	* *	* *	6	* *	6	* *	42	* *	42	* *
Wilson	254	7.9 (65)	55.9 (43)	239	17.6 (11)	53.1 (42)	43	* *	38	* *	228	6.1 (38)	228	10.1 (32)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5B
2008 - 2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Health Indicators by County
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)														
County	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Woodson	28	* *	* *	26	* *	* *	7	* *	12	* *	26	* *	26	* *
Wyandotte	6,341	4.3 (31)	55.7 (40)	5,681	19.6 (24)	51.9 (36)	981	30.9 (13)	1,141	26.0 (20)	5,503	4.5 (15)	5,503	8.3 (19)
Unknown Counties	21	* *	* *	20	* *	* *	1	* *	3	* *	22	* *	22	* *
Kansas	63,394	4.5	54.2	56,431	20.2	51.1	9,783	29.3	13,286	23.5	54,852	5.7	54,836	9.2

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6B
2008 - 2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Behavioral Indicators by County
Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2, 3)										
County	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Pregpregnancy (4)		During Pregpregnancy (5)	
	Number	% (Rank)	%	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Allen	246	92.7 (1)	1.6 (15)		271	38.7 (40)	244	14.3 (40)	214	52.3 (47)
Anderson	125	87.2 (9)	4.8 (37)		146	48.6 (10)	144	20.1 (25)	124	63.7 (22)
Atchison	359	88.6 (8)	2.5 (19)		389	37.3 (47)	376	8.8 (57)	318	58.5 (38)
Barber	117	81.2 (32)	8.5 (53)		133	45.1 (20)	126	34.1 (5)	105	67.6 (12)
Barton	693	84.4 (23)	1.2 (11)		728	51.9 (6)	669	20.3 (22)	548	56.6 (41)
Bourbon	311	90.0 (5)	1.6 (15)		357	45.1 (20)	321	10.9 (49)	276	59.1 (37)
Brown	236	75.8 (43)	11.0 (55)		284	39.4 (38)	254	12.6 (44)	244	64.8 (20)
Butler	750	70.1 (55)	9.9 (54)		760	38.9 (39)	721	14.7 (37)	548	35.6 (56)
Chase	33	* *	* *		35	* *	34	* *	29	* *
Chautauqua	75	* *	* *		78	* *	67	* *	63	* *
Cherokee	437	76.4 (40)	8.2 (52)		560	37.3 (47)	510	12.5 (46)	407	53.6 (44)
Cheyenne	7	* *	* *		13	* *	13	* *	12	* *
Clark	8	* *	* *		8	* *	6	* *	6	* *
Clay	158	84.8 (20)	5.7 (40)		165	43.0 (26)	145	20.0 (26)	134	67.9 (11)
Cloud	230	83.5 (28)	4.8 (37)		238	48.3 (11)	226	20.8 (21)	195	68.2 (9)
Coffey	69	* *	* *		72	* *	72	* *	56	* *
Comanche	25	* *	* *		25	* *	25	* *	20	* *
Cowley	894	69.2 (56)	6.3 (43)		987	44.2 (23)	920	12.6 (44)	782	56.0 (43)
Crawford	930	86.0 (14)	3.0 (21)		1,052	42.2 (28)	975	12.3 (47)	844	60.1 (36)
Decatur	24	* *	* *		30	* *	29	* *	21	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6B
2008 - 2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Behavioral Indicators by County
Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2, 3)										
County	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Pregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	%	(Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Dickinson	349	86.8 (12)	1.4	(14)	370	44.3 (22)	357	20.2 (23)	282	68.4 (7)
Doniphan	116	77.6 (39)	7.8	(49)	123	39.8 (34)	117	35.0 (4)	92	* *
Douglas	1,274	74.0 (47)	5.7	(40)	1,397	37.6 (46)	1,295	19.3 (28)	1,054	60.4 (34)
Edwards	59	* *	* *		68	* *	69	* *	57	* *
Elk	48	* *	* *		53	* *	49	* *	41	* *
Ellis	423	85.6 (16)	1.7	(17)	449	34.5 (54)	433	25.9 (12)	360	68.3 (8)
Ellsworth	93	* *	* *		86	* *	86	* *	67	* *
Finney	1,762	59.0 (61)	24.7	(61)	1,957	35.4 (52)	1,834	11.1 (48)	1,539	39.0 (54)
Ford	1,666	71.6 (54)	14.4	(58)	1,915	45.3 (19)	1,730	9.7 (52)	1,538	51.8 (49)
Franklin	203	76.4 (40)	3.0	(21)	498	38.2 (43)	463	5.4 (61)	370	62.4 (25)
Geary	876	72.1 (50)	1.3	(12)	1,097	35.3 (53)	1,009	19.5 (27)	846	56.9 (40)
Gove	25	* *	* *		27	* *	25	* *	18	* *
Graham	42	* *	* *		49	* *	41	* *	36	* *
Grant	172	68.6 (57)	16.3	(60)	186	33.3 (60)	173	9.2 (54)	135	45.9 (52)
Gray	122	84.4 (23)	3.3	(26)	133	46.6 (16)	123	52.8 (2)	107	68.2 (9)
Greeley	32	* *	* *		34	* *	34	* *	27	* *
Greenwood	119	56.3 (62)	36.1	(62)	132	52.3 (4)	116	14.7 (37)	104	70.2 (4)
Hamilton	66	* *	* *		72	* *	68	* *	60	* *
Harper	129	73.6 (49)	14.7	(59)	144	52.1 (5)	133	26.3 (11)	117	70.1 (5)
Harvey	657	74.0 (47)	12.9	(57)	764	47.8 (12)	705	16.6 (32)	601	66.2 (16)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6B
2008 - 2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Behavioral Indicators by County
Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2, 3)										
County	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Haskell	51	* *	* *	* *	57	* *	53	* *	47	* *
Hodgeman	36	* *	* *	* *	35	* *	34	* *	27	* *
Jackson	118	82.2 (31)	4.2 (31)		225	44.0 (24)	202	10.4 (50)	178	77.0 (1)
Jefferson	160	85.6 (16)	6.9 (47)		174	39.7 (36)	162	6.2 (60)	136	50.7 (50)
Jewell	30	* *	* *	* *	28	* *	24	* *	25	* *
Johnson	4,561	64.9 (60)	3.7 (27)		5,294	32.8 (62)	4,928	13.3 (43)	4,016	62.0 (28)
Kearny	53	* *	* *	* *	57	* *	50	* *	40	* *
Kingman	141	78.0 (38)	4.3 (32)		154	51.9 (6)	147	22.4 (17)	128	67.2 (14)
Kiowa	29	* *	* *	* *	33	* *	30	* *	26	* *
Labette	407	86.5 (13)	3.2 (25)		500	36.6 (51)	462	13.9 (42)	401	61.1 (33)
Lane	31	* *	* *	* *	34	* *	31	* *	29	* *
Leavenworth	931	85.5 (18)	0.6 (4)		965	32.5 (63)	925	20.2 (23)	699	68.5 (6)
Lincoln	58	* *	* *	* *	56	* *	55	* *	45	* *
Linn	175	80.0 (37)	7.4 (48)		180	47.8 (12)	167	21.0 (20)	148	61.5 (30)
Logan	15	* *	* *	* *	37	* *	35	* *	30	* *
Lyon	844	83.9 (25)	0.8 (5)		881	40.4 (33)	810	15.9 (34)	671	62.4 (25)
Marion	373	87.1 (11)	0.5 (3)		383	38.1 (45)	343	23.0 (16)	294	67.3 (13)
Marshall	128	71.9 (52)	3.9 (28)		141	41.8 (29)	135	32.6 (6)	100	65.0 (18)
Mcperson	148	92.6 (3)	0.0 (1)		153	41.8 (29)	133	30.8 (7)	108	65.7 (17)
Meade	114	71.9 (52)	7.9 (51)		116	57.8 (3)	108	37.0 (3)	97	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6B
2008 - 2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Behavioral Indicators by County
Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2, 3)										
County	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	% (Rank)		Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Miami	389	81.2 (32)	1.0 (8)		391	34.3 (56)	371	16.7 (31)	296	53.4 (45)
Mitchell	123	92.7 (1)	0.0 (1)		123	34.1 (58)	114	24.6 (14)	97	* *
Montgomery	712	85.4 (19)	4.4 (33)		944	33.3 (60)	875	14.3 (40)	739	63.6 (23)
Morris	76	* *	* *		87	* *	76	* *	64	* *
Nemaha	109	87.2 (9)	5.5 (39)		110	38.2 (43)	97	* *	86	* *
Neosho	386	88.9 (7)	4.4 (33)		433	38.6 (41)	404	9.2 (54)	343	67.1 (15)
Ness	26	* *	* *		30	* *	32	* *	24	* *
Norton	87	* *	* *		85	* *	84	* *	69	* *
Osage	208	91.3 (4)	1.0 (8)		240	50.0 (9)	231	19.0 (29)	196	38.8 (55)
Osborne	49	* *	* *		54	* *	46	* *	40	* *
Ottawa	72	* *	* *		73	* *	68	* *	59	* *
Pawnee	136	84.6 (22)	2.2 (18)		154	50.6 (8)	148	27.7 (9)	124	61.3 (32)
Phillips	97	* *	* *		102	58.8 (2)	97	* *	89	* *
Pottawatom	92	* *	* *		107	38.3 (42)	93	* *	83	* *
Pratt	202	80.7 (35)	6.4 (44)		212	43.4 (25)	208	7.2 (59)	158	61.4 (31)
Rawlins	11	* *	* *		28	* *	25	* *	23	* *
Reno	1,262	83.6 (26)	2.5 (19)		1,320	39.8 (34)	1,243	21.1 (19)	1,043	62.0 (28)
Republic	89	* *	* *		96	* *	83	* *	72	* *
Rice	203	83.3 (29)	1.0 (8)		220	42.3 (27)	206	16.0 (33)	171	44.4 (53)
Riley	2,092	83.6 (26)	0.8 (5)		2,561	34.4 (55)	2,435	18.7 (30)	1,924	62.2 (27)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6B
2008 - 2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Behavioral Indicators by County
Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2, 3)										
County	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	% (Rank)		Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Rooks	120	82.5 (30)	5.8 (42)		129	39.5 (37)	117	30.8 (7)	100	65.0 (18)
Rush	30	* *	* *		29	* *	27	* *	24	* *
Russell	114	89.5 (6)	0.9 (7)		115	47.0 (15)	109	10.1 (51)	94	* *
Saline	1,234	74.9 (45)	4.1 (30)		1,451	34.1 (58)	1,348	5.0 (62)	1,151	52.2 (48)
Scott	104	80.8 (34)	6.7 (45)		112	45.5 (18)	105	79.0 (1)	95	* *
Sedgwick	12,080	74.9 (45)	4.6 (35)		12,454	28.3 (65)	11,578	15.4 (35)	9,583	47.5 (51)
Seward	1,183	65.4 (59)	11.7 (56)		1,349	34.3 (56)	1,227	8.6 (58)	1,021	64.8 (20)
Shawnee	3,460	76.1 (42)	3.9 (28)		4,068	47.2 (14)	3,799	9.2 (54)	3,247	52.9 (46)
Sheridan	22	* *	* *		22	* *	20	* *	17	* *
Sherman	118	84.7 (21)	6.8 (46)		190	46.3 (17)	171	25.7 (13)	146	70.5 (3)
Smith	38	* *	* *		47	* *	44	* *	42	* *
Stafford	100	72.0 (51)	3.0 (21)		109	60.6 (1)	101	23.8 (15)	87	* *
Stanton	50	* *	* *		51	* *	41	* *	31	* *
Stevens	194	75.3 (44)	3.1 (24)		206	32.0 (64)	197	22.3 (18)	161	60.2 (35)
Sumner	435	80.2 (36)	4.6 (35)		440	40.5 (32)	405	15.1 (36)	325	62.5 (24)
Thomas	96	* *	* *		146	37.0 (50)	133	27.1 (10)	113	76.1 (2)
Trego	39	* *	* *		41	* *	38	* *	34	* *
Wabaunsee	52	* *	* *		52	* *	52	* *	41	* *
Wallace	1	* *	* *		0	* *	1	* *	0	* *
Washington	66	* *	* *		68	* *	64	* *	52	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6B
2008 - 2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Behavioral Indicators by County
Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2, 3)										
County	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Pregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	%	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Wichita	43	* *	* *	* *	45	* *	42	* *	33	* *
Wilson	236	86.0 (14)	1.3	(12)	246	41.5 (31)	226	14.6 (39)	198	56.1 (42)
Woodson	28	* *	* *	* *	27	* *	26	* *	21	* *
Wyandotte	5,629	65.5 (58)	7.8	(49)	6,006	37.2 (49)	5,557	9.3 (53)	4,735	57.5 (39)
Unknown Counties	20	* *	* *	* *	23	* *	23	* *	18	* *
Kansas	54,276	75.1	5.7		60,184	37.0	55,958	14.7	46,711	56.0

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7B
2008 - 2010 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by County
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Allen	276	48.2 (52)	228	30.3 (46)	86	* *	242	39.7 (49)	228	38.2 (48)
Anderson	152	50.7 (59)	132	29.5 (45)	56	* *	133	23.3 (33)	134	20.1 (33)
Atchison	301	48.8 (56)	200	31.5 (49)	73	* *	256	27.3 (37)	204	22.5 (38)
Barber	140	48.6 (55)	114	31.6 (50)	46	* *	117	41.0 (52)	115	44.3 (54)
Barton	780	36.8 (22)	687	24.0 (24)	206	26.7 (21)	634	11.4 (13)	688	5.8 (6)
Bourbon	356	46.3 (47)	277	27.8 (41)	97	* *	296	38.9 (48)	281	34.9 (46)
Brown	285	46.0 (45)	215	26.5 (35)	71	* *	244	42.2 (55)	217	39.6 (51)
Butler	818	49.4 (57)	685	30.7 (47)	252	32.1 (15)	618	13.4 (17)	690	8.4 (14)
Chase	39	* *	34	* *	17	* *	33	* *	34	* *
Chautauqua	71	* *	46	* *	16	* *	62	* *	45	* *
Cherokee	585	40.3 (30)	502	27.7 (40)	157	23.6 (26)	454	33.7 (46)	511	34.2 (45)
Cheyenne	13	* *	10	* *	2	* *	12	* *	10	* *
Clark	3	* *	2	* *	0	* *	5	* *	4	* *
Clay	175	47.4 (51)	154	25.3 (27)	64	* *	155	18.1 (26)	156	10.3 (19)
Cloud	252	35.7 (19)	224	19.2 (15)	63	* *	216	7.4 (4)	226	2.2 (1)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7B
2008 - 2010 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by County
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Coffey	77	* *	68	* *	26	* *	60	* *	68	* *
Comanche	27	* *	24	* *	10	* *	21	* *	24	* *
Cowley	1,041	42.4 (36)	819	28.4 (42)	274	26.3 (23)	864	20.9 (31)	822	20.3 (34)
Crawford	1,045	43.7 (39)	814	25.7 (29)	289	36.3 (6)	909	39.8 (50)	817	39.3 (50)
Decatur	33	* *	28	* *	6	* *	23	* *	29	* *
Dickinson	385	44.2 (41)	337	26.7 (37)	117	32.5 (14)	307	9.8 (10)	340	7.1 (10)
Doniphan	117	41.9 (34)	85	* *	24	* *	87	* *	88	* *
Douglas	1,468	40.9 (32)	1,234	19.8 (16)	385	42.1 (3)	1,179	10.2 (11)	1,236	6.4 (9)
Edwards	77	* *	65	* *	19	* *	63	* *	66	* *
Elk	53	* *	43	* *	13	* *	43	* *	42	* *
Ellis	477	36.1 (21)	408	14.0 (9)	120	46.7 (1)	386	7.0 (2)	410	5.6 (5)
Ellsworth	97	* *	90	* *	29	* *	73	* *	91	* *
Finney	2,076	11.5 (5)	1,861	6.0 (4)	162	38.9 (4)	1,730	15.4 (22)	1,861	14.1 (25)
Ford	1,956	9.0 (4)	1,639	4.5 (3)	105	31.4 (16)	1,704	8.9 (6)	1,637	8.4 (14)
Franklin	533	44.1 (40)	468	28.4 (42)	142	27.5 (20)	403	13.4 (17)	470	9.6 (17)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7B
2008 - 2010 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by County
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Geary	1,134	36.9 (23)	905	20.7 (17)	257	33.9 (10)	942	37.4 (47)	908	34.1 (44)
Gove	28	* *	26	* *	2	* *	19	* *	26	* *
Graham	51	* *	41	* *	19	* *	43	* *	42	* *
Grant	202	6.9 (2)	191	2.6 (1)	8	* *	158	9.5 (9)	191	11.0 (20)
Gray	146	8.2 (3)	119	6.7 (5)	6	* *	125	10.4 (12)	120	7.5 (12)
Greeley	37	* *	32	* *	7	* *	27	* *	32	* *
Greenwood	118	56.8 (63)	60	* *	27	* *	110	30.9 (44)	87	* *
Hamilton	75	* *	68	* *	6	* *	63	* *	68	* *
Harper	139	44.6 (43)	99	* *	33	* *	116	21.6 (32)	92	* *
Harvey	802	37.2 (26)	687	21.7 (18)	209	34.9 (7)	667	7.0 (2)	689	4.2 (3)
Haskell	61	* *	56	* *	8	* *	53	* *	56	* *
Hodgeman	37	* *	32	* *	14	* *	29	* *	31	* *
Jackson	231	48.5 (53)	203	31.0 (48)	80	* *	197	20.8 (30)	205	19.5 (32)
Jefferson	48	* *	21	* *	4	* *	48	* *	25	* *
Jewell	30	* *	28	* *	11	* *	27	* *	28	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7B
2008 - 2010 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by County
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Johnson	5,652	23.9 (9)	4,959	10.6 (7)	821	43.0 (2)	4,490	18.4 (27)	4,972	18.1 (31)
Kearny	61	* *	57	* *	9	* *	48	* *	57	* *
Kingman	149	47.0 (50)	122	27.0 (39)	39	* *	136	41.9 (54)	131	38.9 (49)
Kiowa	31	* *	22	* *	5	* *	25	* *	24	* *
Labette	520	36.0 (20)	458	25.1 (26)	129	17.1 (28)	439	30.5 (43)	463	30.7 (42)
Lane	11	* *	2	* *	0	* *	12	* *	3	* *
Leavenworth	1,059	46.8 (49)	892	25.8 (31)	288	34.7 (8)	788	41.5 (53)	895	40.1 (52)
Lincoln	61	* *	50	* *	21	* *	51	* *	50	* *
Linn	188	56.4 (62)	170	37.6 (54)	84	* *	159	15.1 (20)	172	12.8 (23)
Logan	39	* *	33	* *	11	* *	34	* *	33	* *
Lyon	949	29.7 (14)	840	16.7 (14)	183	32.8 (12)	761	11.8 (14)	841	8.6 (16)
Marion	411	37.7 (27)	371	24.0 (24)	121	29.8 (18)	339	7.4 (4)	372	3.8 (2)
Marshall	153	22.2 (8)	131	15.3 (10)	22	* *	113	5.3 (1)	131	4.6 (4)
Mcperson	159	50.3 (58)	134	32.1 (51)	53	* *	132	53.8 (59)	136	51.5 (56)
Meade	117	24.8 (10)	101	15.8 (12)	20	* *	104	25.0 (34)	103	22.3 (37)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7B
2008 - 2010 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by County
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Miami	395	44.8 (44)	335	25.7 (29)	113	33.6 (11)	306	20.3 (29)	334	16.8 (28)
Mitchell	131	38.2 (28)	120	26.7 (37)	40	* *	109	9.2 (8)	122	8.2 (13)
Montgomery	981	40.8 (31)	826	28.5 (44)	265	17.4 (27)	816	33.2 (45)	830	30.5 (41)
Morris	93	* *	80	* *	29	* *	75	* *	79	* *
Nemaha	108	46.3 (47)	87	* *	37	* *	91	* *	88	* *
Neosho	395	41.8 (33)	237	23.6 (23)	77	* *	346	28.3 (39)	244	17.6 (30)
Ness	29	* *	25	* *	6	* *	23	* *	25	* *
Norton	94	* *	83	* *	31	* *	75	* *	85	* *
Osage	252	46.0 (45)	207	26.6 (36)	70	* *	209	29.7 (41)	208	26.4 (39)
Osborne	58	* *	51	* *	20	* *	43	* *	51	* *
Ottawa	73	* *	46	* *	19	* *	64	* *	59	* *
Pawnee	157	36.9 (23)	124	25.8 (31)	35	* *	133	17.3 (24)	125	14.4 (26)
Phillips	83	* *	50	* *	14	* *	67	* *	50	* *
Pottawatom	114	27.2 (12)	97	* *	23	* *	95	* *	101	5.9 (7)
Pratt	228	38.2 (28)	194	25.3 (27)	60	* *	174	40.2 (51)	194	37.6 (47)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7B
2008 - 2010 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by County
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Rawlins	29	* *	25	* *	6	* *	25	* *	25	* *
Reno	1,416	43.6 (38)	1,224	23.4 (22)	433	36.5 (5)	1,157	16.7 (23)	1,230	15.0 (27)
Republic	101	48.5 (53)	83	* *	29	* *	81	* *	85	* *
Rice	230	42.6 (37)	198	25.8 (31)	61	* *	186	17.7 (25)	200	13.5 (24)
Riley	2,681	28.5 (13)	1,947	15.3 (10)	381	30.7 (17)	2,125	15.1 (20)	2,036	12.3 (22)
Rooks	132	62.1 (64)	105	39.0 (56)	55	* *	110	29.1 (40)	111	33.3 (43)
Rush	31	* *	29	* *	16	* *	28	* *	29	* *
Russell	126	44.4 (42)	111	37.8 (55)	43	* *	103	53.4 (58)	112	52.7 (57)
Saline	1,479	42.0 (35)	1,233	26.4 (34)	430	26.5 (22)	1,235	12.6 (16)	1,237	9.9 (18)
Scott	119	26.1 (11)	95	* *	24	* *	103	30.1 (42)	97	* *
Sedgwick	13,267	31.1 (15)	12,097	16.3 (13)	2,844	34.7 (8)	10,766	9.0 (7)	12,083	7.3 (11)
Seward	1,418	6.8 (1)	1,264	3.2 (2)	66	* *	1,177	13.6 (19)	1,284	11.7 (21)
Shawnee	4,028	35.1 (18)	3,171	22.6 (21)	840	25.8 (25)	3,337	11.8 (14)	3,194	6.1 (8)
Sheridan	24	* *	21	* *	4	* *	20	* *	21	* *
Sherman	198	34.8 (17)	169	21.9 (19)	50	* *	161	25.5 (35)	168	20.8 (36)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7B
2008 - 2010 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by County
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Smith	46	* *	35	* *	12	* *	41	* *	37	* *
Stafford	111	36.9 (23)	93	* *	30	* *	92	* *	94	* *
Stanton	51	* *	47	* *	4	* *	40	* *	47	* *
Stevens	223	19.3 (6)	189	8.5 (6)	27	* *	179	25.7 (36)	191	20.4 (35)
Sumner	466	53.9 (60)	414	33.8 (52)	175	29.1 (19)	367	47.7 (56)	416	49.0 (55)
Thomas	156	34.6 (16)	135	22.2 (20)	34	* *	122	27.9 (38)	136	26.5 (40)
Trego	42	* *	32	* *	13	* *	38	* *	33	* *
Wabaunsee	60	* *	57	* *	18	* *	44	* *	57	* *
Wallace	0	* *	0	* *	0	* *	0	* *	0	* *
Washington	74	* *	65	* *	11	* *	58	* *	64	* *
Wichita	46	* *	37	* *	8	* *	36	* *	37	* *
Wilson	252	56.0 (61)	230	34.3 (53)	110	32.7 (13)	220	47.7 (56)	230	43.0 (53)
Woodson	25	* *	19	* *	2	* *	20	* *	20	* *
Wyandotte	6,335	21.1 (7)	5,508	12.4 (8)	870	26.3 (23)	5,304	19.6 (28)	5,600	17.3 (29)
Unknown Counties	22	* *	22	* *	11	* *	18	* *	23	* *

- (1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7B
2008 - 2010 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by County
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Kansas	62,786	31.6	53,620	17.9	12,870	31.8	51,403	17.6	54,019	15.1

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8B
2008 - 2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Infant Health Indicators by County
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)									
County	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Allen	257	8.6 (48)	3.5 (3)	246	9.3 (27)	218	6.0 (48)	260	57.7 (57)
Anderson	122	8.2 (44)	8.2 (51)	131	8.4 (15)	105	6.7 (50)	136	57.4 (58)
Atchison	352	7.4 (27)	5.1 (19)	345	6.1 (3)	312	4.8 (38)	365	52.3 (60)
Barber	109	11.0 (58)	4.6 (13)	103	14.6 (58)	87	* *	110	77.3 (11)
Barton	662	6.3 (15)	4.4 (11)	624	9.5 (30)	559	4.1 (29)	658	75.4 (15)
Bourbon	318	9.7 (56)	3.8 (4)	306	10.8 (45)	268	6.3 (49)	320	60.9 (49)
Brown	233	9.0 (54)	7.7 (47)	249	10.0 (41)	204	4.9 (44)	251	49.8 (61)
Butler	762	7.6 (33)	6.3 (31)	687	9.5 (30)	612	3.4 (15)	744	69.4 (33)
Chase	33	* *	* *	29	* *	28	* *	32	* *
Chautauqua	71	* *	* *	70	* *	62	* *	71	* *
Cherokee	539	7.4 (27)	6.5 (37)	493	8.7 (18)	447	2.9 (10)	528	58.0 (56)
Cheyenne	9	* *	* *	9	* *	7	* *	9	* *
Clark	9	* *	* *	7	* *	6	* *	8	* *
Clay	157	7.6 (33)	6.4 (35)	146	13.0 (56)	127	3.9 (24)	156	61.5 (44)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8B
2008 - 2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Infant Health Indicators by County
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)										
County	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)							
	Number	% (Rank)	%	(Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Cloud	225	8.0 (41)	3.1 (2)		213	11.7 (53)	187	4.8 (38)	220	71.4 (28)
Coffey	67	* *	* *		58	* *	55	* *	68	* *
Comanche	23	* *	* *		20	* *	18	* *	23	* *
Cowley	928	7.5 (30)	7.5 (46)		871	11.4 (50)	748	4.8 (38)	938	58.5 (54)
Crawford	931	8.1 (43)	4.8 (15)		896	10.4 (44)	785	4.1 (29)	940	58.9 (53)
Decatur	28	* *	* *		27	* *	25	* *	27	* *
Dickinson	332	6.6 (18)	9.6 (54)		317	9.5 (30)	286	2.4 (6)	334	74.0 (22)
Doniphan	116	12.1 (59)	10.3 (56)		107	13.1 (57)	92	* *	114	61.4 (46)
Douglas	1,264	6.7 (20)	7.0 (40)		1,184	8.9 (21)	1,063	4.0 (26)	1,254	77.7 (9)
Edwards	62	* *	* *		55	* *	43	* *	61	* *
Elk	51	* *	* *		48	* *	44	* *	51	* *
Ellis	384	6.3 (15)	6.0 (26)		382	8.6 (16)	322	2.8 (8)	401	77.6 (10)
Ellsworth	89	* *	* *		77	* *	72	* *	88	* *
Finney	1,827	6.7 (20)	6.4 (35)		1,750	9.8 (37)	1,540	3.6 (21)	1,575	78.0 (7)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8B
2008 - 2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Infant Health Indicators by County
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)										
County	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)							
	Number	% (Rank)	%	(Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Ford	1,694	4.8 (5)	6.0	(26)	1,651	8.9 (21)	1,465	2.0 (5)	1,602	74.6 (18)
Franklin	462	5.6 (8)	5.4	(21)	426	8.2 (14)	389	3.9 (24)	457	61.5 (44)
Geary	927	6.9 (24)	6.3	(31)	874	8.9 (21)	794	3.3 (13)	896	70.5 (31)
Gove	24	* * *	* *	*	22	* *	19	* *	22	* *
Graham	38	* * *	* *	*	38	* *	32	* *	39	* *
Grant	187	7.0 (25)	4.3	(8)	173	8.1 (13)	156	4.5 (36)	163	74.2 (19)
Gray	97	* * *	* *	*	108	11.1 (48)	81	* *	117	82.1 (2)
Greeley	32	* * *	* *	*	29	* *	26	* *	32	* *
Greenwood	119	5.0 (6)	5.0	(17)	114	8.8 (20)	104	4.8 (38)	117	60.7 (50)
Hamilton	64	* * *	* *	*	63	* *	57	* *	57	* *
Harper	125	6.4 (17)	9.6	(54)	123	8.9 (21)	106	2.8 (8)	132	75.8 (13)
Harvey	685	7.9 (38)	6.1	(29)	654	10.2 (43)	579	3.8 (22)	695	80.7 (3)
Haskell	53	* * *	* *	*	51	* *	47	* *	43	* *
Hodgeman	36	* * *	* *	*	32	* *	29	* *	34	* *

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one county's rate to other counties. Rank 1 = best rate.
- (4) <2500 g.
- (5) >4000 g.
- (6) <37 weeks gestation.
- (7) 37 or more weeks gestation and <2500 g.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8B
2008 - 2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Infant Health Indicators by County
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)										
County	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)							
	Number	% (Rank)	%	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Jackson	207	8.7 (50)	12.6	(59)	205	9.8 (37)	181	5.0 (45)	206	75.7 (14)
Jefferson	169	8.3 (46)	6.5	(37)	155	12.9 (54)	134	5.2 (47)	166	65.1 (40)
Jewell	28	*	*	* *	28	* *	21	* *	28	* *
Johnson	4,730	6.8 (23)	7.1	(44)	4,622	7.6 (11)	4,020	3.4 (15)	4,877	80.2 (4)
Kearny	55	*	*	* *	53	* *	44	* *	46	* *
Kingman	98	*	*	* *	129	7.0 (8)	85	* *	137	83.2 (1)
Kiowa	28	*	*	* *	29	* *	24	* *	29	* *
Labette	468	5.6 (8)	7.9	(49)	436	9.9 (39)	393	3.3 (13)	451	54.5 (59)
Lane	30	*	*	* *	28	* *	23	* *	31	* *
Leavenworth	897	7.9 (38)	7.0	(40)	823	8.9 (21)	739	4.3 (34)	910	63.5 (42)
Lincoln	57	*	*	* *	50	* *	45	* *	54	* *
Linn	163	9.2 (55)	4.9	(16)	161	11.2 (49)	137	7.3 (52)	167	74.9 (17)
Logan	33	*	*	* *	31	* *	29	* *	32	* *
Lyon	826	7.6 (33)	6.2	(30)	760	8.9 (21)	691	4.1 (29)	810	74.2 (19)

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one county's rate to other counties. Rank 1 = best rate.
- (4) <2500 g.
- (5) >4000 g.
- (6) <37 weeks gestation.
- (7) 37 or more weeks gestation and <2500 g.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8B
2008 - 2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Infant Health Indicators by County
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)										
County	Birthweight				Preterm ⁽⁶⁾		Full Term LBW ⁽⁷⁾		Ever Breastfed	
	Low ⁽⁴⁾		High ⁽⁵⁾		Number	% ^(Rank)	Number	% ^(Rank)	Number	% ^(Rank)
	Number	% ^(Rank)	% ^(Rank)	Number						
Marion	360	4.4 (4)	5.6 (22)	336	6.3 (4)	311	1.3 (3)	361	73.4 (24)	
Marshall	117	5.1 (7)	7.7 (47)	108	7.4 (10)	97	* *	119	79.8 (6)	
Mcperson	139	8.6 (48)	4.3 (8)	135	9.6 (34)	118	5.1 (46)	140	60.0 (52)	
Meade	100	6.0 (13)	5.0 (17)	99	* *	80	* *	105	80.0 (5)	
Miami	386	6.2 (14)	8.5 (52)	349	7.7 (12)	320	3.8 (22)	358	74.0 (22)	
Mitchell	104	5.8 (11)	5.8 (24)	112	7.1 (9)	90	* *	115	73.0 (26)	
Montgomery	877	8.0 (41)	3.8 (4)	816	9.7 (36)	731	4.0 (26)	859	61.8 (43)	
Morris	74	* *	* *	75	* *	61	* *	80	* *	
Nemaha	74	* *	* *	101	9.9 (39)	58	* *	104	60.6 (51)	
Neosho	319	6.6 (18)	5.3 (20)	389	10.8 (45)	265	3.4 (15)	417	58.3 (55)	
Ness	28	* *	* *	25	* *	21	* *	30	* *	
Norton	82	* *	* *	74	* *	62	* *	79	* *	
Osage	238	6.7 (20)	3.8 (4)	222	17.6 (59)	175	3.4 (15)	237	61.2 (47)	
Osborne	48	* *	* *	45	* *	42	* *	47	* *	

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8B
2008 - 2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Infant Health Indicators by County
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)										
County	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)							
	Number	% (Rank)	%	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Ottawa	72	* * * *			68	* *	63	* *	69	* *
Pawnee	133	7.5 (30)	4.5	(12)	127	8.7 (18)	116	1.7 (4)	132	72.7 (27)
Phillips	92	* * * *			87	* *	77	* *	90	* *
Pottawatom	100	4.0 (3)	7.0	(40)	93	* *	85	* *	99	* *
Pratt	193	2.1 (1)	10.4	(57)	176	5.7 (2)	164	1.2 (2)	190	68.4 (36)
Rawlins	25	* * * *			24	* *	23	* *	25	* *
Reno	1,254	7.7 (36)	6.0	(26)	1,160	9.5 (30)	1,037	4.4 (35)	1,243	65.2 (39)
Republic	85	* * * *			81	* *	68	* *	88	* *
Rice	193	7.8 (37)	10.4	(57)	183	11.5 (52)	158	3.2 (12)	192	73.4 (24)
Riley	2,247	5.7 (10)	8.0	(50)	2,052	6.8 (6)	1,891	2.7 (7)	2,240	77.8 (8)
Rooks	117	8.5 (47)	4.3	(8)	112	6.3 (4)	105	4.8 (38)	116	69.0 (35)
Rush	30	* * * *			28	* *	25	* *	30	* *
Russell	108	7.4 (27)	2.8	(1)	98	* *	84	* *	105	61.0 (48)
Saline	1,384	7.9 (38)	6.3	(31)	1,307	9.6 (34)	1,159	4.2 (32)	1,395	63.9 (41)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8B
2008 - 2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Infant Health Indicators by County
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)									
County	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Scott	99	* * * *	*	95	* *	89	* *	98	* *
Sedgwick	11,991	8.9 (53)	7.0 (40)	11,196	8.6 (16)	10,146	4.8 (38)	11,566	69.1 (34)
Seward	1,212	5.8 (11)	6.7 (39)	1,190	9.4 (28)	1,022	3.1 (11)	1,266	70.8 (30)
Shawnee	3,693	7.5 (30)	5.7 (23)	3,512	10.8 (45)	3,086	4.0 (26)	3,697	70.2 (32)
Sheridan	23	* * * *	*	20	* *	16	* *	23	* *
Sherman	170	8.8 (51)	4.7 (14)	159	6.9 (7)	148	3.4 (15)	166	74.1 (21)
Smith	42	* * * *	*	38	* *	32	* *	43	* *
Stafford	92	* * * *	*	91	* *	79	* *	89	* *
Stanton	47	* * * *	*	47	* *	42	* *	37	* *
Stevens	117	3.4 (2)	8.5 (52)	172	5.2 (1)	101	1.0 (1)	189	76.7 (12)
Sumner	422	10.9 (57)	7.1 (44)	394	11.4 (50)	340	6.8 (51)	427	68.4 (36)
Thomas	142	7.0 (25)	4.2 (7)	128	9.4 (28)	116	3.4 (15)	138	75.4 (15)
Trego	38	* * * *	*	36	* *	31	* *	38	* *
Wabaunsee	57	* * * *	*	50	* *	48	* *	56	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8B
2008 - 2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Infant Health Indicators by County
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)										
County	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)							
	Number	% (Rank)	% (Rank)		Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Wallace	1	* * * *	* *		0	* *	0	* *	1	* *
Washington	60	* * * *	* *		60	* *	48	* *	63	* *
Wichita	41	* * * *	* *		39	* *	33	* *	41	* *
Wilson	240	8.8 (51)	5.8 (24)		232	12.9 (54)	200	4.5 (36)	226	66.4 (38)
Woodson	25	* * * *	* *		27	* *	23	* *	28	* *
Wyandotte	5,495	8.2 (44)	6.3 (31)		5,247	10.0 (41)	4,570	4.2 (32)	5,549	71.2 (29)
Unknown Counties	23	* * * *	* *		23	* *	23	* *	22	* *
Kansas	55,301	7.5	6.5		52,556	9.2	46,451	4.0	54,920	70.7

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

