

Table 3B
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Racial and Ethnic Distribution by County

Comparison of Racial and Ethnic Distribution ⁽²⁾								
County	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Allen	71	*	*	*	*	*	*	*
Anderson	47	*	*	*	*	*	*	*
Atchison	148	83.8	12.2	0.7	0.7	0.0	2.7	0.0
Barber	37	*	*	*	*	*	*	*
Barton	259	68.7	0.0	30.1	0.0	0.0	1.2	0.0
Bourbon	111	90.1	4.5	1.8	1.8	0.9	0.9	0.0
Brown	89	*	*	*	*	*	*	*
Butler	253	86.6	0.8	7.9	1.6	0.0	3.2	0.0
Chase	16	*	*	*	*	*	*	*
Chautauqua	32	*	*	*	*	*	*	*
Cherokee	168	89.9	1.8	3.0	1.8	2.4	1.2	0.0
Cheyenne	20	*	*	*	*	*	*	*
Clark	4	*	*	*	*	*	*	*
Clay	50	*	*	*	*	*	*	*
Cloud	60	*	*	*	*	*	*	*
Coffey	26	*	*	*	*	*	*	*
Comanche	9	*	*	*	*	*	*	*
Cowley	361	76.7	1.4	18.0	0.8	1.9	1.1	0.0
Crawford	368	85.1	3.0	8.7	1.1	1.4	0.8	0.0
Decatur	16	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3B
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Racial and Ethnic Distribution by County

Comparison of Racial and Ethnic Distribution ⁽²⁾								
County	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Dickinson	122	91.8	0.8	4.9	0.0	1.6	0.8	0.0
Doniphan	40	*	*	*	*	*	*	*
Douglas	525	64.8	8.0	10.7	8.6	5.9	2.1	0.0
Edwards	21	*	*	*	*	*	*	*
Elk	12	*	*	*	*	*	*	*
Ellis	154	76.6	1.9	18.8	1.3	0.6	0.6	0.0
Ellsworth	34	*	*	*	*	*	*	*
Finney	684	26.8	1.3	66.8	0.0	5.0	0.1	0.0
Ford	674	19.6	3.0	76.0	0.3	0.9	0.3	0.0
Franklin	189	92.1	1.1	3.7	2.1	0.5	0.5	0.0
Geary	406	55.9	22.2	15.3	0.7	2.7	3.2	0.0
Gove	13	*	*	*	*	*	*	*
Graham	21	*	*	*	*	*	*	*
Grant	72	*	*	*	*	*	*	*
Gray	54	*	*	*	*	*	*	*
Greeley	14	*	*	*	*	*	*	*
Greenwood	35	*	*	*	*	*	*	*
Hamilton	19	*	*	*	*	*	*	*
Harper	60	*	*	*	*	*	*	*
Harvey	251	73.3	3.2	21.5	0.8	0.4	0.8	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3B
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Racial and Ethnic Distribution by County

Comparison of Racial and Ethnic Distribution ⁽²⁾								
County	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Haskell	22	*	*	*	*	*	*	*
Hodgeman	12	*	*	*	*	*	*	*
Jackson	77	*	*	*	*	*	*	*
Jefferson	57	*	*	*	*	*	*	*
Jewell	6	*	*	*	*	*	*	*
Johnson	1,898	47.5	14.6	33.6	0.3	2.8	1.2	0.0
Kearny	16	*	*	*	*	*	*	*
Kingman	47	*	*	*	*	*	*	*
Kiowa	15	*	*	*	*	*	*	*
Labette	160	80.6	10.6	4.4	2.5	0.0	1.9	0.0
Lane	16	*	*	*	*	*	*	*
Leavenworth	367	66.8	16.3	11.2	1.4	1.1	3.3	0.0
Lincoln	23	*	*	*	*	*	*	*
Linn	69	*	*	*	*	*	*	*
Logan	11	*	*	*	*	*	*	*
Lyon	267	57.3	6.0	34.8	0.4	1.1	0.4	0.0
Marion	140	85.0	2.1	10.7	0.0	0.7	1.4	0.0
Marshall	53	*	*	*	*	*	*	*
Mcperson	60	*	*	*	*	*	*	*
Meade	36	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3B
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Racial and Ethnic Distribution by County

Comparison of Racial and Ethnic Distribution ⁽²⁾								
County	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaska Native	Asian/Pacific Islander	Multiple Races	All Other/Unknown
		%	%	%	%	%	%	%
Miami	158	91.1	1.3	5.7	0.0	0.6	1.3	0.0
Mitchell	43	*	*	*	*	*	*	*
Montgomery	320	75.9	8.4	12.8	1.3	0.9	0.6	0.0
Morris	23	*	*	*	*	*	*	*
Nemaha	42	*	*	*	*	*	*	*
Neosho	154	92.2	1.3	5.2	0.0	0.0	1.3	0.0
Ness	8	*	*	*	*	*	*	*
Norton	50	*	*	*	*	*	*	*
Osage	102	96.1	1.0	2.0	0.0	0.0	1.0	0.0
Osborne	17	*	*	*	*	*	*	*
Ottawa	14	*	*	*	*	*	*	*
Pawnee	49	*	*	*	*	*	*	*
Phillips	32	*	*	*	*	*	*	*
Pottawatom	21	*	*	*	*	*	*	*
Pratt	98	*	*	*	*	*	*	*
Rawlins	8	*	*	*	*	*	*	*
Reno	460	79.6	3.0	14.3	0.2	0.2	2.6	0.0
Republic	28	*	*	*	*	*	*	*
Rice	73	*	*	*	*	*	*	*
Riley	1,018	68.2	11.3	12.5	2.0	4.5	1.6	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3B
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Racial and Ethnic Distribution by County

Comparison of Racial and Ethnic Distribution ⁽²⁾								
County	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Rooks	50	*	*	*	*	*	*	*
Rush	11	*	*	*	*	*	*	*
Russell	47	*	*	*	*	*	*	*
Saline	486	71.8	5.6	19.3	0.2	0.6	2.5	0.0
Scott	48	*	*	*	*	*	*	*
Sedgwick	4,536	45.2	17.0	30.2	1.4	3.8	2.4	0.0
Seward	447	15.7	2.2	81.0	0.4	0.4	0.2	0.0
Shawnee	1,459	53.7	17.0	23.8	1.6	1.0	2.9	0.0
Sheridan	8	*	*	*	*	*	*	*
Sherman	57	*	*	*	*	*	*	*
Smith	18	*	*	*	*	*	*	*
Stafford	24	*	*	*	*	*	*	*
Stanton	11	*	*	*	*	*	*	*
Stevens	72	*	*	*	*	*	*	*
Sumner	136	83.8	1.5	8.8	2.9	0.0	2.9	0.0
Thomas	56	*	*	*	*	*	*	*
Trego	12	*	*	*	*	*	*	*
Wabaunsee	19	*	*	*	*	*	*	*
Washington	26	*	*	*	*	*	*	*
Wichita	8	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3B
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Racial and Ethnic Distribution by County

Comparison of Racial and Ethnic Distribution ⁽²⁾								
County	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Wilson	84	*	*	*	*	*	*	*
Woodson	14	*	*	*	*	*	*	*
Wyandotte	2,101	26.2	27.0	39.3	0.6	6.1	0.9	0.0
Unknown Counties	25	*	*	*	*	*	*	*
Kansas	21,370	56.0	11.3	27.2	1.2	2.6	1.7	0.0
Nation (Prior Year)	1,162,961	41.6	22.7	29.2	1.1	3.4	1.3	0.7

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4B
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Age Distribution by County

Comparison of Age Distribution ⁽²⁾								
County	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Allen	71	*	*	*	*	*	*	*
Anderson	47	*	*	*	*	*	*	*
Atchison	148	0.0	8.1	16.9	61.5	13.5	0.0	0.0
Barber	37	*	*	*	*	*	*	*
Barton	259	0.0	7.3	16.2	64.1	11.2	1.2	0.0
Bourbon	111	0.0	4.5	18.0	59.5	17.1	0.0	0.9
Brown	89	*	*	*	*	*	*	*
Butler	253	0.0	3.6	13.0	68.0	14.2	1.2	0.0
Chase	16	*	*	*	*	*	*	*
Chautauqua	32	*	*	*	*	*	*	*
Cherokee	168	0.6	6.0	14.9	66.1	11.9	0.6	0.0
Cheyenne	20	*	*	*	*	*	*	*
Clark	4	*	*	*	*	*	*	*
Clay	50	*	*	*	*	*	*	*
Cloud	60	*	*	*	*	*	*	*
Coffey	26	*	*	*	*	*	*	*
Comanche	9	*	*	*	*	*	*	*
Cowley	361	0.0	6.6	15.5	62.9	14.1	0.8	0.0
Crawford	368	0.5	2.7	16.3	62.8	16.3	1.4	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4B
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Age Distribution by County

Comparison of Age Distribution ⁽²⁾								
County	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Decatur	16	*	*	*	*	*	*	*
Dickinson	122	0.0	4.9	17.2	65.6	12.3	0.0	0.0
Doniphan	40	*	*	*	*	*	*	*
Douglas	525	0.2	3.2	9.3	63.8	21.3	2.1	0.0
Edwards	21	*	*	*	*	*	*	*
Elk	12	*	*	*	*	*	*	*
Ellis	154	0.6	2.6	16.9	64.3	14.9	0.6	0.0
Ellsworth	34	*	*	*	*	*	*	*
Finney	684	0.4	5.6	16.7	55.6	20.3	1.5	0.0
Ford	674	0.0	5.2	12.8	58.8	20.9	2.4	0.0
Franklin	189	0.0	4.8	13.8	61.4	19.0	1.1	0.0
Geary	406	0.2	3.7	16.0	66.3	12.8	1.0	0.0
Gove	13	*	*	*	*	*	*	*
Graham	21	*	*	*	*	*	*	*
Grant	72	*	*	*	*	*	*	*
Gray	54	*	*	*	*	*	*	*
Greeley	14	*	*	*	*	*	*	*
Greenwood	35	*	*	*	*	*	*	*
Hamilton	19	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4B
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Age Distribution by County

Comparison of Age Distribution ⁽²⁾								
County	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Harper	60	*	*	*	*	*	*	*
Harvey	251	0.4	6.8	13.9	54.2	22.7	2.0	0.0
Haskell	22	*	*	*	*	*	*	*
Hodgeman	12	*	*	*	*	*	*	*
Jackson	77	*	*	*	*	*	*	*
Jefferson	57	*	*	*	*	*	*	*
Jewell	6	*	*	*	*	*	*	*
Johnson	1,898	0.1	3.0	9.5	62.2	23.4	1.7	0.1
Kearny	16	*	*	*	*	*	*	*
Kingman	47	*	*	*	*	*	*	*
Kiowa	15	*	*	*	*	*	*	*
Labette	160	1.3	6.9	11.9	66.3	13.1	0.6	0.0
Lane	16	*	*	*	*	*	*	*
Leavenworth	367	0.3	3.5	15.8	66.8	12.8	0.8	0.0
Lincoln	23	*	*	*	*	*	*	*
Linn	69	*	*	*	*	*	*	*
Logan	11	*	*	*	*	*	*	*
Lyon	267	0.4	7.5	11.6	66.3	13.1	1.1	0.0
Marion	140	0.7	4.3	15.0	62.1	16.4	1.4	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4B
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Age Distribution by County

Comparison of Age Distribution ⁽²⁾								
County	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Marshall	53	*	*	*	*	*	*	*
Mcpherson	60	*	*	*	*	*	*	*
Meade	36	*	*	*	*	*	*	*
Miami	158	0.0	3.8	17.1	70.9	8.2	0.0	0.0
Mitchell	43	*	*	*	*	*	*	*
Montgomery	320	0.6	6.9	14.7	60.3	16.3	0.9	0.3
Morris	23	*	*	*	*	*	*	*
Nemaha	42	*	*	*	*	*	*	*
Neosho	154	0.0	4.5	14.9	66.9	11.7	0.6	1.3
Ness	8	*	*	*	*	*	*	*
Norton	50	*	*	*	*	*	*	*
Osage	102	0.0	1.0	13.7	62.7	21.6	1.0	0.0
Osborne	17	*	*	*	*	*	*	*
Ottawa	14	*	*	*	*	*	*	*
Pawnee	49	*	*	*	*	*	*	*
Phillips	32	*	*	*	*	*	*	*
Pottawatom	21	*	*	*	*	*	*	*
Pratt	98	*	*	*	*	*	*	*
Rawlins	8	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4B
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Age Distribution by County

Comparison of Age Distribution ⁽²⁾								
County	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Reno	460	0.4	4.6	15.9	65.7	13.0	0.4	0.0
Republic	28	*	*	*	*	*	*	*
Rice	73	*	*	*	*	*	*	*
Riley	1,018	0.0	1.3	10.8	74.8	13.1	0.1	0.0
Rooks	50	*	*	*	*	*	*	*
Rush	11	*	*	*	*	*	*	*
Russell	47	*	*	*	*	*	*	*
Saline	486	0.6	7.2	13.8	61.3	15.6	1.4	0.0
Scott	48	*	*	*	*	*	*	*
Sedgwick	4,536	0.4	5.2	12.9	60.6	19.3	1.5	0.1
Seward	447	0.2	9.4	12.5	53.5	22.6	1.6	0.2
Shawnee	1,459	0.5	6.0	14.3	62.8	15.4	1.0	0.0
Sheridan	8	*	*	*	*	*	*	*
Sherman	57	*	*	*	*	*	*	*
Smith	18	*	*	*	*	*	*	*
Stafford	24	*	*	*	*	*	*	*
Stanton	11	*	*	*	*	*	*	*
Stevens	72	*	*	*	*	*	*	*
Sumner	136	0.0	3.7	13.2	64.7	16.9	1.5	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4B
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Age Distribution by County

Comparison of Age Distribution ⁽²⁾								
County	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Thomas	56	*	*	*	*	*	*	*
Trego	12	*	*	*	*	*	*	*
Wabaunsee	19	*	*	*	*	*	*	*
Washington	26	*	*	*	*	*	*	*
Wichita	8	*	*	*	*	*	*	*
Wilson	84	*	*	*	*	*	*	*
Woodson	14	*	*	*	*	*	*	*
Wyandotte	2,101	0.4	6.9	12.9	57.8	20.4	1.6	0.0
Unknown Counties	25	*	*	*	*	*	*	*
Kansas	21,370	0.3	5.1	13.2	62.1	17.9	1.3	0.1
Nation (Prior Year)	1,162,961	0.3	5.3	11.5	59.4	21.7	1.7	0.2

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5B
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Health Indicators by County

Comparison of Maternal Health Indicators ^(2, 3)																
County	Prepregnancy BMI (4)				Weight Gain (5)				Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight		< Ideal		> Ideal		3rd Trimester		Postpartum					
	Number	%	%	%	Number	%	%	%	Number	%	Number	%	Number	%	Number	%
	(Rank)	(Rank)	(Rank)	(Rank)	(Rank)	(Rank)	(Rank)	(Rank)	(Rank)	(Rank)	(Rank)	(Rank)	(Rank)	(Rank)	(Rank)	
Allen	70	* *	* *	* *	62	* *	* *	* *	9	* *	20	* *	64	* *	66	* *
Anderson	46	* *	* *	* *	38	* *	* *	* *	4	* *	11	* *	39	* *	39	* *
Atchison	142	8.5 (32)	47.9 (1)		120	28.3 (31)	46.7 (10)		16	* *	26	* *	120	5.0 (7)	120	7.5(4)
Barber	37	* *	* *	* *	29	* *	* *	* *	3	* *	4	* *	30	* *	30	* *
Barton	256	4.3 (19)	57.8 (26)		226	26.5 (29)	42.9 (3)		24	* *	30	* *	231	6.9 (16)	231	12.6(22)
Bourbon	110	2.7 (3)	60.9 (31)		101	21.8 (10)	52.5 (29)		16	* *	15	* *	101	14.9 (31)	101	14.9(28)
Brown	87	* *	* *	* *	72	* *	* *	* *	12	* *	12	* *	70	* *	70	* *
Butler	248	4.8 (23)	55.2 (16)		224	24.6 (23)	48.7 (16)		36	* *	38	* *	231	6.1 (12)	231	13.4(24)
Chase	16	* *	* *	* *	15	* *	* *	* *	3	* *	2	* *	15	* *	15	* *
Chautauqua	32	* *	* *	* *	28	* *	* *	* *	7	* *	4	* *	28	* *	28	* *
Cherokee	159	6.9 (28)	53.5 (10)		142	22.5 (18)	49.3 (19)		21	* *	25	* *	145	4.1 (5)	145	11.7(19)
Cheyenne	19	* *	* *	* *	16	* *	* *	* *	3	* *	8	* *	17	* *	17	* *
Clark	4	* *	* *	* *	4	* *	* *	* *	1	* *	1	* *	4	* *	4	* *
Clay	49	* *	* *	* *	42	* *	* *	* *	7	* *	9	* *	44	* *	44	* *
Cloud	58	* *	* *	* *	53	* *	* *	* *	3	* *	12	* *	55	* *	56	* *
Coffey	26	* *	* *	* *	22	* *	* *	* *	5	* *	4	* *	22	* *	22	* *
Comanche	9	* *	* *	* *	9	* *	* *	* *	0	* *	3	* *	9	* *	9	* *

- (1) Reporting period is January 1 through December 31.
 - (2) Excludes records with unknown data or errors.
 - (3) Rank compares one county's rate to other counties. Rank 1 = best rate.
 - (4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
 - (5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
 - (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5B
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Health Indicators by County

Comparison of Maternal Health Indicators ^(2, 3)														
County	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	%	%	Number	%	%	Number	%	Number	%	Number	%	Number	%
	(Rank)	(Rank)		(Rank)	(Rank)		(Rank)		(Rank)		(Rank)		(Rank)	
Cowley	354	3.1 (4)	59.0 (28)	306	21.9 (11)	47.4 (13)	41	* *	47	* *	316	6.0 (11)	316	12.0(20)
Crawford	362	6.6 (27)	56.6 (22)	316	22.2 (17)	47.5 (14)	46	* *	57	* *	321	10.3 (28)	324	13.3(23)
Decatur	16	* *	* *	15	* *	* *	3	* *	4	* *	15	* *	15	* *
Dickinson	121	3.3 (7)	57.0 (23)	104	26.0 (28)	45.2 (9)	27	* *	19	* *	105	4.8 (6)	105	12.4(21)
Doniphan	38	* *	* *	26	* *	* *	5	* *	10	* *	29	* *	30	* *
Douglas	516	3.3 (7)	52.9 (6)	432	20.1 (6)	51.9 (27)	67	* *	124	29.0 (5)	426	9.9 (27)	426	7.7(5)
Edwards	21	* *	* *	19	* *	* *	1	* *	4	* *	18	* *	18	* *
Elk	11	* *	* *	10	* *	* *	2	* *	1	* *	11	* *	11	* *
Ellis	150	3.3 (7)	50.0 (3)	125	20.0 (5)	41.6 (2)	24	* *	35	* *	132	1.5 (1)	131	5.3(2)
Ellsworth	34	* *	* *	31	* *	* *	4	* *	1	* *	30	* *	30	* *
Finney	671	7.7 (31)	53.4 (9)	598	27.1 (30)	38.5 (1)	82	* *	94	* *	606	8.9 (23)	606	5.9(3)
Ford	669	2.4 (2)	61.1 (32)	589	25.0 (24)	47.5 (14)	77	* *	70	* *	593	7.4 (19)	591	8.6(9)
Franklin	183	3.3 (7)	58.5 (27)	157	22.9 (19)	51.6 (25)	21	* *	44	* *	137	7.3 (17)	137	13.9(25)
Geary	387	3.6 (13)	51.4 (5)	310	21.9 (11)	51.0 (24)	52	* *	64	* *	323	6.2 (13)	322	8.7(10)
Gove	12	* *	* *	10	* *	* *	3	* *	3	* *	11	* *	11	* *
Graham	20	* *	* *	18	* *	* *	0	* *	1	* *	19	* *	19	* *
Grant	72	* *	* *	67	* *	* *	13	* *	17	* *	67	* *	67	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5B
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Health Indicators by County

Comparison of Maternal Health Indicators ^(2, 3)														
County	Prepregnancy BMI ⁽⁴⁾			Weight Gain ⁽⁵⁾			Anemia (Low Hb/Hct) ⁽⁶⁾				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Gray	54	* *	* *	44	* *	* *	3	* *	10	* *	44	* *	44	* *
Greeley	14	* *	* *	12	* *	* *	2	* *	3	* *	12	* *	12	* *
Greenwood	35	* *	* *	33	* *	* *	4	* *	4	* *	33	* *	33	* *
Hamilton	19	* *	* *	17	* *	* *	3	* *	4	* *	17	* *	17	* *
Harper	59	* *	* *	50	* *	* *	5	* *	11	* *	52	* *	52	* *
Harvey	246	2.0 (1)	57.3 (24)	218	22.9 (19)	47.2 (12)	31	* *	43	* *	226	3.1 (2)	226	15.9(29)
Haskell	21	* *	* *	18	* *	* *	3	* *	2	* *	19	* *	19	* *
Hodgeman	12	* *	* *	12	* *	* *	2	* *	2	* *	12	* *	12	* *
Jackson	77	* *	* *	64	* *	* *	8	* *	17	* *	66	* *	66	* *
Jefferson	56	* *	* *	50	* *	* *	9	* *	18	* *	51	* *	50	* *
Jewell	6	* *	* *	5	* *	* *	1	* *	4	* *	5	* *	5	* *
Johnson	1,836	3.1 (4)	54.7 (13)	1,616	22.1 (16)	47.0 (11)	263	21.3 (1)	446	16.6 (1)	1,665	10.9 (30)	1,664	7.8(6)
Kearny	16	* *	* *	14	* *	* *	4	* *	3	* *	14	* *	14	* *
Kingman	46	* *	* *	34	* *	* *	4	* *	11	* *	36	* *	36	* *
Kiowa	15	* *	* *	14	* *	* *	3	* *	6	* *	14	* *	14	* *
Labette	157	5.1 (24)	56.1 (20)	140	25.7 (27)	44.3 (7)	22	* *	32	* *	139	9.4 (25)	140	19.3(30)
Lane	15	* *	* *	11	* *	* *	1	* *	1	* *	12	* *	12	* *

- (1) Reporting period is January 1 through December 31.
 - (2) Excludes records with unknown data or errors.
 - (3) Rank compares one county's rate to other counties. Rank 1 = best rate.
 - (4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
 - (5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
 - (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5B
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Health Indicators by County

Comparison of Maternal Health Indicators ^(2, 3)														
County	Prepregnancy BMI ⁽⁴⁾			Weight Gain ⁽⁵⁾			Anemia (Low Hb/Hct) ⁽⁶⁾				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Leavenworth	350	4.3 (19)	52.9 (6)	290	17.9 (3)	49.3 (19)	54	* *	80	* *	302	5.6 (8)	301	11.0(16)
Lincoln	23	* *	* *	19	* *	* *	1	* *	2	* *	19	* *	19	* *
Linn	69	* *	* *	60	* *	* *	4	* *	6	* *	61	* *	61	* *
Logan	11	* *	* *	10	* *	* *	2	* *	2	* *	10	* *	10	* *
Lyon	264	3.8 (15)	59.8 (30)	230	23.9 (21)	48.7 (16)	26	* *	49	* *	232	10.3 (28)	231	14.7(26)
Marion	136	5.9 (25)	52.9 (6)	118	21.2 (8)	44.9 (8)	14	* *	37	* *	124	3.2 (3)	125	10.4(15)
Marshall	53	* *	* *	42	* *	* *	8	* *	5	* *	45	* *	45	* *
Mcperson	57	* *	* *	51	* *	* *	9	* *	18	* *	55	* *	55	* *
Meade	36	* *	* *	36	* *	* *	5	* *	8	* *	36	* *	35	* *
Miami	155	3.2 (6)	54.8 (14)	141	12.1 (1)	60.3 (31)	28	* *	35	* *	148	3.4 (4)	148	11.5(17)
Mitchell	42	* *	* *	37	* *	* *	7	* *	3	* *	37	* *	38	* *
Montgomery	311	7.4 (30)	49.2 (2)	284	25.0 (24)	43.7 (5)	61	* *	71	* *	287	9.8 (26)	286	14.7(26)
Morris	21	* *	* *	21	* *	* *	0	* *	9	* *	21	* *	22	* *
Nemaha	42	* *	* *	36	* *	* *	4	* *	6	* *	36	* *	36	* *
Neosho	147	3.4 (11)	53.7 (11)	132	22.0 (14)	49.2 (18)	23	* *	24	* *	141	7.8 (20)	141	9.9(13)
Ness	8	* *	* *	7	* *	* *	0	* *	3	* *	7	* *	7	* *
Norton	48	* *	* *	42	* *	* *	3	* *	12	* *	44	* *	44	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5B
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Health Indicators by County

Comparison of Maternal Health Indicators ^(2, 3)														
County	Prepregnancy BMI ⁽⁴⁾			Weight Gain ⁽⁵⁾			Anemia (Low Hb/Hct) ⁽⁶⁾				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Osage	100	7.0 (29)	55.0 (15)	88	*	*	13	*	25	93	93	*	*	
Osborne	17	*	15	*	3	2	15	*	15	*	15	*	*	
Ottawa	14	*	12	*	1	3	12	*	12	*	12	*	*	
Pawnee	48	*	43	*	9	9	45	*	45	*	45	*	*	
Phillips	32	*	26	*	8	6	26	*	26	*	26	*	*	
Pottawatom	21	*	16	*	2	6	17	*	17	*	17	*	*	
Pratt	95	*	86	*	14	22	89	*	89	*	89	*	*	
Rawlins	7	*	5	*	0	2	5	*	5	*	5	*	*	
Reno	451	3.8 (15)	56.1 (20)	393	19.1 (4)	52.4 (28)	54	*	69	402	6.2 (13)	402	10.2(14)	
Republic	27	*	23	*	0	8	25	*	25	*	25	*	*	
Rice	73	*	64	*	11	15	65	*	65	*	65	*	*	
Riley	972	3.7 (14)	50.6 (4)	809	20.6 (7)	51.8 (26)	149	41.6 (4)	225	27.6 (4)	844	8.9 (23)	849	9.7(12)
Rooks	48	*	44	*	4	11	43	*	43	*	43	*	*	
Rush	10	*	7	*	1	2	8	*	8	*	8	*	*	
Russell	45	*	42	*	11	9	43	*	43	*	43	*	*	
Saline	478	4.4 (22)	55.6 (18)	434	15.9 (2)	53.7 (30)	67	*	65	439	7.3 (17)	438	9.6(11)	
Scott	48	*	47	*	11	14	45	*	45	*	45	*	*	

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5B
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Health Indicators by County

Comparison of Maternal Health Indicators ^(2, 3)														
County	Prepregnancy BMI ⁽⁴⁾			Weight Gain ⁽⁵⁾			Anemia (Low Hb/Hct) ⁽⁶⁾				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Sedgwick	4,402	3.9 (17)	55.9 (19)	3,986	21.9 (11)	49.5 (22)	836	29.3 (2)	1,016	22.3 (2)	4,112	8.5 (22)	4,113	11.5(17)
Seward	440	3.4 (11)	59.5 (29)	402	25.1 (26)	43.8 (6)	74	* *	55	* *	408	8.3 (21)	407	4.7(1)
Shawnee	1,431	4.3 (19)	54.6 (12)	1,260	21.7 (9)	49.8 (23)	146	47.9 (5)	205	29.8 (6)	1,287	6.4 (15)	1,287	8.1(8)
Sheridan	7	* *	* *	7	* *	* *	2	* *	0	* *	8	* *	8	* *
Sherman	55	* *	* *	48	* *	* *	5	* *	15	* *	50	* *	50	* *
Smith	18	* *	* *	17	* *	* *	3	* *	2	* *	17	* *	17	* *
Stafford	24	* *	* *	18	* *	* *	2	* *	3	* *	18	* *	18	* *
Stanton	11	* *	* *	10	* *	* *	1	* *	4	* *	10	* *	10	* *
Stevens	70	* *	* *	62	* *	* *	15	* *	18	* *	62	* *	63	* *
Sumner	132	6.1 (26)	57.6 (25)	119	24.4 (22)	42.9 (3)	20	* *	32	* *	123	5.7 (9)	123	19.5(31)
Thomas	55	* *	* *	44	* *	* *	9	* *	7	* *	46	* *	46	* *
Trego	12	* *	* *	10	* *	* *	3	* *	2	* *	8	* *	10	* *
Wabaunsee	18	* *	* *	17	* *	* *	1	* *	10	* *	18	* *	18	* *
Washington	26	* *	* *	19	* *	* *	6	* *	4	* *	19	* *	19	* *
Wichita	8	* *	* *	8	* *	* *	1	* *	0	* *	8	* *	8	* *
Wilson	81	* *	* *	80	* *	* *	12	* *	13	* *	82	* *	82	* *
Woodson	14	* *	* *	14	* *	* *	2	* *	3	* *	14	* *	14	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5B
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Health Indicators by County

Comparison of Maternal Health Indicators ^(2, 3)														
County	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Wyandotte	2,034	4.0 (18)	55.5 (17)	1,821	22.0 (14)	49.4 (21)	310	36.5 (3)	371	22.4 (3)	1,851	5.7 (9)	1,849	8.0(7)
Unknown Counties	25	* *	* *	24	* *	* *	2	* *	3	* *	24	* *	24	* *
Kansas	20,811	4.1	55.5	18,364	22.5	48.6	3,063	30.9	4,062	22.1	18,787	7.6	18,793	10.1
Nation (Prior Year)	1,106,370	4.5	53.4	835,927	21.5	48.0	164,201	33.9	346,981	28.1	419,956	5.2	467,675	6.3

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6B
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Behavioral Indicators by County

Comparison of Maternal Behavioral Indicators ^(2, 3)										
County	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	%	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Allen	60	* *	* *	* *	71	* *	71	* *	58	* *
Anderson	37	* *	* *	* *	47	* *	47	* *	43	* *
Atchison	121	90.9 (1)	3.3 (17)		147	46.3 (2)	148	12.8 (22)	122	77.9 (1)
Barber	30	* *	* *	* *	37	* *	37	* *	33	* *
Barton	230	89.6 (6)	0.4 (3)		257	45.1 (6)	259	17.8 (12)	199	69.8 (9)
Bourbon	97	* *	* *	* *	111	38.7 (15)	111	16.2 (14)	96	* *
Brown	54	* *	* *	* *	88	* *	89	* *	79	* *
Butler	231	86.1 (12)	0.9 (6)		252	36.9 (19)	253	19.0 (10)	198	67.2 (14)
Chase	15	* *	* *	* *	16	* *	16	* *	15	* *
Chautauqua	28	* *	* *	* *	31	* *	32	* *	29	* *
Cherokee	106	84.9 (14)	4.7 (22)		168	37.5 (17)	168	10.1 (28)	136	63.2 (18)
Cheyenne	17	* *	* *	* *	20	* *	20	* *	16	* *
Clark	4	* *	* *	* *	4	* *	4	* *	4	* *
Clay	44	* *	* *	* *	50	* *	50	* *	44	* *
Cloud	56	* *	* *	* *	60	* *	60	* *	48	* *
Coffey	21	* *	* *	* *	26	* *	26	* *	23	* *
Comanche	8	* *	* *	* *	9	* *	9	* *	8	* *
Cowley	270	72.2 (25)	5.9 (24)		357	42.3 (11)	361	16.1 (16)	292	59.6 (24)
Crawford	275	86.2 (11)	4.0 (18)		365	45.8 (5)	368	13.6 (20)	304	55.9 (27)
Decatur	15	* *	* *	* *	16	* *	16	* *	14	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6B
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Behavioral Indicators by County

Comparison of Maternal Behavioral Indicators ^(2, 3)										
County	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Pregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	%	(Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Dickinson	105	82.9 (16)	1.9	(10)	120	40.8 (13)	122	21.3 (7)	104	68.3 (10)
Doniphan	31	* *	* *		40	* *	40	* *	30	* *
Douglas	391	70.6 (26)	11.0	(27)	522	42.7 (10)	525	18.7 (11)	429	64.1 (17)
Edwards	17	* *	* *		21	* *	21	* *	17	* *
Elk	11	* *	* *		12	* *	12	* *	9	* *
Ellis	125	88.0 (9)	3.2	(16)	153	28.8 (32)	154	27.9 (1)	129	61.2 (19)
Ellsworth	31	* *	* *		34	* *	34	* *	33	* *
Finney	557	49.9 (30)	29.4	(30)	682	38.6 (16)	684	7.9 (31)	555	41.3 (30)
Ford	562	76.3 (23)	9.6	(26)	667	46.0 (4)	674	12.6 (23)	588	60.4 (21)
Franklin	127	83.5 (15)	7.1	(25)	188	44.1 (7)	189	13.8 (18)	150	65.3 (15)
Geary	309	79.0 (21)	1.0	(7)	405	42.0 (12)	406	20.4 (9)	334	59.9 (22)
Gove	12	* *	* *		13	* *	13	* *	8	* *
Graham	19	* *	* *		20	* *	21	* *	18	* *
Grant	56	* *	* *		70	* *	72	* *	61	* *
Gray	43	* *	* *		52	* *	54	* *	44	* *
Greeley	7	* *	* *		14	* *	14	* *	11	* *
Greenwood	31	* *	* *		35	* *	35	* *	28	* *
Hamilton	15	* *	* *		19	* *	19	* *	15	* *
Harper	52	* *	* *		59	* *	60	* *	48	* *
Harvey	224	81.7 (19)	4.0	(18)	251	48.2 (1)	251	23.5 (4)	212	67.5 (12)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6B
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Behavioral Indicators by County

Comparison of Maternal Behavioral Indicators ^(2, 3)										
County	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Pregpregnancy (4)		During Pregpregnancy (5)	
	Number	% (Rank)	%	(Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Haskell	17	* *	* *	* *	22	* *	22	* *	18	* *
Hodgeman	12	* *	* *	* *	12	* *	12	* *	11	* *
Jackson	39	* *	* *	* *	77	* *	77	* *	59	* *
Jefferson	50	* *	* *	* *	57	* *	57	* *	49	* *
Jewell	6	* *	* *	* *	6	* *	6	* *	5	* *
Johnson	1,576	67.1 (27)	2.4 (13)		1,892	35.1 (25)	1,898	15.6 (17)	1,552	60.6 (20)
Kearny	13	* *	* *	* *	16	* *	16	* *	15	* *
Kingman	36	* *	* *	* *	47	* *	47	* *	39	* *
Kiowa	14	* *	* *	* *	15	* *	15	* *	11	* *
Labette	101	90.1 (4)	2.0 (11)		160	36.3 (21)	160	11.9 (26)	131	71.0 (5)
Lane	12	* *	* *	* *	16	* *	16	* *	13	* *
Leavenwort	302	88.1 (8)	1.3 (8)		365	31.0 (30)	367	21.5 (5)	292	74.3 (2)
Lincoln	19	* *	* *	* *	23	* *	23	* *	22	* *
Linn	61	* *	* *	* *	69	* *	69	* *	64	* *
Logan	9	* *	* *	* *	11	* *	11	* *	11	* *
Lyon	219	90.0 (5)	0.5 (4)		267	46.1 (3)	267	21.3 (7)	226	58.8 (25)
Marion	123	87.0 (10)	2.4 (13)		140	35.7 (24)	140	24.3 (3)	107	70.1 (8)
Marshall	45	* *	* *	* *	53	* *	53	* *	48	* *
Mcperson	54	* *	* *	* *	60	* *	60	* *	50	* *
Meade	34	* *	* *	* *	35	* *	36	* *	33	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6B
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Behavioral Indicators by County

Comparison of Maternal Behavioral Indicators ^(2, 3)										
County	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	%	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Miami	146	82.2 (18)	0.0	(1)	158	34.2 (26)	158	12.0 (25)	120	64.2 (16)
Mitchell	38	* *	* *		43	* *	43	* *	35	* *
Montgomery	217	89.4 (7)	1.4	(9)	316	32.3 (29)	320	13.8 (18)	270	71.5 (4)
Morris	16	* *	* *		22	* *	23	* *	16	* *
Nemaha	36	* *	* *		42	* *	42	* *	38	* *
Neosho	130	90.8 (2)	2.3	(12)	153	39.9 (14)	154	16.2 (14)	125	70.4 (6)
Ness	7	* *	* *		7	* *	8	* *	7	* *
Norton	44	* *	* *		49	* *	50	* *	41	* *
Osage	69	* *	* *		102	36.3 (21)	102	8.8 (29)	83	* *
Osborne	14	* *	* *		17	* *	17	* *	15	* *
Ottawa	12	* *	* *		14	* *	14	* *	11	* *
Pawnee	44	* *	* *		49	* *	49	* *	41	* *
Phillips	26	* *	* *		30	* *	32	* *	28	* *
Pottawatom	16	* *	* *		21	* *	21	* *	17	* *
Pratt	89	* *	* *		98	* *	98	* *	81	* *
Rawlins	5	* *	* *		8	* *	8	* *	7	* *
Reno	397	82.9 (16)	2.8	(15)	457	42.9 (9)	460	21.5 (5)	365	59.7 (23)
Republic	25	* *	* *		27	* *	28	* *	26	* *
Rice	65	* *	* *		72	* *	73	* *	62	* *
Riley	785	85.1 (13)	0.8	(5)	1,017	35.9 (23)	1,018	26.2 (2)	822	68.0 (11)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6B
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Behavioral Indicators by County

Comparison of Maternal Behavioral Indicators ^(2, 3)										
County	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	%	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Rooks	45	* *	* *	* *	50	* *	50	* *	39	* *
Rush	8	* *	* *	* *	11	* *	11	* *	8	* *
Russell	43	* *	* *	* *	46	* *	47	* *	40	* *
Saline	305	79.7 (20)	5.2 (23)		486	34.2 (26)	486	8.6 (30)	406	67.5 (12)
Scott	47	* *	* *	* *	48	* *	48	* *	44	* *
Sedgwick	3,910	76.0 (24)	4.3 (20)		4,528	29.3 (31)	4,536	16.7 (13)	3,699	52.1 (29)
Seward	355	63.4 (28)	18.0 (28)		446	37.4 (18)	447	12.1 (24)	380	70.3 (7)
Shawnee	1,148	76.8 (22)	4.6 (21)		1,455	43.2 (8)	1,459	13.4 (21)	1,247	53.6 (28)
Sheridan	8	* *	* *	* *	8	* *	8	* *	4	* *
Sherman	49	* *	* *	* *	57	* *	57	* *	44	* *
Smith	15	* *	* *	* *	18	* *	18	* *	16	* *
Stafford	16	* *	* *	* *	24	* *	24	* *	21	* *
Stanton	7	* *	* *	* *	11	* *	11	* *	9	* *
Stevens	62	* *	* *	* *	71	* *	72	* *	56	* *
Sumner	123	90.2 (3)	0.0 (1)		136	33.1 (28)	136	11.8 (27)	114	71.9 (3)
Thomas	45	* *	* *	* *	56	* *	56	* *	49	* *
Trego	10	* *	* *	* *	12	* *	12	* *	11	* *
Wabaunsee	18	* *	* *	* *	19	* *	19	* *	13	* *
Washington	19	* *	* *	* *	26	* *	26	* *	24	* *
Wichita	8	* *	* *	* *	8	* *	8	* *	7	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6B
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Behavioral Indicators by County

Comparison of Maternal Behavioral Indicators ^(2, 3)										
County	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Pregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	%	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Wilson	75	* *	* *	* *	83	* *	84	* *	73	* *
Woodson	14	* *	* *	* *	14	* *	14	* *	13	* *
Wyandotte	1,452	57.9 (29)	19.5	(29)	2,093	36.8 (20)	2,101	5.9 (32)	1,782	57.5 (26)
Unknown Counties	17	* *	* *	* *	25	* *	25	* *	22	* *
Kansas	17,166	76.1	6.3		21,285	37.4	21,370	16.4	17,671	59.5
Nation (Prior Year)	429,770	83.1	4.1		1,074,863	36.0	380,154	19.2	418,448	85.0

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7B
2011 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by County

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Allen	69	* *	56	* *	13	* *	58	* *	57	* *
Anderson	47	* *	36	* *	11	* *	42	* *	36	* *
Atchison	102	41.2 (18)	54	* *	18	* *	90	* *	54	* *
Barber	37	* *	30	* *	13	* *	33	* *	30	* *
Barton	258	38.8 (15)	230	23.0 (15)	62	* *	195	13.8 (18)	230	7.4 (13)
Bourbon	109	44.0 (24)	96	* *	38	* *	95	* *	96	* *
Brown	85	* *	56	* *	20	* *	77	* *	59	* *
Butler	251	53.4 (31)	229	33.6 (29)	102	28.4 (6)	196	7.1 (6)	229	2.6 (4)
Chase	16	* *	14	* *	9	* *	15	* *	14	* *
Chautauqua	30	* *	26	* *	8	* *	28	* *	26	* *
Cherokee	161	43.5 (23)	141	28.4 (25)	47	* *	132	36.4 (27)	141	34.8 (25)
Cheyenne	20	* *	17	* *	4	* *	16	* *	17	* *
Clark	4	* *	4	* *	0	* *	4	* *	4	* *
Clay	50	* *	44	* *	21	* *	44	* *	44	* *
Cloud	60	* *	56	* *	15	* *	48	* *	56	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7B
2011 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by County

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Coffey	26	* *	21	* *	9	* *	23	* *	21	* *
Comanche	9	* *	9	* *	2	* *	8	* *	9	* *
Cowley	351	38.5 (14)	257	26.1 (21)	85	* *	284	19.4 (20)	239	18.8 (21)
Crawford	351	44.4 (25)	269	25.7 (20)	91	* *	297	41.1 (29)	269	40.1 (29)
Decatur	14	* *	11	* *	1	* *	13	* *	11	* *
Dickinson	121	52.9 (30)	103	25.2 (19)	40	* *	103	4.9 (2)	103	4.9 (8)
Doniphan	35	* *	26	* *	4	* *	26	* *	26	* *
Douglas	511	33.5 (11)	396	17.4 (9)	106	46.2 (2)	422	11.1 (15)	414	8.7 (14)
Edwards	21	* *	17	* *	4	* *	17	* *	18	* *
Elk	12	* *	11	* *	6	* *	9	* *	11	* *
Ellis	152	42.1 (22)	127	22.8 (13)	44	* *	129	8.5 (8)	127	7.1 (12)
Ellsworth	33	* *	28	* *	13	* *	33	* *	28	* *
Finney	680	10.4 (3)	581	4.5 (2)	44	* *	553	10.1 (13)	580	10.5 (17)
Ford	646	7.3 (2)	529	5.1 (3)	27	* *	565	7.4 (7)	529	5.5 (9)
Franklin	188	41.5 (20)	159	28.9 (26)	50	* *	150	10.0 (12)	161	6.8 (11)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7B
2011 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by County

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Geary	399	40.6 (17)	307	19.5 (11)	95	* *	332	30.4 (23)	314	22.3 (23)
Gove	12	* *	10	* *	1	* *	8	* *	10	* *
Graham	20	* *	18	* *	10	* *	18	* *	18	* *
Grant	72	* *	65	* *	10	* *	61	* *	65	* *
Gray	54	* *	44	* *	4	* *	44	* *	44	* *
Greeley	13	* *	11	* *	1	* *	11	* *	11	* *
Greenwood	33	* *	30	* *	10	* *	28	* *	31	* *
Hamilton	18	* *	15	* *	1	* *	15	* *	15	* *
Harper	59	* *	50	* *	18	* *	46	* *	50	* *
Harvey	249	37.8 (13)	215	22.8 (13)	64	* *	210	2.9 (1)	214	1.9 (2)
Haskell	22	* *	19	* *	2	* *	18	* *	19	* *
Hodgeman	12	* *	12	* *	4	* *	11	* *	12	* *
Jackson	76	* *	62	* *	22	* *	58	* *	63	* *
Jefferson	49	* *	28	* *	11	* *	43	* *	29	* *
Jewell	5	* *	2	* *	0	* *	4	* *	3	* *

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one county's rate to other counties. Rank 1 = best rate.
- (4) Among women who smoked 3 months prior to pregnancy.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7B
2011 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by County

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Johnson	1,883	23.3 (5)	1,654	9.1 (4)	285	50.9 (1)	1,537	17.6 (19)	1,655	18.2 (20)
Kearny	16	* *	13	* *	3	* *	15	* *	13	* *
Kingman	46	* *	32	* *	8	* *	38	* *	33	* *
Kiowa	15	* *	12	* *	3	* *	11	* *	14	* *
Labette	154	41.6 (21)	135	31.1 (27)	49	* *	126	34.9 (26)	136	36.8 (28)
Lane	10	* *	1	* *	0	* *	9	* *	2	* *
Leavenworth	356	44.4 (25)	284	24.3 (16)	94	* *	287	34.8 (25)	286	35.3 (26)
Lincoln	22	* *	18	* *	12	* *	22	* *	18	* *
Linn	68	* *	59	* *	23	* *	63	* *	60	* *
Logan	11	* *	10	* *	2	* *	11	* *	10	* *
Lyon	260	30.0 (8)	224	12.9 (6)	56	* *	224	8.9 (10)	208	2.4 (3)
Marion	140	32.9 (10)	125	21.6 (12)	31	* *	106	5.7 (3)	125	1.6 (1)
Marshall	53	* *	43	* *	10	* *	48	* *	43	* *
Mcperson	60	* *	53	* *	20	* *	50	* *	53	* *
Meade	34	* *	32	* *	11	* *	32	* *	34	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7B
2011 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by County

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Miami	157	46.5 (27)	143	32.2 (28)	48	* *	115	20.9 (21)	143	16.1 (19)
Mitchell	42	* *	37	* *	8	* *	35	* *	37	* *
Montgomery	312	37.5 (12)	280	26.8 (23)	86	* *	268	31.0 (24)	278	33.5 (24)
Morris	22	* *	21	* *	4	* *	16	* *	22	* *
Nemaha	17	* *	11	* *	4	* *	11	* *	13	* *
Neosho	145	41.4 (19)	103	26.2 (22)	34	* *	120	25.8 (22)	103	20.4 (22)
Ness	8	* *	7	* *	3	* *	6	* *	7	* *
Norton	48	* *	34	* *	10	* *	40	* *	36	* *
Osage	97	* *	86	* *	32	* *	80	* *	87	* *
Osborne	16	* *	9	* *	2	* *	15	* *	9	* *
Ottawa	13	* *	10	* *	2	* *	10	* *	10	* *
Pawnee	49	* *	40	* *	11	* *	41	* *	40	* *
Phillips	30	* *	21	* *	7	* *	25	* *	22	* *
Pottawatom	21	* *	17	* *	4	* *	17	* *	17	* *
Pratt	96	* *	85	* *	31	* *	76	* *	80	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7B
2011 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by County

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Rawlins	7	* *	5	* *	1	* *	7	* *	5	* *
Reno	453	48.1 (28)	391	26.9 (24)	140	37.9 (4)	359	12.5 (16)	393	8.9 (16)
Republic	28	* *	24	* *	10	* *	26	* *	25	* *
Rice	72	* *	64	* *	28	* *	62	* *	63	* *
Riley	991	28.1 (6)	814	13.6 (7)	191	42.9 (3)	813	7.0 (5)	820	4.3 (5)
Rooks	46	* *	36	* *	13	* *	34	* *	37	* *
Rush	11	* *	8	* *	4	* *	8	* *	8	* *
Russell	46	* *	43	* *	14	* *	40	* *	42	* *
Saline	476	40.1 (16)	429	24.5 (17)	135	25.9 (9)	401	6.7 (4)	431	4.6 (6)
Scott	40	* *	32	* *	9	* *	36	* *	32	* *
Sedgwick	4,500	29.2 (7)	4,055	16.7 (8)	905	33.8 (5)	3,670	8.7 (9)	4,063	5.8 (10)
Seward	442	4.3 (1)	398	1.8 (1)	12	* *	376	10.4 (14)	402	8.7 (14)
Shawnee	1,370	32.3 (9)	1,016	19.1 (10)	249	26.5 (8)	1,163	9.9 (11)	1,001	4.8 (7)
Sheridan	7	* *	5	* *	1	* *	4	* *	7	* *
Sherman	56	* *	50	* *	10	* *	43	* *	50	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7B
2011 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by County

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Smith	18	* *	13	* *	4	* *	16	* *	13	* *
Stafford	23	* *	17	* *	3	* *	20	* *	16	* *
Stanton	11	* *	10	* *	1	* *	9	* *	9	* *
Stevens	71	* *	62	* *	15	* *	56	* *	63	* *
Sumner	135	49.6 (29)	121	24.8 (18)	49	* *	111	37.8 (28)	120	35.8 (27)
Thomas	55	* *	45	* *	12	* *	46	* *	46	* *
Trego	12	* *	10	* *	4	* *	10	* *	10	* *
Wabaunsee	19	* *	18	* *	3	* *	13	* *	18	* *
Washington	26	* *	18	* *	6	* *	24	* *	19	* *
Wichita	8	* *	6	* *	1	* *	7	* *	6	* *
Wilson	83	* *	81	* *	30	* *	72	* *	80	* *
Woodson	14	* *	14	* *	5	* *	13	* *	14	* *
Wyandotte	2,062	19.4 (4)	1,800	11.9 (5)	271	27.7 (7)	1,760	12.5 (16)	1,845	11.2 (18)
Unknown Counties	25	* *	23	* *	8	* *	22	* *	22	* *
Kansas	20,850	30.4	17,728	17.5	4,187	32.7	17,287	15.1	17,791	12.6

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7B
2011 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by County

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal	Postpartum			
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)		
Nation (Prior Year)	814,631	23.0	715,034	12.1	137,883	38.1	660,570	16.3	562,126	15.6

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8B
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Infant Health Indicators by County

Comparison of Infant Health Indicators ^(2, 3)										
County	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)		Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
	Number	% (Rank)	% (Rank)	Number						
Allen	59	* * * *	* * * *	65	* *	58	* *	65	* *	
Anderson	36	* * * *	* * * *	38	* *	33	* *	38	* *	
Atchison	114	13.2 (30)	1.8 (1)	125	10.4 (24)	102	8.8 (29)	125	52.0 (30)	
Barber	27	* * * *	* * * *	28	* *	25	* *	27	* *	
Barton	220	5.0 (4)	5.0 (7)	220	6.4 (6)	204	3.4 (8)	221	74.2 (15)	
Bourbon	99	* * * *	* * * *	99	* *	87	* *	99	* *	
Brown	69	* * * *	* * * *	75	* *	60	* *	75	* *	
Butler	220	7.7 (18)	5.0 (7)	223	7.6 (11)	202	5.4 (24)	223	76.7 (11)	
Chase	15	* * * *	* * * *	15	* *	15	* *	15	* *	
Chautauqua	25	* * * *	* * * *	25	* *	23	* *	26	* *	
Cherokee	146	8.9 (27)	13.0 (30)	147	10.9 (27)	130	5.4 (24)	147	63.9 (27)	
Cheyenne	15	* * * *	* * * *	15	* *	12	* *	15	* *	
Clark	4	* * * *	* * * *	4	* *	4	* *	4	* *	
Clay	41	* * * *	* * * *	42	* *	36	* *	42	* *	

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8B
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Infant Health Indicators by County

Comparison of Infant Health Indicators ^(2, 3)										
County	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)							
	Number	% (Rank)	%	(Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Cloud	52	* * * *	* *	* *	53	* *	45	* *	53	* *
Coffey	21	* * * *	* *	* *	21	* *	21	* *	21	* *
Comanche	7	* * * *	* *	* *	8	* *	6	* *	8	* *
Cowley	283	6.7 (7)	5.7 (13)		303	8.9 (15)	253	4.7 (19)	303	64.0 (26)
Crawford	302	7.9 (21)	5.0 (7)		313	9.6 (20)	269	3.3 (7)	316	66.5 (23)
Decatur	14	* * * *	* *	* *	14	* *	13	* *	14	* *
Dickinson	100	7.0 (11)	10.0 (29)		98	* *	90	* *	100	81.0 (5)
Doniphan	28	* * * *	* *	* *	28	* *	23	* *	27	* *
Douglas	411	5.6 (5)	6.3 (17)		422	7.1 (8)	379	3.2 (6)	423	80.4 (6)
Edwards	18	* * * *	* *	* *	19	* *	13	* *	17	* *
Elk	10	* * * *	* *	* *	11	* *	9	* *	11	* *
Ellis	121	7.4 (16)	7.4 (23)		127	3.1 (1)	116	5.2 (23)	128	81.3 (4)
Ellsworth	31	* * * *	* *	* *	31	* *	30	* *	31	* *
Finney	593	6.6 (6)	5.1 (10)		600	10.2 (23)	533	3.6 (10)	596	75.3 (14)

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one county's rate to other counties. Rank 1 = best rate.
- (4) <2500 g.
- (5) >4000 g.
- (6) <37 weeks gestation.
- (7) 37 or more weeks gestation and <2500 g.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8B
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Infant Health Indicators by County

Comparison of Infant Health Indicators ^(2, 3)										
County	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)							
	Number	% (Rank)	%	(Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Ford	549	7.3 (15)	4.9	(6)	558	8.4 (13)	496	4.2 (14)	557	77.6 (9)
Franklin	156	7.1 (12)	4.5	(4)	158	12.7 (28)	135	3.0 (5)	159	64.2 (25)
Geary	314	8.0 (23)	9.6	(28)	316	6.0 (4)	294	5.4 (24)	315	75.6 (12)
Gove	12	*	*	*	12	*	*	12	*	*
Graham	16	*	*	*	17	*	*	13	*	*
Grant	65	*	*	*	63	*	*	60	*	*
Gray	41	*	*	*	40	*	*	37	*	*
Greeley	12	*	*	*	12	*	*	11	*	*
Greenwood	31	*	*	*	32	*	*	30	*	*
Hamilton	17	*	*	*	17	*	*	16	*	*
Harper	50	*	*	*	50	*	*	45	*	*
Harvey	217	7.8 (20)	7.8	(26)	219	10.0 (22)	195	5.1 (22)	219	88.1 (1)
Haskell	17	*	*	*	17	*	*	15	*	*
Hodgeman	11	*	*	*	11	*	*	10	*	*

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one county's rate to other counties. Rank 1 = best rate.
- (4) <2500 g.
- (5) >4000 g.
- (6) <37 weeks gestation.
- (7) 37 or more weeks gestation and <2500 g.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8B
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Infant Health Indicators by County

Comparison of Infant Health Indicators ^(2, 3)										
County	Birthweight				Preterm ⁽⁶⁾		Full Term LBW ⁽⁷⁾		Ever Breastfed	
	Low ⁽⁴⁾		High ⁽⁵⁾		Number	%	Number	%	Number	%
	Number	% (Rank)	% (Rank)	(Rank)						
Jackson	60	* * * *	* * * *	63	* *	56	* *	63	* *	
Jefferson	49	* * * *	* * * *	49	* *	40	* *	49	* *	
Jewell	6	* * * *	* * * *	6	* *	5	* *	6	* *	
Johnson	1,566	6.7 (7)	7.2 (22)	1,630	6.3 (5)	1,459	3.9 (13)	1,627	82.7 (2)	
Kearny	14	* * * *	* * * *	14	* *	10	* *	14	* *	
Kingman	24	* * * *	* * * *	32	* *	22	* *	32	* *	
Kiowa	14	* * * *	* * * *	14	* *	12	* *	13	* *	
Labette	139	7.2 (13)	5.8 (14)	139	12.9 (29)	121	5.0 (21)	138	55.1 (29)	
Lane	10	* * * *	* * * *	10	* *	9	* *	10	* *	
Leavenworth	281	10.0 (29)	5.3 (11)	304	7.9 (12)	257	6.6 (28)	304	66.1 (24)	
Lincoln	17	* * * *	* * * *	18	* *	14	* *	18	* *	
Linn	61	* * * *	* * * *	61	* *	51	* *	61	* *	
Logan	10	* * * *	* * * *	10	* *	9	* *	10	* *	
Lyon	229	7.9 (21)	7.4 (23)	229	9.6 (20)	207	2.9 (4)	228	79.4 (8)	

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8B
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Infant Health Indicators by County

Comparison of Infant Health Indicators ^(2, 3)											
County	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed		
	Low (4)		High (5)								
	Number	% (Rank)	%	(Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	
Marion	118	3.4 (1)	2.5 (2)		120	4.2 (2)		113	1.8 (2)	120	77.5 (10)
Marshall	44	* *	* *		44	* *		39	* *	44	* *
Mcperson	52	* *	* *		52	* *		47	* *	52	* *
Meade	32	* *	* *		34	* *		27	* *	34	* *
Miami	147	6.8 (9)	7.5 (25)		148	9.5 (19)		133	3.8 (11)	147	66.7 (22)
Mitchell	34	* *	* *		37	* *		34	* *	37	* *
Montgomery	279	9.0 (28)	2.9 (3)		280	7.1 (8)		257	6.2 (27)	281	60.5 (28)
Morris	21	* *	* *		21	* *		15	* *	22	* *
Nemaha	32	* *	* *		36	* *		31	* *	36	* *
Neosho	130	4.6 (3)	6.2 (16)		139	5.8 (3)		121	1.7 (1)	139	72.7 (17)
Ness	6	* *	* *		6	* *		4	* *	7	* *
Norton	39	* *	* *		39	* *		35	* *	39	* *
Osage	86	* *	* *		91	* *		79	* *	91	* *
Osborne	14	* *	* *		14	* *		14	* *	14	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8B
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Infant Health Indicators by County

Comparison of Infant Health Indicators ^(2, 3)										
County	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)							
	Number	% (Rank)	%	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Ottawa	10	* * * *	* *	* *	12	* *	9	* *	12	* *
Pawnee	41	* * * *	* *	* *	43	* *	35	* *	42	* *
Phillips	24	* * * *	* *	* *	23	* *	22	* *	24	* *
Pottawatom	15	* * * *	* *	* *	15	* *	14	* *	15	* *
Pratt	87	* * * *	* *	* *	87	* *	81	* *	87	* *
Rawlins	6	* * * *	* *	* *	6	* *	6	* *	6	* *
Reno	385	6.8 (9)	6.8 (20)		386	7.3 (10)	354	4.2 (14)	387	72.6 (18)
Republic	23	* * * *	* *	* *	23	* *	19	* *	24	* *
Rice	63	* * * *	* *	* *	62	* *	58	* *	63	* *
Riley	825	7.2 (13)	6.8 (20)		830	6.5 (7)	771	4.7 (19)	825	82.4 (3)
Rooks	45	* * * *	* *	* *	46	* *	40	* *	45	* *
Rush	7	* * * *	* *	* *	7	* *	7	* *	7	* *
Russell	42	* * * *	* *	* *	41	* *	33	* *	42	* *
Saline	400	8.0 (23)	4.5 (4)		430	10.7 (25)	355	3.4 (8)	429	70.4 (20)

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one county's rate to other counties. Rank 1 = best rate.
- (4) <2500 g.
- (5) >4000 g.
- (6) <37 weeks gestation.
- (7) 37 or more weeks gestation and <2500 g.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8B
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Infant Health Indicators by County

Comparison of Infant Health Indicators ^(2, 3)										
County	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)							
	Number	% (Rank)	%	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Scott	45	* * * *	* *	* *	46	* *	39	* *	46	* *
Sedgwick	3,998	8.6 (26)	6.5 (18)		4,010	8.7 (14)	3,645	4.6 (18)	4,004	73.2 (16)
Seward	368	4.1 (2)	8.4 (27)		381	9.2 (17)	333	2.1 (3)	382	80.1 (7)
Shawnee	1,243	8.5 (25)	5.5 (12)		1,252	9.3 (18)	1,123	4.5 (17)	1,249	69.9 (21)
Sheridan	8	* * * *	* *	* *	8	* *	8	* *	8	* *
Sherman	47	* * * *	* *	* *	47	* *	43	* *	47	* *
Smith	17	* * * *	* *	* *	17	* *	16	* *	17	* *
Stafford	16	* * * *	* *	* *	16	* *	13	* *	16	* *
Stanton	10	* * * *	* *	* *	10	* *	10	* *	10	* *
Stevens	36	* * * *	* *	* *	62	* *	32	* *	63	* *
Sumner	119	7.6 (17)	6.7 (19)		122	10.7 (25)	106	3.8 (11)	122	75.4 (13)
Thomas	43	* * * *	* *	* *	43	* *	40	* *	43	* *
Trego	8	* * * *	* *	* *	10	* *	7	* *	10	* *
Wabaunsee	18	* * * *	* *	* *	18	* *	17	* *	18	* *

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one county's rate to other counties. Rank 1 = best rate.
- (4) <2500 g.
- (5) >4000 g.
- (6) <37 weeks gestation.
- (7) 37 or more weeks gestation and <2500 g.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8B
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Infant Health Indicators by County

Comparison of Infant Health Indicators ^(2, 3)										
County	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)							
	Number	% (Rank)	%	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Washington	17	* * * *	* *	* *	17	* *	17	* *	17	* *
Wichita	8	* * * *	* *	* *	8	* *	8	* *	8	* *
Wilson	75	* * * *	* *	* *	77	* *	68	* *	77	* *
Woodson	14	* * * *	* *	* *	14	* *	13	* *	14	* *
Wyandotte	1,786	7.7 (18)	5.8	(14)	1,832	9.1 (16)	1,619	4.3 (16)	1,817	72.2 (19)
Unknown Counties	24	* * * *	* *	* *	24	* *	24	* *	24	* *
Kansas	17,976	7.6	6.2		18,356	8.5	16,366	4.3	18,328	74.2
Nation (Prior Year)	880,933	8.0	6.7		948,286	10.7	749,107	3.4	871,811	68.2

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3B
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Racial and Ethnic Distribution by County
Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾								
County	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Allen	246	91.9	2.8	1.2	1.6	0.0	2.4	0.0
Anderson	153	96.7	0.0	3.3	0.0	0.0	0.0	0.0
Atchison	443	86.9	9.5	1.1	0.9	0.0	1.6	0.0
Barber	139	95.0	1.4	1.4	0.0	1.4	0.7	0.0
Barton	771	67.4	1.3	29.8	0.1	0.5	0.8	0.0
Bourbon	355	92.1	2.5	2.5	1.1	0.6	1.1	0.0
Brown	279	74.2	1.8	5.7	16.5	0.0	1.8	0.0
Butler	820	88.3	1.1	7.3	1.0	0.2	2.1	0.0
Chase	41	*	*	*	*	*	*	*
Chautauqua	83	*	*	*	*	*	*	*
Cherokee	563	89.7	1.4	2.7	1.8	3.0	1.4	0.0
Cheyenne	33	*	*	*	*	*	*	*
Clark	10	*	*	*	*	*	*	*
Clay	160	93.8	0.6	1.9	1.9	0.0	1.9	0.0
Cloud	242	92.6	1.7	2.1	2.1	0.8	0.8	0.0
Coffey	84	*	*	*	*	*	*	*
Comanche	31	*	*	*	*	*	*	*
Cowley	1,087	75.1	2.1	19.3	0.8	1.7	1.0	0.0
Crawford	1,108	84.2	3.2	9.4	0.8	1.2	1.2	0.0
Decatur	40	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3B
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Racial and Ethnic Distribution by County
Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾								
County	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaska Native	Asian/Pacific Islander	Multiple Races	All Other/Unknown
		%	%	%	%	%	%	%
Dickinson	411	91.2	0.2	5.6	1.0	1.0	1.0	0.0
Doniphan	130	90.8	2.3	4.6	0.8	0.0	1.5	0.0
Douglas	1,562	65.2	9.3	10.8	8.1	4.7	1.9	0.0
Edwards	71	*	*	*	*	*	*	*
Elk	56	*	*	*	*	*	*	*
Ellis	492	81.5	1.0	14.8	1.2	0.6	0.8	0.0
Ellsworth	104	90.4	0.0	6.7	1.0	0.0	1.9	0.0
Finney	2,105	28.5	1.1	65.9	0.0	4.1	0.3	0.0
Ford	2,057	21.0	2.1	75.9	0.2	0.7	0.1	0.0
Franklin	561	92.2	0.5	4.3	1.6	0.4	1.1	0.0
Geary	1,220	54.7	22.3	15.3	0.8	3.1	3.8	0.0
Gove	35	*	*	*	*	*	*	*
Graham	58	*	*	*	*	*	*	*
Grant	209	23.9	0.0	75.6	0.0	0.0	0.5	0.0
Gray	145	73.1	0.0	26.2	0.7	0.0	0.0	0.0
Greeley	41	*	*	*	*	*	*	*
Greenwood	131	93.1	0.8	4.6	0.8	0.0	0.8	0.0
Hamilton	77	*	*	*	*	*	*	*
Harper	165	80.6	0.0	17.0	0.6	0.0	1.8	0.0
Harvey	818	71.5	2.9	21.9	1.3	0.9	1.5	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3B
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Racial and Ethnic Distribution by County
Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾								
County	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Haskell	59	*	*	*	*	*	*	*
Hodgeman	39	*	*	*	*	*	*	*
Jackson	240	74.6	0.4	6.3	15.8	0.0	2.9	0.0
Jefferson	183	94.0	0.5	2.2	2.7	0.0	0.5	0.0
Jewell	22	*	*	*	*	*	*	*
Johnson	5,804	47.0	13.6	34.5	0.4	3.2	1.4	0.0
Kearny	56	*	*	*	*	*	*	*
Kingman	155	92.3	0.0	5.8	0.6	1.3	0.0	0.0
Kiowa	39	*	*	*	*	*	*	*
Labette	530	85.7	7.2	3.2	2.1	0.2	1.7	0.0
Lane	40	*	*	*	*	*	*	*
Leavenworth	1,090	69.2	14.2	11.4	0.8	1.2	3.2	0.0
Lincoln	64	*	*	*	*	*	*	*
Linn	206	95.6	0.0	2.4	1.5	0.0	0.5	0.0
Logan	38	*	*	*	*	*	*	*
Lyon	884	57.0	5.1	34.8	0.5	1.7	0.9	0.0
Marion	409	85.8	1.7	10.3	0.5	0.7	1.0	0.0
Marshall	161	91.9	0.0	7.5	0.0	0.6	0.0	0.0
Mcperson	160	89.4	0.6	3.1	1.9	0.0	5.0	0.0
Meade	118	65.3	0.8	33.1	0.0	0.0	0.8	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3B
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Racial and Ethnic Distribution by County
Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾								
County	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Miami	455	91.2	2.4	4.0	0.2	0.7	1.5	0.0
Mitchell	133	90.2	0.0	7.5	0.0	1.5	0.8	0.0
Montgomery	1,009	76.1	9.1	10.8	2.3	0.8	0.9	0.0
Morris	85	*	*	*	*	*	*	*
Nemaha	123	90.2	0.0	7.3	0.8	0.8	0.8	0.0
Neosho	466	91.8	1.9	4.5	0.4	0.0	1.3	0.0
Ness	35	*	*	*	*	*	*	*
Norton	120	90.0	1.7	6.7	0.0	0.8	0.8	0.0
Osage	282	96.5	0.4	1.8	0.4	0.0	1.1	0.0
Osborne	57	*	*	*	*	*	*	*
Ottawa	71	*	*	*	*	*	*	*
Pawnee	169	78.7	4.1	14.2	1.8	0.6	0.6	0.0
Phillips	109	93.6	0.0	3.7	0.0	1.8	0.9	0.0
Pottawatom	95	*	*	*	*	*	*	*
Pratt	265	84.9	0.8	14.0	0.0	0.0	0.4	0.0
Rawlins	27	*	*	*	*	*	*	*
Reno	1,412	79.2	2.6	15.4	0.4	0.5	1.9	0.0
Republic	98	*	*	*	*	*	*	*
Rice	237	75.9	2.1	20.3	0.8	0.8	0.0	0.0
Riley	2,981	70.7	11.0	12.5	0.9	3.9	1.0	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3B
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Racial and Ethnic Distribution by County
Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾								
County	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Rooks	150	94.7	1.3	2.0	1.3	0.7	0.0	0.0
Rush	30	*	*	*	*	*	*	*
Russell	133	89.5	4.5	5.3	0.8	0.0	0.0	0.0
Saline	1,563	70.9	4.7	20.3	0.2	1.7	2.2	0.0
Scott	135	61.5	0.0	38.5	0.0	0.0	0.0	0.0
Sedgwick	13,808	45.2	17.7	29.8	1.5	3.7	2.1	0.0
Seward	1,407	15.2	2.0	81.7	0.5	0.4	0.1	0.0
Shawnee	4,473	54.6	16.0	24.2	2.2	0.9	2.1	0.0
Sheridan	26	*	*	*	*	*	*	*
Sherman	189	75.1	0.5	23.3	0.0	0.5	0.5	0.0
Smith	53	*	*	*	*	*	*	*
Stafford	100	65.0	1.0	33.0	0.0	1.0	0.0	0.0
Stanton	41	*	*	*	*	*	*	*
Stevens	231	46.8	0.4	51.5	1.3	0.0	0.0	0.0
Sumner	457	86.9	1.3	7.0	2.0	0.2	2.6	0.0
Thomas	166	83.7	1.2	13.3	0.6	1.2	0.0	0.0
Trego	44	*	*	*	*	*	*	*
Wabaunsee	61	*	*	*	*	*	*	*
Washington	75	*	*	*	*	*	*	*
Wichita	44	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3B
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Racial and Ethnic Distribution by County
Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾								
County	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Wilson	260	94.2	1.5	1.5	0.4	0.0	2.3	0.0
Woodson	40	*	*	*	*	*	*	*
Wyandotte	6,487	26.4	28.8	39.3	0.5	4.2	0.8	0.0
Unknown Counties	48	*	*	*	*	*	*	*
Kansas	65,683	56.6	11.2	27.2	1.2	2.3	1.5	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4B
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Age Distribution by County
Combined 3 Years

Comparison of Age Distribution ⁽²⁾								
County	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Allen	246	0.4	8.5	16.3	63.4	10.6	0.8	0.0
Anderson	153	0.7	4.6	17.6	61.4	15.7	0.0	0.0
Atchison	443	0.0	7.7	15.6	61.9	14.7	0.2	0.0
Barber	139	0.0	5.8	20.1	59.7	13.7	0.7	0.0
Barton	771	0.0	7.8	14.4	62.3	13.7	1.8	0.0
Bourbon	355	0.3	6.8	17.5	59.2	15.5	0.6	0.3
Brown	279	0.4	7.2	11.5	65.6	14.3	1.1	0.0
Butler	820	0.0	5.7	16.0	63.8	13.7	0.6	0.2
Chase	41	*	*	*	*	*	*	*
Chautauqua	83	*	*	*	*	*	*	*
Cherokee	563	0.4	8.3	19.2	60.0	11.4	0.5	0.2
Cheyenne	33	*	*	*	*	*	*	*
Clark	10	*	*	*	*	*	*	*
Clay	160	1.3	9.4	18.1	56.9	14.4	0.0	0.0
Cloud	242	0.4	6.6	15.3	59.5	15.7	2.5	0.0
Coffey	84	*	*	*	*	*	*	*
Comanche	31	*	*	*	*	*	*	*
Cowley	1,087	0.5	7.1	16.4	62.2	13.2	0.6	0.0
Crawford	1,108	0.3	4.7	15.7	64.8	13.8	0.6	0.1

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4B
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Age Distribution by County
Combined 3 Years

Comparison of Age Distribution ⁽²⁾								
County	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Decatur	40	*	*	*	*	*	*	*
Dickinson	411	0.0	5.6	14.8	65.0	14.4	0.2	0.0
Doniphan	130	0.8	2.3	11.5	66.2	18.5	0.0	0.8
Douglas	1,562	0.1	3.8	11.1	63.8	19.8	1.5	0.0
Edwards	71	*	*	*	*	*	*	*
Elk	56	*	*	*	*	*	*	*
Ellis	492	0.6	4.1	14.8	65.9	13.6	0.8	0.2
Ellsworth	104	0.0	2.9	11.5	57.7	24.0	3.8	0.0
Finney	2,105	0.5	7.1	15.5	55.7	19.3	1.9	0.0
Ford	2,057	0.2	6.4	12.9	55.5	22.5	2.4	0.0
Franklin	561	0.2	5.2	16.6	61.0	16.0	1.1	0.0
Geary	1,220	0.2	4.2	14.4	68.2	12.1	0.7	0.1
Gove	35	*	*	*	*	*	*	*
Graham	58	*	*	*	*	*	*	*
Grant	209	0.0	5.3	12.9	62.7	17.7	1.4	0.0
Gray	145	0.0	4.1	12.4	60.7	21.4	1.4	0.0
Greeley	41	*	*	*	*	*	*	*
Greenwood	131	0.0	4.6	16.0	62.6	16.0	0.0	0.8
Hamilton	77	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4B
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Age Distribution by County
Combined 3 Years

Comparison of Age Distribution ⁽²⁾								
County	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Harper	165	0.0	5.5	12.1	67.9	13.9	0.6	0.0
Harvey	818	0.4	6.2	15.3	59.5	17.6	1.0	0.0
Haskell	59	*	*	*	*	*	*	*
Hodgeman	39	*	*	*	*	*	*	*
Jackson	240	0.0	5.4	18.3	63.8	11.7	0.8	0.0
Jefferson	183	0.0	3.8	16.4	56.3	23.0	0.5	0.0
Jewell	22	*	*	*	*	*	*	*
Johnson	5,804	0.1	3.3	10.1	60.8	23.8	1.8	0.1
Kearny	56	*	*	*	*	*	*	*
Kingman	155	0.0	6.5	16.8	64.5	10.3	1.9	0.0
Kiowa	39	*	*	*	*	*	*	*
Labette	530	0.4	8.1	17.4	61.3	12.1	0.8	0.0
Lane	40	*	*	*	*	*	*	*
Leavenworth	1,090	0.2	5.2	14.4	66.4	13.1	0.6	0.0
Lincoln	64	*	*	*	*	*	*	*
Linn	206	0.0	6.3	13.1	62.1	18.0	0.5	0.0
Logan	38	*	*	*	*	*	*	*
Lyon	884	0.7	5.5	13.6	64.5	14.4	1.4	0.0
Marion	409	0.5	4.2	12.5	64.5	17.1	1.2	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4B
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Age Distribution by County
Combined 3 Years

Comparison of Age Distribution ⁽²⁾								
County	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Marshall	161	0.0	1.2	16.1	59.6	19.9	3.1	0.0
Mcperson	160	0.0	5.0	17.5	63.8	13.1	0.6	0.0
Meade	118	0.0	10.2	11.0	55.9	22.0	0.8	0.0
Miami	455	0.4	4.2	17.6	65.3	11.9	0.7	0.0
Mitchell	133	0.0	1.5	14.3	67.7	16.5	0.0	0.0
Montgomery	1,009	0.5	5.3	16.4	62.9	13.7	1.1	0.2
Morris	85	*	*	*	*	*	*	*
Nemaha	123	0.0	3.3	11.4	65.9	13.0	5.7	0.8
Neosho	466	0.0	6.4	16.3	63.9	12.0	0.9	0.4
Ness	35	*	*	*	*	*	*	*
Norton	120	0.0	1.7	15.8	57.5	24.2	0.8	0.0
Osage	282	0.0	3.9	18.4	61.0	14.9	1.4	0.4
Osborne	57	*	*	*	*	*	*	*
Ottawa	71	*	*	*	*	*	*	*
Pawnee	169	0.6	5.3	13.0	69.2	11.2	0.6	0.0
Phillips	109	0.9	1.8	14.7	66.1	15.6	0.9	0.0
Pottawatom	95	*	*	*	*	*	*	*
Pratt	265	0.4	7.2	14.0	64.2	13.6	0.8	0.0
Rawlins	27	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4B
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Age Distribution by County
Combined 3 Years

Comparison of Age Distribution ⁽²⁾								
County	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Reno	1,412	0.2	6.4	15.8	63.7	13.4	0.4	0.0
Republic	98	*	*	*	*	*	*	*
Rice	237	0.0	7.6	12.2	62.0	17.3	0.8	0.0
Riley	2,981	0.0	1.7	11.4	74.3	12.3	0.2	0.0
Rooks	150	0.0	4.7	14.0	64.0	15.3	1.3	0.7
Rush	30	*	*	*	*	*	*	*
Russell	133	0.0	7.5	15.0	60.2	15.8	1.5	0.0
Saline	1,563	0.3	6.6	15.0	62.3	14.7	1.2	0.1
Scott	135	0.0	3.7	13.3	64.4	18.5	0.0	0.0
Sedgwick	13,808	0.3	6.3	13.4	60.5	18.1	1.3	0.0
Seward	1,407	0.1	8.2	12.7	55.4	22.0	1.4	0.1
Shawnee	4,473	0.4	6.3	14.7	61.1	16.3	1.1	0.0
Sheridan	26	*	*	*	*	*	*	*
Sherman	189	0.0	3.2	18.5	61.4	15.3	1.6	0.0
Smith	53	*	*	*	*	*	*	*
Stafford	100	0.0	5.0	11.0	68.0	14.0	2.0	0.0
Stanton	41	*	*	*	*	*	*	*
Stevens	231	0.0	6.9	15.2	56.7	19.5	1.7	0.0
Sumner	457	0.0	6.1	15.5	61.5	15.5	1.3	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4B
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Age Distribution by County
Combined 3 Years

Comparison of Age Distribution ⁽²⁾								
County	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Thomas	166	0.0	10.2	15.7	60.8	12.0	1.2	0.0
Trego	44	*	*	*	*	*	*	*
Wabaunsee	61	*	*	*	*	*	*	*
Washington	75	*	*	*	*	*	*	*
Wichita	44	*	*	*	*	*	*	*
Wilson	260	0.4	7.7	14.2	65.8	11.5	0.4	0.0
Woodson	40	*	*	*	*	*	*	*
Wyandotte	6,487	0.4	7.8	13.3	57.1	19.8	1.6	0.0
Unknown Counties	48	*	*	*	*	*	*	*
Kansas	65,683	0.3	5.8	13.8	61.5	17.4	1.2	0.1

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5B
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Health Indicators by County
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)														
County	Prepregnancy BMI ⁽⁴⁾			Weight Gain ⁽⁵⁾			Anemia (Low Hb/Hct) ⁽⁶⁾				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Allen	241	6.2 (58)	58.1 (45)	214	19.2 (15)	58.4 (61)	37	* *	67	* *	221	5.4 (16)	223	15.2(58)
Anderson	149	6.0 (54)	49.7 (4)	132	25.0 (54)	48.5 (21)	17	* *	28	* *	137	5.1 (15)	137	11.7(36)
Atchison	427	6.6 (60)	52.2 (11)	372	18.5 (9)	55.4 (55)	57	* *	79	* *	359	5.8 (21)	359	11.7(36)
Barber	137	3.6 (20)	51.8 (9)	109	31.2 (62)	48.6 (22)	16	* *	26	* *	114	0.0 (1)	114	14.0(50)
Barton	752	4.9 (41)	57.0 (43)	652	22.1 (33)	48.8 (24)	75	* *	133	21.1(12)	670	6.4 (27)	668	13.3(47)
Bourbon	349	2.3 (6)	58.7 (49)	308	24.0 (46)	51.3 (37)	42	* *	56	* *	310	9.7 (59)	311	14.1(51)
Brown	274	2.2 (4)	58.4 (46)	233	24.0 (46)	45.5 (11)	31	* *	56	* *	232	6.9 (36)	232	17.2(60)
Butler	791	5.9 (52)	53.6 (22)	711	21.2 (28)	51.9 (41)	118	44.1 (18)	137	30.7(25)	746	6.4 (27)	746	12.5(41)
Chase	40	* *	* *	35	* *	* *	4	* *	2	* *	37	* *	37	* *
Chautauqua	82	* *	* *	75	* *	* *	13	* *	12	* *	76	* *	76	* *
Cherokee	521	7.5 (64)	52.4 (13)	463	18.8 (12)	53.8 (51)	69	* *	105	28.6(22)	492	5.7 (19)	492	12.4(40)
Cheyenne	31	* *	* *	25	* *	* *	6	* *	10	* *	27	* *	27	* *
Clark	10	* *	* *	9	* *	* *	3	* *	1	* *	9	* *	9	* *
Clay	156	5.1 (44)	53.2 (19)	134	21.6 (30)	53.0 (48)	19	* *	39	* *	142	7.0 (38)	142	10.6(27)
Cloud	237	5.5 (48)	52.3 (12)	214	23.8 (45)	50.5 (30)	25	* *	62	* *	216	7.4 (45)	221	10.4(25)
Coffey	80	* *	* *	64	* *	* *	11	* *	18	* *	67	* *	67	* *
Comanche	31	* *	* *	29	* *	* *	5	* *	7	* *	29	* *	29	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5B
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Health Indicators by County
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)														
County	Prepregnancy BMI ⁽⁴⁾			Weight Gain ⁽⁵⁾			Anemia (Low Hb/Hct) ⁽⁶⁾				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Cowley	1,065	3.8 (23)	55.6 (35)	926	21.8 (32)	50.5 (30)	133	33.8 (13)	149	22.1 (13)	947	6.9 (36)	949	11.3 (29)
Crawford	1,088	6.8 (62)	54.0 (24)	935	21.7 (31)	50.3 (29)	145	22.8 (5)	154	15.6 (3)	959	9.1 (56)	962	13.2 (46)
Decatur	40	* *	* *	38	* *	* *	8	* *	9	* *	38	* *	38	* *
Dickinson	398	2.8 (12)	60.1 (55)	342	22.2 (34)	48.8 (24)	59	* *	79	* *	357	5.9 (23)	353	12.5 (41)
Doniphan	125	0.0 (1)	59.2 (50)	101	16.8 (6)	58.4 (61)	20	* *	29	* *	108	6.5 (29)	109	8.3 (13)
Douglas	1,520	3.3 (16)	49.8 (5)	1,276	18.8 (12)	52.7 (46)	199	34.2 (14)	391	27.4 (21)	1,297	8.0 (48)	1,297	8.8 (15)
Edwards	71	* *	* *	65	* *	* *	10	* *	11	* *	64	* *	64	* *
Elk	54	* *	* *	48	* *	* *	6	* *	10	* *	49	* *	49	* *
Ellis	471	4.2 (29)	51.0 (8)	400	18.3 (8)	46.5 (14)	94	* *	105	16.2 (4)	428	2.1 (3)	427	5.9 (2)
Ellsworth	102	3.9 (26)	63.7 (62)	94	* *	* *	15	* *	9	* *	94	* *	94	* *
Finney	2,062	6.7 (61)	52.4 (13)	1,842	25.4 (55)	42.8 (3)	273	26.4 (6)	293	18.1 (7)	1,885	8.3 (51)	1,883	6.9 (6)
Ford	2,033	2.4 (8)	58.6 (48)	1,788	25.4 (55)	45.8 (13)	241	19.1 (2)	222	15.3 (1)	1,790	6.8 (35)	1,786	9.0 (16)
Franklin	535	3.6 (20)	54.2 (25)	465	19.8 (21)	52.9 (47)	60	* *	148	15.5 (2)	404	5.7 (19)	404	11.6 (34)
Geary	1,170	4.4 (37)	53.2 (19)	928	19.4 (18)	52.0 (42)	168	38.7 (17)	211	35.1 (26)	969	4.5 (11)	967	10.2 (23)
Gove	29	* *	* *	24	* *	* *	4	* *	12	* *	30	* *	30	* *
Graham	57	* *	* *	48	* *	* *	3	* *	6	* *	49	* *	49	* *
Grant	205	3.4 (17)	65.4 (65)	189	27.5 (60)	43.4 (5)	27	* *	56	* *	195	6.7 (32)	193	6.7 (4)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5B
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Health Indicators by County
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)														
County	Prepregnancy BMI ⁽⁴⁾			Weight Gain ⁽⁵⁾			Anemia (Low Hb/Hct) ⁽⁶⁾				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Gray	141	5.0 (42)	52.5 (15)	117	23.1 (39)	56.4 (59)	11	* *	30	* *	121	9.1 (56)	121	11.6(34)
Greeley	39	* *	* *	33	* *	* *	3	* *	6	* *	35	* *	35	* *
Greenwood	127	2.4 (8)	64.6 (64)	113	19.5 (20)	54.9 (54)	14	* *	18	* *	113	6.2 (25)	114	17.5(61)
Hamilton	74	* *	* *	68	* *	* *	16	* *	21	* *	69	* *	69	* *
Harper	159	3.1 (14)	56.0 (39)	137	15.3 (4)	49.6 (26)	16	* *	24	* *	139	6.5 (29)	139	9.4(20)
Harvey	799	3.9 (26)	56.6 (41)	703	22.8 (38)	47.9 (19)	109	37.6 (16)	119	20.2(11)	723	4.6 (13)	722	13.0(45)
Haskell	58	* *	* *	51	* *	* *	6	* *	10	* *	52	* *	52	* *
Hodgeman	37	* *	* *	35	* *	* *	6	* *	7	* *	36	* *	36	* *
Jackson	231	4.3 (31)	58.4 (46)	203	19.2 (15)	56.2 (57)	32	* *	53	* *	208	8.7 (54)	209	14.8(56)
Jefferson	179	3.4 (17)	55.3 (31)	159	18.2 (7)	54.7 (53)	30	* *	51	* *	163	6.7 (32)	162	12.3(39)
Jewell	22	* *	* *	21	* *	* *	3	* *	7	* *	21	* *	21	* *
Johnson	5,564	3.6 (20)	54.8 (30)	4,876	20.1 (23)	49.7 (27)	851	22.7 (4)	1,500	17.4 (5)	5,052	9.0 (55)	5,042	7.3(8)
Kearny	56	* *	* *	52	* *	* *	8	* *	8	* *	52	* *	52	* *
Kingman	154	5.2 (45)	49.4 (3)	127	26.0 (58)	47.2 (18)	13	* *	26	* *	130	5.4 (16)	130	14.6(54)
Kiowa	37	* *	* *	33	* *	* *	6	* *	12	* *	35	* *	35	* *
Labette	519	6.0 (54)	55.9 (38)	472	25.8 (57)	45.6 (12)	88	* *	114	23.7(14)	472	7.0 (38)	473	14.6(54)
Lane	38	* *	* *	32	* *	* *	3	* *	6	* *	34	* *	34	* *

- (1) Reporting period is January 1 through December 31.
 - (2) Excludes records with unknown data or errors.
 - (3) Rank compares one county's rate to other counties. Rank 1 = best rate.
 - (4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
 - (5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
 - (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5B
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Health Indicators by County
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)														
County	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	%	%	Number	%	%	Number	%	Number	%	Number	%	Number	%
	(Rank)	(Rank)		(Rank)	(Rank)		(Rank)	(Rank)	(Rank)	(Rank)	(Rank)		(Rank)	
Leavenworth	1,045	4.3 (31)	55.4 (32)	879	19.2 (15)	52.3 (44)	166	27.1 (7)	247	19.0 (8)	909	6.7 (32)	906	11.4(31)
Lincoln	61	* *	* *	53	* *	* *	11	* *	12	* *	55	* *	55	* *
Linn	206	6.3 (59)	60.7 (57)	185	24.3 (50)	48.1 (20)	25	* *	25	* *	185	7.6 (46)	186	7.0(7)
Logan	37	* *	* *	32	* *	* *	6	* *	5	* *	32	* *	32	* *
Lyon	858	2.7 (11)	59.9 (54)	746	24.3 (50)	45.4 (10)	105	15.2 (1)	157	19.7(10)	766	8.0 (48)	765	14.1(51)
Marion	400	5.0 (42)	53.5 (21)	349	23.2 (40)	45.0 (8)	46	* *	99	* *	361	5.8 (21)	364	9.9(21)
Marshall	156	5.8 (49)	50.6 (7)	126	19.8 (21)	46.8 (16)	21	* *	30	* *	137	1.5 (2)	138	6.5(3)
Mcperson	155	5.2 (45)	55.5 (33)	141	27.7 (61)	43.3 (4)	19	* *	40	* *	146	9.6 (58)	145	13.8(48)
Meade	117	1.7 (3)	63.2 (61)	107	18.7 (11)	48.6 (22)	10	* *	20	* *	106	11.3 (62)	105	10.5(26)
Miami	439	4.3 (31)	55.8 (37)	386	14.5 (1)	56.2 (57)	80	* *	113	26.5(20)	411	6.3 (26)	411	11.9(38)
Mitchell	129	3.1 (14)	62.8 (60)	116	35.3 (63)	44.0 (7)	13	* *	24	* *	119	10.1 (60)	119	17.6(62)
Montgomery	981	6.1 (57)	52.8 (17)	869	23.2 (40)	47.0 (17)	193	34.2 (14)	228	24.6(17)	881	8.1 (50)	880	12.8(44)
Morris	81	* *	* *	75	* *	* *	8	* *	22	* *	77	* *	78	* *
Nemaha	112	0.9 (2)	60.7 (57)	100	19.0 (14)	56.0 (56)	22	* *	22	* *	110	4.5 (11)	110	9.1(18)
Neosho	451	5.8 (49)	52.5 (15)	407	22.4 (35)	51.1 (35)	71	* *	70	* *	422	7.6 (46)	421	14.3(53)
Ness	33	* *	* *	28	* *	* *	3	* *	9	* *	29	* *	29	* *
Norton	118	3.4 (17)	48.3 (2)	106	22.6 (37)	43.4 (5)	15	* *	32	* *	108	4.6 (13)	108	14.8(56)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5B
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Health Indicators by County
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)														
County	Prepregnancy BMI ⁽⁴⁾			Weight Gain ⁽⁵⁾			Anemia (Low Hb/Hct) ⁽⁶⁾				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Osage	276	5.8 (49)	51.8 (9)	243	15.2 (3)	56.4 (59)	33	* *	59	* *	253	4.3 (8)	253	11.5(33)
Osborne	56	* *	* *	47	* *	* *	7	* *	12	* *	48	* *	48	* *
Ottawa	67	* *	* *	59	* *	* *	9	* *	9	* *	66	* *	66	* *
Pawnee	164	3.0 (13)	59.8 (52)	133	24.8 (53)	51.1 (35)	18	* *	29	* *	143	11.2 (61)	142	11.3(29)
Phillips	105	3.8 (23)	61.0 (59)	84	* *	* *	15	* *	21	* *	92	* *	92	* *
Pottawatom	94	* *	* *	83	* *	* *	7	* *	35	* *	85	* *	85	* *
Pratt	258	5.4 (47)	56.6 (41)	224	21.4 (29)	51.3 (37)	35	* *	53	* *	232	3.0 (5)	232	8.2(10)
Rawlins	26	* *	* *	22	* *	* *	2	* *	7	* *	22	* *	22	* *
Reno	1,381	4.3 (31)	55.5 (33)	1,209	19.4 (18)	52.4 (45)	170	30.6 (10)	209	26.3(19)	1,236	4.3 (8)	1,236	13.8(48)
Republic	95	* *	* *	77	* *	* *	10	* *	23	* *	81	* *	81	* *
Rice	232	4.7 (40)	63.8 (63)	203	23.2 (40)	50.2 (28)	30	* *	49	* *	211	6.6 (31)	210	9.0(16)
Riley	2,830	4.3 (31)	50.5 (6)	2,290	18.6 (10)	53.4 (49)	461	33.0 (11)	682	25.5(18)	2,400	6.1 (24)	2,405	10.9(28)
Rooks	148	8.1 (65)	45.9 (1)	131	24.4 (52)	39.7 (1)	16	* *	26	* *	128	7.0 (38)	128	10.2(23)
Rush	29	* *	* *	26	* *	* *	3	* *	8	* *	27	* *	27	* *
Russell	130	2.3 (6)	53.8 (23)	118	23.7 (43)	53.4 (49)	24	* *	22	* *	120	3.3 (6)	120	6.7(4)
Saline	1,532	4.5 (38)	55.7 (36)	1,374	16.4 (5)	54.2 (52)	252	22.6 (3)	247	19.0 (8)	1,392	7.2 (43)	1,394	8.2(10)
Scott	134	6.0 (54)	53.0 (18)	120	22.5 (36)	46.7 (15)	21	* *	25	* *	118	2.5 (4)	118	7.6(9)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5B
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Health Indicators by County
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)														
County	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight	Over- weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
							Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)
Sedgwick	13,402	4.1 (28)	54.6 (28)	12,083	20.8 (27)	50.8 (34)	2,584	30.1 (9)	2,990	23.7(14)	12,460	7.1 (42)	12,457	11.4(31)
Seward	1,381	2.5 (10)	59.2 (50)	1,244	24.1 (49)	45.0 (8)	230	27.4 (8)	188	28.7(23)	1,266	8.5 (52)	1,267	5.8(1)
Shawnee	4,356	4.5 (38)	54.4 (27)	3,779	20.6 (25)	51.3 (37)	433	46.9 (19)	680	29.4(24)	3,872	7.2 (43)	3,874	8.2(10)
Sheridan	25	* *	* *	23	* *	* *	4	* *	3	* *	25	* *	25	* *
Sherman	183	3.8 (23)	54.6 (28)	156	26.3 (59)	51.3 (37)	17	* *	34	* *	162	4.3 (8)	162	9.9(21)
Smith	53	* *	* *	46	* *	* *	9	* *	7	* *	45	* *	45	* *
Stafford	99	* *	* *	84	* *	* *	10	* *	13	* *	84	* *	85	* *
Stanton	40	* *	* *	38	* *	* *	3	* *	19	* *	39	* *	38	* *
Stevens	225	2.2 (4)	60.4 (56)	198	20.2 (24)	50.5 (30)	44	* *	61	* *	201	15.9 (63)	202	8.4(14)
Sumner	440	5.9 (52)	54.3 (26)	397	23.7 (43)	41.3 (2)	63	* *	112	17.9 (6)	413	7.0 (38)	414	17.9(63)
Thomas	162	4.3 (31)	58.0 (44)	142	14.8 (2)	59.9 (63)	25	* *	26	* *	147	4.1 (7)	147	15.6(59)
Trego	42	* *	* *	34	* *	* *	8	* *	5	* *	34	* *	36	* *
Wabaunsee	58	* *	* *	56	* *	* *	8	* *	28	* *	59	* *	59	* *
Washington	72	* *	* *	59	* *	* *	9	* *	20	* *	62	* *	62	* *
Wichita	42	* *	* *	38	* *	* *	6	* *	2	* *	40	* *	40	* *
Wilson	256	7.0 (63)	59.8 (52)	242	24.0 (46)	52.1 (43)	42	* *	37	* *	244	8.6 (53)	244	12.7(43)
Woodson	40	* *	* *	38	* *	* *	9	* *	15	* *	38	* *	38	* *

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one county's rate to other counties. Rank 1 = best rate.
- (4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
- (5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
- (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5B
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Health Indicators by County
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)															
County	Prepregnancy BMI (4)				Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy	Hypertension During Pregnancy		
	Under-weight		Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)			Number	% (Rank)
Wyandotte	6,299	4.2 (29)	56.1 (40)		5,579	20.7 (26)	50.7 (33)	971	33.1 (12)	1,154	24.1 (16)	5,666	5.4 (16)	5,663	9.1 (18)
Unknown Counties	46	* *	* *		44	* *	* *	3	* *	6	* *	46	* *	46	* *
Kansas	63,724	4.3	54.9		55,948	21.2	50.1	9,633	29.8	12,815	22.8	57,413	6.9	57,405	10.4

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6B
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Behavioral Indicators by County
Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2, 3)										
County	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Pregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	%	(Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Allen	204	92.6 (2)	2.0	(17)	246	35.4 (52)	246	17.5 (34)	208	48.1 (55)
Anderson	114	85.1 (23)	7.0	(50)	153	48.4 (5)	153	21.6 (24)	138	67.4 (23)
Atchison	376	89.6 (5)	2.7	(23)	441	39.0 (39)	443	11.1 (57)	369	67.8 (21)
Barber	114	81.6 (37)	7.9	(52)	138	46.4 (12)	139	35.3 (6)	117	75.2 (2)
Barton	669	87.4 (14)	0.9	(7)	768	48.3 (6)	771	19.6 (31)	615	59.3 (44)
Bourbon	293	88.4 (10)	1.7	(14)	355	42.0 (29)	355	13.5 (50)	294	59.9 (42)
Brown	194	79.9 (41)	10.8	(57)	277	39.0 (39)	279	15.4 (41)	240	65.8 (28)
Butler	727	75.1 (52)	5.8	(44)	819	36.1 (48)	820	16.5 (38)	629	45.2 (57)
Chase	37	* *	* *		39	* *	41	* *	36	* *
Chautauqua	76	* *	* *		81	* *	83	* *	75	* *
Cherokee	366	79.0 (43)	6.6	(47)	561	33.5 (60)	563	12.4 (54)	435	57.9 (46)
Cheyenne	24	* *	* *		33	* *	33	* *	28	* *
Clark	9	* *	* *		10	* *	10	* *	9	* *
Clay	143	86.0 (20)	3.5	(29)	160	43.8 (22)	160	21.3 (25)	146	68.5 (18)
Cloud	222	84.7 (24)	4.5	(37)	240	47.1 (11)	242	25.6 (14)	207	68.6 (17)
Coffey	66	* *	* *		84	* *	84	* *	71	* *
Comanche	28	* *	* *		31	* *	31	* *	26	* *
Cowley	850	70.2 (57)	6.1	(46)	1,075	42.0 (29)	1,087	14.1 (45)	902	57.9 (46)
Crawford	875	85.5 (21)	4.0	(34)	1,105	42.7 (27)	1,108	13.6 (48)	936	58.3 (45)
Decatur	32	* *	* *		40	* *	40	* *	31	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6B
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Behavioral Indicators by County
Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2, 3)										
County	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Pregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	%	(Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Dickinson	360	85.3 (22)	1.4	(13)	406	41.4 (32)	411	20.9 (26)	331	70.7 (11)
Doniphan	111	82.0 (34)	4.5	(37)	130	33.8 (59)	130	36.2 (5)	96	* *
Douglas	1,262	74.2 (53)	7.0	(50)	1,557	38.2 (42)	1,562	20.2 (29)	1,260	63.9 (33)
Edwards	58	* *	* *		69	* *	71	* *	61	* *
Elk	48	* *	* *		56	* *	56	* *	46	* *
Ellis	420	86.2 (19)	2.1	(19)	490	32.9 (62)	492	26.8 (11)	405	67.2 (25)
Ellsworth	96	* *	* *		104	48.1 (7)	104	23.1 (19)	90	* *
Finney	1,741	53.8 (63)	28.1	(62)	2,088	35.9 (49)	2,105	11.1 (57)	1,737	41.6 (60)
Ford	1,679	72.8 (56)	12.9	(59)	2,044	44.6 (19)	2,057	11.3 (55)	1,790	55.0 (51)
Franklin	235	79.1 (42)	6.0	(45)	557	39.5 (37)	561	8.9 (62)	429	60.6 (40)
Geary	911	74.2 (53)	1.3	(12)	1,218	35.1 (53)	1,220	20.4 (28)	997	59.7 (43)
Gove	30	* *	* *		35	* *	35	* *	23	* *
Graham	49	* *	* *		57	* *	58	* *	49	* *
Grant	174	63.8 (60)	19.5	(61)	205	34.1 (57)	209	10.0 (60)	169	44.4 (58)
Gray	116	87.9 (11)	1.7	(14)	142	47.2 (10)	145	61.4 (2)	127	74.8 (3)
Greeley	29	* *	* *		41	* *	41	* *	31	* *
Greenwood	110	60.9 (62)	30.9	(63)	131	43.5 (24)	131	16.8 (37)	111	74.8 (3)
Hamilton	66	* *	* *		77	* *	77	* *	64	* *
Harper	140	77.1 (46)	10.0	(55)	160	45.6 (14)	165	30.3 (9)	136	73.5 (5)
Harvey	669	76.7 (49)	10.0	(55)	813	47.8 (8)	818	19.6 (31)	685	67.4 (23)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6B
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Behavioral Indicators by County
Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2, 3)										
County	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Pregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	%	(Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Haskell	47	* *	* *	* *	58	* *	59	* *	50	* *
Hodgeman	36	* *	* *	* *	39	* *	39	* *	34	* *
Jackson	115	81.7 (36)	3.5 (29)		239	43.5 (24)	240	12.9 (52)	195	76.4 (1)
Jefferson	158	83.5 (27)	9.5 (54)		183	44.3 (21)	183	9.8 (61)	152	41.4 (61)
Jewell	22	* *	* *	* *	22	* *	22	* *	20	* *
Johnson	4,612	66.4 (58)	3.4 (27)		5,780	33.3 (61)	5,804	14.4 (43)	4,677	61.8 (36)
Kearny	47	* *	* *	* *	56	* *	56	* *	46	* *
Kingman	128	80.5 (39)	4.7 (40)		154	54.5 (3)	155	21.9 (22)	131	70.2 (13)
Kiowa	35	* *	* *	* *	39	* *	39	* *	29	* *
Labette	380	87.6 (12)	3.2 (26)		528	35.6 (51)	530	13.2 (51)	443	64.1 (32)
Lane	34	* *	* *	* *	40	* *	40	* *	33	* *
Leavenworth	913	86.7 (15)	0.8 (3)		1,086	31.2 (63)	1,090	22.1 (20)	840	71.8 (9)
Lincoln	55	* *	* *	* *	64	* *	64	* *	58	* *
Linn	188	80.3 (40)	8.0 (53)		206	47.6 (9)	206	23.3 (17)	181	65.2 (29)
Logan	18	* *	* *	* *	38	* *	38	* *	34	* *
Lyon	751	86.7 (15)	0.8 (3)		883	41.0 (34)	884	18.0 (33)	728	62.4 (35)
Marion	358	86.6 (17)	1.1 (11)		409	37.4 (43)	409	24.9 (15)	334	69.5 (15)
Marshall	133	78.9 (44)	2.3 (21)		160	39.4 (38)	161	39.1 (4)	127	67.7 (22)
Mcperson	146	93.2 (1)	0.0 (1)		160	44.4 (20)	160	45.6 (3)	133	69.9 (14)
Meade	104	77.9 (45)	6.7 (48)		116	56.0 (2)	118	33.9 (7)	105	73.3 (7)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6B
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Behavioral Indicators by County
Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2, 3)										
County	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Pregpregnancy (4)		During Pregpregnancy (5)	
	Number	% (Rank)	%	(Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Miami	406	82.3 (31)	1.0	(10)	455	34.3 (56)	455	14.9 (42)	354	57.9 (46)
Mitchell	118	91.5 (3)	0.8	(3)	132	37.1 (45)	133	23.3 (17)	110	61.8 (36)
Montgomery	675	86.4 (18)	2.7	(23)	1,003	30.2 (64)	1,009	14.1 (45)	849	68.2 (19)
Morris	66	* *	* *		83	* *	85	* *	67	* *
Nemaha	109	89.0 (8)	3.7	(31)	123	35.8 (50)	123	13.8 (47)	109	65.1 (30)
Neosho	387	89.4 (6)	4.1	(36)	464	39.0 (39)	466	12.7 (53)	391	70.3 (12)
Ness	25	* *	* *		34	* *	35	* *	30	* *
Norton	108	90.7 (4)	0.9	(7)	119	46.2 (13)	120	17.5 (34)	102	13.7 (62)
Osage	201	89.1 (7)	2.0	(17)	277	43.7 (23)	282	14.2 (44)	242	42.1 (59)
Osborne	45	* *	* *		57	* *	57	* *	45	* *
Ottawa	65	* *	* *		71	* *	71	* *	60	* *
Pawnee	139	82.0 (34)	2.2	(20)	167	45.5 (15)	169	24.9 (15)	140	62.9 (34)
Phillips	92	* *	* *		107	52.3 (4)	109	22.0 (21)	96	* *
Pottawatom	80	* *	* *		95	* *	95	* *	79	* *
Pratt	231	82.7 (30)	3.9	(33)	264	41.3 (33)	265	7.2 (65)	210	68.1 (20)
Rawlins	11	* *	* *		27	* *	27	* *	23	* *
Reno	1,209	83.4 (28)	2.8	(25)	1,403	40.7 (35)	1,412	20.7 (27)	1,161	60.9 (39)
Republic	81	* *	* *		97	* *	98	* *	84	* *
Rice	209	82.3 (31)	0.5	(2)	233	42.1 (28)	237	16.9 (36)	192	45.3 (56)
Riley	2,237	84.4 (26)	0.8	(3)	2,979	34.0 (58)	2,981	21.9 (22)	2,378	66.4 (27)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6B
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Behavioral Indicators by County
Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2, 3)										
County	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	% (Rank)		Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Rooks	130	84.6 (25)	4.6 (39)		149	45.0 (17)	150	32.7 (8)	121	60.3 (41)
Rush	27	* *	* *		29	* *	30	* *	25	* *
Russell	120	83.3 (29)	2.5 (22)		131	45.0 (17)	133	8.3 (63)	113	70.8 (10)
Saline	1,126	76.4 (50)	4.7 (40)		1,559	35.0 (54)	1,563	6.0 (66)	1,315	57.6 (49)
Scott	118	77.1 (46)	6.8 (49)		132	43.2 (26)	135	84.4 (1)	121	50.4 (53)
Sedgwick	12,117	75.5 (51)	4.8 (42)		13,771	28.6 (65)	13,808	16.4 (40)	11,304	49.8 (54)
Seward	1,111	64.5 (59)	13.8 (60)		1,397	34.4 (55)	1,407	10.2 (59)	1,163	66.7 (26)
Shawnee	3,446	77.0 (48)	3.8 (32)		4,454	45.3 (16)	4,473	11.3 (55)	3,755	52.2 (52)
Sheridan	25	* *	* *		26	* *	26	* *	20	* *
Sherman	120	80.8 (38)	5.0 (43)		189	40.2 (36)	189	25.9 (13)	152	73.0 (8)
Smith	41	* *	* *		52	* *	53	* *	48	* *
Stafford	83	* *	* *		100	58.0 (1)	100	20.0 (30)	85	* *
Stanton	36	* *	* *		41	* *	41	* *	28	* *
Stevens	201	73.6 (55)	4.0 (34)		228	28.1 (66)	231	26.0 (12)	188	69.1 (16)
Sumner	413	82.1 (33)	3.4 (27)		457	37.2 (44)	457	13.6 (48)	371	64.7 (31)
Thomas	105	87.6 (12)	1.9 (16)		166	36.7 (46)	166	28.3 (10)	136	73.5 (5)
Trego	36	* *	* *		44	* *	44	* *	39	* *
Wabaunsee	54	* *	* *		61	* *	61	* *	47	* *
Washington	61	* *	* *		74	* *	75	* *	62	* *
Wichita	40	* *	* *		44	* *	44	* *	35	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6B
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Behavioral Indicators by County
Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2, 3)										
County	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Pregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	%	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Wilson	235	88.9 (9)	0.9	(7)	258	41.5 (31)	260	16.5 (38)	224	61.6 (38)
Woodson	40	* *	*	*	40	* *	40	* *	34	* *
Wyandotte	5,159	63.0 (61)	11.4	(58)	6,442	36.2 (47)	6,487	7.7 (64)	5,470	57.4 (50)
Unknown Counties	37	* *	*	*	48	* *	48	* *	40	* *
Kansas	53,378	75.5	6.0		65,378	36.4	65,683	15.8	54,117	57.7

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7B
2009 - 2011 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by County
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Allen	234	45.3 (45)	184	26.6 (29)	61	* *	204	41.2 (53)	187	35.8 (47)
Anderson	151	47.0 (53)	125	27.2 (33)	50	* *	136	23.5 (35)	127	20.5 (35)
Atchison	329	44.1 (40)	216	29.6 (44)	71	* *	279	29.0 (41)	220	24.1 (38)
Barber	138	47.1 (54)	111	31.5 (51)	45	* *	117	47.9 (59)	112	49.1 (57)
Barton	766	36.4 (21)	668	23.2 (23)	188	28.2 (19)	611	12.6 (19)	666	6.0 (10)
Bourbon	337	46.6 (48)	271	28.0 (39)	101	25.7 (25)	280	40.7 (52)	274	37.6 (49)
Brown	265	43.8 (39)	186	31.2 (50)	65	* *	230	45.2 (56)	191	42.9 (55)
Butler	807	49.6 (55)	716	30.6 (48)	276	31.5 (17)	623	11.1 (14)	723	6.8 (12)
Chase	41	* *	36	* *	19	* *	36	* *	36	* *
Chautauqua	71	* *	50	* *	18	* *	65	* *	50	* *
Cherokee	545	41.7 (34)	480	27.5 (36)	154	23.4 (26)	426	35.9 (48)	483	36.0 (48)
Cheyenne	33	* *	27	* *	6	* *	28	* *	27	* *
Clark	6	* *	6	* *	0	* *	8	* *	8	* *
Clay	160	50.0 (57)	141	29.1 (43)	60	* *	146	14.4 (22)	141	5.7 (9)
Cloud	242	36.4 (21)	219	20.1 (18)	67	* *	207	8.7 (7)	221	2.7 (1)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7B
2009 - 2011 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by County
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Coffey	80	* *	66	* *	27	* *	69	* *	66	* *
Comanche	31	* *	29	* *	12	* *	26	* *	29	* *
Cowley	1,053	40.4 (32)	784	27.6 (37)	259	26.6 (24)	878	20.4 (31)	766	19.6 (33)
Crawford	1,035	44.7 (43)	798	26.6 (29)	283	31.8 (16)	896	40.2 (51)	800	39.8 (52)
Decatur	38	* *	32	* *	6	* *	30	* *	33	* *
Dickinson	406	46.6 (48)	353	26.3 (28)	125	36.0 (9)	326	7.7 (4)	354	5.6 (8)
Doniphan	121	38.0 (26)	89	* *	19	* *	90	* *	90	* *
Douglas	1,532	37.5 (24)	1,249	18.7 (15)	373	42.4 (2)	1,243	10.5 (10)	1,268	6.9 (13)
Edwards	71	* *	62	* *	18	* *	60	* *	63	* *
Elk	52	* *	43	* *	17	* *	43	* *	42	* *
Ellis	489	39.9 (30)	414	18.1 (14)	130	40.8 (3)	404	7.7 (4)	416	5.3 (7)
Ellsworth	103	46.6 (48)	91	* *	32	* *	90	* *	91	* *
Finney	2,082	10.6 (5)	1,815	5.2 (4)	138	39.9 (4)	1,727	13.1 (21)	1,815	12.5 (25)
Ford	1,990	8.8 (2)	1,635	5.0 (3)	103	29.1 (18)	1,747	8.8 (8)	1,639	8.1 (15)
Franklin	558	43.2 (37)	480	29.6 (44)	145	21.4 (27)	427	12.9 (20)	484	8.7 (16)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7B
2009 - 2011 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by County
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Geary	1,200	36.9 (23)	942	20.2 (19)	254	35.0 (11)	990	34.3 (47)	949	28.9 (42)
Gove	33	* *	29	* *	3	* *	22	* *	29	* *
Graham	57	* *	47	* *	22	* *	49	* *	48	* *
Grant	209	10.0 (4)	194	4.1 (2)	13	* *	169	11.8 (15)	194	12.4 (24)
Gray	145	9.7 (3)	116	7.8 (5)	7	* *	126	7.9 (6)	117	5.1 (6)
Greeley	40	* *	33	* *	5	* *	31	* *	33	* *
Greenwood	110	53.6 (61)	57	* *	23	* *	102	31.4 (44)	83	* *
Hamilton	75	* *	68	* *	5	* *	64	* *	68	* *
Harper	153	44.4 (41)	124	27.4 (35)	41	* *	121	26.4 (37)	116	25.0 (40)
Harvey	804	37.9 (25)	692	21.1 (21)	212	37.7 (6)	674	6.1 (2)	692	2.9 (3)
Haskell	59	* *	52	* *	7	* *	50	* *	52	* *
Hodgeman	37	* *	35	* *	15	* *	32	* *	34	* *
Jackson	238	49.6 (55)	201	32.3 (53)	75	* *	193	19.7 (30)	203	19.7 (34)
Jefferson	81	* *	40	* *	14	* *	75	* *	43	* *
Jewell	21	* *	17	* *	8	* *	19	* *	18	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7B
2009 - 2011 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by County
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Johnson	5,734	23.5 (8)	5,016	9.8 (6)	829	46.4 (1)	4,627	18.0 (28)	5,021	17.7 (32)
Kearny	56	* *	50	* *	10	* *	46	* *	50	* *
Kingman	145	46.9 (52)	112	32.1 (52)	38	* *	127	39.4 (50)	116	33.6 (46)
Kiowa	38	* *	30	* *	6	* *	28	* *	33	* *
Labette	513	38.4 (27)	454	28.2 (42)	145	18.6 (29)	434	31.8 (45)	456	32.9 (44)
Lane	18	* *	3	* *	0	* *	17	* *	5	* *
Leavenworth	1,068	46.0 (47)	879	24.5 (26)	287	36.2 (8)	832	37.7 (49)	880	38.4 (51)
Lincoln	63	* *	51	* *	27	* *	58	* *	51	* *
Linn	203	52.7 (60)	182	32.4 (54)	86	* *	179	14.5 (23)	183	10.4 (21)
Logan	37	* *	33	* *	9	* *	34	* *	33	* *
Lyon	871	29.6 (13)	760	15.8 (11)	176	35.8 (10)	725	10.6 (11)	743	6.1 (11)
Marion	408	35.0 (20)	364	20.9 (20)	105	33.3 (13)	331	6.9 (3)	364	2.7 (1)
Marshall	160	23.8 (9)	134	17.2 (13)	26	* *	127	3.9 (1)	134	4.5 (4)
Mcperson	156	44.9 (44)	135	26.7 (32)	49	* *	129	46.5 (58)	136	41.2 (53)
Meade	107	24.3 (10)	96	* *	23	* *	101	24.8 (36)	100	23.0 (37)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7B
2009 - 2011 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by County
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Miami	438	42.9 (35)	380	26.6 (29)	122	27.0 (21)	336	20.8 (32)	380	16.1 (30)
Mitchell	131	33.6 (17)	116	23.3 (24)	34	* *	110	10.9 (13)	117	9.4 (19)
Montgomery	975	40.1 (31)	833	28.0 (39)	267	20.6 (28)	835	33.8 (46)	831	32.9 (44)
Morris	84	* *	74	* *	22	* *	65	* *	74	* *
Nemaha	88	* *	64	* *	24	* *	73	* *	67	* *
Neosho	396	43.4 (38)	231	27.7 (38)	82	* *	346	30.9 (43)	237	21.9 (36)
Ness	30	* *	25	* *	7	* *	26	* *	25	* *
Norton	116	39.7 (29)	94	* *	35	* *	100	27.0 (39)	98	* *
Osage	264	46.6 (48)	217	27.2 (33)	78	* *	225	28.0 (40)	219	26.5 (41)
Osborne	56	* *	42	* *	15	* *	45	* *	42	* *
Ottawa	68	* *	41	* *	17	* *	59	* *	54	* *
Pawnee	164	34.8 (19)	123	19.5 (17)	31	* *	138	18.1 (29)	124	14.5 (28)
Phillips	96	* *	68	* *	18	* *	75	* *	69	* *
Pottawatom	95	* *	81	* *	18	* *	79	* *	84	* *
Pratt	258	39.5 (28)	221	28.1 (41)	74	* *	204	42.6 (54)	214	37.9 (50)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7B
2009 - 2011 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by County
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Rawlins	26	* *	22	* *	6	* *	23	* *	22	* *
Reno	1,391	44.4 (41)	1,198	24.1 (25)	422	38.9 (5)	1,144	14.9 (25)	1,204	12.1 (23)
Republic	97	* *	76	* *	26	* *	81	* *	78	* *
Rice	232	43.1 (36)	204	29.9 (46)	71	* *	189	14.8 (24)	205	10.7 (22)
Riley	2,889	28.2 (12)	2,183	14.0 (9)	463	37.6 (7)	2,346	12.3 (18)	2,208	8.7 (16)
Rooks	143	55.2 (62)	115	36.5 (55)	51	* *	114	29.8 (42)	119	31.9 (43)
Rush	30	* *	27	* *	16	* *	25	* *	27	* *
Russell	132	45.5 (46)	118	39.8 (56)	46	* *	113	54.9 (60)	117	53.0 (58)
Saline	1,522	40.9 (33)	1,305	25.4 (27)	435	26.9 (22)	1,289	11.8 (15)	1,308	9.1 (18)
Scott	125	29.6 (13)	98	* *	26	* *	110	22.7 (34)	100	16.0 (29)
Sedgwick	13,698	29.8 (15)	12,370	16.4 (12)	2,816	34.2 (12)	11,239	9.4 (9)	12,371	6.9 (13)
Seward	1,378	5.3 (1)	1,232	2.7 (1)	46	* *	1,153	12.0 (17)	1,247	9.9 (20)
Shawnee	4,167	34.0 (18)	3,198	21.7 (22)	830	26.9 (22)	3,473	10.7 (12)	3,206	4.9 (5)
Sheridan	24	* *	21	* *	4	* *	20	* *	23	* *
Sherman	185	28.1 (11)	161	14.9 (10)	40	* *	147	22.4 (33)	161	13.7 (26)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7B
2009 - 2011 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by County
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Smith	50	* *	37	* *	13	* *	45	* *	38	* *
Stafford	95	* *	80	* *	24	* *	80	* *	79	* *
Stanton	40	* *	37	* *	3	* *	28	* *	36	* *
Stevens	228	21.5 (7)	199	11.1 (7)	32	* *	186	26.9 (38)	202	24.3 (39)
Sumner	451	52.5 (59)	400	30.8 (49)	167	32.3 (15)	363	43.0 (55)	402	43.5 (56)
Thomas	164	32.3 (16)	144	18.8 (16)	36	* *	132	16.7 (26)	146	16.4 (31)
Trego	42	* *	32	* *	13	* *	37	* *	32	* *
Wabaunsee	61	* *	58	* *	16	* *	47	* *	58	* *
Washington	75	* *	59	* *	12	* *	62	* *	60	* *
Wichita	43	* *	34	* *	7	* *	34	* *	34	* *
Wilson	256	51.2 (58)	238	30.3 (47)	105	33.3 (13)	222	45.9 (57)	237	42.2 (54)
Woodson	37	* *	31	* *	7	* *	32	* *	32	* *
Wyandotte	6,373	19.9 (6)	5,554	11.6 (8)	841	27.8 (20)	5,418	16.7 (26)	5,656	13.9 (27)
Unknown Counties	47	* *	45	* *	19	* *	40	* *	45	* *
Kansas	63,865	30.9	54,208	17.6	12,825	32.5	52,872	16.8	54,494	14.0

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8B
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Infant Health Indicators by County
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)									
County	Birthweight			Preterm ⁽⁶⁾		Full Term LBW ⁽⁷⁾		Ever Breastfed	
	Low ⁽⁴⁾	High ⁽⁵⁾							
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Allen	209	8.6 (45)	2.4 (3)	220	8.2 (24)	191	5.8 (49)	219	62.1 (52)
Anderson	112	10.7 (56)	6.3 (32)	132	9.8 (42)	102	5.9 (50)	131	61.1 (58)
Atchison	365	9.6 (54)	3.8 (7)	386	7.8 (20)	335	6.3 (52)	385	51.9 (62)
Barber	105	7.6 (31)	4.8 (20)	105	11.4 (56)	92	* *	105	80.0 (5)
Barton	639	5.6 (7)	4.1 (11)	642	7.0 (13)	591	4.2 (27)	642	74.8 (22)
Bourbon	300	8.7 (47)	3.0 (4)	304	10.9 (53)	267	5.6 (48)	303	61.7 (53)
Brown	216	8.8 (50)	6.5 (33)	236	11.0 (55)	190	5.3 (47)	236	55.1 (60)
Butler	718	6.8 (21)	6.1 (30)	730	7.7 (19)	661	4.1 (23)	727	70.4 (38)
Chase	36	* *	* *	34	* *	33	* *	36	* *
Chautauqua	70	* *	* *	72	* *	65	* *	74	* *
Cherokee	491	7.7 (32)	8.8 (51)	493	8.1 (23)	449	3.6 (14)	493	58.6 (59)
Cheyenne	24	* *	* *	24	* *	19	* *	24	* *
Clark	9	* *	* *	9	* *	8	* *	9	* *
Clay	138	8.7 (47)	9.4 (55)	139	11.5 (57)	122	1.6 (3)	139	67.6 (42)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8B
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Infant Health Indicators by County
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)									
County	Birthweight			Preterm ⁽⁶⁾		Full Term LBW ⁽⁷⁾		Ever Breastfed	
	Low ⁽⁴⁾	High ⁽⁵⁾		Number	%	Number	%	Number	%
	Number	% ^(Rank)	% ^(Rank)						
Cloud	214	8.4 (43)	4.2 (12)	214	10.3 (47)	190	4.7 (37)	214	78.0 (13)
Coffey	65	* *	* *	65	* *	64	* *	65	* *
Comanche	25	* *	* *	26	* *	22	* *	26	* *
Cowley	891	7.5 (29)	7.1 (41)	929	10.4 (49)	786	4.7 (37)	930	61.4 (55)
Crawford	909	7.8 (34)	4.5 (16)	937	10.5 (51)	808	4.0 (20)	938	61.2 (57)
Decatur	35	* *	* *	35	* *	32	* *	35	* *
Dickinson	345	6.4 (16)	9.3 (54)	340	10.0 (45)	306	3.3 (8)	345	75.4 (21)
Doniphan	107	12.1 (59)	10.3 (57)	108	13.9 (60)	92	* *	107	64.5 (50)
Douglas	1,268	6.8 (21)	6.7 (37)	1,293	7.5 (17)	1,169	4.1 (23)	1,288	79.2 (9)
Edwards	60	* *	* *	60	* *	47	* *	59	* *
Elk	49	* *	* *	51	* *	45	* *	51	* *
Ellis	387	5.9 (10)	8.0 (49)	413	6.5 (10)	358	3.4 (10)	414	77.5 (14)
Ellsworth	92	* *	* *	92	* *	88	* *	92	* *
Finney	1,802	7.0 (25)	5.9 (27)	1,836	9.8 (42)	1,616	4.0 (20)	1,830	77.0 (15)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8B
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Infant Health Indicators by County
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)										
County	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)							
	Number	% (Rank)	%	(Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Ford	1,681	5.6 (7)	5.7	(24)	1,726	8.5 (27)	1,529	2.9 (5)	1,719	75.6 (20)
Franklin	473	5.9 (10)	5.1	(21)	474	9.1 (34)	426	3.8 (17)	477	64.6 (49)
Geary	954	7.8 (34)	7.7	(47)	956	8.3 (25)	873	4.2 (27)	950	72.6 (29)
Gove	29	* *	* *	*	29	* *	27	* *	27	* *
Graham	43	* *	* *	*	45	* *	37	* *	44	* *
Grant	190	6.8 (21)	4.2	(12)	189	5.8 (5)	175	4.0 (20)	192	73.4 (26)
Gray	101	5.0 (4)	10.9	(58)	114	8.8 (32)	92	* *	114	78.1 (12)
Greeley	34	* *	* *	*	34	* *	30	* *	34	* *
Greenwood	110	3.6 (2)	4.5	(16)	111	4.5 (2)	105	3.8 (17)	111	69.4 (41)
Hamilton	68	* *	* *	*	69	* *	63	* *	69	* *
Harper	138	8.0 (38)	10.1	(56)	136	7.4 (16)	123	4.1 (23)	139	74.8 (22)
Harvey	688	7.4 (26)	6.8	(38)	693	8.4 (26)	625	4.5 (33)	695	83.7 (2)
Haskell	47	* *	* *	*	48	* *	43	* *	48	* *
Hodgeman	35	* *	* *	*	35	* *	32	* *	35	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8B
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Infant Health Indicators by County
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)										
County	Birthweight				Preterm ⁽⁶⁾		Full Term LBW ⁽⁷⁾		Ever Breastfed	
	Low ⁽⁴⁾		High ⁽⁵⁾		Number	% ^(Rank)	Number	% ^(Rank)	Number	% ^(Rank)
	Number	% ^(Rank)	% ^(Rank)							
Jackson	201	7.5 (29)	11.9 (59)		208	7.2 (14)	186	4.8 (42)	207	72.9 (27)
Jefferson	159	6.3 (13)	4.4 (15)		160	15.0 (62)	135	3.7 (15)	158	67.1 (43)
Jewell	21	*	*	*	22	*	18	*	22	*
Johnson	4,728	6.8 (21)	7.2 (42)		4,982	6.9 (12)	4,379	3.8 (17)	4,973	81.5 (4)
Kearny	50	*	*	*	51	*	41	*	51	*
Kingman	79	*	*	*	120	5.0 (3)	74	*	121	79.3 (7)
Kiowa	34	*	*	*	35	*	29	*	34	*
Labette	458	6.3 (13)	6.6 (34)		458	10.3 (47)	411	4.4 (31)	456	54.4 (61)
Lane	30	*	*	*	31	*	25	*	31	*
Leavenworth	879	8.1 (40)	7.5 (44)		911	8.6 (30)	801	5.1 (45)	910	64.8 (47)
Lincoln	52	*	*	*	53	*	45	*	52	*
Linn	178	6.7 (20)	3.9 (8)		183	10.9 (53)	158	4.4 (31)	183	76.5 (16)
Logan	33	*	*	*	33	*	31	*	33	*
Lyon	754	8.4 (43)	6.6 (34)		753	9.2 (36)	684	4.7 (37)	749	76.2 (17)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8B
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Infant Health Indicators by County
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)										
County	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)							
	Number	% (Rank)	%	(Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Marion	345	4.3 (3)	4.6	(18)	350	6.0 (7)	324	1.2 (1)	349	74.5 (24)
Marshall	122	7.4 (26)	6.6	(34)	124	7.3 (15)	112	4.5 (33)	124	85.5 (1)
Mcperson	137	8.8 (50)	2.2	(2)	141	9.9 (44)	123	6.5 (53)	140	64.3 (51)
Meade	94	* * *	* *		103	9.7 (41)	82	* *	103	78.6 (11)
Miami	407	6.6 (18)	8.8	(51)	410	7.6 (18)	377	3.7 (15)	407	70.0 (39)
Mitchell	98	* * *	* *		116	4.3 (1)	94	* *	115	75.7 (19)
Montgomery	868	7.8 (34)	3.3	(5)	871	7.9 (22)	795	4.7 (37)	870	61.3 (56)
Morris	66	* * *	* *		73	* *	57	* *	74	* *
Nemaha	73	* * *	* *		109	6.4 (9)	68	* *	104	61.5 (54)
Neosho	330	6.4 (16)	5.2	(22)	416	8.9 (33)	299	3.0 (7)	414	64.7 (48)
Ness	26	* * *	* *		28	* *	23	* *	29	* *
Norton	101	10.9 (57)	4.0	(10)	101	13.9 (60)	87	* *	99	* *
Osage	241	8.7 (47)	3.7	(6)	249	12.9 (59)	205	5.9 (50)	254	65.4 (46)
Osborne	45	* * *	* *		46	* *	43	* *	45	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8B
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Infant Health Indicators by County
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)										
County	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)							
	Number	% (Rank)	%	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Ottawa	63	* * * *	* *	* *	66	* *	59	* *	66	* *
Pawnee	135	7.4 (26)	5.9 (27)		135	10.4 (49)	119	3.4 (10)	136	70.6 (37)
Phillips	87	* * * *	* *	* *	87	* *	78	* *	86	* *
Pottawatom	81	* * * *	* *	* *	81	* *	74	* *	81	* *
Pratt	223	3.1 (1)	9.0 (53)		224	6.7 (11)	207	1.4 (2)	225	71.6 (31)
Rawlins	22	* * * *	* *	* *	22	* *	21	* *	22	* *
Reno	1,201	8.0 (38)	5.7 (24)		1,208	8.5 (27)	1,092	4.9 (43)	1,212	69.7 (40)
Republic	76	* * * *	* *	* *	78	* *	70	* *	79	* *
Rice	197	6.6 (18)	7.6 (46)		196	9.2 (36)	175	2.9 (5)	200	76.0 (18)
Riley	2,373	6.2 (12)	8.0 (49)		2,398	6.1 (8)	2,226	3.5 (13)	2,386	79.3 (7)
Rooks	129	7.8 (34)	4.7 (19)		129	7.8 (20)	119	4.2 (27)	129	72.9 (27)
Rush	26	* * * *	* *	* *	25	* *	23	* *	26	* *
Russell	114	5.3 (5)	5.3 (23)		112	15.2 (63)	95	* *	114	65.8 (45)
Saline	1,337	7.7 (32)	6.2 (31)		1,384	9.5 (40)	1,205	4.3 (30)	1,380	66.4 (44)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8B
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Infant Health Indicators by County
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)										
County	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)							
	Number	% (Rank)	%	(Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Scott	116	8.6 (45)	1.7	(1)	115	8.7 (31)	103	4.9 (43)	117	70.9 (34)
Sedgwick	12,156	9.0 (52)	6.8	(38)	12,213	8.5 (27)	11,100	5.1 (45)	12,187	70.8 (36)
Seward	1,156	5.4 (6)	7.9	(48)	1,215	9.1 (34)	1,040	3.4 (10)	1,216	73.6 (25)
Shawnee	3,765	8.1 (40)	5.7	(24)	3,801	10.0 (45)	3,370	4.6 (35)	3,788	70.9 (34)
Sheridan	25	* * *	* *	*	25	* *	22	* *	25	* *
Sherman	155	9.0 (52)	3.9	(8)	155	5.8 (5)	146	4.1 (23)	155	80.0 (5)
Smith	44	* * *	* *	*	44	* *	39	* *	45	* *
Stafford	78	* * *	* *	*	80	* *	69	* *	77	* *
Stanton	36	* * *	* *	*	38	* *	36	* *	38	* *
Stevens	121	5.8 (9)	7.4	(43)	196	5.1 (4)	112	2.7 (4)	197	78.7 (10)
Sumner	395	11.6 (58)	6.8	(38)	407	12.5 (58)	344	6.7 (54)	406	71.7 (30)
Thomas	141	9.9 (55)	4.3	(14)	141	9.2 (36)	128	4.7 (37)	141	81.6 (3)
Trego	35	* * *	* *	*	37	* *	32	* *	37	* *
Wabaunsee	58	* * *	* *	*	58	* *	57	* *	58	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8B
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Infant Health Indicators by County
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)										
County	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)		Number	%	Number	%	Number	%
	Number	% (Rank)	% (Rank)	Number						
Washington	54	* * * *	* *	57	* *	48	* *	57	* *	
Wichita	38	* * * *	* *	38	* *	34	* *	38	* *	
Wilson	239	6.3 (13)	7.5 (44)	242	10.7 (52)	212	3.3 (8)	241	71.0 (33)	
Woodson	37	* * * *	* *	40	* *	35	* *	40	* *	
Wyandotte	5,426	8.1 (40)	6.0 (29)	5,594	9.2 (36)	4,902	4.6 (35)	5,581	71.4 (32)	
Unknown Counties	47	* * * *	* *	47	* *	47	* *	46	* *	
Kansas	55,067	7.6	6.4	56,409	8.6	50,073	4.4	56,310	72.2	

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

