

Table 3F
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Racial and Ethnic Distribution by Local Agency

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Local Agency	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaska Native	Asian/Pacific Islander	Multiple Races	All Other/Unknown
		%	%	%	%	%	%	%
Barton	452	72.6	2.0	24.8	0.2	0.0	0.4	0.0
Butler	286	92.7	1.0	4.2	0.0	0.0	2.1	0.0
Chautauqua	45	*	*	*	*	*	*	*
Cherokee	715	85.2	5.7	6.2	1.4	0.7	0.8	0.0
Cowley	356	76.1	2.5	18.0	0.8	2.0	0.6	0.0
Crawford	629	86.3	4.0	7.5	0.2	1.0	1.1	0.0
Doniphan	43	*	*	*	*	*	*	*
Douglas	463	65.7	6.0	10.2	10.2	5.2	2.8	0.0
Ford	768	29.0	1.0	69.5	0.1	0.1	0.1	0.0
Franklin	166	89.8	2.4	4.2	1.8	0.0	1.8	0.0
Geary	340	53.5	23.5	15.0	0.0	2.9	5.0	0.0
Graham	35	*	*	*	*	*	*	*
Greeley	10	*	*	*	*	*	*	*
Greenwood	44	*	*	*	*	*	*	*
Harvey	251	69.3	1.6	27.5	0.4	0.0	1.2	0.0
Jefferson	65	*	*	*	*	*	*	*
Johnson	1,844	45.7	12.7	36.3	0.6	2.9	1.7	0.0
Leavenworth	356	66.0	17.4	8.7	0.6	2.5	4.8	0.0
Linn	54	*	*	*	*	*	*	*
Lyon	382	62.3	4.7	31.4	0.3	0.8	0.5	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3F
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Racial and Ethnic Distribution by Local Agency

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Local Agency	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Marion	183	90.2	0.0	8.2	1.1	0.0	0.5	0.0
Marshall	63	*	*	*	*	*	*	*
Miami	136	90.4	2.2	4.4	0.0	1.5	1.5	0.0
NEK	298	81.9	3.4	3.4	8.4	0.0	3.0	0.0
Nemaha	37	*	*	*	*	*	*	*
Neosho	153	91.5	2.6	3.9	0.7	0.7	0.7	0.0
Osborne	314	87.9	2.5	8.0	0.6	1.0	0.0	0.0
Reno	639	82.9	1.9	12.8	0.8	0.6	0.9	0.0
Riley	866	71.9	11.9	11.2	0.9	3.9	0.1	0.0
Saline	819	81.2	2.4	13.3	0.4	1.5	1.2	0.0
Scott	49	*	*	*	*	*	*	*
Sedgwick	4,149	45.0	18.7	29.5	1.5	3.8	1.5	0.0
SEK	50	*	*	*	*	*	*	*
Seward	496	16.1	1.0	77.8	3.4	1.2	0.4	0.0
Shawnee	1,458	57.4	14.3	24.0	1.8	0.8	1.8	0.0
Sheridan	7	*	*	*	*	*	*	*
Sherman	153	81.0	0.7	15.0	0.0	1.3	2.0	0.0
SKCPH	257	81.3	0.8	16.0	0.4	1.2	0.4	0.0
Stevens	66	*	*	*	*	*	*	*
Sumner	158	90.5	0.0	7.6	1.3	0.6	0.0	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3F
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Racial and Ethnic Distribution by Local Agency

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Local Agency	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
SW KS WIC	842	31.0	0.8	65.8	0.0	2.0	0.4	0.0
Wichita	11	*	*	*	*	*	*	*
Wilson	83	*	*	*	*	*	*	*
Wyandotte	2,112	27.3	31.0	39.3	0.2	1.8	0.4	0.0
Kansas	20,703	56.9	11.3	27.3	1.2	2.0	1.2	0.0
Nation (Prior Year)	1,252,728	41.7	23.3	27.9	1.1	3.2	1.0	1.8

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4F
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Age Distribution by Local Agency

Comparison of Age Distribution ⁽²⁾								
Local Agency	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Barton	452	0.4	6.0	17.3	58.6	17.0	0.7	0.0
Butler	286	0.3	7.0	16.1	61.9	14.0	0.7	0.0
Chautauqua	45	*	*	*	*	*	*	*
Cherokee	715	0.1	7.3	18.7	61.7	11.2	0.7	0.3
Cowley	356	0.6	9.0	15.4	61.2	13.5	0.3	0.0
Crawford	629	0.2	7.5	15.3	64.1	12.4	0.5	0.2
Doniphan	43	*	*	*	*	*	*	*
Douglas	463	0.0	4.1	13.6	63.5	17.1	1.7	0.0
Ford	768	0.3	7.2	12.1	55.9	23.4	1.2	0.0
Franklin	166	0.6	7.2	18.1	58.4	15.1	0.0	0.6
Geary	340	0.6	5.6	17.9	65.0	10.3	0.6	0.0
Graham	35	*	*	*	*	*	*	*
Greeley	10	*	*	*	*	*	*	*
Greenwood	44	*	*	*	*	*	*	*
Harvey	251	0.0	5.6	8.0	66.9	16.7	2.8	0.0
Jefferson	65	*	*	*	*	*	*	*
Johnson	1,844	0.3	4.6	11.7	60.7	21.4	1.2	0.1
Leavenworth	356	0.0	5.9	19.4	61.5	11.0	2.2	0.0
Linn	54	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4F
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Age Distribution by Local Agency

Comparison of Age Distribution ⁽²⁾								
Local Agency	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Lyon	382	0.3	8.4	14.9	59.9	13.1	3.4	0.0
Marion	183	0.5	4.4	14.2	63.4	16.4	1.1	0.0
Marshall	63	*	*	*	*	*	*	*
Miami	136	0.0	7.4	11.0	67.6	14.0	0.0	0.0
NEK	298	0.0	7.4	16.4	64.1	11.1	1.0	0.0
Nemaha	37	*	*	*	*	*	*	*
Neosho	153	0.7	3.9	20.9	60.8	13.1	0.7	0.0
Osborne	314	0.3	4.5	15.6	64.0	13.7	1.9	0.0
Reno	639	0.3	5.3	15.3	65.3	12.8	0.9	0.0
Riley	866	0.0	1.7	12.8	73.1	11.7	0.7	0.0
Saline	819	0.0	6.2	17.0	61.1	14.8	1.0	0.0
Scott	49	*	*	*	*	*	*	*
Sedgwick	4,149	0.3	6.7	14.3	61.4	15.9	1.4	0.0
SEK	50	*	*	*	*	*	*	*
Seward	496	0.4	6.7	15.1	57.3	19.2	1.4	0.0
Shawnee	1,458	0.2	6.4	14.2	62.1	15.9	1.1	0.0
Sheridan	7	*	*	*	*	*	*	*
Sherman	153	0.0	6.5	12.4	62.7	15.7	2.6	0.0
SKCPH	257	0.4	7.0	12.8	68.1	11.7	0.0	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4F
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Age Distribution by Local Agency

Comparison of Age Distribution ⁽²⁾								
Local Agency	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Stevens	66	*	*	*	*	*	*	*
Sumner	158	0.0	5.7	17.1	63.3	13.9	0.0	0.0
SW KS WIC	842	0.1	9.6	15.9	55.2	17.5	1.4	0.2
Wichita	11	*	*	*	*	*	*	*
Wilson	83	*	*	*	*	*	*	*
Wyandotte	2,112	0.7	8.6	13.4	58.0	18.1	1.2	0.0
Kansas	20,703	0.3	6.5	14.6	61.6	15.9	1.2	0.0
Nation (Prior Year)	1,252,728	0.4	6.5	12.7	59.9	19.0	1.4	0.1

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5F
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Health Indicators by Local Agency

Comparison of Maternal Health Indicators ^(2, 3)														
Local Agency	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	%	%	Number	%	%	Number	%	Number	%	Number	%	Number	%
	(Rank)	(Rank)		(Rank)	(Rank)		(Rank)		(Rank)		(Rank)		(Rank)	
Barton	446	10.8 (11)	48.7 (22)	413	28.3 (29)	40.7 (2)	45	* *	97	* *	349	3.7 (17)	349	6.3(12)
Butler	283	12.7 (22)	41.3 (2)	269	21.6 (7)	46.8 (19)	32	* *	89	* *	208	2.4 (5)	208	9.6(25)
Chautauqua	45	* *	* *	39	* *	* *	2	* *	7	* *	34	* *	34	* *
Cherokee	711	13.8 (25)	47.0 (16)	644	26.1 (25)	44.1 (10)	102	25.5 (2)	169	24.3 (6)	558	2.9 (12)	558	7.0(16)
Cowley	355	9.9 (5)	49.9 (28)	332	21.7 (9)	45.8 (16)	40	* *	58	* *	286	2.8 (9)	286	7.7(20)
Crawford	625	9.9 (5)	49.8 (27)	567	23.6 (19)	45.5 (14)	95	* *	138	21.0 (4)	506	5.1 (25)	505	8.3(22)
Doniphan	43	* *	* *	35	* *	* *	8	* *	6	* *	28	* *	28	* *
Douglas	459	13.1 (23)	41.8 (3)	409	20.3 (4)	53.1 (28)	67	* *	117	27.4 (8)	347	3.2 (14)	346	4.0(2)
Ford	758	5.8 (1)	49.3 (25)	687	24.3 (23)	45.3 (13)	106	22.6 (1)	113	10.6 (1)	625	2.6 (7)	625	4.8(4)
Franklin	164	11.0 (15)	45.1 (12)	145	16.6 (1)	42.8 (9)	13	* *	32	* *	119	2.5 (6)	119	8.4(23)
Geary	336	11.0 (15)	44.3 (8)	283	21.2 (5)	48.1 (25)	55	* *	50	* *	256	5.1 (25)	256	9.8(26)
Graham	35	* *	* *	31	* *	* *	6	* *	3	* *	31	* *	31	* *
Greeley	10	* *	* *	10	* *	* *	2	* *	2	* *	10	* *	10	* *
Greenwood	44	* *	* *	44	* *	* *	3	* *	7	* *	40	* *	40	* *
Harvey	250	10.8 (11)	44.4 (9)	228	27.2 (27)	41.7 (8)	30	* *	46	* *	192	3.1 (13)	190	6.8(15)
Jefferson	65	* *	* *	60	* *	* *	10	* *	18	* *	53	* *	53	* *
Johnson	1,819	10.8 (11)	44.6 (11)	1,684	21.6 (7)	44.4 (11)	314	25.8 (4)	400	16.5 (2)	1,428	4.0 (20)	1,428	4.3(3)

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.
- (4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).
- (5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).
- (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5F
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Health Indicators by Local Agency

Comparison of Maternal Health Indicators ^(2, 3)															
Local Agency	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy		
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum						
	Number	%	%	Number	%	%	Number	%	Number	%	Number	%	Number	%	Number
	(Rank)	(Rank)		(Rank)	(Rank)		(Rank)	(Rank)	(Rank)	(Rank)	(Rank)	(Rank)	(Rank)	(Rank)	(Rank)
Leavenworth	352	10.5 (8)	48.3 (18)	323	21.7 (9)	49.8 (26)	55	* *	99	* *	253	1.6 (1)	254	5.1(5)	
Linn	52	* *	* *	49	* *	* *	3	* *	10	* *	45	* *	45	* *	
Lyon	379	11.6 (19)	52.0 (29)	347	21.3 (6)	52.4 (27)	53	* *	73	* *	294	2.0 (4)	294	10.9(28)	
Marion	182	9.9 (5)	41.2 (1)	164	23.2 (17)	47.0 (21)	28	* *	36	* *	136	4.4 (22)	136	8.1(21)	
Marshall	62	* *	* *	55	* *	* *	11	* *	13	* *	46	* *	46	* *	
Miami	135	14.1 (26)	48.1 (17)	130	19.2 (2)	53.8 (29)	32	* *	42	* *	104	4.8 (24)	104	5.8(8)	
NEK	296	10.8 (11)	48.6 (21)	272	23.9 (21)	47.4 (22)	47	* *	64	* *	246	1.6 (1)	246	9.8(26)	
Nemaha	36	* *	* *	36	* *	* *	9	* *	16	* *	31	* *	31	* *	
Neosho	151	14.6 (27)	48.3 (18)	142	19.7 (3)	40.1 (1)	23	* *	25	* *	117	1.7 (3)	117	6.0(11)	
Osborne	313	11.8 (20)	44.4 (9)	284	23.2 (17)	40.8 (5)	53	* *	75	* *	235	3.4 (15)	235	7.2(18)	
Reno	638	12.5 (21)	45.9 (15)	600	25.5 (24)	40.7 (2)	96	* *	132	17.4 (3)	496	2.8 (9)	497	7.2(18)	
Riley	860	9.3 (3)	42.9 (5)	736	21.7 (9)	47.7 (23)	113	25.7 (3)	192	32.8(11)	597	3.9 (19)	597	5.9(9)	
Saline	816	10.5 (8)	49.5 (26)	768	22.5 (15)	46.0 (18)	150	26.0 (5)	206	23.3 (5)	673	4.0 (20)	673	5.9(9)	
Scott	48	* *	* *	44	* *	* *	9	* *	9	* *	38	* *	38	* *	
Sedgwick	4,121	11.2 (17)	45.8 (14)	3,907	22.7 (16)	45.6 (15)	822	33.8 (8)	1,156	25.2 (7)	3,347	3.7 (17)	3,346	6.3(12)	
SEK	50	* *	* *	47	* *	* *	9	* *	8	* *	38	* *	38	* *	
Seward	493	8.7 (2)	45.6 (13)	450	22.0 (13)	44.4 (11)	82	* *	81	* *	393	5.3 (27)	391	6.6(14)	

- (1) Reporting period is January 1 through December 31.
 - (2) Excludes records with unknown data or errors.
 - (3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.
 - (4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).
 - (5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).
 - (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5F
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Health Indicators by Local Agency

Comparison of Maternal Health Indicators ^(2, 3)														
Local Agency	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	%	%	Number	%	%	Number	%	Number	%	Number	%	Number	%
	(Rank)	(Rank)		(Rank)	(Rank)		(Rank)	(Rank)	(Rank)	(Rank)	(Rank)		(Rank)	
Shawnee	1,443	10.7 (10)	49.2 (24)	1,310	22.0 (13)	47.7 (23)	152	32.9 (7)	261	29.5 (10)	1,146	5.3 (27)	1,149	5.7 (7)
Sheridan	7	* *	* *	5	* *	* *	3	* *	1	* *	5	* *	5	* *
Sherman	151	15.2 (28)	48.3 (18)	133	24.1 (22)	41.4 (6)	14	* *	38	* *	114	3.5 (16)	114	7.0 (16)
SKCPH	257	15.2 (28)	42.0 (4)	220	27.7 (28)	41.4 (6)	29	* *	64	* *	179	4.5 (23)	178	11.8 (29)
Stevens	66	* *	* *	55	* *	* *	8	* *	17	* *	44	* *	44	* *
Sumner	158	11.4 (18)	43.7 (7)	148	23.6 (19)	45.9 (17)	25	* *	43	* *	117	2.6 (7)	117	9.4 (24)
SW KS WIC	833	13.3 (24)	43.1 (6)	777	26.6 (26)	40.7 (2)	123	37.4 (9)	144	34.7 (12)	696	5.9 (29)	696	3.2 (1)
Wichita	11	* *	* *	11	* *	* *	1	* *	4	* *	10	* *	10	* *
Wilson	81	* *	* *	79	* *	* *	13	* *	14	* *	68	* *	68	* *
Wyandotte	2,077	9.5 (4)	48.9 (23)	1,924	21.9 (12)	46.9 (20)	320	29.7 (6)	358	28.5 (9)	1,688	2.8 (9)	1,689	5.2 (6)
Kansas	20,516	10.9	46.6	18,896	22.9	45.6	3,213	29.3	4,533	24.0	16,226	3.7	16,224	6.2
Nation (Prior Year)	1,175,574	10.9	44.5	922,658	25.0	42.8	180,176	33.5	414,735	29.8	230,547	3.9	280,574	4.3

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.
(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).
(5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6F
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Behavioral Indicators by Local Agency

Comparison of Maternal Behavioral Indicators ^(2, 3)											
Local Agency	Medical Care				WIC Enrollment			Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester			Prepregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	% (Rank)		Number	% (Rank)		Number	% (Rank)	Number	% (Rank)
Barton	406	82.0 (12)	1.7 (8)		359	56.0 (1)		265	20.8 (3)	229	57.6 (10)
Butler	254	70.5 (22)	13.4 (27)		193	48.2 (8)		154	12.3 (13)	117	37.6 (22)
Chautauqua	38	* *	* *		37	* *		21	* *	21	* *
Cherokee	559	83.0 (9)	6.4 (19)		556	45.0 (13)		393	13.0 (10)	357	53.2 (14)
Cowley	314	69.1 (24)	6.4 (19)		269	50.6 (4)		194	10.8 (18)	172	52.3 (16)
Crawford	547	89.8 (1)	0.9 (5)		521	45.1 (12)		381	7.6 (20)	354	61.0 (6)
Doniphan	36	* *	* *		33	* *		27	* *	26	* *
Douglas	403	70.0 (23)	6.9 (21)		362	42.3 (17)		258	12.4 (11)	223	48.0 (18)
Ford	657	74.1 (18)	12.8 (26)		626	48.7 (7)		412	9.2 (19)	390	49.5 (17)
Franklin	95	* *	* *		129	41.1 (18)		91	* *	91	* *
Geary	274	73.0 (21)	0.7 (4)		284	45.8 (11)		195	15.9 (5)	183	47.0 (20)
Graham	32	* *	* *		26	* *		13	* *	14	* *
Greeley	10	* *	* *		7	* *		7	* *	7	* *
Greenwood	40	* *	* *		36	* *		20	* *	21	* *
Harvey	212	73.6 (20)	12.7 (25)		202	48.0 (9)		138	11.6 (16)	128	61.7 (4)
Jefferson	52	* *	* *		48	* *		36	* *	33	* *
Johnson	1,525	62.8 (28)	3.3 (12)		1,406	34.1 (27)		1,022	11.5 (17)	891	60.6 (7)
Leavenworth	320	84.4 (6)	0.9 (5)		244	36.1 (26)		202	12.4 (11)	151	61.6 (5)
Linn	48	* *	* *		43	* *		30	* *	31	* *
Lyon	347	82.4 (10)	0.6 (2)		291	44.7 (14)		216	14.4 (8)	185	56.2 (11)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6F
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Behavioral Indicators by Local Agency

Comparison of Maternal Behavioral Indicators ^(2, 3)														
Local Agency	Medical Care				WIC Enrollment			Multivitamin Consumption						
	1st Trimester		No Care		1st Trimester			Prepregnancy (4)		During Pregnancy (5)				
	Number	% (Rank)	%	% (Rank)	Number	%	% (Rank)	Number	%	% (Rank)				
Marion	160	83.1 (8)	2.5	(10)	144	53.5	(2)	109	14.7	(6)	89	*	*	
Marshall	56	*	*	*	*	53	*	*	33	*	*	25	*	*
Miami	129	79.1 (15)	0.0	(1)	94	*	*	74	*	*	62	*	*	
NEK	242	81.8 (13)	6.2	(18)	253	46.2	(10)	184	5.4	(24)	196	58.7	(8)	
Nemaha	36	*	*	*	*	29	*	*	16	*	*	15	*	*
Neosho	129	89.1 (2)	3.1	(11)	122	38.5	(22)	92	*	*	77	*	*	
Osborne	284	86.6 (3)	1.4	(7)	239	42.7	(16)	184	23.4	(1)	172	67.4	(1)	
Reno	588	84.7 (5)	1.7	(8)	488	39.8	(20)	365	21.9	(2)	314	63.4	(3)	
Riley	668	82.2 (11)	0.6	(2)	632	39.1	(21)	491	14.7	(6)	389	47.8	(19)	
Saline	719	79.8 (14)	3.6	(13)	630	36.3	(25)	451	14.0	(9)	399	53.9	(13)	
Scott	42	*	*	*	*	38	*	*	25	*	*	27	*	*
Sedgwick	3,873	74.0 (19)	3.8	(14)	3,211	28.5	(28)	2,306	11.9	(15)	1,978	42.7	(21)	
SEK	48	*	*	*	*	40	*	*	38	*	*	29	*	*
Seward	427	66.0 (27)	11.5	(24)	398	37.4	(23)	267	6.0	(22)	238	64.3	(2)	
Shawnee	1,254	75.4 (16)	4.4	(16)	1,144	49.9	(5)	846	6.3	(21)	783	56.2	(11)	
Sheridan	5	*	*	*	*	4	*	*	2	*	*	1	*	*
Sherman	102	86.3 (4)	4.9	(17)	119	51.3	(3)	82	*	*	82	*	*	
SKCPH	214	75.2 (17)	10.3	(23)	200	49.5	(6)	160	18.1	(4)	138	52.9	(15)	
Stevens	55	*	*	*	*	49	*	*	38	*	*	29	*	*
Sumner	145	83.4 (7)	4.1	(15)	119	44.5	(15)	84	*	*	68	*	*	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6F
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Behavioral Indicators by Local Agency

Comparison of Maternal Behavioral Indicators ^(2, 3)										
Local Agency	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Pregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	%	(Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
SW KS WIC	708	67.2 (25)	17.8	(28)	675	36.9 (24)	496	6.0 (22)	431	31.1 (23)
Wichita	11	* *	* *		9	* *	6	* *	5	* *
Wilson	78	* *	* *		72	* *	50	* *	47	* *
Wyandotte	1,922	66.7 (26)	7.0	(22)	1,657	40.5 (19)	1,171	12.3 (13)	1,047	57.8 (9)
Kansas	18,064	74.9	5.2		16,091	39.8	11,645	12.0	10,265	53.0
Nation (Prior Year)	612,462	79.4	4.7		1,034,046	31.3	98,170	15.7	238,509	75.9

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7F
2008 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by Local Agency

Comparison of Smoking Indicators (2, 3)										
Local Agency	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Barton	442	40.7 (14)	395	25.8 (20)	128	26.6 (8)	359	12.8 (8)	399	9.0 (7)
Butler	262	52.7 (27)	198	34.3 (27)	78	* *	191	14.7 (11)	196	7.7 (4)
Chautauqua	43	* *	33	* *	8	* *	34	* *	32	* *
Cherokee	680	38.1 (12)	575	26.6 (22)	167	15.0 (12)	540	29.6 (22)	589	28.0 (22)
Cowley	339	44.5 (24)	292	28.8 (24)	100	27.0 (7)	270	21.1 (19)	295	21.0 (19)
Crawford	600	43.0 (20)	487	25.5 (19)	169	33.7 (4)	517	37.3 (25)	487	37.6 (25)
Doniphan	31	* *	22	* *	9	* *	23	* *	24	* *
Douglas	447	44.1 (21)	381	20.7 (10)	118	44.9 (1)	358	10.1 (4)	382	7.3 (3)
Ford	724	10.2 (2)	626	5.3 (2)	43	* *	609	9.5 (3)	620	7.9 (6)
Franklin	163	44.2 (22)	147	25.2 (18)	47	* *	126	11.1 (5)	147	9.5 (8)
Geary	333	41.4 (17)	270	21.1 (12)	98	* *	284	39.8 (26)	273	38.8 (26)
Graham	33	* *	29	* *	11	* *	28	* *	30	* *
Greeley	10	* *	10	* *	3	* *	7	* *	10	* *
Greenwood	41	* *	33	* *	14	* *	36	* *	35	* *
Harvey	247	35.2 (9)	210	24.8 (17)	61	* *	203	5.9 (1)	211	6.2 (1)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7F
2008 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by Local Agency

Comparison of Smoking Indicators (2, 3)										
Local Agency	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Jefferson	16	* *	9	* *	1	* *	16	* *	11	* *
Johnson	1,801	24.6 (5)	1,597	11.5 (4)	277	40.8 (2)	1,400	18.9 (17)	1,606	19.6 (18)
Leavenworth	347	47.0 (25)	297	28.3 (23)	95	* *	243	46.5 (27)	301	40.5 (27)
Linn	53	* *	47	* *	21	* *	43	* *	49	* *
Lyon	375	32.5 (7)	339	17.7 (8)	78	* *	286	14.3 (9)	341	12.6 (12)
Marion	177	39.5 (13)	154	22.1 (13)	49	* *	144	11.8 (6)	156	7.7 (4)
Marshall	63	* *	52	* *	24	* *	53	* *	53	* *
Miami	114	54.4 (29)	98	* *	39	* *	85	* *	97	* *
NEK	248	49.6 (26)	187	29.9 (26)	73	* *	220	30.0 (23)	188	26.6 (21)
Nemaha	37	* *	34	* *	17	* *	29	* *	34	* *
Neosho	144	36.8 (11)	109	17.4 (7)	29	* *	120	18.3 (16)	110	10.9 (11)
Osborne	290	41.4 (17)	238	21.0 (11)	81	* *	228	18.9 (17)	243	17.3 (17)
Reno	621	44.4 (23)	549	26.0 (21)	198	28.8 (6)	486	15.6 (14)	552	14.3 (15)
Riley	823	28.9 (6)	611	18.2 (9)	118	22.0 (11)	625	14.7 (11)	682	13.8 (13)
Saline	746	41.8 (19)	643	24.7 (16)	221	29.4 (5)	595	11.8 (6)	650	9.8 (9)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7F
2008 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by Local Agency

Comparison of Smoking Indicators (2, 3)										
Local Agency	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Scott	37	* *	29	* *	7	* *	33	* *	29	* *
Sedgwick	4,069	33.4 (8)	3,782	16.4 (6)	933	35.4 (3)	3,197	7.1 (2)	3,775	6.9 (2)
SEK	48	* *	43	* *	17	* *	39	* *	43	* *
Seward	482	8.9 (1)	430	3.3 (1)	32	* *	400	15.3 (13)	439	14.1 (14)
Shawnee	1,334	36.3 (10)	1,082	22.5 (14)	288	24.7 (9)	1,101	14.4 (10)	1,082	10.1 (10)
Sheridan	7	* *	5	* *	1	* *	4	* *	5	* *
Sherman	148	41.2 (15)	119	29.4 (25)	36	* *	119	26.1 (21)	118	30.5 (23)
SKCPH	240	41.3 (16)	186	22.6 (15)	57	* *	198	30.8 (24)	197	31.5 (24)
Stevens	66	* *	52	* *	10	* *	49	* *	52	* *
Sumner	150	54.0 (28)	135	34.8 (28)	57	* *	115	53.0 (28)	134	53.7 (28)
SW KS WIC	824	13.1 (3)	767	6.9 (3)	82	* *	679	17.7 (15)	766	16.2 (16)
Wichita	11	* *	9	* *	2	* *	9	* *	9	* *
Wilson	81	* *	75	* *	35	* *	71	* *	75	* *
Wyandotte	2,024	23.2 (4)	1,754	14.2 (5)	300	23.3 (10)	1,646	21.6 (20)	1,789	21.7 (20)
Kansas	19,771	32.9	17,140	18.5	4,232	30.8	15,818	17.6	17,316	16.1

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7F
2008 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by Local Agency

Comparison of Smoking Indicators (2, 3)										
Local Agency	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)		
Nation (Prior Year)	895,227	26.6	716,629	15.8	161,864	35.1	572,120	20.9	486,636	22.6

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8F
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Infant Health Indicators by Local Agency

Comparison of Infant Health Indicators ^(2, 3)										
Local Agency	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)							
	Number	% (Rank)	%	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Barton	390	7.7 (17)	7.2	(24)	326	13.5 (24)	279	2.5 (9)	377	75.1 (8)
Butler	264	9.8 (29)	5.7	(9)	180	14.4 (27)	153	3.3 (17)	240	72.9 (12)
Chautauqua	38	*	*	* *	31	* *	28	* *	34	* *
Cherokee	631	7.8 (19)	6.0	(11)	489	12.9 (23)	424	3.1 (14)	585	60.3 (23)
Cowley	320	6.9 (16)	7.2	(24)	245	11.8 (19)	215	5.1 (25)	311	55.3 (26)
Crawford	548	7.7 (17)	5.8	(10)	464	9.3 (9)	419	3.8 (20)	540	58.5 (24)
Doniphan	37	*	*	* *	27	* *	23	* *	34	* *
Douglas	407	5.4 (5)	7.1	(22)	313	12.1 (20)	273	2.6 (11)	389	75.6 (5)
Ford	653	4.7 (1)	6.1	(14)	557	10.2 (13)	492	1.2 (2)	533	75.6 (5)
Franklin	145	6.2 (12)	5.5	(7)	110	10.9 (16)	98	* *	139	54.0 (27)
Geary	287	5.2 (3)	5.2	(5)	234	7.7 (2)	215	2.3 (7)	261	69.0 (16)
Graham	29	*	*	* *	24	* *	18	* *	30	* *
Greeley	10	*	*	* *	7	* *	7	* *	10	* *
Greenwood	40	*	*	* *	35	* *	29	* *	38	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8F
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Infant Health Indicators by Local Agency

Comparison of Infant Health Indicators ^(2, 3)										
Local Agency	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)							
	Number	% (Rank)	%	(Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Harvey	214	9.3 (26)	5.6	(8)	180	17.2 (28)	149	2.7 (12)	219	78.5 (3)
Jefferson	59	* * *	* *	*	44	* *	39	* *	57	* *
Johnson	1,568	6.7 (15)	7.0	(20)	1,270	8.8 (5)	1,100	2.5 (9)	1,531	78.8 (1)
Leavenworth	299	9.4 (28)	4.0	(2)	216	8.8 (5)	195	4.1 (22)	304	62.2 (22)
Linn	46	* * *	* *	*	39	* *	30	* *	45	* *
Lyon	336	5.7 (9)	6.0	(11)	260	8.5 (3)	236	0.8 (1)	324	71.9 (13)
Marion	155	5.8 (10)	7.7	(27)	126	9.5 (10)	113	3.5 (19)	156	74.4 (9)
Marshall	54	* * *	* *	*	46	* *	42	* *	52	* *
Miami	126	5.6 (7)	6.3	(15)	87	* *	76	* *	98	* *
NEK	253	7.9 (20)	8.3	(28)	232	9.1 (8)	204	3.4 (18)	257	56.0 (25)
Nemaha	33	* * *	* *	*	28	* *	21	* *	36	* *
Neosho	119	5.0 (2)	6.7	(18)	112	11.6 (18)	87	* *	142	53.5 (28)
Osborne	268	9.3 (26)	3.0	(1)	211	10.9 (16)	181	5.0 (24)	258	75.6 (5)
Reno	571	5.4 (5)	7.0	(20)	444	9.0 (7)	399	2.3 (7)	550	62.5 (21)

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.
- (4) <2500 g.
- (5) >4000 g.
- (6) <37 weeks gestation.
- (7) 37 or more weeks gestation and <2500 g.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8F
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Infant Health Indicators by Local Agency

Comparison of Infant Health Indicators ^(2, 3)									
Local Agency	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Riley	733	5.9 (11)	6.5 (17)	511	9.6 (12)	461	2.0 (4)	712	77.9 (4)
Saline	747	7.9 (20)	5.0 (3)	590	13.7 (25)	505	3.2 (15)	729	65.0 (19)
Scott	38	* * *	* *	33	* *	32	* *	37	* *
Sedgwick	3,833	8.1 (23)	7.1 (22)	2,993	9.5 (10)	2,691	3.2 (15)	3,383	67.8 (17)
SEK	46	* * *	* *	37	* *	36	* *	43	* *
Seward	424	5.2 (3)	5.0 (3)	356	10.4 (14)	315	1.3 (3)	432	71.1 (15)
Shawnee	1,271	6.5 (14)	5.4 (6)	1,037	12.4 (22)	896	2.1 (5)	1,248	67.4 (18)
Sheridan	6	* * *	* *	3	* *	2	* *	6	* *
Sherman	133	8.3 (24)	6.0 (11)	103	8.7 (4)	94	* *	123	73.2 (11)
SKCPH	201	8.5 (25)	10.0 (29)	167	13.8 (26)	137	4.4 (23)	206	74.3 (10)
Stevens	32	* * *	* *	38	* *	21	* *	55	* *
Sumner	146	6.2 (12)	7.5 (26)	109	6.4 (1)	102	3.9 (21)	143	65.0 (19)
SW KS WIC	756	5.6 (7)	6.3 (15)	627	10.5 (15)	556	2.2 (6)	411	78.6 (2)
Wichita	11	* * *	* *	9	* *	7	* *	11	* *

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.
- (4) <2500 g.
- (5) >4000 g.
- (6) <37 weeks gestation.
- (7) 37 or more weeks gestation and <2500 g.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8F
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Infant Health Indicators by Local Agency

Comparison of Infant Health Indicators ^(2, 3)									
Local Agency	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Wilson	78	* * *	* *	68	* *	57	* *	64	* *
Wyandotte	1,855	8.0 (22)	6.8 (19)	1,485	12.1 (20)	1,287	2.9 (13)	1,785	71.8 (14)
Kansas	18,210	7.2	6.5	14,503	10.7	12,744	2.8	16,938	69.4
Nation (Prior Year)	1,039,036	8.2	6.7	947,705	11.9	856,467	3.7	936,194	64.6

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3F
2006 - 2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Racial and Ethnic Distribution by Local Agency
Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Local Agency	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Barton	1,395	75.8	1.4	22.2	0.1	0.2	0.2	0.0
Butler	802	91.4	2.0	4.5	0.4	0.1	1.6	0.0
Chautauqua	142	91.5	0.0	6.3	1.4	0.0	0.7	0.0
Cherokee	2,201	84.5	6.7	5.8	1.7	0.7	0.7	0.0
Cowley	1,008	80.6	1.9	14.2	1.0	1.9	0.5	0.0
Crawford	1,882	88.5	3.5	5.8	0.2	1.3	0.7	0.0
Doniphan	123	86.2	4.1	7.3	1.6	0.0	0.8	0.0
Douglas	1,433	64.3	8.4	10.7	9.6	4.7	2.2	0.0
Ford	2,208	29.8	0.7	68.6	0.1	0.6	0.1	0.0
Franklin	507	90.1	2.4	4.3	1.2	0.4	1.6	0.0
Geary	948	54.0	24.3	13.8	0.5	3.7	3.7	0.0
Graham	133	92.5	1.5	5.3	0.0	0.0	0.8	0.0
Greeley	23	*	*	*	*	*	*	*
Greenwood	130	93.8	0.8	3.8	0.0	0.8	0.8	0.0
Harvey	772	69.4	2.1	26.2	0.6	0.0	1.7	0.0
Jefferson	199	95.5	0.0	1.5	2.0	0.5	0.5	0.0
Johnson	5,464	46.7	12.6	36.2	0.5	3.1	1.0	0.0
Leavenwort	1,055	70.0	16.0	8.9	0.7	1.7	2.7	0.0
Linn	179	96.1	0.0	2.8	0.0	0.0	1.1	0.0
Lyon	1,207	58.4	4.9	35.0	0.2	1.0	0.4	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3F
2006 - 2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Racial and Ethnic Distribution by Local Agency
Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Local Agency	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Marion	576	92.4	0.5	5.6	1.0	0.2	0.3	0.0
Marshall	617	40.0	1.5	57.1	0.8	0.2	0.5	0.0
Miami	389	93.1	2.3	3.1	0.0	0.5	1.0	0.0
NEK	926	82.1	5.1	2.8	8.4	0.2	1.4	0.0
Nemaha	107	92.5	0.0	5.6	0.0	0.9	0.9	0.0
Neosho	452	91.4	2.4	4.9	0.4	0.7	0.2	0.0
Osborne	977	90.2	1.8	6.6	0.3	1.0	0.1	0.0
Reno	1,881	84.5	2.6	10.9	0.8	0.7	0.5	0.0
Riley	2,810	76.2	10.5	8.4	0.6	4.1	0.2	0.0
Saline	2,341	81.9	2.7	13.0	0.3	1.3	0.8	0.0
Scott	181	69.6	0.6	27.1	1.7	0.0	1.1	0.0
Sedgwick	12,379	47.5	17.8	28.2	1.6	3.6	1.3	0.0
SEK	159	93.1	0.6	4.4	0.6	1.3	0.0	0.0
Seward	1,033	19.5	1.4	75.4	2.3	1.2	0.3	0.0
Shawnee	4,574	58.0	15.9	21.9	2.1	0.7	1.5	0.0
Sheridan	29	*	*	*	*	*	*	*
Sherman	485	83.3	0.6	13.6	0.6	1.0	0.8	0.0
SKCPH	785	85.4	0.5	12.6	0.8	0.5	0.3	0.0
Stevens	169	45.6	0.0	53.8	0.0	0.6	0.0	0.0
Sumner	427	94.4	0.5	4.2	0.5	0.2	0.2	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3F
2006 - 2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Racial and Ethnic Distribution by Local Agency
Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Local Agency	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
SW KS WIC	2,560	34.5	0.9	62.9	0.0	1.6	0.1	0.0
Wichita	57	*	*	*	*	*	*	*
Wilson	322	95.7	0.3	3.1	0.0	0.3	0.6	0.0
Wyandotte	6,483	25.8	31.6	39.8	0.2	2.3	0.3	0.0
Kansas	62,530	58.3	11.4	26.2	1.2	2.0	0.9	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4F
2006 - 2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Age Distribution by Local Agency
Combined 3 Years

Comparison of Age Distribution ⁽²⁾								
Local Agency	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Barton	1,395	0.3	5.7	16.5	61.1	15.7	0.8	0.0
Butler	802	0.1	6.4	16.5	62.0	14.2	0.9	0.0
Chautauqua	142	0.0	4.9	17.6	63.4	9.9	4.2	0.0
Cherokee	2,201	0.4	7.5	17.5	62.8	11.1	0.5	0.1
Cowley	1,008	0.5	8.3	17.3	59.4	14.0	0.5	0.0
Crawford	1,882	0.3	7.1	16.5	63.8	11.5	0.6	0.2
Doniphan	123	0.0	4.9	17.1	65.0	13.0	0.0	0.0
Douglas	1,433	0.1	4.0	12.9	64.2	17.3	1.4	0.0
Ford	2,208	0.3	6.7	11.7	58.2	21.8	1.1	0.2
Franklin	507	1.0	7.7	16.8	61.5	12.4	0.4	0.2
Geary	948	0.4	5.8	16.2	66.2	10.8	0.5	0.0
Graham	133	0.0	2.3	11.3	72.2	13.5	0.8	0.0
Greeley	23	*	*	*	*	*	*	*
Greenwood	130	0.0	6.9	16.9	63.1	13.1	0.0	0.0
Harvey	772	0.1	6.0	11.4	65.7	14.9	1.9	0.0
Jefferson	199	0.0	10.6	21.6	56.3	10.6	1.0	0.0
Johnson	5,464	0.2	4.3	11.7	62.5	20.1	1.2	0.1
Leavenworth	1,055	0.3	5.8	18.3	63.9	10.6	1.1	0.0
Linn	179	0.0	6.1	16.8	63.7	11.7	1.7	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4F
2006 - 2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Age Distribution by Local Agency
Combined 3 Years

Comparison of Age Distribution ⁽²⁾								
Local Agency	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Lyon	1,207	0.4	6.4	14.1	61.6	15.4	2.0	0.1
Marion	576	0.2	6.3	12.5	64.2	15.6	1.0	0.2
Marshall	617	0.3	7.1	12.5	60.3	18.8	0.6	0.3
Miami	389	0.0	7.7	15.4	65.8	10.5	0.5	0.0
NEK	926	0.0	7.2	16.8	64.5	10.6	0.6	0.2
Nemaha	107	0.0	5.6	15.9	61.7	15.9	0.9	0.0
Neosho	452	0.7	5.8	20.8	61.3	11.1	0.4	0.0
Osborne	977	0.3	4.8	14.4	65.3	13.4	1.6	0.1
Reno	1,881	0.3	4.9	15.0	66.1	13.0	0.6	0.1
Riley	2,810	0.1	2.2	12.0	73.3	11.7	0.6	0.1
Saline	2,341	0.2	6.7	16.3	61.2	14.7	0.9	0.0
Scott	181	0.6	6.1	7.2	69.6	14.4	1.7	0.6
Sedgwick	12,379	0.3	6.8	13.9	61.5	16.0	1.3	0.1
SEK	159	0.0	8.2	12.6	66.7	11.9	0.6	0.0
Seward	1,033	0.5	7.4	13.7	56.3	20.3	1.7	0.0
Shawnee	4,574	0.5	6.3	14.3	63.2	14.8	0.8	0.1
Sheridan	29	*	*	*	*	*	*	*
Sherman	485	0.2	5.4	15.5	61.4	15.3	2.1	0.2
SKCPH	785	0.1	6.9	14.3	65.2	13.4	0.1	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4F
2006 - 2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Age Distribution by Local Agency
Combined 3 Years

Comparison of Age Distribution ⁽²⁾								
Local Agency	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Stevens	169	0.6	5.9	14.8	62.7	15.4	0.6	0.0
Sumner	427	0.2	6.1	16.9	63.5	13.1	0.2	0.0
SW KS WIC	2,560	0.5	9.6	13.8	55.7	19.0	1.3	0.1
Wichita	57	*	*	*	*	*	*	*
Wilson	322	0.0	11.5	13.0	63.0	12.4	0.0	0.0
Wyandotte	6,483	0.7	8.2	14.3	58.5	17.1	1.2	0.1
Kansas	62,530	0.3	6.4	14.3	62.2	15.6	1.1	0.1

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5F
2006 - 2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Health Indicators by Local Agency
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)														
Local Agency	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	%	%	Number	%	%	Number	%	Number	%	Number	%	Number	%
	(Rank)	(Rank)		(Rank)	(Rank)		(Rank)	(Rank)	(Rank)	(Rank)	(Rank)	(Rank)	(Rank)	
Barton	1,353	10.3 (13)	47.5 (32)	1,213	25.8 (37)	43.0 (4)	157	20.4 (3)	238	18.1 (6)	1,200	1.7 (21)	1,200	3.3(24)
Butler	772	13.5 (36)	45.5 (18)	716	20.5 (10)	48.7 (33)	90	* *	212	28.8(23)	686	1.2 (10)	686	5.2(36)
Chautauqua	138	10.1 (12)	53.6 (40)	111	25.2 (36)	43.2 (6)	6	* *	18	* *	120	1.7 (21)	120	1.7(5)
Cherokee	2,133	12.7 (34)	44.8 (12)	1,761	23.6 (29)	46.5 (22)	327	32.4 (16)	520	24.2(15)	1,917	1.2 (10)	1,917	3.1(20)
Cowley	976	11.8 (28)	45.0 (13)	885	23.2 (27)	44.3 (13)	138	31.2 (14)	124	21.8(10)	875	1.4 (14)	875	3.2(22)
Crawford	1,807	12.6 (31)	46.5 (27)	1,563	21.5 (16)	47.0 (25)	259	18.5 (1)	348	19.5 (7)	1,635	2.3 (32)	1,634	3.4(25)
Doniphan	118	5.1 (1)	56.8 (41)	90	* *	* *	25	* *	24	* *	98	* *	98	* *
Douglas	1,378	11.7 (27)	43.1 (7)	1,157	18.8 (6)	51.4 (39)	192	31.8 (15)	351	29.3(25)	1,168	1.7 (21)	1,167	2.0(7)
Ford	2,151	7.0 (3)	45.6 (20)	1,936	24.8 (34)	43.2 (6)	304	22.4 (5)	260	16.9 (4)	1,945	1.2 (10)	1,945	2.3(11)
Franklin	484	11.4 (25)	46.3 (26)	423	17.5 (1)	48.2 (30)	48	* *	115	17.4 (5)	425	0.7 (4)	425	4.7(33)
Geary	916	10.9 (18)	45.4 (16)	771	20.4 (9)	48.5 (32)	150	38.0 (20)	137	28.5(22)	775	2.5 (34)	775	4.6(32)
Graham	127	8.7 (7)	48.0 (34)	108	30.6 (39)	45.4 (14)	17	* *	26	* *	123	0.8 (6)	123	2.4(12)
Greeley	23	* *	* *	22	* *	* *	3	* *	6	* *	22	* *	22	* *
Greenwood	127	7.9 (4)	52.0 (39)	105	21.9 (19)	46.7 (23)	15	* *	27	* *	118	0.8 (6)	118	3.4(25)
Harvey	755	10.7 (15)	47.5 (32)	687	22.6 (23)	46.7 (23)	108	28.7 (11)	126	26.2(18)	678	2.1 (30)	676	2.4(12)
Jefferson	191	14.7 (38)	48.2 (35)	161	18.0 (3)	48.4 (31)	26	* *	51	* *	173	2.9 (36)	173	1.2(3)
Johnson	5,232	10.5 (14)	44.1 (10)	4,534	20.8 (13)	45.8 (17)	896	23.8 (7)	1,423	16.3 (3)	4,686	3.9 (38)	4,686	2.2(10)

- (1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.
(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).
(5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5F
2006 - 2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Health Indicators by Local Agency
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)														
Local Agency	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight	Over- weight	%	< Ideal	> Ideal	3rd Trimester		Postpartum						
						Number	% (Rank)	%	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Leavenworth	1,006	11.0 (20)	46.0 (22)	884	20.9 (14)	50.1 (38)	161	41.0 (22)	210	22.4 (13)	867	0.9 (8)	868	2.0 (7)
Linn	173	11.0 (20)	49.1 (36)	160	28.8 (38)	43.8 (9)	23	* *	33	* *	162	0.6 (3)	162	3.1 (20)
Lyon	1,170	8.5 (5)	50.5 (37)	1,073	22.6 (23)	47.8 (28)	168	19.0 (2)	209	13.9 (2)	1,061	1.4 (14)	1,061	5.6 (37)
Marion	553	10.7 (15)	43.0 (6)	493	23.5 (28)	43.6 (8)	71	* *	115	20.9 (9)	499	1.6 (19)	499	3.2 (22)
Marshall	573	9.9 (9)	46.2 (24)	492	18.7 (4)	44.1 (11)	108	27.8 (10)	60	* *	553	0.4 (2)	553	0.7 (2)
Miami	368	14.7 (38)	45.4 (16)	343	20.1 (8)	49.6 (36)	78	* *	89	* *	334	1.8 (25)	334	1.8 (6)
NEK	893	11.1 (22)	46.8 (29)	713	23.1 (26)	47.7 (26)	154	35.7 (18)	161	25.5 (17)	818	1.1 (9)	818	4.8 (34)
Nemaha	104	8.7 (7)	34.6 (1)	98	* *	* *	25	* *	33	* *	97	* *	97	* *
Neosho	439	15.3 (40)	46.2 (24)	401	17.5 (1)	46.4 (20)	77	* *	68	* *	400	1.3 (13)	400	2.0 (7)
Osborne	946	11.6 (26)	42.1 (5)	819	22.6 (23)	40.3 (2)	155	20.6 (4)	220	13.2 (1)	843	1.8 (25)	843	3.7 (27)
Reno	1,837	12.2 (30)	46.8 (29)	1,661	22.3 (22)	45.4 (14)	282	27.7 (9)	429	22.1 (11)	1,623	1.5 (18)	1,624	2.6 (16)
Riley	2,681	10.0 (11)	42.0 (4)	2,189	18.7 (4)	48.8 (34)	377	22.5 (6)	486	27.2 (20)	2,236	1.6 (19)	2,236	2.5 (14)
Saline	2,284	10.9 (18)	46.0 (22)	2,078	21.5 (16)	46.4 (20)	364	25.3 (8)	508	20.3 (8)	2,064	1.9 (29)	2,064	3.0 (19)
Scott	177	15.8 (41)	41.2 (3)	153	19.0 (7)	49.0 (35)	28	* *	35	* *	158	0.0 (1)	158	0.6 (1)
Sedgwick	11,992	11.2 (24)	45.2 (14)	11,054	20.6 (12)	46.3 (19)	2,467	32.4 (16)	3,230	25.4 (16)	10,966	1.8 (25)	10,965	2.9 (18)
SEK	151	13.9 (37)	41.1 (2)	142	24.6 (32)	43.0 (4)	22	* *	28	* *	144	0.7 (4)	144	6.9 (39)
Seward	1,010	8.6 (6)	45.2 (14)	923	22.0 (20)	45.7 (16)	164	29.9 (12)	176	23.9 (14)	890	2.9 (36)	888	4.4 (31)

- (1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.
(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).
(5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5F
2006 - 2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Health Indicators by Local Agency
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)															
Local Agency	Prepregnancy BMI (4)				Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight		Over-weight		< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Shawnee	4,442	10.7 (15)	45.9 (21)	3,908	21.6 (18)	47.9 (29)	458	39.1 (21)	778	29.0(24)	3,977	2.8 (35)	3,980	2.6(16)	
Sheridan	28	* *	* *	22	* *	* *	3	* *	5	* *	24	* *	24	* *	
Sherman	464	13.4 (35)	46.6 (28)	407	23.8 (30)	41.5 (3)	51	* *	84	* *	418	2.2 (31)	418	5.0(35)	
SKCPH	753	12.6 (31)	44.6 (11)	623	24.7 (33)	43.8 (9)	92	* *	162	22.2(12)	649	1.8 (25)	648	4.3(30)	
Stevens	169	5.3 (2)	51.5 (38)	151	20.5 (10)	49.7 (37)	26	* *	43	* *	142	4.2 (39)	142	5.6(37)	
Sumner	415	12.0 (29)	43.1 (7)	285	21.4 (15)	47.7 (26)	87	* *	93	* *	357	1.7 (21)	357	3.9(29)	
SW KS WIC	2,497	11.1 (22)	45.5 (18)	2,304	25.0 (35)	40.2 (1)	343	30.0 (13)	416	28.4(21)	2,298	2.3 (32)	2,298	1.3(4)	
Wichita	57	* *	* *	54	* *	* *	7	* *	10	* *	53	* *	53	* *	
Wilson	310	12.6 (31)	43.5 (9)	254	24.0 (31)	44.1 (11)	52	* *	48	* *	292	1.4 (14)	292	3.8(28)	
Wyandotte	6,225	9.9 (9)	46.8 (29)	5,748	22.1 (21)	46.1 (18)	1,103	36.0 (19)	1,132	26.4(19)	5,747	1.4 (14)	5,748	2.5(14)	
Kansas	60,398	10.9	45.5	53,675	21.7	46.1	9,707	30.3	12,867	23.4	54,316	1.9	54,314	2.9	

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.
(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).
(5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6F
2006 - 2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Behavioral Indicators by Local Agency
Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2, 3)										
Local Agency	Medical Care			WIC Enrollment			Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)		
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	
Barton	1,231	80.8 (20)	1.4 (14)	1,126	55.4 (2)	265	20.8 (3)	580	83.3 (13)	
Butler	715	76.8 (26)	6.7 (36)	557	44.2 (18)	154	12.3 (13)	317	77.0 (28)	
Chautauqua	121	84.3 (14)	0.8 (9)	113	50.4 (5)	21	* *	58	* *	
Cherokee	1,779	85.0 (13)	3.1 (26)	1,744	44.0 (19)	393	13.0 (10)	1,016	83.6 (12)	
Cowley	882	73.1 (37)	3.6 (27)	797	44.5 (15)	194	10.8 (18)	416	80.3 (23)	
Crawford	1,640	90.1 (2)	0.9 (10)	1,498	45.3 (14)	381	7.6 (20)	878	84.3 (9)	
Doniphan	105	80.0 (22)	4.8 (32)	98	* *	27	* *	66	* *	
Douglas	1,199	74.1 (34)	5.8 (33)	1,122	44.0 (19)	258	12.4 (11)	609	81.0 (18)	
Ford	1,948	75.8 (31)	8.5 (38)	1,831	47.4 (11)	412	9.2 (19)	910	78.4 (25)	
Franklin	386	81.3 (19)	1.6 (16)	346	39.9 (29)	91	* *	161	86.3 (6)	
Geary	752	73.7 (35)	0.4 (3)	769	45.6 (13)	195	15.9 (5)	431	77.5 (27)	
Graham	124	86.3 (9)	0.0 (1)	102	56.9 (1)	13	* *	52	* *	
Greeley	22	* *	* *	16	* *	7	* *	10	* *	
Greenwood	105	73.3 (36)	18.1 (41)	108	51.9 (3)	20	* *	56	* *	
Harvey	665	74.3 (33)	9.8 (40)	633	47.9 (10)	138	11.6 (16)	312	84.3 (9)	
Jefferson	164	87.8 (5)	1.2 (12)	138	31.9 (38)	36	* *	87	* *	
Johnson	4,618	66.4 (41)	2.1 (18)	4,197	32.0 (37)	1,022	11.5 (17)	2,340	85.0 (8)	
Leavenworth	923	85.5 (12)	0.7 (5)	738	39.0 (32)	202	12.4 (11)	429	86.5 (5)	
Linn	162	79.6 (23)	3.7 (28)	146	43.8 (21)	30	* *	82	* *	
Lyon	1,102	82.9 (16)	0.7 (5)	905	44.4 (16)	216	14.4 (8)	504	83.9 (11)	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6F
2006 - 2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Behavioral Indicators by Local Agency
Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2, 3)										
Local Agency	Medical Care			WIC Enrollment		Multivitamin Consumption				
	1st Trimester		No Care	1st Trimester		Pregpregnancy (4)		During Pregpregnancy (5)		
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	
Marion	513	82.5 (17)	2.7 (23)	423	50.6 (4)	109	14.7 (6)	226	82.7 (16)	
Marshall	555	75.5 (32)	0.7 (5)	481	36.4 (34)	33	* *	318	96.5 (1)	
Miami	353	76.8 (26)	0.3 (2)	263	35.4 (36)	74	* *	138	76.8 (29)	
NEK	785	86.5 (8)	3.8 (29)	748	41.0 (26)	184	5.4 (24)	424	80.9 (19)	
Nemaha	102	88.2 (4)	3.9 (30)	80	* *	16	* *	35	* *	
Neosho	397	90.4 (1)	2.0 (17)	369	44.4 (16)	92	* *	181	81.2 (17)	
Osborne	872	86.9 (7)	0.9 (10)	766	46.9 (12)	184	23.4 (1)	475	88.2 (3)	
Reno	1,698	85.9 (11)	1.4 (14)	1,450	42.6 (22)	365	21.9 (2)	868	86.8 (4)	
Riley	1,940	82.4 (18)	0.6 (4)	2,066	39.9 (29)	491	14.7 (6)	1,212	83.3 (13)	
Saline	2,092	80.1 (21)	2.8 (24)	1,791	41.2 (25)	451	14.0 (9)	903	79.6 (24)	
Scott	160	78.8 (24)	6.3 (35)	144	42.4 (23)	25	* *	83	* *	
Sedgwick	11,318	76.7 (28)	2.6 (21)	9,638	28.8 (39)	2,306	11.9 (15)	5,099	77.8 (26)	
SEK	154	87.7 (6)	1.3 (13)	121	49.6 (7)	38	* *	66	* *	
Seward	913	68.2 (40)	7.1 (37)	817	38.1 (33)	267	6.0 (22)	333	74.5 (30)	
Shawnee	3,989	76.5 (29)	2.9 (25)	3,615	48.3 (9)	846	6.3 (21)	1,771	80.6 (22)	
Sheridan	23	* *	* *	21	* *	2	* *	14	* *	
Sherman	380	84.2 (15)	2.1 (18)	375	50.4 (5)	82	* *	246	90.7 (2)	
SKCPH	670	76.3 (30)	6.0 (34)	618	48.9 (8)	160	18.1 (4)	338	80.8 (20)	
Stevens	153	77.8 (25)	2.6 (21)	130	39.2 (31)	38	* *	47	* *	
Sumner	348	86.2 (10)	2.3 (20)	320	40.6 (27)	84	* *	167	86.2 (7)	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6F
2006 - 2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Behavioral Indicators by Local Agency
Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2, 3)									
Local Agency	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Pregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
SW KS WIC	2,271	72.3 (38)	8.7 (39)	2,068	40.0 (28)	496	6.0 (22)	1,108	73.2 (31)
Wichita	54	* *	* *	46	* *	6	* *	30	* *
Wilson	295	88.5 (3)	0.7 (5)	273	41.4 (24)	50	* *	150	80.7 (21)
Wyandotte	5,883	69.3 (39)	4.2 (31)	5,137	35.7 (35)	1,171	12.3 (13)	2,640	83.3 (13)
Kansas	54,561	76.9	3.2	48,744	39.3	11,645	12.0	26,186	81.6

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7F
2006 - 2008 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by Local Agency
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
Local Agency	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Barton	1,366	42.4 (18)	1,211	26.5 (24)	399	28.1 (25)	1,116	21.1 (19)	1,216	12.2 (9)
Butler	762	53.8 (35)	646	33.6 (34)	248	30.6 (19)	539	19.5 (11)	645	9.5 (5)
Chautauqua	140	40.0 (15)	113	29.2 (29)	31	* *	107	43.9 (37)	113	39.8 (35)
Cherokee	2,097	41.1 (17)	1,795	25.8 (22)	530	21.3 (28)	1,700	31.4 (27)	1,855	26.2 (29)
Cowley	976	47.1 (29)	857	31.3 (31)	325	25.5 (27)	780	21.5 (20)	863	17.5 (18)
Crawford	1,809	45.4 (27)	1,548	27.8 (27)	544	30.1 (20)	1,460	36.2 (33)	1,547	33.1 (32)
Doniphan	99	* *	68	* *	29	* *	79	* *	70	* *
Douglas	1,390	44.0 (21)	1,151	20.7 (11)	362	44.2 (3)	1,096	12.6 (4)	1,157	7.1 (2)
Ford	2,126	12.5 (2)	1,792	5.7 (2)	163	38.0 (7)	1,808	10.5 (2)	1,804	8.3 (3)
Franklin	499	46.7 (28)	434	27.2 (25)	128	35.2 (11)	335	20.9 (17)	447	14.1 (10)
Geary	914	43.5 (19)	752	21.3 (12)	268	38.8 (5)	756	39.9 (34)	755	41.1 (36)
Graham	129	51.2 (33)	110	30.0 (30)	44	* *	100	34.0 (31)	112	33.9 (33)
Greeley	23	* *	22	* *	4	* *	16	* *	22	* *
Greenwood	121	47.1 (29)	73	* *	30	* *	107	33.6 (29)	107	22.4 (26)
Harvey	759	35.2 (12)	660	23.5 (16)	188	27.1 (26)	621	6.8 (1)	663	4.8 (1)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7F
2006 - 2008 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by Local Agency
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
Local Agency	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Jefferson	127	54.3 (36)	87	* *	33	* *	102	44.1 (38)	96	* *
Johnson	5,311	26.2 (7)	4,684	11.1 (5)	837	47.2 (2)	4,065	20.2 (14)	4,697	18.7 (20)
Leavenworth	1,028	48.7 (32)	867	27.8 (27)	283	28.6 (23)	738	42.4 (36)	878	42.6 (37)
Linn	177	59.3 (40)	159	44.0 (37)	77	* *	144	20.8 (16)	161	14.9 (12)
Lyon	1,183	32.8 (10)	1,081	17.3 (10)	252	36.1 (9)	889	19.0 (10)	1,085	15.4 (13)
Marion	557	44.0 (21)	497	25.2 (21)	167	31.1 (17)	416	14.7 (5)	504	8.3 (3)
Marshall	609	24.5 (6)	543	13.6 (7)	97	* *	472	30.1 (26)	546	24.9 (28)
Miami	344	54.9 (37)	282	32.3 (33)	106	35.8 (10)	245	27.8 (23)	280	20.0 (21)
NEK	821	44.6 (23)	660	24.4 (18)	213	28.2 (24)	694	34.3 (32)	671	28.9 (30)
Nemaha	106	47.2 (31)	99	* *	36	* *	80	* *	98	* *
Neosho	441	45.1 (26)	382	25.1 (20)	136	33.8 (12)	366	27.9 (25)	383	21.4 (25)
Osborne	937	43.6 (20)	800	23.5 (16)	284	39.4 (4)	746	23.1 (22)	808	21.3 (24)
Reno	1,848	44.9 (25)	1,648	24.6 (19)	566	32.9 (13)	1,429	19.8 (12)	1,657	14.4 (11)
Riley	2,634	29.9 (8)	2,061	16.4 (8)	417	38.1 (6)	2,001	18.2 (8)	2,215	16.5 (16)
Saline	2,204	44.7 (24)	1,877	26.2 (23)	654	30.1 (20)	1,722	17.7 (7)	1,882	11.6 (8)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7F
2006 - 2008 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by Local Agency
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
Local Agency	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Scott	163	30.1 (9)	137	21.9 (13)	28	* *	133	27.8 (23)	140	24.3 (27)
Sedgwick	12,115	33.8 (11)	10,995	16.9 (9)	2,710	37.3 (8)	9,460	12.4 (3)	11,020	10.3 (6)
SEK	153	57.5 (38)	140	33.6 (34)	59	* *	119	33.6 (29)	141	20.6 (22)
Seward	1,000	9.1 (1)	894	4.4 (1)	59	* *	813	16.6 (6)	902	16.5 (16)
Shawnee	4,312	37.5 (13)	3,596	21.9 (13)	967	29.9 (22)	3,478	18.2 (8)	3,591	11.4 (7)
Sheridan	25	* *	18	* *	3	* *	20	* *	18	* *
Sherman	464	40.7 (16)	402	27.4 (26)	120	18.3 (29)	364	20.9 (17)	402	15.4 (13)
SKCPH	730	38.2 (14)	546	22.0 (15)	166	31.9 (15)	585	31.6 (28)	598	29.1 (31)
Stevens	168	20.8 (4)	145	9.0 (4)	23	* *	128	20.3 (15)	146	15.8 (15)
Sumner	388	51.8 (34)	297	31.3 (31)	116	31.0 (18)	306	47.7 (39)	299	49.5 (38)
SW KS WIC	2,511	15.5 (3)	2,313	6.9 (3)	272	49.3 (1)	2,056	19.9 (13)	2,319	17.7 (19)
Wichita	57	* *	52	* *	7	* *	47	* *	52	* *
Wilson	316	57.6 (39)	282	33.7 (36)	137	31.4 (16)	267	42.3 (35)	287	38.7 (34)
Wyandotte	6,269	24.2 (5)	5,632	13.3 (6)	1,006	32.4 (14)	5,036	22.4 (21)	5,688	21.0 (23)
Kansas	60,208	34.1	52,408	18.6	13,124	34.0	47,541	20.5	52,940	17.1

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8F
2006 - 2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Infant Health Indicators by Local Agency
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)									
Local Agency	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Barton	1,201	7.7 (20)	5.0 (7)	990	12.1 (23)	865	3.6 (26)	1,161	70.8 (16)
Butler	721	9.2 (36)	5.8 (15)	504	14.9 (36)	426	3.5 (24)	692	73.4 (13)
Chautauqua	111	9.0 (34)	4.5 (4)	95	* *	73	* *	113	69.9 (19)
Cherokee	1,950	7.9 (24)	5.4 (12)	1,539	12.2 (24)	1,343	3.2 (21)	1,871	57.0 (35)
Cowley	894	7.7 (20)	7.5 (31)	710	11.0 (19)	622	4.5 (31)	869	55.9 (37)
Crawford	1,610	8.4 (29)	5.0 (7)	1,310	12.6 (26)	1,131	4.0 (29)	1,616	58.0 (34)
Doniphan	89	* *	* *	82	* *	57	* *	101	46.5 (41)
Douglas	1,200	6.5 (10)	6.9 (26)	918	11.3 (21)	802	2.6 (11)	1,145	77.6 (4)
Ford	1,885	5.4 (3)	6.1 (18)	1,628	10.6 (14)	1,416	2.2 (5)	1,686	73.5 (12)
Franklin	435	6.9 (16)	5.3 (10)	287	13.2 (31)	246	2.8 (13)	419	53.9 (38)
Geary	797	6.6 (14)	7.0 (29)	628	8.3 (1)	571	2.3 (8)	734	68.5 (20)
Graham	93	* *	* *	89	* *	63	* *	114	78.1 (3)
Greeley	15	* *	* *	15	* *	11	* *	21	* *
Greenwood	115	8.7 (33)	3.5 (1)	99	* *	83	* *	109	62.4 (29)

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.
- (4) <2500 g.
- (5) >4000 g.
- (6) <37 weeks gestation.
- (7) 37 or more weeks gestation and <2500 g.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8F
2006 - 2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Infant Health Indicators by Local Agency
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)									
Local Agency	Birthweight			Preterm ⁽⁶⁾		Full Term LBW ⁽⁷⁾		Ever Breastfed	
	Low ⁽⁴⁾	High ⁽⁵⁾		Number	%	Number	%	Number	%
	Number	% (Rank)	% (Rank)						
Harvey	592	9.1 (35)	5.9 (17)	572	13.6 (33)	419	3.6 (26)	681	78.3 (2)
Jefferson	169	6.5 (10)	8.3 (36)	121	13.2 (31)	104	3.8 (28)	172	57.0 (35)
Johnson	4,646	6.4 (8)	7.7 (32)	3,675	9.0 (4)	3,233	2.5 (10)	4,540	79.4 (1)
Leavenworth	865	8.3 (28)	4.5 (4)	625	9.3 (6)	553	2.9 (17)	893	61.4 (30)
Linn	160	6.9 (16)	5.0 (7)	132	13.6 (33)	113	4.4 (30)	156	60.3 (32)
Lyon	1,090	6.1 (6)	6.7 (22)	812	9.5 (9)	732	2.0 (2)	1,049	71.8 (15)
Marion	496	6.5 (10)	6.3 (20)	373	10.7 (15)	322	2.8 (13)	495	76.6 (6)
Marshall	518	7.7 (20)	7.9 (33)	426	10.1 (12)	364	2.2 (5)	551	67.2 (21)
Miami	352	6.5 (10)	5.7 (13)	233	12.4 (25)	202	4.5 (31)	305	67.2 (21)
NEK	704	7.7 (20)	6.7 (22)	670	11.0 (19)	532	2.8 (13)	765	58.2 (33)
Nemaha	81	* * * *		75	* *	49	* *	101	65.3 (25)
Neosho	336	8.0 (27)	7.1 (30)	341	13.8 (35)	240	3.3 (22)	419	51.6 (39)
Osborne	836	7.4 (18)	4.2 (3)	679	10.9 (17)	583	2.7 (12)	835	75.6 (8)
Reno	1,659	6.4 (8)	8.1 (35)	1,282	10.5 (13)	1,133	2.1 (3)	1,622	66.8 (23)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8F
2006 - 2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Infant Health Indicators by Local Agency
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)										
Local Agency	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)							
	Number	% (Rank)	% (Rank)		Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Riley	2,332	5.6 (4)	8.4 (37)		1,637	9.5 (9)	1,473	2.1 (3)	2,265	77.1 (5)
Saline	2,017	8.5 (30)	5.3 (10)		1,606	12.7 (27)	1,354	3.5 (24)	2,055	64.0 (27)
Scott	128	3.9 (1)	6.3 (20)		126	8.7 (3)	101	3.0 (18)	146	70.5 (17)
Sedgwick	11,211	7.9 (24)	6.8 (25)		8,765	9.3 (6)	7,879	3.1 (20)	10,274	65.8 (24)
SEK	149	6.7 (15)	4.7 (6)		115	11.3 (21)	102	5.9 (35)	144	50.7 (40)
Seward	864	4.9 (2)	6.7 (22)		734	9.3 (6)	637	1.9 (1)	903	72.6 (14)
Shawnee	3,860	7.6 (19)	5.7 (13)		3,204	12.8 (28)	2,680	2.8 (13)	3,911	65.1 (26)
Sheridan	24	* * *	* *		17	* *	15	* *	23	* *
Sherman	429	9.3 (37)	5.8 (15)		330	9.1 (5)	300	5.3 (34)	412	75.5 (9)
SKCPH	573	8.6 (31)	8.0 (34)		523	13.0 (29)	396	3.3 (22)	630	73.8 (11)
Stevens	99	* * *	* *		114	9.6 (11)	64	* *	148	75.7 (7)
Sumner	334	6.0 (5)	6.9 (26)		280	8.6 (2)	224	2.2 (5)	350	62.9 (28)
SW KS WIC	2,304	6.3 (7)	6.2 (19)		1,898	10.7 (15)	1,677	2.4 (9)	1,746	74.7 (10)
Wichita	37	* * *	* *		43	* *	22	* *	53	* *

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.
- (4) <2500 g.
- (5) >4000 g.
- (6) <37 weeks gestation.
- (7) 37 or more weeks gestation and <2500 g.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8F
2006 - 2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Infant Health Indicators by Local Agency
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)									
Local Agency	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Wilson	290	8.6 (31)	4.1 (2)	252	13.1 (30)	215	4.7 (33)	266	60.9 (31)
Wyandotte	5,745	7.9 (24)	6.9 (26)	4,637	10.9 (17)	4,080	3.0 (18)	5,600	70.2 (18)
Kansas	54,016	7.3	6.6	43,191	10.7	37,507	2.9	52,161	68.6

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

