

Table 3F
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Racial and Ethnic Distribution by Local Agency

Comparison of Racial and Ethnic Distribution (2)								
Local Agency	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaska Native	Asian/Pacific Islander	Multiple Races	All Other/Unknown
		%	%	%	%	%	%	%
Barton	451	69.0	1.3	27.9	0.2	0.7	0.9	0.0
Butler	282	88.3	1.1	7.1	0.7	0.7	2.1	0.0
Chautauqua	52	*	*	*	*	*	*	*
Cherokee	750	82.7	5.9	6.3	2.8	1.3	1.1	0.0
Cloud	218	90.8	1.8	4.6	0.9	0.5	1.4	0.0
Cowley	363	73.0	2.8	20.9	0.8	1.4	1.1	0.0
Crawford	609	87.0	2.3	7.4	0.5	1.0	1.8	0.0
Doniphan	37	*	*	*	*	*	*	*
Douglas	507	63.1	10.5	11.6	8.1	5.3	1.4	0.0
Ford	686	21.6	1.0	76.7	0.1	0.6	0.0	0.0
Franklin	187	90.4	0.5	7.0	0.5	0.0	1.6	0.0
Geary	393	49.9	24.9	15.5	0.8	3.6	5.3	0.0
Graham	47	*	*	*	*	*	*	*
Greeley	15	*	*	*	*	*	*	*
Greenwood	49	*	*	*	*	*	*	*
Harvey	284	71.5	2.5	22.2	1.8	0.7	1.4	0.0
Jefferson	68	*	*	*	*	*	*	*
Johnson	1,916	45.5	13.3	35.8	0.6	3.4	1.4	0.0
Leavenworth	359	71.6	13.1	10.6	0.3	0.8	3.6	0.0
Lincoln	145	93.1	0.7	6.2	0.0	0.0	0.0	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3F
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Racial and Ethnic Distribution by Local Agency

Comparison of Racial and Ethnic Distribution (2)								
Local Agency	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Linn	66	*	*	*	*	*	*	*
Lyon	364	58.2	4.7	33.0	0.5	2.2	1.4	0.0
Marion	204	92.2	0.0	4.9	1.0	1.0	1.0	0.0
Marshall	48	*	*	*	*	*	*	*
Meade	109	69.7	0.0	28.4	0.0	0.0	1.8	0.0
Miami	146	92.5	3.4	2.1	0.0	0.7	1.4	0.0
NEK	329	81.2	4.0	3.0	9.4	0.0	2.4	0.0
Nemaha	43	*	*	*	*	*	*	*
Neosho	146	91.1	3.4	4.1	0.0	0.0	1.4	0.0
Osborne	384	89.6	0.5	8.3	1.0	0.5	0.0	0.0
Reno	643	82.4	2.3	13.5	0.3	0.3	1.1	0.0
Riley	991	74.6	10.3	11.2	0.4	2.9	0.6	0.0
Saline	531	68.4	4.9	21.3	0.4	2.6	2.4	0.0
Scott	64	*	*	*	*	*	*	*
Sedgwick	4,721	44.6	18.0	29.8	1.8	4.0	1.8	0.0
SEK	58	*	*	*	*	*	*	*
Seward	532	15.0	1.7	82.1	0.6	0.4	0.2	0.0
Shawnee	1,630	59.0	14.5	22.5	2.0	0.7	1.3	0.0
Sheridan	10	*	*	*	*	*	*	*
Sherman	150	83.3	0.7	15.3	0.0	0.0	0.7	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3F
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Racial and Ethnic Distribution by Local Agency

Comparison of Racial and Ethnic Distribution (2)								
Local Agency	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaska Native	Asian/Pacific Islander	Multiple Races	All Other/Unknown
		%	%	%	%	%	%	%
SKCPH	297	88.6	0.3	10.1	0.7	0.0	0.3	0.0
Stevens	89	*	*	*	*	*	*	*
Sumner	157	88.5	1.3	4.5	1.9	0.0	3.8	0.0
SW KS WIC	873	29.8	0.5	66.9	0.1	2.4	0.3	0.0
Wichita	18	*	*	*	*	*	*	*
Wilson	101	95.0	1.0	3.0	1.0	0.0	0.0	0.0
Wyandotte	2,286	26.2	29.9	40.3	0.3	2.8	0.4	0.0
Unknown Local Agencies	3	*	*	*	*	*	*	*
Kansas	22,411	56.4	11.3	27.6	1.3	2.2	1.3	0.0
Nation (Prior Year)	1,307,887	42.0	24.2	28.1	1.0	3.1	1.1	0.5

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4F
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Age Distribution by Local Agency

Comparison of Age Distribution (2)								
Local Agency	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Barton	451	0.2	8.0	13.5	63.4	13.5	1.3	0.0
Butler	282	0.0	7.8	19.9	59.9	11.7	0.4	0.4
Chautauqua	52	*	*	*	*	*	*	*
Cherokee	750	0.1	5.9	18.1	63.5	11.6	0.5	0.3
Cloud	218	0.5	6.4	18.3	60.6	12.8	1.4	0.0
Cowley	363	0.8	8.0	16.5	61.2	12.7	0.8	0.0
Crawford	609	0.2	7.1	17.2	62.1	12.8	0.5	0.2
Doniphan	37	*	*	*	*	*	*	*
Douglas	507	0.0	3.9	11.4	63.1	20.1	1.4	0.0
Ford	686	0.4	7.9	14.0	53.9	21.7	1.9	0.1
Franklin	187	0.0	5.3	21.9	56.7	14.4	1.6	0.0
Geary	393	0.3	4.8	14.5	67.9	12.2	0.3	0.0
Graham	47	*	*	*	*	*	*	*
Greeley	15	*	*	*	*	*	*	*
Greenwood	49	*	*	*	*	*	*	*
Harvey	284	0.7	7.4	18.3	59.9	13.4	0.4	0.0
Jefferson	68	*	*	*	*	*	*	*
Johnson	1,916	0.2	3.8	10.9	59.5	24.0	1.5	0.1
Leavenwort	359	0.0	6.7	17.3	62.7	12.8	0.6	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4F
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Age Distribution by Local Agency

Comparison of Age Distribution (2)								
Local Agency	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Lincoln	145	0.0	2.1	15.9	62.8	18.6	0.7	0.0
Linn	66	*	*	*	*	*	*	*
Lyon	364	0.5	4.4	13.7	64.3	15.7	1.4	0.0
Marion	204	0.0	8.3	14.7	62.3	13.2	1.5	0.0
Marshall	48	*	*	*	*	*	*	*
Meade	109	0.0	8.3	9.2	57.8	23.9	0.9	0.0
Miami	146	0.7	4.8	17.1	63.0	13.7	0.7	0.0
NEK	329	0.0	6.4	16.4	63.8	13.1	0.3	0.0
Nemaha	43	*	*	*	*	*	*	*
Neosho	146	0.0	9.6	19.2	59.6	11.0	0.7	0.0
Osborne	384	0.0	2.9	13.3	68.8	13.5	1.6	0.0
Reno	643	0.0	6.5	15.4	63.9	13.8	0.3	0.0
Riley	991	0.0	2.1	12.2	73.7	11.7	0.3	0.0
Saline	531	0.2	7.2	14.5	64.6	12.4	1.1	0.0
Scott	64	*	*	*	*	*	*	*
Sedgwick	4,721	0.3	6.9	13.6	60.6	17.2	1.2	0.0
SEK	58	*	*	*	*	*	*	*
Seward	532	0.0	7.7	14.3	54.7	21.8	1.5	0.0
Shawnee	1,630	0.3	6.8	16.4	60.1	15.1	1.2	0.1

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4F
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Age Distribution by Local Agency

Comparison of Age Distribution (2)								
Local Agency	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Sheridan	10	*	*	*	*	*	*	*
Sherman	150	0.0	7.3	16.7	59.3	15.3	1.3	0.0
SKCPH	297	0.0	6.7	17.8	61.6	12.8	1.0	0.0
Stevens	89	*	*	*	*	*	*	*
Sumner	157	0.0	5.7	19.7	60.5	12.7	1.3	0.0
SW KS WIC	873	0.5	8.4	15.5	55.6	18.6	1.6	0.0
Wichita	18	*	*	*	*	*	*	*
Wilson	101	0.0	6.9	19.8	61.4	11.9	0.0	0.0
Wyandotte	2,286	0.4	7.6	13.7	57.9	18.9	1.4	0.0
Unknown Local Agencies	3	*	*	*	*	*	*	*
Kansas	22,411	0.3	6.3	14.6	61.1	16.6	1.1	0.1
Nation (Prior Year)	1,307,887	0.3	6.3	12.6	59.7	19.4	1.4	0.3

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5F
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Maternal Health Indicators by Local Agency

Comparison of Maternal Health Indicators (2, 3)														
Local Agency	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Barton	440	3.9 (7)	57.7 (28)	373	19.8 (19)	49.6 (11)	45	* *	81	* *	389	6.2 (17)	386	11.1(15)
Butler	263	6.8 (29)	51.7 (5)	234	17.9 (7)	49.1 (8)	41	* *	55	* *	255	8.6 (26)	255	12.9(22)
Chautauqua	51	* *	* *	50	* *	* *	5	* *	8	* *	51	* *	51	* *
Cherokee	719	7.1 (32)	56.2 (23)	641	20.3 (22)	49.1 (8)	116	31.9 (5)	175	25.1 (7)	661	5.4 (10)	662	11.3(18)
Cloud	214	6.1 (25)	52.3 (9)	184	23.9 (28)	50.0 (13)	23	* *	62	* *	188	4.3 (4)	191	13.1(24)
Cowley	356	4.8 (18)	52.8 (10)	313	21.4 (23)	53.4 (23)	52	* *	45	* *	316	5.7 (12)	317	12.0(21)
Crawford	592	6.4 (27)	55.2 (19)	515	19.0 (12)	56.3 (28)	90	* *	99	* *	531	7.3 (22)	531	13.9(27)
Doniphan	35	* *	* *	31	* *	* *	5	* *	9	* *	33	* *	33	* *
Douglas	491	3.7 (6)	48.1 (1)	421	16.9 (3)	55.1 (26)	70	* *	137	21.2 (5)	432	6.7 (19)	432	9.0(6)
Ford	677	2.5 (5)	55.5 (20)	604	22.8 (25)	45.5 (3)	80	* *	61	* *	601	6.3 (18)	600	9.8(8)
Franklin	180	2.2 (4)	55.6 (21)	160	18.1 (8)	56.9 (30)	19	* *	56	* *	128	4.7 (7)	128	10.9(14)
Geary	378	5.3 (23)	51.1 (4)	301	15.9 (1)	52.5 (20)	55	* *	83	* *	314	3.5 (3)	314	10.5(11)
Graham	42	* *	* *	36	* *	* *	5	* *	9	* *	41	* *	41	* *
Greeley	13	* *	* *	12	* *	* *	1	* *	2	* *	14	* *	14	* *
Greenwood	47	* *	* *	43	* *	* *	5	* *	5	* *	44	* *	45	* *
Harvey	277	4.7 (17)	53.1 (11)	241	26.6 (31)	45.2 (2)	40	* *	37	* *	246	6.1 (15)	246	11.8(19)
Jefferson	67	* *	* *	61	* *	* *	11	* *	20	* *	62	* *	62	* *

- (1) Reporting period is January 1 through December 31.
 - (2) Excludes records with unknown data or errors.
 - (3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.
 - (4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
 - (5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
 - (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5F
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Maternal Health Indicators by Local Agency

Comparison of Maternal Health Indicators (2, 3)														
Local Agency	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Johnson	1,812	4.5 (14)	54.8 (18)	1,609	19.4 (15)	51.0 (14)	307	24.1 (2)	592	17.9 (4)	1,686	8.3 (25)	1,683	7.3(4)
Leavenwort	347	4.6 (16)	57.3 (25)	297	19.2 (13)	51.5 (16)	61	* *	75	* *	303	8.9 (28)	302	9.9(9)
Lincoln	139	4.3 (12)	62.6 (33)	130	24.6 (30)	53.8 (24)	16	* *	29	* *	134	9.0 (29)	134	11.9(20)
Linn	66	* *	* *	62	* *	* *	10	* *	10	* *	61	* *	61	* *
Lyon	344	1.7 (1)	62.2 (32)	299	23.1 (26)	45.5 (3)	47	* *	60	* *	316	7.0 (20)	316	13.6(26)
Marion	194	6.2 (26)	52.1 (8)	172	18.6 (11)	54.7 (25)	21	* *	55	* *	184	4.3 (4)	180	10.6(12)
Marshall	47	* *	* *	41	* *	* *	5	* *	10	* *	42	* *	42	* *
Meade	108	1.9 (2)	57.4 (27)	95	* *	* *	10	* *	17	* *	96	* *	96	* *
Miami	137	6.6 (28)	57.7 (28)	120	18.3 (9)	49.2 (10)	22	* *	44	* *	131	9.2 (31)	131	13.0(23)
NEK	314	5.1 (21)	57.3 (25)	282	17.4 (5)	56.7 (29)	46	* *	74	* *	285	6.0 (13)	284	15.5(30)
Nemaha	37	* *	* *	34	* *	* *	8	* *	10	* *	40	* *	40	* *
Neosho	140	8.6 (33)	49.3 (3)	128	21.9 (24)	52.3 (19)	22	* *	19	* *	133	9.0 (29)	133	14.3(28)
Osborne	368	4.1 (8)	51.9 (7)	317	19.2 (13)	47.9 (7)	66	* *	85	* *	335	4.8 (8)	335	10.7(13)
Reno	631	5.1 (21)	54.5 (17)	552	19.7 (18)	49.8 (12)	72	* *	99	* *	566	3.2 (1)	566	15.0(29)
Riley	935	4.9 (19)	48.3 (2)	750	16.4 (2)	55.5 (27)	161	21.1 (1)	226	17.7 (3)	800	4.3 (4)	799	10.4(10)
Saline	516	4.1 (8)	53.9 (15)	473	16.9 (3)	53.1 (22)	89	* *	105	14.3 (1)	482	5.6 (11)	486	6.2(1)
Scott	63	* *	* *	57	* *	* *	8	* *	8	* *	57	* *	57	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5F
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Maternal Health Indicators by Local Agency

Comparison of Maternal Health Indicators (2, 3)														
Local Agency	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Sedgwick	4,574	4.4 (13)	53.5 (13)	4,150	20.1 (20)	51.6 (17)	914	31.4 (4)	988	25.3 (8)	4,278	6.1 (15)	4,281	11.2(16)
SEK	57	* *	* *	51	* *	* *	9	* *	10	* *	53	* *	53	* *
Seward	518	1.9 (2)	58.7 (31)	468	24.1 (29)	46.2 (5)	93	* *	64	* *	478	8.8 (27)	482	6.4(2)
Shawnee	1,579	4.5 (14)	53.8 (14)	1,379	19.6 (17)	52.2 (18)	164	42.1 (7)	264	31.8 (9)	1,413	7.2 (21)	1,415	7.8(5)
Sheridan	10	* *	* *	10	* *	* *	2	* *	2	* *	10	* *	10	* *
Sherman	146	5.5 (24)	58.2 (30)	130	18.5 (10)	58.5 (31)	19	* *	22	* *	133	6.0 (13)	133	13.5(25)
SKCPH	288	4.9 (19)	53.1 (11)	243	17.7 (6)	52.7 (21)	39	* *	52	* *	251	3.2 (1)	251	11.2(16)
Stevens	86	* *	* *	76	* *	* *	14	* *	26	* *	79	* *	79	* *
Sumner	147	4.1 (8)	54.4 (16)	134	20.1 (20)	41.0 (1)	20	* *	37	* *	144	7.6 (23)	144	16.7(31)
SW KS WIC	851	6.8 (29)	51.7 (5)	766	23.1 (26)	46.9 (6)	115	27.0 (3)	154	17.5 (2)	786	7.9 (24)	782	7.0(3)
Wichita	18	* *	* *	15	* *	* *	3	* *	2	* *	15	* *	15	* *
Wilson	101	6.9 (31)	56.4 (24)	93	* *	* *	17	* *	14	* *	93	* *	93	* *
Wyandotte	2,228	4.2 (11)	55.7 (22)	1,973	19.5 (16)	51.2 (15)	351	31.9 (5)	393	22.4 (6)	2,004	5.2 (9)	2,005	9.5(7)
Unknown Local Agencies	3	* *	* *	3	* *	* *	0	* *	1	* *	3	* *	3	* *
Kansas	21,646	4.6	54.3	19,129	19.9	51.1	3,394	29.5	4,491	22.7	19,697	6.4	19,699	10.3
Nation (Prior Year)	1,224,766	4.7	52.1	963,624	21.4	48.2	188,530	33.8	416,710	29.6	346,704	4.4	395,960	5.2

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6F
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Maternal Behavioral Indicators by Local Agency

Comparison of Maternal Behavioral Indicators (2, 3)											
Local Agency	Medical Care				WIC Enrollment			Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester			Prepregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	% (Rank)		Number	% (Rank)		Number	% (Rank)	Number	% (Rank)
Barton	383	82.5 (14)	1.8 (8)		447	48.8 (1)		451	20.0 (10)	374	51.1 (27)
Butler	241	64.7 (27)	13.3 (26)		282	38.3 (16)		282	16.7 (14)	207	31.9 (31)
Chautauqua	50	* *	* *		51	* *		52	* *	46	* *
Cherokee	535	82.8 (12)	5.2 (18)		745	33.6 (24)		750	14.9 (19)	609	59.8 (18)
Cloud	191	85.9 (6)	3.1 (11)		217	48.4 (2)		218	20.2 (9)	188	68.6 (7)
Cowley	283	66.1 (25)	6.0 (21)		358	42.2 (11)		363	12.9 (24)	307	58.0 (20)
Crawford	503	85.5 (7)	3.2 (13)		609	42.2 (11)		609	13.8 (23)	537	56.8 (22)
Doniphan	33	* *	* *		37	* *		37	* *	25	* *
Douglas	428	75.7 (18)	4.7 (16)		507	33.1 (28)		507	20.5 (7)	397	63.0 (12)
Ford	584	70.0 (22)	16.4 (28)		682	43.0 (8)		686	7.4 (31)	611	52.7 (25)
Franklin	32	* *	* *		184	42.9 (9)		187	4.3 (33)	152	53.3 (24)
Geary	285	67.0 (23)	1.4 (5)		392	33.2 (27)		393	18.6 (12)	321	60.7 (15)
Graham	39	* *	* *		47	* *		47	* *	37	* *
Greeley	14	* *	* *		15	* *		15	* *	10	* *
Greenwood	42	* *	* *		49	* *		49	* *	40	* *
Harvey	225	74.7 (20)	14.7 (27)		281	47.0 (4)		284	16.2 (16)	238	68.9 (5)
Jefferson	60	* *	* *		68	* *		68	* *	54	* *
Johnson	1,518	66.5 (24)	3.8 (14)		1,905	30.6 (32)		1,916	14.0 (22)	1,527	64.2 (9)
Leavenworth	304	88.5 (3)	0.0 (1)		357	34.7 (22)		359	22.6 (5)	275	72.4 (2)
Lincoln	133	92.5 (1)	0.8 (4)		144	37.5 (18)		145	24.8 (4)	120	60.8 (14)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6F
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Maternal Behavioral Indicators by Local Agency

Comparison of Maternal Behavioral Indicators (2, 3)										
Local Agency	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Linn	62	* *	* *	* *	66	* *	66	* *	57	* *
Lyon	313	87.9 (4)	0.6 (3)		363	39.1 (14)	364	15.1 (18)	294	63.3 (11)
Marion	180	81.7 (15)	1.7 (7)		203	42.9 (9)	204	22.5 (6)	152	71.7 (3)
Marshall	42	* *	* *	* *	48	* *	48	* *	40	* *
Meade	92	* *	* *	* *	107	43.9 (7)	109	44.0 (1)	91	* *
Miami	129	85.3 (9)	1.6 (6)		146	30.8 (31)	146	14.4 (20)	107	54.2 (23)
NEK	260	83.5 (11)	5.8 (20)		327	33.3 (26)	329	11.2 (26)	274	69.3 (4)
Nemaha	40	* *	* *	* *	43	* *	43	* *	39	* *
Neosho	124	88.7 (2)	6.5 (23)		146	37.7 (17)	146	8.9 (29)	127	68.5 (8)
Osborne	325	86.8 (5)	2.5 (10)		382	39.0 (15)	384	27.6 (3)	311	62.4 (13)
Reno	550	84.9 (10)	1.8 (8)		639	39.6 (13)	643	20.5 (7)	535	60.7 (15)
Riley	746	85.4 (8)	0.5 (2)		990	33.1 (28)	991	16.5 (15)	793	59.1 (19)
Saline	419	74.0 (21)	5.7 (19)		529	32.9 (30)	531	4.9 (32)	431	51.5 (26)
Scott	56	* *	* *	* *	62	* *	64	* *	55	* *
Sedgwick	4,225	75.4 (19)	4.1 (15)		4,705	28.0 (33)	4,721	14.4 (20)	3,847	47.1 (29)
SEK	44	* *	* *	* *	58	* *	58	* *	51	* *
Seward	420	64.0 (28)	10.2 (25)		528	34.8 (21)	532	10.2 (28)	438	63.5 (10)
Shawnee	1,261	79.3 (16)	3.1 (11)		1,617	47.2 (3)	1,630	8.5 (30)	1,373	49.2 (28)
Sheridan	10	* *	* *	* *	10	* *	10	* *	8	* *
Sherman	39	* *	* *	* *	150	34.7 (22)	150	28.7 (2)	126	77.8 (1)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6F
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Maternal Behavioral Indicators by Local Agency

Comparison of Maternal Behavioral Indicators (2, 3)										
Local Agency	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Pregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	% (Rank)		Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
SKCPH	247	82.6 (13)	6.1 (22)		291	45.7 (6)	297	19.9 (11)	243	68.7 (6)
Stevens	79	* *	* *		88	* *	89	* *	72	* *
Sumner	143	79.0 (17)	4.9 (17)		157	35.7 (20)	157	15.9 (17)	121	60.3 (17)
SW KS WIC	712	61.4 (29)	24.0 (29)		866	33.5 (25)	873	12.7 (25)	712	36.5 (30)
Wichita	15	* *	* *		18	* *	18	* *	15	* *
Wilson	92	* *	* *		101	46.5 (5)	101	16.8 (13)	88	* *
Wyandotte	1,981	65.6 (26)	8.0 (24)		2,267	35.9 (19)	2,286	10.4 (27)	1,910	56.9 (21)
Unknown Local Agencies	3	* *	* *		3	* *	3	* *	1	* *
Kansas	18,492	75.4	5.7		22,287	35.7	22,411	14.6	18,386	55.7
Nation (Prior Year)	580,485	80.1	3.1		1,187,387	31.7	266,567	19.3	269,548	80.6

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7F
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by Local Agency

Comparison of Smoking Indicators (2, 3)										
Local Agency	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Barton	445	40.0 (15)	377	24.9 (18)	125	28.8 (6)	370	13.8 (8)	377	8.0 (6)
Butler	275	44.0 (22)	234	26.5 (21)	77	* *	204	12.7 (6)	236	7.6 (5)
Chautauqua	42	* *	29	* *	10	* *	37	* *	28	* *
Cherokee	730	38.9 (14)	626	28.0 (25)	192	18.8 (11)	598	32.1 (23)	629	31.5 (25)
Cloud	217	47.0 (28)	185	23.8 (15)	69	* *	186	21.5 (19)	188	14.4 (14)
Cowley	347	41.8 (17)	249	28.5 (26)	86	* *	297	22.2 (20)	249	22.1 (22)
Crawford	560	44.1 (23)	432	25.0 (19)	149	30.2 (5)	505	41.8 (29)	433	40.2 (28)
Doniphan	34	* *	26	* *	7	* *	24	* *	27	* *
Douglas	497	44.3 (24)	422	21.6 (12)	150	41.3 (2)	392	10.2 (2)	425	5.6 (2)
Ford	673	11.1 (2)	568	5.5 (2)	44	* *	601	10.5 (4)	570	10.5 (9)
Franklin	186	47.8 (30)	165	30.9 (28)	61	* *	151	13.9 (9)	166	8.4 (8)
Geary	386	35.0 (10)	307	19.2 (10)	82	* *	320	37.5 (27)	307	29.6 (23)
Graham	44	* *	36	* *	12	* *	35	* *	37	* *
Greeley	15	* *	14	* *	3	* *	10	* *	14	* *
Greenwood	38	* *	14	* *	6	* *	35	* *	31	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7F
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by Local Agency

Comparison of Smoking Indicators (2, 3)										
Local Agency	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Harvey	275	36.0 (11)	234	19.7 (11)	70	* *	231	9.5 (1)	236	5.1 (1)
Jefferson	17	* *	1	* *	0	* *	19	* *	2	* *
Johnson	1,888	23.2 (5)	1,665	9.8 (4)	266	42.9 (1)	1,517	18.4 (18)	1,666	17.5 (19)
Leavenworth	351	45.9 (27)	294	25.2 (20)	95	* *	274	40.1 (28)	295	41.0 (29)
Lincoln	143	42.7 (20)	121	22.3 (13)	41	* *	120	14.2 (10)	124	15.3 (16)
Linn	66	* *	60	* *	26	* *	57	* *	60	* *
Lyon	357	30.5 (8)	313	16.6 (8)	73	* *	293	14.3 (11)	312	11.9 (11)
Marion	203	38.4 (13)	179	26.8 (24)	51	* *	149	10.7 (5)	179	6.7 (3)
Marshall	45	* *	37	* *	8	* *	38	* *	38	* *
Meade	99	* *	86	* *	17	* *	85	* *	89	* *
Miami	138	42.0 (18)	116	26.7 (23)	38	* *	102	16.7 (15)	116	14.7 (15)
NEK	279	47.0 (28)	214	30.8 (27)	72	* *	232	32.8 (24)	217	32.3 (26)
Nemaha	40	* *	33	* *	15	* *	34	* *	33	* *
Neosho	112	45.5 (26)	54	* *	21	* *	100	33.0 (25)	58	* *
Osborne	374	40.9 (16)	312	24.7 (17)	99	* *	296	22.3 (21)	316	20.6 (21)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7F
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by Local Agency

Comparison of Smoking Indicators (2, 3)										
Local Agency	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Reno	633	42.2 (19)	544	23.5 (14)	186	36.0 (3)	527	16.1 (14)	548	13.5 (13)
Riley	975	27.4 (6)	737	14.8 (6)	138	28.3 (8)	789	14.8 (13)	756	12.4 (12)
Saline	503	43.1 (21)	406	26.6 (22)	135	24.4 (9)	410	17.8 (17)	408	15.4 (17)
Scott	54	* *	45	* *	9	* *	47	* *	47	* *
Sedgwick	4,680	31.0 (9)	4,250	16.2 (7)	1,000	35.3 (4)	3,827	10.3 (3)	4,253	8.3 (7)
SEK	56	* *	45	* *	20	* *	50	* *	46	* *
Seward	514	5.8 (1)	457	3.1 (1)	17	* *	435	14.7 (12)	463	11.7 (10)
Shawnee	1,527	37.7 (12)	1,201	23.9 (16)	351	28.8 (6)	1,294	13.5 (7)	1,212	7.3 (4)
Sheridan	9	* *	10	* *	1	* *	8	* *	10	* *
Sherman	148	29.7 (7)	132	18.9 (9)	32	* *	123	34.1 (26)	132	30.3 (24)
SKCPH	283	45.2 (25)	227	32.6 (29)	84	* *	229	44.5 (30)	224	38.8 (27)
Stevens	87	* *	77	* *	13	* *	70	* *	79	* *
Sumner	153	52.9 (31)	135	32.6 (29)	55	* *	117	46.2 (31)	138	49.3 (30)
SW KS WIC	859	11.4 (3)	758	5.5 (2)	63	* *	707	17.4 (16)	759	16.2 (18)
Wichita	18	* *	12	* *	2	* *	15	* *	12	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7F
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by Local Agency

Comparison of Smoking Indicators (2, 3)										
Local Agency	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Wilson	99	* *	88	* *	40	* *	87	* *	88	* *
Wyandotte	2,255	19.8 (4)	2,000	12.1 (5)	299	23.7 (10)	1,897	22.5 (22)	2,004	19.4 (20)
Unknown Local Agencies	3	* *	3	* *	1	* *	1	* *	3	* *
Kansas	21,732	31.5	18,530	17.8	4,411	31.7	17,945	18.8	18,640	15.9
Nation (Prior Year)	999,199	23.1	856,953	14.2	146,933	35.5	655,095	18.9	578,852	19.9

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8F
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Infant Health Indicators by Local Agency

Comparison of Infant Health Indicators (2, 3)										
Local Agency	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)							
	Number	% (Rank)	%	(Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Barton	365	6.6 (14)	3.3	(1)	365	10.4 (26)	323	3.4 (7)	364	72.3 (10)
Butler	246	5.7 (6)	7.3	(17)	250	5.2 (1)	233	3.4 (7)	248	70.2 (15)
Chautauqua	49	*	*	*	*	*	*	*	*	*
Cherokee	645	6.4 (11)	5.4	(10)	643	8.2 (9)	589	3.6 (10)	642	57.9 (30)
Cloud	183	6.0 (8)	3.8	(2)	184	10.3 (25)	163	1.8 (3)	185	66.5 (23)
Cowley	306	9.2 (27)	7.5	(18)	309	11.7 (29)	266	5.6 (27)	310	63.5 (25)
Crawford	509	10.2 (28)	4.3	(5)	523	11.1 (27)	451	5.5 (26)	521	56.0 (31)
Doniphan	31	*	*	*	*	*	*	*	*	*
Douglas	422	6.4 (11)	5.2	(8)	430	7.7 (8)	390	3.8 (13)	427	77.8 (4)
Ford	568	4.6 (1)	6.0	(11)	588	10.0 (24)	510	1.6 (1)	587	72.1 (12)
Franklin	164	5.5 (5)	4.9	(6)	163	9.2 (16)	146	3.4 (7)	165	68.5 (20)
Geary	314	8.9 (24)	5.1	(7)	313	9.9 (21)	282	5.3 (24)	311	71.4 (14)
Graham	39	*	*	*	*	*	*	*	*	*
Greeley	13	*	*	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8F
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Infant Health Indicators by Local Agency

Comparison of Infant Health Indicators (2, 3)										
Local Agency	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)							
	Number	% (Rank)	%	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Greenwood	40	* * * *			40	* *	38	* *	40	* *
Harvey	234	6.0 (8)	8.5 (26)		234	8.5 (11)	211	3.8 (13)	235	80.0 (1)
Jefferson	61	* * * *			62	* *	49	* *	61	* *
Johnson	1,561	6.6 (14)	7.7 (20)		1,662	7.0 (4)	1,444	3.9 (15)	1,663	79.9 (2)
Leavenwort	298	9.1 (26)	8.4 (23)		303	8.6 (13)	272	6.3 (28)	304	62.2 (27)
Lincoln	116	6.0 (8)	10.3 (29)		127	6.3 (3)	109	3.7 (11)	126	66.7 (22)
Linn	59	* * * *			60	* *	57	* *	60	* *
Lyon	305	7.9 (20)	6.2 (12)		304	9.2 (16)	276	5.1 (22)	301	72.8 (9)
Marion	163	6.7 (16)	11.0 (30)		167	7.2 (5)	150	4.0 (16)	166	72.3 (10)
Marshall	40	* * * *			41	* *	36	* *	41	* *
Meade	84	* * * *			89	* *	74	* *	88	* *
Miami	128	10.2 (28)	8.6 (27)		130	8.5 (11)	118	6.8 (29)	128	68.8 (19)
NEK	281	7.8 (19)	6.8 (15)		292	7.2 (5)	260	4.6 (19)	292	58.6 (29)
Nemaha	26	* * * *			40	* *	24	* *	39	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8F
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Infant Health Indicators by Local Agency

Comparison of Infant Health Indicators (2, 3)										
Local Agency	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)		Number	%	Number	%	Number	%
	Number	% (Rank)	% (Rank)	Number						
Neosho	70	* * * *			133	12.8 (31)	61	* *	132	62.1 (28)
Osborne	308	6.5 (13)	7.8 (21)		323	9.9 (21)	275	3.3 (5)	322	75.8 (7)
Reno	550	7.5 (18)	4.2 (4)		550	8.9 (14)	498	4.2 (18)	554	67.1 (21)
Riley	790	5.4 (4)	7.8 (21)		793	5.2 (1)	748	3.3 (5)	793	76.2 (6)
Saline	466	8.8 (23)	8.4 (23)		478	9.8 (20)	418	5.3 (24)	476	63.2 (26)
Scott	55	* * * *			54	* *	49	* *	56	* *
Sedgwick	4,182	9.0 (25)	6.8 (15)		4,198	8.2 (9)	3,828	5.1 (22)	4,190	68.9 (18)
SEK	41	* * * *			51	* *	37	* *	51	* *
Seward	442	5.9 (7)	8.4 (23)		466	9.9 (21)	394	4.1 (17)	464	64.0 (24)
Shawnee	1,382	8.5 (22)	5.2 (8)		1,393	11.6 (28)	1,210	5.0 (21)	1,396	71.7 (13)
Sheridan	10	* * * *			10	* *	8	* *	10	* *
Sherman	131	4.6 (1)	3.8 (2)		131	7.6 (7)	121	1.7 (2)	131	73.3 (8)
SKCPH	225	5.3 (3)	7.6 (19)		236	9.3 (18)	197	2.0 (4)	241	76.3 (5)
Stevens	46	* * * *			75	* *	42	* *	76	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8F
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Infant Health Indicators by Local Agency

Comparison of Infant Health Indicators (2, 3)										
Local Agency	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)							
	Number	% (Rank)	%	(Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Sumner	135	11.1 (30)	8.9	(28)	140	12.1 (30)	118	6.8 (29)	140	70.0 (17)
SW KS WIC	746	7.0 (17)	6.2	(12)	764	9.7 (19)	667	3.7 (11)	761	77.9 (3)
Wichita	15	* * * *	*	*	15	* *	12	* *	15	* *
Wilson	93	* * * *	*	*	94	* *	81	* *	93	* *
Wyandotte	1,925	8.3 (21)	6.4	(14)	1,966	8.9 (14)	1,742	4.8 (20)	1,970	70.2 (15)
Unknown Local Agencies	3	* * * *	*	*	3	* *	3	* *	3	* *
Kansas	18,865	7.7	6.5		19,327	8.8	17,104	4.4	19,312	70.3
Nation (Prior Year)	1,085,134	8.2	6.6		1,025,278	11.5	890,981	3.6	952,980	65.4

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3F
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Racial and Ethnic Distribution by Local Agency
Combined 3 Years

Comparison of Racial and Ethnic Distribution (2)								
Local Agency	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Barton	1,359	72.7	1.4	25.0	0.1	0.3	0.4	0.0
Butler	815	90.8	1.3	5.4	0.4	0.4	1.7	0.0
Chautauqua	137	93.4	0.0	5.1	0.7	0.0	0.7	0.0
Cherokee	2,172	83.7	6.5	5.8	2.3	0.9	0.8	0.0
Cloud	611	92.5	1.0	4.4	0.7	0.8	0.7	0.0
Cowley	1,020	76.5	2.4	17.8	1.0	1.6	0.8	0.0
Crawford	1,845	87.4	3.3	6.7	0.3	1.1	1.2	0.0
Doniphan	122	86.1	4.1	7.4	0.8	0.0	1.6	0.0
Douglas	1,442	63.9	9.1	10.9	9.2	5.2	1.8	0.0
Ford	1,930	22.8	0.9	75.4	0.2	0.7	0.1	0.0
Franklin	509	90.6	1.8	4.7	1.2	0.4	1.4	0.0
Geary	1,056	53.1	24.4	15.2	0.6	2.8	3.9	0.0
Graham	132	89.4	1.5	8.3	0.0	0.0	0.8	0.0
Greeley	31	*	*	*	*	*	*	*
Greenwood	140	92.9	0.7	5.0	0.0	0.7	0.7	0.0
Harvey	790	69.9	2.5	24.7	1.1	0.3	1.5	0.0
Jefferson	193	96.4	0.5	1.0	2.1	0.0	0.0	0.0
Johnson	5,514	46.4	12.8	35.9	0.5	3.1	1.2	0.0
Leavenworth	1,045	69.4	15.6	9.8	0.7	1.4	3.2	0.0
Lincoln	398	94.7	0.3	4.5	0.0	0.0	0.5	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3F
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Racial and Ethnic Distribution by Local Agency
Combined 3 Years

Comparison of Racial and Ethnic Distribution (2)								
Local Agency	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaska Native	Asian/Pacific Islander	Multiple Races	All Other/Unknown
		%	%	%	%	%	%	%
Linn	186	96.2	0.0	2.7	0.0	0.0	1.1	0.0
Lyon	1,136	57.8	4.8	35.1	0.3	1.2	0.7	0.0
Marion	565	91.7	0.0	6.4	1.1	0.4	0.5	0.0
Marshall	171	91.8	0.0	3.5	2.9	0.0	1.8	0.0
Meade	343	66.8	0.3	32.1	0.0	0.0	0.9	0.0
Miami	409	92.7	2.4	2.9	0.0	0.7	1.2	0.0
NEK	926	81.5	4.5	3.3	8.6	0.0	1.9	0.0
Nemaha	116	89.7	0.0	7.8	0.0	0.9	1.7	0.0
Neosho	438	90.6	2.5	5.3	0.5	0.5	0.7	0.0
Osborne	1,063	89.6	1.6	7.3	0.7	0.8	0.0	0.0
Reno	1,840	83.4	2.6	12.0	0.7	0.6	0.8	0.0
Riley	2,769	74.2	10.9	10.1	0.6	3.8	0.4	0.0
Saline	1,437	71.7	4.4	19.7	0.4	2.3	1.5	0.0
Scott	181	64.6	0.6	33.1	1.7	0.0	0.0	0.0
Sedgwick	12,755	45.6	18.4	29.1	1.7	3.8	1.4	0.0
SEK	180	92.8	0.6	5.0	0.6	1.1	0.0	0.0
Seward	1,478	17.4	1.4	78.3	1.8	0.7	0.3	0.0
Shawnee	4,629	58.5	14.7	22.8	2.1	0.7	1.3	0.0
Sheridan	27	*	*	*	*	*	*	*
Sherman	427	81.7	0.7	15.7	0.2	0.7	0.9	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3F
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Racial and Ethnic Distribution by Local Agency
Combined 3 Years

Comparison of Racial and Ethnic Distribution (2)								
Local Agency	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
SKCPH	788	86.2	0.5	11.8	0.9	0.4	0.3	0.0
Stevens	250	45.2	0.0	53.6	0.8	0.4	0.0	0.0
Sumner	443	91.2	0.9	5.0	1.1	0.2	1.6	0.0
SW KS WIC	2,487	29.9	0.8	66.9	0.0	2.1	0.2	0.0
Wichita	46	*	*	*	*	*	*	*
Wilson	281	96.4	0.7	2.1	0.4	0.0	0.4	0.0
Wyandotte	6,467	26.0	31.1	39.9	0.2	2.5	0.3	0.0
Unknown Local Agencies	3	*	*	*	*	*	*	*
Kansas	63,102	57.2	11.4	27.0	1.3	2.1	1.0	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4F
2007 - 2009 Pregnancy Nutrition Surveillance
Kansas
Comparison of Age Distribution by Local Agency
Combined 3 Years

(1)

Comparison of Age Distribution (2)								
Local Agency	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Barton	1,359	0.3	6.2	16.1	60.9	15.6	0.9	0.0
Butler	815	0.1	7.6	17.2	61.7	12.8	0.5	0.1
Chautauqua	137	0.0	3.6	21.2	59.1	13.1	2.2	0.7
Cherokee	2,172	0.2	7.0	18.1	62.7	11.2	0.6	0.2
Cloud	611	0.2	5.9	17.7	60.9	14.1	1.3	0.0
Cowley	1,020	0.8	8.1	16.4	60.9	13.4	0.4	0.0
Crawford	1,845	0.2	7.3	16.9	62.9	12.0	0.5	0.2
Doniphan	122	0.8	4.1	14.8	63.9	16.4	0.0	0.0
Douglas	1,442	0.1	3.8	13.2	63.9	17.7	1.4	0.0
Ford	1,930	0.4	7.6	12.5	56.1	22.1	1.3	0.1
Franklin	509	0.8	6.3	19.8	58.2	14.1	0.6	0.2
Geary	1,056	0.5	4.9	15.4	67.2	11.6	0.4	0.0
Graham	132	0.0	3.0	15.2	66.7	14.4	0.8	0.0
Greeley	31	*	*	*	*	*	*	*
Greenwood	140	0.0	5.7	17.1	62.1	15.0	0.0	0.0
Harvey	790	0.3	6.5	13.4	63.3	15.3	1.3	0.0
Jefferson	193	0.0	8.3	17.6	60.1	13.5	0.5	0.0
Johnson	5,514	0.2	4.4	11.6	60.8	21.7	1.3	0.1
Leavenworth	1,045	0.1	5.9	18.4	63.3	11.3	1.1	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4F
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Age Distribution by Local Agency
Combined 3 Years

Comparison of Age Distribution (2)								
Local Agency	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Lincoln	398	0.0	2.5	14.8	62.1	20.1	0.5	0.0
Linn	186	0.0	5.4	17.2	62.9	13.4	1.1	0.0
Lyon	1,136	0.4	6.3	14.5	61.8	14.9	2.1	0.1
Marion	565	0.2	6.5	14.7	63.2	14.3	1.1	0.0
Marshall	171	0.0	5.3	9.9	69.0	15.8	0.0	0.0
Meade	343	0.0	6.4	10.2	58.9	22.7	1.7	0.0
Miami	409	0.2	7.3	14.4	65.5	12.0	0.5	0.0
NEK	926	0.0	6.7	16.4	63.8	12.4	0.6	0.0
Nemaha	116	0.0	5.2	12.9	62.9	15.5	2.6	0.9
Neosho	438	0.2	7.1	19.6	61.4	11.2	0.5	0.0
Osborne	1,063	0.1	3.8	14.2	66.2	14.1	1.5	0.1
Reno	1,840	0.2	5.5	15.7	65.8	12.3	0.5	0.0
Riley	2,769	0.0	2.2	11.8	73.8	11.7	0.5	0.0
Saline	1,437	0.2	7.7	15.1	63.1	12.9	1.0	0.1
Scott	181	0.6	5.0	9.4	65.7	17.1	2.2	0.0
Sedgwick	12,755	0.3	6.9	13.9	60.8	16.7	1.3	0.0
SEK	180	0.6	8.9	11.7	67.8	10.6	0.6	0.0
Seward	1,478	0.3	7.7	13.9	55.8	20.6	1.6	0.0
Shawnee	4,629	0.3	6.7	15.0	61.8	15.0	1.0	0.1

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4F
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Age Distribution by Local Agency
Combined 3 Years

Comparison of Age Distribution (2)								
Local Agency	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Sheridan	27	*	*	*	*	*	*	*
Sherman	427	0.2	6.6	16.4	60.2	14.5	2.1	0.0
SKCPH	788	0.1	6.7	16.0	63.2	13.5	0.5	0.0
Stevens	250	0.0	6.8	14.8	59.2	18.8	0.4	0.0
Sumner	443	0.0	5.0	18.7	62.8	12.9	0.7	0.0
SW KS WIC	2,487	0.3	9.2	14.8	55.2	18.9	1.4	0.1
Wichita	46	*	*	*	*	*	*	*
Wilson	281	0.0	8.9	16.0	62.6	12.5	0.0	0.0
Wyandotte	6,467	0.6	8.2	13.9	58.0	18.0	1.3	0.1
Unknown Local Agencies	3	*	*	*	*	*	*	*
Kansas	63,102	0.3	6.5	14.5	61.6	16.1	1.1	0.1

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5F
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Maternal Health Indicators by Local Agency
Combined 3 Years

Comparison of Maternal Health Indicators (2, 3)														
Local Agency	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Barton	1,339	4.3 (14)	54.4 (29)	1,195	22.2 (34)	48.4 (9)	151	22.5 (5)	260	20.0 (9)	1,156	3.7 (26)	1,153	6.8(21)
Butler	791	6.3 (38)	50.2 (4)	728	18.7 (15)	52.5 (30)	103	38.8 (20)	216	28.7 (26)	692	4.2 (33)	692	8.8(37)
Chautauqua	136	4.4 (15)	58.8 (39)	123	24.4 (42)	45.5 (2)	8	* *	21	* *	121	2.5 (4)	121	5.8(15)
Cherokee	2,133	6.0 (35)	53.6 (20)	1,895	20.7 (28)	51.4 (18)	333	30.3 (14)	542	24.5 (16)	1,869	3.0 (10)	1,870	7.1(26)
Cloud	605	5.0 (26)	53.4 (18)	536	20.1 (25)	53.0 (32)	68	* *	147	27.9 (24)	518	3.5 (22)	521	6.9(22)
Cowley	1,009	5.1 (28)	52.9 (16)	915	20.8 (30)	52.0 (24)	145	32.4 (18)	129	23.3 (13)	877	3.3 (17)	878	7.4(30)
Crawford	1,809	5.2 (30)	54.2 (25)	1,594	18.9 (17)	55.3 (41)	274	19.7 (2)	360	21.4 (11)	1,583	4.5 (35)	1,582	8.0(33)
Doniphan	119	2.5 (4)	63.9 (44)	97	* *	* *	25	* *	24	* *	98	* *	98	* *
Douglas	1,421	4.4 (15)	49.3 (1)	1,246	17.3 (5)	56.1 (43)	202	31.2 (16)	403	24.8 (17)	1,199	3.9 (29)	1,198	5.2(10)
Ford	1,906	2.4 (3)	54.7 (33)	1,727	21.1 (31)	48.2 (7)	252	21.8 (3)	200	16.0 (3)	1,690	3.3 (17)	1,689	5.1(8)
Franklin	499	3.2 (8)	54.3 (28)	437	16.7 (1)	55.4 (42)	49	* *	133	16.5 (5)	381	2.4 (2)	381	8.4(35)
Geary	1,031	4.6 (19)	51.6 (12)	860	17.6 (8)	54.3 (37)	165	38.8 (20)	193	29.5 (27)	854	3.4 (21)	854	7.8(31)
Graham	126	7.1 (40)	50.8 (9)	111	21.6 (33)	48.6 (10)	17	* *	27	* *	120	3.3 (17)	120	5.0(6)
Greeley	29	* *	* *	27	* *	* *	4	* *	6	* *	29	* *	29	* *
Greenwood	137	2.9 (5)	60.6 (43)	128	22.7 (36)	52.3 (27)	16	* *	25	* *	126	4.0 (31)	127	9.4(38)
Harvey	779	4.2 (12)	54.0 (24)	707	22.8 (37)	48.8 (11)	114	30.7 (15)	133	25.6 (22)	679	3.5 (22)	677	6.6(19)
Jefferson	192	7.3 (42)	55.2 (34)	172	16.9 (4)	55.2 (40)	29	* *	58	* *	169	4.7 (36)	169	6.5(17)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5F
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Maternal Health Indicators by Local Agency
Combined 3 Years

Comparison of Maternal Health Indicators (2, 3)														
Local Agency	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Johnson	5,359	4.2 (12)	53.9 (23)	4,828	18.2 (10)	51.3 (17)	920	24.9 (7)	1,632	17.7 (6)	4,702	5.3 (41)	4,699	4.6(5)
Leavenwort	1,019	4.9 (24)	53.5 (19)	905	18.6 (14)	55.0 (39)	171	29.2 (11)	246	19.5 (8)	857	3.7 (26)	857	5.5(13)
Lincoln	391	3.8 (10)	58.8 (39)	373	23.3 (40)	52.0 (24)	50	* *	98	* *	357	5.3 (41)	357	7.3(28)
Linn	184	5.4 (32)	58.2 (38)	171	29.8 (43)	43.9 (1)	24	* *	30	* *	168	3.0 (10)	168	4.2(4)
Lyon	1,109	3.3 (9)	58.8 (39)	1,008	20.1 (25)	51.7 (20)	153	14.4 (1)	222	16.2 (4)	978	3.2 (15)	978	9.9(41)
Marion	552	5.8 (34)	50.0 (3)	503	20.3 (27)	51.5 (19)	71	* *	136	21.3 (10)	489	3.3 (17)	485	7.0(23)
Marshall	167	7.2 (41)	50.3 (5)	140	22.9 (38)	47.9 (6)	23	* *	35	* *	137	0.7 (1)	137	5.1(8)
Meade	339	2.9 (5)	57.8 (37)	295	22.4 (35)	51.9 (22)	43	* *	77	* *	287	3.1 (12)	287	7.3(28)
Miami	395	8.1 (43)	54.2 (25)	362	18.2 (10)	53.3 (34)	78	* *	116	25.0 (19)	352	5.1 (39)	352	6.5(17)
NEK	904	4.8 (22)	54.2 (25)	798	18.2 (10)	54.9 (38)	140	30.0 (12)	198	24.2 (15)	803	2.7 (6)	802	10.1(42)
Nemaha	109	1.8 (1)	53.2 (17)	104	19.2 (18)	51.9 (22)	25	* *	37	* *	106	2.8 (7)	106	2.8(1)
Neosho	424	8.5 (44)	50.7 (8)	400	17.5 (7)	50.3 (14)	72	* *	71	* *	387	3.9 (29)	387	7.0(23)
Osborne	1,041	4.7 (20)	50.3 (5)	924	20.7 (28)	47.4 (5)	177	22.0 (4)	261	15.7 (2)	911	3.2 (15)	911	7.2(27)
Reno	1,818	5.5 (33)	53.7 (22)	1,647	19.9 (23)	50.7 (15)	242	28.5 (10)	406	23.4 (14)	1,573	2.4 (2)	1,574	8.0(33)
Riley	2,693	4.5 (18)	49.5 (2)	2,239	16.8 (2)	53.7 (35)	398	23.9 (6)	627	25.0 (19)	2,171	2.9 (9)	2,170	6.3(16)
Saline	1,415	4.7 (20)	53.6 (20)	1,307	18.4 (13)	52.3 (27)	264	27.7 (8)	340	19.4 (7)	1,277	3.5 (22)	1,281	5.0(6)
Scott	176	4.0 (11)	54.5 (32)	155	17.4 (6)	54.2 (36)	26	* *	31	* *	156	2.6 (5)	156	3.2(2)

- (1) Reporting period is January 1 through December 31.
 - (2) Excludes records with unknown data or errors.
 - (3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.
 - (4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
 - (5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
 - (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5F
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Maternal Health Indicators by Local Agency
Combined 3 Years

Comparison of Maternal Health Indicators (2, 3)														
Local Agency	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Sedgwick	12,538	4.9 (24)	52.4 (14)	11,593	19.4 (21)	51.8 (21)	2,553	32.3 (17)	3,421	25.4 (21)	11,241	3.8 (28)	11,243	7.0(23)
SEK	178	5.1 (28)	51.7 (13)	166	24.1 (41)	46.4 (3)	30	* *	32	* *	161	3.1 (12)	161	11.8(43)
Seward	1,450	2.9 (5)	54.4 (29)	1,328	21.4 (32)	50.1 (13)	249	27.7 (8)	218	24.8 (17)	1,294	5.2 (40)	1,296	5.3(12)
Shawnee	4,543	4.8 (22)	52.8 (15)	4,050	19.3 (20)	52.0 (24)	482	39.0 (22)	848	29.8 (28)	3,973	4.9 (37)	3,978	5.2(10)
Sheridan	26	* *	* *	23	* *	* *	5	* *	3	* *	24	* *	24	* *
Sherman	418	6.5 (39)	55.7 (35)	372	19.6 (22)	53.2 (33)	48	* *	84	* *	361	3.6 (25)	361	9.4(38)
SKCPH	776	5.0 (26)	50.3 (5)	660	20.0 (24)	51.2 (16)	97	* *	176	22.7 (12)	635	2.8 (7)	634	8.7(36)
Stevens	247	2.0 (2)	60.3 (42)	220	16.8 (2)	49.1 (12)	38	* *	63	* *	213	8.9 (43)	213	6.6(19)
Sumner	433	5.3 (31)	51.0 (10)	385	19.2 (18)	48.3 (8)	77	* *	113	12.4 (1)	378	4.0 (31)	378	9.8(40)
SW KS WIC	2,440	6.1 (36)	51.5 (11)	2,256	22.9 (38)	46.4 (3)	340	30.0 (12)	436	28.2 (25)	2,221	4.9 (37)	2,217	3.8(3)
Wichita	46	* *	* *	42	* *	* *	7	* *	10	* *	41	* *	41	* *
Wilson	279	6.1 (36)	56.6 (36)	265	18.1 (9)	52.8 (31)	46	* *	50	* *	255	4.3 (34)	255	7.8(31)
Wyandotte	6,336	4.4 (15)	54.4 (29)	5,782	18.7 (15)	52.3 (27)	1,060	34.3 (19)	1,196	26.3 (23)	5,623	3.1 (12)	5,625	5.7(14)
Unknown Local Agencies	3	* *	* *	3	* *	* *	0	* *	1	* *	3	* *	3	* *
Kansas	61,869	4.7	53.2	55,802	19.5	51.6	9,864	30.1	14,091	23.5	54,295	3.8	54,295	6.4

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6F
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Maternal Behavioral Indicators by Local Agency
Combined 3 Years

Comparison of Maternal Behavioral Indicators (2, 3)													
Local Agency	Medical Care				WIC Enrollment			Multivitamin Consumption					
	1st Trimester		No Care		1st Trimester			Prepregnancy (4)		During Pregnancy (5)			
	Number	% (Rank)	%	% (Rank)	Number	%	% (Rank)	Number	%	% (Rank)			
Barton	1,194	79.8 (22)	1.7	(9)	1,177	52.8	(3)	716	20.3	(6)	683	59.0	(29)
Butler	716	70.8 (37)	10.9	(40)	638	43.9	(16)	436	15.1	(18)	364	41.2	(36)
Chautauqua	121	80.2 (20)	5.0	(28)	120	58.3	(1)	73	*	*	75	*	*
Cherokee	1,671	82.5 (19)	4.8	(26)	1,855	39.8	(26)	1,143	14.3	(21)	1,136	63.7	(20)
Cloud	543	85.6 (10)	3.1	(18)	530	51.5	(4)	325	17.8	(11)	303	67.3	(11)
Cowley	866	68.4 (39)	5.5	(31)	870	44.9	(13)	557	12.2	(24)	535	60.6	(27)
Crawford	1,575	88.8 (4)	1.9	(11)	1,607	44.1	(15)	990	11.4	(25)	1,014	63.5	(21)
Doniphan	106	78.3 (27)	5.7	(33)	103	39.8	(26)	64	*	*	72	*	*
Douglas	1,243	72.3 (34)	6.9	(36)	1,242	39.3	(30)	765	17.8	(11)	719	63.4	(22)
Ford	1,668	72.1 (35)	14.0	(42)	1,712	45.7	(10)	1,033	6.8	(31)	1,074	57.3	(32)
Franklin	254	79.5 (25)	3.1	(18)	414	41.8	(23)	278	3.2	(35)	292	68.2	(9)
Geary	822	70.7 (38)	0.7	(2)	923	40.4	(25)	588	17.7	(13)	569	60.8	(26)
Graham	119	88.2 (5)	0.8	(5)	107	46.7	(9)	60	*	*	63	*	*
Greeley	29	*	*	*	*	*	*	22	*	*	17	*	*
Greenwood	118	61.9 (44)	29.7	(44)	124	54.8	(2)	69	*	*	73	*	*
Harvey	659	72.1 (35)	13.8	(41)	688	47.2	(7)	422	14.7	(20)	417	70.5	(3)
Jefferson	162	90.7 (1)	1.9	(11)	153	39.2	(32)	104	6.7	(32)	105	54.3	(34)
Johnson	4,545	64.0 (43)	3.0	(17)	4,617	31.6	(42)	2,938	13.2	(23)	2,727	67.1	(12)
Leavenworth	915	85.4 (12)	0.7	(2)	823	36.5	(37)	561	18.9	(9)	485	72.4	(2)
Lincoln	375	85.6 (10)	2.4	(15)	323	39.3	(30)	218	31.7	(2)	201	66.7	(15)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6F
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Maternal Behavioral Indicators by Local Agency
Combined 3 Years

Comparison of Maternal Behavioral Indicators (2, 3)										
Local Agency	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	% (Rank)		Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Linn	170	79.4 (26)	6.5 (34)		164	45.1 (12)	96	* *	109	66.1 (16)
Lyon	1,025	82.7 (17)	1.0 (7)		922	42.2 (21)	580	14.8 (19)	524	63.9 (19)
Marion	507	80.1 (21)	3.4 (20)		470	47.0 (8)	313	19.8 (7)	271	69.7 (5)
Marshall	147	90.5 (2)	0.7 (2)		138	42.0 (22)	81	* *	72	* *
Meade	296	79.7 (23)	4.1 (25)		285	45.6 (11)	174	38.5 (1)	171	69.0 (7)
Miami	371	77.9 (29)	0.8 (5)		323	32.5 (41)	220	15.5 (16)	195	58.5 (30)
NEK	757	84.7 (13)	5.5 (31)		811	39.5 (29)	513	9.2 (28)	533	69.0 (7)
Nemaha	111	86.5 (8)	5.4 (30)		96	* *	59	* *	59	* *
Neosho	377	88.9 (3)	4.0 (24)		382	42.4 (20)	238	6.7 (32)	223	66.8 (14)
Osborne	940	87.1 (7)	1.7 (9)		902	43.1 (18)	573	26.5 (3)	588	70.4 (4)
Reno	1,639	84.1 (14)	2.0 (14)		1,529	40.6 (24)	1,008	21.0 (5)	959	66.1 (16)
Riley	2,047	83.2 (16)	0.6 (1)		2,283	36.4 (38)	1,482	15.9 (15)	1,416	62.8 (23)
Saline	1,209	74.0 (33)	4.8 (26)		1,205	33.9 (40)	802	5.2 (34)	775	57.0 (33)
Scott	158	79.7 (23)	8.2 (38)		156	39.7 (28)	89	* *	102	69.6 (6)
Sedgwick	11,643	74.7 (31)	3.6 (22)		10,832	28.2 (43)	7,027	13.6 (22)	6,444	50.8 (35)
SEK	161	88.2 (5)	1.9 (11)		153	43.8 (17)	96	* *	95	* *
Seward	1,260	66.4 (41)	8.4 (39)		1,279	37.1 (34)	799	8.8 (29)	740	66.9 (13)
Shawnee	3,871	76.3 (30)	3.7 (23)		3,968	48.5 (5)	2,476	7.7 (30)	2,457	57.6 (31)
Sheridan	23	* *	* *		20	* *	12	* *	12	* *
Sherman	238	83.6 (15)	5.0 (28)		363	44.4 (14)	227	26.4 (4)	246	79.7 (1)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6F
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Maternal Behavioral Indicators by Local Agency
Combined 3 Years

Comparison of Maternal Behavioral Indicators (2, 3)													
Local Agency	Medical Care				WIC Enrollment			Multivitamin Consumption					
	1st Trimester		No Care		1st Trimester			Prepregnancy (4)		During Pregnancy (5)			
	Number	% (Rank)	%	% (Rank)	Number	%	% (Rank)	Number	%	% (Rank)			
SKCPH	660	78.3 (27)	7.6	(37)	675	47.7	(6)	457	19.3	(8)	442	68.1	(10)
Stevens	224	74.6 (32)	2.7	(16)	210	35.2	(39)	127	18.9	(9)	115	60.0	(28)
Sumner	398	82.7 (17)	3.5	(21)	362	38.1	(33)	241	16.6	(14)	209	66.0	(18)
SW KS WIC	2,125	66.1 (42)	17.1	(43)	2,126	36.8	(36)	1,369	10.3	(27)	1,270	41.0	(37)
Wichita	42	* *	* *	* *	40	* *	* *	24	* *	* *	24	* *	* *
Wilson	260	85.8 (9)	1.5	(8)	254	42.9	(19)	151	15.2	(17)	156	62.8	(23)
Wyandotte	5,808	66.5 (40)	6.6	(35)	5,566	37.0	(35)	3,457	11.0	(26)	3,286	61.5	(25)
Unknown Local Agencies	3	* *	* *	* *	3	* *	* *	3	* *	* *	1	* *	* *
Kansas	54,171	75.0	4.9		53,570	38.0		34,056	13.7		32,418	60.0	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7F
2007 - 2009 Pregnancy Nutrition Surveillance
Kansas
Comparison of Smoking Indicators by Local Agency
Combined 3 Years

(1)

Comparison of Smoking Indicators (2, 3)										
Local Agency	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Barton	1,335	41.2 (19)	1,172	25.8 (22)	391	29.2 (18)	1,101	14.7 (11)	1,178	10.4 (9)
Butler	771	50.5 (36)	639	31.0 (35)	235	30.6 (14)	558	14.3 (10)	641	8.6 (6)
Chautauqua	125	39.2 (15)	93	* *	27	* *	103	37.9 (34)	92	* *
Cherokee	2,083	39.9 (18)	1,759	27.5 (30)	529	17.4 (30)	1,695	29.9 (29)	1,797	28.4 (31)
Cloud	599	43.7 (24)	514	24.7 (18)	180	28.3 (22)	496	20.4 (21)	519	16.0 (18)
Cowley	981	44.5 (27)	802	30.3 (32)	297	26.6 (26)	810	19.4 (20)	808	18.8 (22)
Crawford	1,751	44.9 (29)	1,426	26.8 (26)	499	31.5 (12)	1,494	38.4 (35)	1,427	37.0 (35)
Doniphan	98	* *	62	* *	23	* *	76	* *	65	* *
Douglas	1,407	43.7 (24)	1,195	20.6 (13)	392	42.9 (2)	1,121	10.1 (4)	1,200	6.5 (2)
Ford	1,855	10.0 (2)	1,531	4.9 (2)	105	28.6 (19)	1,618	8.9 (2)	1,532	8.3 (4)
Franklin	504	46.0 (31)	444	27.3 (29)	136	31.6 (11)	378	14.8 (12)	446	10.5 (10)
Geary	1,032	39.2 (15)	842	20.4 (12)	265	36.6 (5)	850	39.8 (36)	844	37.0 (35)
Graham	126	46.8 (32)	104	30.8 (34)	38	* *	98	* *	107	29.9 (33)
Greeley	31	* *	29	* *	6	* *	22	* *	29	* *
Greenwood	124	54.8 (40)	72	* *	31	* *	110	31.8 (32)	104	26.0 (30)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7F
2007 - 2009 Pregnancy Nutrition Surveillance
Kansas
Comparison of Smoking Indicators by Local Agency
Combined 3 Years

(1)

Comparison of Smoking Indicators (2, 3)										
Local Agency	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Harvey	773	35.1 (12)	658	22.5 (14)	189	30.2 (15)	638	5.8 (1)	662	4.1 (1)
Jefferson	70	* *	28	* *	9	* *	69	* *	32	* *
Johnson	5,398	24.9 (7)	4,746	10.7 (4)	812	44.5 (1)	4,213	18.4 (19)	4,761	18.3 (21)
Leavenworth	1,010	47.6 (33)	853	27.1 (28)	281	29.9 (16)	742	42.7 (37)	862	40.5 (37)
Lincoln	384	42.4 (20)	342	24.6 (17)	116	31.9 (10)	300	14.0 (7)	348	12.4 (12)
Linn	185	56.2 (41)	166	40.4 (40)	76	* *	155	14.2 (9)	168	10.7 (11)
Lyon	1,114	31.3 (10)	1,005	16.6 (10)	225	36.9 (4)	846	15.1 (13)	1,008	13.2 (13)
Marion	553	39.8 (17)	492	25.0 (19)	150	30.7 (13)	415	10.1 (4)	496	6.9 (3)
Marshall	166	50.6 (37)	132	32.6 (37)	46	* *	128	52.3 (40)	137	49.6 (40)
Meade	318	21.4 (4)	268	13.4 (7)	41	* *	263	16.3 (17)	274	13.9 (15)
Miami	359	47.9 (35)	287	30.3 (32)	99	* *	265	21.5 (22)	286	16.8 (20)
NEK	784	45.7 (30)	587	27.8 (31)	198	24.7 (28)	674	31.6 (31)	599	29.7 (32)
Nemaha	113	47.8 (34)	101	26.7 (25)	41	* *	88	* *	100	42.0 (39)
Neosho	393	43.0 (22)	292	25.7 (21)	102	29.4 (17)	335	24.2 (26)	297	21.5 (26)
Osborne	1,016	43.6 (23)	852	24.4 (16)	299	35.1 (7)	805	21.7 (23)	860	21.2 (25)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7F
2007 - 2009 Pregnancy Nutrition Surveillance
Kansas
Comparison of Smoking Indicators by Local Agency
Combined 3 Years

(1)

Comparison of Smoking Indicators (2, 3)											
Local Agency	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	
Reno	1,801	44.6 (28)	1,581	25.2 (20)	544	32.7 (8)	1,416	15.3 (14)	1,592	13.6 (14)	
Riley	2,679	27.8 (8)	2,000	16.5 (9)	382	28.0 (23)	2,073	15.3 (14)	2,152	13.9 (15)	
Saline	1,303	44.2 (26)	1,029	26.9 (27)	348	27.3 (24)	1,040	13.2 (6)	1,034	10.3 (8)	
Scott	154	28.6 (9)	124	16.9 (11)	26	* *	136	30.1 (30)	129	23.3 (27)	
Sedgwick	12,530	32.2 (11)	11,365	16.4 (8)	2,701	35.4 (6)	9,927	9.7 (3)	11,365	8.3 (4)	
SEK	172	56.4 (42)	147	32.0 (36)	63	* *	145	29.7 (28)	150	25.3 (29)	
Seward	1,428	7.5 (1)	1,278	3.5 (1)	69	* *	1,188	15.7 (16)	1,292	14.4 (17)	
Shawnee	4,303	37.0 (13)	3,432	22.7 (15)	946	26.8 (25)	3,577	14.1 (8)	3,446	9.1 (7)	
Sheridan	22	* *	19	* *	2	* *	18	* *	19	* *	
Sherman	413	37.8 (14)	353	25.8 (22)	103	21.4 (29)	335	26.0 (27)	353	23.8 (28)	
SKCPH	734	42.6 (21)	567	26.6 (24)	189	28.6 (19)	606	35.6 (33)	605	32.6 (34)	
Stevens	247	21.9 (6)	214	11.2 (5)	36	* *	192	22.4 (25)	217	18.9 (23)	
Sumner	418	53.6 (39)	357	33.9 (39)	147	28.6 (19)	316	49.1 (39)	362	50.0 (41)	
SW KS WIC	2,433	12.9 (3)	2,221	6.5 (3)	209	39.7 (3)	1,975	17.5 (18)	2,223	16.2 (19)	
Wichita	46	* *	37	* *	5	* *	37	* *	37	* *	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7F
2007 - 2009 Pregnancy Nutrition Surveillance
Kansas
Comparison of Smoking Indicators by Local Agency
Combined 3 Years

(1)

Comparison of Smoking Indicators (2, 3)										
Local Agency	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Wilson	276	53.3 (38)	247	33.6 (38)	113	32.7 (8)	239	45.6 (38)	250	41.6 (38)
Wyandotte	6,305	21.6 (5)	5,597	12.7 (6)	906	26.3 (27)	5,183	22.2 (24)	5,641	21.0 (24)
Unknown Local Agencies	3	* *	3	* *	1	* *	1	* *	3	* *
Kansas	60,725	32.6	52,064	18.3	12,628	31.8	48,870	18.4	52,549	16.3

(1) Reporting period is January 1 through December 31.
 (2) Excludes records with unknown data or errors.
 (3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.
 (4) Among women who smoked 3 months prior to pregnancy.
 * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8F
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Infant Health Indicators by Local Agency
Combined 3 Years

Comparison of Infant Health Indicators (2, 3)									
Local Agency	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Barton	1,152	7.3 (19)	5.1 (8)	1,020	12.4 (35)	887	3.3 (17)	1,117	72.8 (13)
Butler	727	8.1 (30)	5.6 (12)	575	10.4 (21)	509	3.7 (23)	700	72.1 (15)
Chautauqua	119	8.4 (34)	4.2 (3)	109	11.9 (32)	94	* *	116	70.7 (19)
Cherokee	1,914	6.9 (18)	5.7 (15)	1,627	10.8 (27)	1,449	3.2 (15)	1,834	58.8 (37)
Cloud	518	6.4 (12)	2.7 (1)	461	14.3 (40)	385	2.9 (8)	521	68.1 (25)
Cowley	894	7.8 (26)	7.3 (33)	771	10.9 (29)	675	4.9 (34)	875	58.3 (38)
Crawford	1,581	9.0 (37)	4.8 (4)	1,402	12.1 (33)	1,216	4.5 (31)	1,570	57.8 (40)
Doniphan	104	5.8 (6)	5.8 (18)	87	* *	78	* *	102	52.0 (44)
Douglas	1,234	6.1 (7)	6.6 (23)	1,060	10.6 (23)	933	3.0 (10)	1,188	77.2 (5)
Ford	1,658	5.1 (3)	5.7 (15)	1,519	10.4 (21)	1,335	1.9 (2)	1,467	72.1 (15)
Franklin	438	6.2 (9)	5.0 (6)	350	10.3 (19)	311	3.2 (15)	425	60.7 (35)
Geary	881	7.4 (21)	6.6 (23)	755	8.7 (6)	686	3.5 (20)	826	69.5 (21)
Graham	111	9.0 (37)	10.8 (43)	94	* *	84	* *	112	75.9 (7)
Greeley	28	* *	* *	24	* *	23	* *	27	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8F
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Infant Health Indicators by Local Agency
Combined 3 Years

Comparison of Infant Health Indicators (2, 3)									
Local Agency	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Greenwood	121	6.6 (14)	3.3 (2)	109	10.1 (17)	97	* *	116	60.3 (36)
Harvey	668	7.3 (19)	7.0 (29)	604	12.3 (34)	518	3.3 (17)	680	78.8 (3)
Jefferson	172	8.1 (30)	7.6 (36)	138	16.7 (41)	114	5.3 (35)	170	61.8 (34)
Johnson	4,628	6.5 (13)	7.5 (34)	4,066	8.0 (4)	3,551	3.0 (10)	4,612	79.2 (2)
Leavenwort	862	9.4 (40)	5.0 (6)	713	9.4 (10)	631	4.8 (33)	892	63.2 (32)
Lincoln	351	6.8 (17)	8.3 (40)	298	8.7 (6)	261	3.1 (12)	344	65.7 (28)
Linn	166	8.4 (34)	4.8 (4)	150	10.0 (16)	133	6.8 (37)	163	68.7 (23)
Lyon	1,002	6.7 (15)	6.8 (25)	809	9.8 (15)	728	2.9 (8)	963	71.1 (18)
Marion	480	6.3 (10)	9.0 (41)	406	7.9 (3)	366	3.3 (17)	480	74.0 (11)
Marshall	138	9.4 (40)	5.1 (8)	113	10.6 (23)	99	* *	140	56.4 (41)
Meade	264	6.1 (7)	6.8 (25)	239	12.6 (37)	195	4.1 (28)	272	81.6 (1)
Miami	370	7.6 (22)	5.9 (19)	290	9.7 (12)	260	5.8 (36)	322	72.7 (14)
NEK	780	7.6 (22)	7.1 (31)	726	9.6 (11)	634	3.9 (25)	789	58.0 (39)
Nemaha	89	* *	* *	91	* *	63	* *	109	64.2 (30)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8F
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Infant Health Indicators by Local Agency
Combined 3 Years

Comparison of Infant Health Indicators (2, 3)									
Local Agency	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)	High (5)							
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Neosho	302	7.9 (27)	6.3 (20)	355	12.7 (38)	231	3.5 (20)	407	55.5 (42)
Osborne	908	7.7 (25)	5.4 (11)	793	10.8 (27)	684	3.1 (12)	902	75.5 (8)
Reno	1,608	6.7 (15)	7.0 (29)	1,345	9.7 (12)	1,201	2.7 (5)	1,576	65.3 (29)
Riley	2,282	5.7 (5)	7.8 (38)	1,828	7.2 (2)	1,689	2.5 (4)	2,222	76.7 (6)
Saline	1,283	9.3 (39)	6.5 (21)	1,095	11.0 (30)	957	4.4 (30)	1,285	62.3 (33)
Scott	146	4.8 (1)	8.2 (39)	135	5.2 (1)	124	2.4 (3)	150	71.3 (17)
Sedgwick	11,520	8.4 (34)	6.9 (28)	9,833	8.8 (8)	8,896	4.0 (27)	10,628	66.7 (27)
SEK	155	9.7 (43)	5.2 (10)	141	10.6 (23)	116	8.6 (38)	159	52.8 (43)
Seward	1,270	5.2 (4)	7.1 (31)	1,148	9.7 (12)	998	2.8 (7)	1,298	69.4 (22)
Shawnee	3,902	7.6 (22)	5.6 (12)	3,500	12.4 (35)	2,963	3.6 (22)	3,950	67.6 (26)
Sheridan	24	* * *	* *	18	* *	15	* *	24	* *
Sherman	372	8.1 (30)	5.6 (12)	315	8.3 (5)	289	4.5 (31)	356	73.3 (12)
SKCPH	598	7.9 (27)	7.5 (34)	557	11.0 (30)	456	3.1 (12)	623	75.0 (9)
Stevens	143	4.9 (2)	9.1 (42)	181	8.8 (8)	104	1.0 (1)	217	77.4 (4)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8F
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Infant Health Indicators by Local Agency
Combined 3 Years

Comparison of Infant Health Indicators (2, 3)										
Local Agency	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)							
	Number	% (Rank)	% (Rank)		Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Sumner	393	8.1 (30)	7.6 (36)		324	10.2 (18)	286	4.2 (29)	375	68.5 (24)
SW KS WIC	2,216	6.3 (10)	6.5 (21)		1,943	10.6 (23)	1,704	2.7 (5)	1,656	74.8 (10)
Wichita	42	* * *	* *		36	* *	27	* *	42	* *
Wilson	261	9.6 (42)	5.7 (15)		240	14.2 (39)	205	3.9 (25)	235	63.8 (31)
Wyandotte	5,635	8.0 (29)	6.8 (25)		4,952	10.3 (19)	4,358	3.7 (23)	5,555	70.7 (19)
Unknown Local Agencies	3	* * *	* *		3	* *	3	* *	3	* *
Kansas	54,513	7.4	6.6		47,348	10.0	41,621	3.5	52,595	69.2

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

