

Table 3F
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Racial and Ethnic Distribution by Local Agency

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Local Agency	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Barton	467	71.7	2.6	24.2	0.6	0.9	0.0	0.0
Butler	285	89.8	1.4	7.0	0.7	0.0	1.1	0.0
Chautauqua	43	*	*	*	*	*	*	*
Cherokee	704	83.0	6.7	5.8	1.7	1.3	1.6	0.0
Cloud	193	91.7	1.0	4.1	2.1	0.5	0.5	0.0
Cowley	363	75.5	2.2	19.0	0.8	1.7	0.8	0.0
Crawford	550	86.9	3.6	6.5	1.3	0.5	1.1	0.0
Doniphan	53	*	*	*	*	*	*	*
Douglas	530	67.5	9.4	10.2	7.5	3.0	2.3	0.0
Ford	697	21.8	2.3	75.0	0.3	0.6	0.0	0.0
Franklin	185	94.1	0.0	2.2	2.2	0.5	1.1	0.0
Geary	421	58.0	20.0	15.2	1.0	3.1	2.9	0.0
Graham	44	*	*	*	*	*	*	*
Greeley	12	*	*	*	*	*	*	*
Greenwood	47	*	*	*	*	*	*	*
Harvey	283	70.0	3.2	21.9	1.4	1.4	2.1	0.0
Jefferson	58	*	*	*	*	*	*	*
Johnson	1,990	47.8	13.0	34.0	0.5	3.3	1.5	0.0
Leavenworth	364	69.2	13.2	12.4	0.8	1.6	2.7	0.0
Lincoln	129	90.7	0.0	8.5	0.0	0.0	0.8	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3F
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Racial and Ethnic Distribution by Local Agency

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Local Agency	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Linn	71	*	*	*	*	*	*	*
Lyon	336	62.8	4.8	29.5	0.3	1.2	1.5	0.0
Marion	255	90.2	0.4	7.5	0.8	0.4	0.8	0.0
Marshall	52	*	*	*	*	*	*	*
Meade	97	*	*	*	*	*	*	*
Miami	151	90.1	2.6	4.0	0.7	0.7	2.0	0.0
NEK	319	82.8	4.7	3.8	8.2	0.0	0.6	0.0
Nemaha	38	*	*	*	*	*	*	*
Neosho	166	92.2	1.2	4.2	1.2	0.0	1.2	0.0
Osborne	386	88.1	1.8	6.7	1.0	1.0	1.3	0.0
Reno	578	78.5	2.1	15.9	0.7	1.0	1.7	0.0
Riley	1,046	70.6	10.7	13.7	0.3	4.0	0.8	0.0
Saline	546	72.5	3.7	20.3	0.0	1.6	1.8	0.0
Scott	47	*	*	*	*	*	*	*
Sedgwick	4,551	45.9	18.1	29.4	1.3	3.4	1.9	0.0
SEK	74	*	*	*	*	*	*	*
Seward	428	15.0	2.1	82.0	0.5	0.5	0.0	0.0
Shawnee	1,606	57.0	14.3	23.2	2.6	0.9	1.9	0.0
Sheridan	8	*	*	*	*	*	*	*
Sherman	151	78.8	0.7	18.5	0.7	1.3	0.0	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3F
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Racial and Ethnic Distribution by Local Agency

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Local Agency	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
SKCPH	281	85.8	0.7	12.1	0.0	1.1	0.4	0.0
Stevens	70	*	*	*	*	*	*	*
Sumner	164	87.8	1.2	7.9	1.2	0.6	1.2	0.0
SW KS WIC	850	28.1	1.4	66.2	0.0	3.9	0.4	0.0
Wichita	18	*	*	*	*	*	*	*
Wilson	75	*	*	*	*	*	*	*
Wyandotte	2,100	26.8	29.4	38.3	0.5	4.0	1.1	0.0
Unknown Local Agencies	20	*	*	*	*	*	*	*
Kansas	21,902	57.3	11.2	26.7	1.2	2.3	1.4	0.0
Nation (Prior Year)	1,304,320	41.6	24.1	28.3	1.1	3.2	1.2	0.5

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4F
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Age Distribution by Local Agency

Comparison of Age Distribution ⁽²⁾								
Local Agency	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Barton	467	0.0	6.4	12.6	63.0	16.3	1.7	0.0
Butler	285	0.0	5.6	14.7	63.9	15.1	0.4	0.4
Chautauqua	43	*	*	*	*	*	*	*
Cherokee	704	0.4	8.0	19.6	58.5	12.2	1.3	0.0
Cloud	193	1.6	8.3	14.5	55.4	18.1	2.1	0.0
Cowley	363	0.6	6.6	17.1	62.5	12.9	0.3	0.0
Crawford	550	0.2	6.4	14.9	65.5	12.5	0.5	0.0
Doniphan	53	*	*	*	*	*	*	*
Douglas	530	0.0	4.2	12.5	64.3	17.9	1.1	0.0
Ford	697	0.3	6.0	12.1	53.8	24.8	3.0	0.0
Franklin	185	0.5	5.4	14.1	64.9	14.6	0.5	0.0
Geary	421	0.2	4.0	12.8	70.3	11.4	1.0	0.2
Graham	44	*	*	*	*	*	*	*
Greeley	12	*	*	*	*	*	*	*
Greenwood	47	*	*	*	*	*	*	*
Harvey	283	0.0	4.6	13.4	64.0	17.3	0.7	0.0
Jefferson	58	*	*	*	*	*	*	*
Johnson	1,990	0.1	3.0	10.0	60.9	23.8	2.2	0.1
Leavenworth	364	0.3	5.5	10.2	69.8	13.7	0.5	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4F
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Age Distribution by Local Agency

Comparison of Age Distribution ⁽²⁾								
Local Agency	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Lincoln	129	0.0	4.7	12.4	58.9	20.9	3.1	0.0
Linn	71	*	*	*	*	*	*	*
Lyon	336	0.9	6.0	15.8	60.4	15.8	1.2	0.0
Marion	255	0.4	2.4	14.5	67.8	14.1	0.8	0.0
Marshall	52	*	*	*	*	*	*	*
Meade	97	*	*	*	*	*	*	*
Miami	151	0.7	4.0	18.5	61.6	13.9	1.3	0.0
NEK	319	0.3	6.6	14.4	61.1	16.6	0.9	0.0
Nemaha	38	*	*	*	*	*	*	*
Neosho	166	0.0	5.4	15.1	65.1	13.3	1.2	0.0
Osborne	386	0.5	5.2	15.5	60.1	17.4	0.8	0.5
Reno	578	0.3	6.7	14.0	63.0	15.1	0.9	0.0
Riley	1,046	0.1	1.8	10.9	73.4	13.3	0.5	0.0
Saline	546	0.0	5.5	16.5	61.0	15.9	0.9	0.2
Scott	47	*	*	*	*	*	*	*
Sedgwick	4,551	0.3	6.6	13.6	60.2	18.0	1.2	0.0
SEK	74	*	*	*	*	*	*	*
Seward	428	0.2	7.7	11.0	58.2	21.7	1.2	0.0
Shawnee	1,606	0.2	6.0	14.1	60.6	17.7	1.2	0.1

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4F
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Age Distribution by Local Agency

Comparison of Age Distribution ⁽²⁾								
Local Agency	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Sheridan	8	*	*	*	*	*	*	*
Sherman	151	0.0	4.0	15.2	67.5	12.6	0.7	0.0
SKCPH	281	0.4	6.4	12.1	64.4	15.7	1.1	0.0
Stevens	70	*	*	*	*	*	*	*
Sumner	164	0.0	8.5	13.4	59.8	17.1	1.2	0.0
SW KS WIC	850	0.4	6.9	13.8	57.5	19.1	2.4	0.0
Wichita	18	*	*	*	*	*	*	*
Wilson	75	*	*	*	*	*	*	*
Wyandotte	2,100	0.5	9.0	13.3	55.5	20.1	1.7	0.0
Unknown Local Agencies	20	*	*	*	*	*	*	*
Kansas	21,902	0.3	5.9	13.5	61.3	17.7	1.3	0.0
Nation (Prior Year)	1,304,320	0.3	5.8	12.3	59.4	20.4	1.5	0.3

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5F
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Health Indicators by Local Agency

Comparison of Maternal Health Indicators ^(2, 3)														
Local Agency	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Barton	450	5.3 (25)	58.4 (26)	388	21.6 (17)	51.0 (13)	47	* *	98	* *	401	7.5 (18)	401	13.5(26)
Butler	280	6.1 (29)	53.9 (11)	253	21.3 (15)	57.3 (30)	41	* *	44	* *	260	4.6 (6)	260	11.2(14)
Chautauqua	42	* *	* *	35	* *	* *	5	* *	9	* *	35	* *	35	* *
Cherokee	675	5.5 (27)	52.0 (6)	597	23.8 (24)	50.4 (11)	130	33.8 (5)	144	25.7 (5)	613	8.0 (23)	612	13.4(24)
Cloud	186	6.5 (30)	53.2 (9)	163	22.1 (19)	51.5 (17)	24	* *	49	* *	170	10.6 (30)	171	9.4(8)
Cowley	355	3.4 (10)	54.9 (17)	307	22.1 (19)	50.8 (12)	40	* *	57	* *	315	8.9 (26)	316	9.8(10)
Crawford	544	4.8 (21)	54.2 (13)	463	24.6 (25)	47.5 (8)	63	* *	86	* *	473	8.9 (26)	474	13.3(23)
Doniphan	52	* *	* *	44	* *	* *	10	* *	10	* *	46	* *	46	* *
Douglas	513	2.9 (7)	48.3 (1)	423	19.4 (8)	51.3 (16)	62	* *	130	32.3 (9)	439	7.5 (18)	439	9.6(9)
Ford	687	2.2 (3)	59.1 (27)	595	28.4 (30)	44.4 (4)	84	* *	91	* *	596	6.5 (15)	595	8.6(4)
Franklin	172	5.2 (24)	48.3 (1)	148	18.2 (4)	50.0 (10)	20	* *	48	* *	139	5.0 (8)	139	10.1(12)
Geary	405	4.2 (16)	56.8 (23)	317	20.2 (10)	52.7 (23)	61	* *	64	* *	332	3.9 (3)	331	11.5(16)
Graham	42	* *	* *	32	* *	* *	4	* *	8	* *	34	* *	34	* *
Greeley	12	* *	* *	9	* *	* *	0	* *	1	* *	9	* *	9	* *
Greenwood	45	* *	* *	37	* *	* *	5	* *	9	* *	36	* *	36	* *
Harvey	276	4.7 (20)	59.4 (29)	244	18.9 (7)	51.2 (15)	38	* *	39	* *	251	4.4 (5)	250	11.6(17)
Jefferson	56	* *	* *	48	* *	* *	10	* *	13	* *	50	* *	50	* *

- (1) Reporting period is January 1 through December 31.
 - (2) Excludes records with unknown data or errors.
 - (3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.
 - (4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
 - (5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
 - (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5F
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Health Indicators by Local Agency

Comparison of Maternal Health Indicators ^(2, 3)														
Local Agency	Prepregnancy BMI ⁽⁴⁾			Weight Gain ⁽⁵⁾			Anemia (Low Hb/Hct) ⁽⁶⁾				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Johnson	1,916	3.2 (9)	54.8 (16)	1,651	18.7 (6)	51.0 (13)	281	22.4 (1)	462	17.5 (1)	1,701	7.8 (22)	1,695	6.8(2)
Leavenwort	348	4.0 (13)	56.0 (21)	292	20.5 (11)	56.2 (29)	51	* *	92	* *	304	5.6 (12)	303	13.2(22)
Lincoln	123	0.8 (1)	62.6 (31)	109	30.3 (31)	53.2 (25)	21	* *	19	* *	118	13.6 (31)	117	17.1(29)
Linn	71	* *	* *	63	* *	* *	11	* *	9	* *	63	* *	64	* *
Lyon	328	2.4 (5)	56.7 (22)	279	24.7 (26)	45.5 (7)	39	* *	62	* *	285	6.3 (14)	285	13.7(27)
Marion	246	1.2 (2)	61.8 (30)	204	23.0 (21)	44.6 (5)	32	* *	43	* *	216	4.6 (6)	217	11.1(13)
Marshall	51	* *	* *	49	* *	* *	5	* *	12	* *	49	* *	48	* *
Meade	91	* *	* *	77	* *	* *	9	* *	20	* *	80	* *	80	* *
Miami	147	3.4 (10)	55.1 (18)	125	13.6 (1)	58.4 (31)	30	* *	34	* *	132	6.8 (17)	132	11.4(15)
NEK	312	2.2 (3)	53.8 (10)	270	20.7 (13)	55.2 (26)	38	* *	59	* *	258	10.5 (29)	260	15.0(28)
Nemaha	33	* *	* *	30	* *	* *	10	* *	6	* *	34	* *	34	* *
Neosho	164	5.5 (27)	54.3 (15)	147	23.1 (23)	51.7 (20)	26	* *	27	* *	148	6.1 (13)	147	18.4(31)
Osborne	379	5.3 (25)	50.9 (4)	327	22.0 (18)	44.3 (3)	63	* *	68	* *	337	3.9 (3)	337	8.9(5)
Reno	563	3.9 (12)	55.4 (20)	495	21.4 (16)	51.7 (20)	76	* *	103	21.4 (2)	505	5.3 (10)	507	13.4(24)
Riley	996	4.5 (18)	52.7 (7)	798	18.4 (5)	52.9 (24)	156	37.8 (6)	260	30.0 (8)	824	5.2 (9)	825	12.1(21)
Saline	538	5.0 (23)	57.4 (25)	467	16.5 (2)	55.9 (28)	96	* *	77	* *	471	8.7 (25)	470	9.1(6)
Scott	46	* *	* *	37	* *	* *	4	* *	8	* *	38	* *	38	* *

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.
- (4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
- (5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
- (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5F
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Health Indicators by Local Agency

Comparison of Maternal Health Indicators ^(2, 3)														
Local Agency	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Sedgwick	4,426	4.1 (14)	54.2 (13)	3,947	20.6 (12)	51.5 (17)	834	29.4 (3)	986	23.5 (4)	4,070	6.5 (15)	4,063	11.6(17)
SEK	72	* *	* *	67	* *	* *	11	* *	19	* *	69	* *	69	* *
Seward	423	2.4 (5)	59.3 (28)	374	23.0 (21)	44.9 (6)	63	* *	69	* *	380	8.2 (24)	378	6.3(1)
Shawnee	1,562	4.9 (22)	54.0 (12)	1,334	20.1 (9)	52.5 (22)	150	46.0 (7)	263	28.1 (7)	1,373	7.7 (21)	1,373	9.3(7)
Sheridan	8	* *	* *	6	* *	* *	0	* *	1	* *	7	* *	7	* *
Sherman	146	4.1 (14)	50.7 (3)	124	16.9 (3)	55.6 (27)	18	* *	26	* *	129	2.3 (1)	129	11.6(17)
SKCPH	277	2.9 (7)	52.7 (7)	240	25.4 (27)	47.9 (9)	32	* *	46	* *	244	3.7 (2)	244	11.9(20)
Stevens	69	* *	* *	60	* *	* *	15	* *	17	* *	60	* *	60	* *
Sumner	161	7.5 (31)	51.6 (5)	144	26.4 (29)	40.3 (1)	23	* *	43	* *	146	7.5 (18)	147	17.7(30)
SW KS WIC	834	4.4 (17)	55.2 (19)	750	26.1 (28)	42.0 (2)	112	25.9 (2)	129	23.3 (3)	773	9.6 (28)	772	7.0(3)
Wichita	16	* *	* *	15	* *	* *	2	* *	0	* *	17	* *	17	* *
Wilson	74	* *	* *	69	* *	* *	13	* *	10	* *	69	* *	69	* *
Wyandotte	2,037	4.5 (18)	57.1 (24)	1,785	20.8 (14)	51.5 (17)	310	31.0 (4)	390	27.4 (6)	1,811	5.4 (11)	1,809	9.8(10)
Unknown Local Agencies	18	* *	* *	17	* *	* *	1	* *	2	* *	19	* *	19	* *
Kansas	21,267	4.1	55.0	18,455	21.1	50.7	3,176	29.0	4,262	23.7	18,929	6.8	18,913	10.7
Nation (Prior Year)	1,226,875	4.5	52.9	917,032	21.2	48.2	189,944	33.9	412,633	28.8	389,161	5.0	394,368	6.8

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6F
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Behavioral Indicators by Local Agency

Comparison of Maternal Behavioral Indicators ^(2, 3)										
Local Agency	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Prepregnancy ⁽⁴⁾		During Pregnancy ⁽⁵⁾	
	Number	% (Rank)	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Barton	399	86.5 (6)	0.8 (1)		464	47.2 (2)	467	21.4 (12)	375	56.8 (22)
Butler	255	74.9 (23)	3.1 (12)		285	33.3 (24)	285	14.0 (17)	224	37.9 (31)
Chautauqua	35	* *	* *		43	* *	43	* *	37	* *
Cherokee	462	83.8 (11)	3.5 (14)		703	29.3 (29)	704	12.8 (23)	581	65.2 (11)
Cloud	172	86.6 (5)	3.5 (14)		191	42.9 (6)	193	27.5 (1)	169	75.7 (1)
Cowley	297	72.4 (25)	6.4 (25)		360	41.4 (10)	363	13.2 (22)	303	56.1 (23)
Crawford	437	88.6 (3)	3.7 (17)		550	39.6 (15)	550	14.0 (17)	443	58.9 (18)
Doniphan	47	* *	* *		53	* *	53	* *	41	* *
Douglas	443	76.1 (20)	5.6 (22)		528	38.6 (18)	530	21.5 (11)	434	64.5 (14)
Ford	533	72.2 (26)	12.4 (28)		695	44.7 (5)	697	13.9 (19)	591	52.1 (28)
Franklin	76	* *	* *		185	31.4 (26)	185	8.6 (28)	127	63.8 (15)
Geary	317	76.0 (21)	1.6 (8)		421	30.2 (28)	421	22.1 (10)	342	58.5 (20)
Graham	35	* *	* *		44	* *	44	* *	37	* *
Greeley	8	* *	* *		12	* *	12	* *	10	* *
Greenwood	37	* *	* *		47	* *	47	* *	43	* *
Harvey	220	73.6 (24)	11.4 (27)		281	48.4 (1)	283	19.4 (13)	235	66.0 (9)
Jefferson	48	* *	* *		58	* *	58	* *	49	* *
Johnson	1,518	65.5 (28)	4.0 (19)		1,983	34.1 (23)	1,990	13.5 (20)	1,598	60.8 (17)
Leavenworth	307	83.7 (12)	1.0 (3)		364	28.0 (31)	364	22.3 (8)	273	68.5 (6)
Lincoln	117	85.5 (8)	2.6 (9)		129	41.9 (9)	129	26.4 (2)	112	58.9 (18)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6F
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas

Comparison of Maternal Behavioral Indicators by Local Agency

Comparison of Maternal Behavioral Indicators ^(2, 3)										
Local Agency	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Linn	65	* *	* *	* *	71	* *	71	* *	60	* *
Lyon	286	84.3 (10)	1.0 (3)		334	39.8 (13)	336	18.2 (15)	277	65.0 (13)
Marion	213	86.4 (7)	1.4 (6)		251	40.2 (12)	255	24.3 (5)	205	68.3 (7)
Marshall	50	* *	* *	* *	52	* *	52	* *	43	* *
Meade	80	* *	* *	* *	97	* *	97	* *	92	* *
Miami	131	79.4 (16)	1.5 (7)		151	37.7 (20)	151	18.5 (14)	127	55.1 (25)
NEK	211	84.8 (9)	4.7 (20)		318	40.9 (11)	319	12.2 (26)	270	65.2 (11)
Nemaha	33	* *	* *	* *	38	* *	38	* *	32	* *
Neosho	133	88.7 (2)	3.8 (18)		165	39.4 (16)	166	12.7 (25)	139	71.9 (3)
Osborne	336	87.5 (4)	3.0 (11)		384	42.4 (7)	386	24.9 (4)	328	55.2 (24)
Reno	497	83.5 (13)	2.8 (10)		576	39.1 (17)	578	22.3 (8)	488	65.8 (10)
Riley	770	82.3 (14)	1.3 (5)		1,046	33.2 (25)	1,046	22.6 (7)	825	70.2 (4)
Saline	402	76.4 (19)	3.2 (13)		544	37.9 (19)	546	4.6 (31)	478	54.6 (26)
Scott	37	* *	* *	* *	46	* *	47	* *	42	* *
Sedgwick	3,982	75.1 (22)	5.9 (23)		4,538	28.6 (30)	4,551	18.2 (15)	3,758	50.4 (29)
SEK	59	* *	* *	* *	74	* *	74	* *	65	* *
Seward	336	66.4 (27)	13.7 (29)		423	30.7 (27)	428	8.4 (29)	345	67.0 (8)
Shawnee	1,205	76.5 (18)	3.5 (14)		1,599	45.2 (3)	1,606	13.4 (21)	1,328	53.1 (27)
Sheridan	7	* *	* *	* *	8	* *	8	* *	8	* *
Sherman	114	89.5 (1)	0.9 (2)		151	39.7 (14)	151	25.2 (3)	120	73.3 (2)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6F
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Behavioral Indicators by Local Agency

Comparison of Maternal Behavioral Indicators ^(2, 3)													
Local Agency	Medical Care				WIC Enrollment			Multivitamin Consumption					
	1st Trimester		No Care		1st Trimester			Pregnancy (4)		During Pregnancy (5)			
	Number	% (Rank)	%	% (Rank)	Number	%	% (Rank)	Number	%	% (Rank)			
SKCPH	241	80.9 (15)	6.2	(24)	278	45.0	(4)	281	24.2	(6)	230	70.0	(5)
Stevens	60	* *	* *	*	69	* *	*	70	* *	*	60	* *	*
Sumner	147	78.2 (17)	4.8	(21)	164	42.1	(8)	164	12.8	(23)	136	62.5	(16)
SW KS WIC	734	53.5 (30)	28.2	(30)	839	34.3	(22)	850	12.1	(27)	709	48.0	(30)
Wichita	17	* *	* *	*	18	* *	*	18	* *	*	13	* *	*
Wilson	68	* *	* *	*	74	* *	*	75	* *	*	63	* *	*
Wyandotte	1,726	64.1 (29)	8.6	(26)	2,082	36.0	(21)	2,100	6.6	(30)	1,778	57.9	(21)
Unknown Local Agencies	17	* *	* *	*	20	* *	*	20	* *	*	17	* *	*
Kansas	17,720	75.1	6.1		21,806	36.1		21,902	16.2		18,060	58.1	
Nation (Prior Year)	527,184	82.2	3.1		1,213,982	34.1		360,369	18.7		372,743	83.5	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7F
2010 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by Local Agency

Comparison of Smoking Indicators (2, 3)										
Local Agency	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Barton	451	33.9 (11)	384	22.4 (17)	101	29.7 (8)	367	14.2 (13)	385	8.1 (9)
Butler	281	51.6 (30)	253	31.6 (28)	97	* *	223	13.0 (12)	258	9.7 (12)
Chautauqua	39	* *	27	* *	11	* *	34	* *	27	* *
Cherokee	676	41.4 (23)	585	27.5 (23)	192	22.9 (12)	571	36.1 (28)	586	35.3 (27)
Cloud	193	35.2 (13)	168	20.2 (12)	44	* *	168	15.5 (17)	168	10.1 (14)
Cowley	355	40.8 (22)	278	28.1 (24)	88	* *	297	19.5 (20)	278	18.0 (22)
Crawford	517	48.4 (29)	400	30.8 (27)	154	29.2 (10)	425	39.8 (30)	406	36.7 (28)
Doniphan	52	* *	37	* *	8	* *	40	* *	37	* *
Douglas	524	34.9 (12)	431	17.2 (8)	117	40.2 (3)	429	10.3 (6)	429	6.3 (3)
Ford	671	8.0 (2)	538	4.3 (2)	32	* *	581	8.4 (2)	540	8.0 (8)
Franklin	184	40.2 (20)	156	28.8 (26)	34	* *	126	15.1 (15)	157	10.8 (18)
Geary	415	35.2 (13)	328	21.6 (15)	77	* *	338	35.2 (27)	328	34.5 (26)
Graham	44	* *	34	* *	11	* *	37	* *	34	* *
Greeley	12	* *	8	* *	1	* *	10	* *	8	* *
Greenwood	39	* *	13	* *	7	* *	39	* *	21	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7F
2010 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by Local Agency

Comparison of Smoking Indicators (2, 3)										
Local Agency	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Harvey	280	40.0 (18)	243	21.0 (14)	78	* *	233	5.6 (1)	242	1.7 (1)
Jefferson	15	* *	11	* *	3	* *	13	* *	12	* *
Johnson	1,963	23.9 (5)	1,697	10.4 (4)	278	45.3 (1)	1,573	17.9 (19)	1,700	17.4 (20)
Leavenworth	361	47.6 (28)	301	23.9 (19)	98	* *	271	38.4 (29)	299	38.8 (29)
Lincoln	128	45.3 (26)	100	32.0 (29)	42	* *	112	19.6 (21)	111	15.3 (19)
Linn	69	* *	63	* *	37	* *	59	* *	63	* *
Lyon	333	32.7 (8)	290	18.3 (10)	75	* *	275	12.4 (11)	290	7.9 (6)
Marion	251	37.8 (15)	215	23.7 (18)	68	* *	202	9.9 (4)	215	7.9 (6)
Marshall	51	* *	45	* *	21	* *	41	* *	45	* *
Meade	92	* *	75	* *	9	* *	91	* *	76	* *
Miami	143	39.9 (17)	121	19.8 (11)	36	* *	119	24.4 (22)	121	17.4 (20)
NEK	290	46.9 (27)	217	28.1 (24)	79	* *	245	29.4 (24)	221	23.5 (24)
Nemaha	31	* *	20	* *	5	* *	28	* *	21	* *
Neosho	139	43.9 (25)	74	* *	27	* *	126	34.1 (25)	76	* *
Osborne	376	43.4 (24)	314	24.5 (20)	120	31.7 (7)	318	27.0 (23)	319	23.2 (23)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7F
2010 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by Local Agency

Comparison of Smoking Indicators (2, 3)										
Local Agency	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Reno	573	40.0 (18)	502	20.9 (13)	170	41.2 (2)	483	11.8 (10)	502	9.0 (11)
Riley	997	29.1 (6)	696	14.1 (6)	148	37.8 (4)	806	15.3 (16)	699	9.9 (13)
Saline	543	39.4 (16)	470	25.3 (22)	165	29.7 (8)	478	10.9 (7)	469	7.7 (5)
Scott	39	* *	23	* *	8	* *	35	* *	24	* *
Sedgwick	4,518	29.2 (7)	4,065	16.1 (7)	911	33.3 (5)	3,742	9.2 (3)	4,055	6.6 (4)
SEK	71	* *	61	* *	21	* *	63	* *	63	* *
Seward	422	5.7 (1)	377	3.2 (1)	17	* *	342	10.2 (5)	382	8.9 (10)
Shawnee	1,479	33.3 (9)	1,152	22.0 (16)	289	27.7 (11)	1,195	11.1 (9)	1,165	5.2 (2)
Sheridan	8	* *	6	* *	2	* *	8	* *	6	* *
Sherman	148	33.8 (10)	128	18.0 (9)	36	* *	118	16.9 (18)	129	10.1 (14)
SKCPH	268	40.7 (21)	227	24.7 (21)	71	* *	225	35.1 (26)	225	32.4 (25)
Stevens	70	* *	60	* *	4	* *	60	* *	60	* *
Sumner	163	54.6 (31)	144	34.0 (30)	63	* *	135	44.4 (31)	144	44.4 (30)
SW KS WIC	843	9.6 (3)	755	5.2 (3)	52	* *	706	11.0 (8)	755	10.3 (16)
Wichita	17	* *	16	* *	4	* *	12	* *	16	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7F
2010 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by Local Agency

Comparison of Smoking Indicators (2, 3)										
Local Agency	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Wilson	74	* *	69	* *	35	* *	63	* *	69	* *
Wyandotte	2,056	20.6 (4)	1,754	10.9 (5)	271	32.5 (6)	1,761	14.7 (14)	1,807	10.6 (17)
Unknown Local Agencies	19	* *	19	* *	10	* *	17	* *	20	* *
Kansas	21,283	30.7	17,950	17.5	4,227	33.0	17,640	16.5	18,063	13.3
Nation (Prior Year)	1,040,568	22.9	838,781	13.0	156,709	36.7	650,648	17.1	559,995	16.8

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8F
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Infant Health Indicators by Local Agency

Comparison of Infant Health Indicators ^(2, 3)										
Local Agency	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)							
	Number	% (Rank)	%	(Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Barton	383	5.7 (6)	5.2 (6)		387	6.2 (3)	356	4.2 (15)	390	75.6 (12)
Butler	252	7.1 (13)	6.0 (14)		257	10.1 (28)	226	3.5 (9)	256	65.2 (25)
Chautauqua	35	* * * *	* *		37	* *	34	* *	37	* *
Cherokee	608	7.6 (18)	5.3 (7)		613	8.0 (14)	558	3.8 (13)	611	58.6 (29)
Cloud	166	8.4 (26)	6.0 (14)		169	8.9 (21)	150	4.7 (24)	168	75.0 (13)
Cowley	302	6.6 (11)	7.9 (24)		317	10.7 (30)	267	3.7 (12)	317	56.8 (31)
Crawford	449	7.6 (18)	2.7 (1)		461	10.4 (29)	401	5.2 (27)	459	63.4 (26)
Doniphan	48	* * * *	* *		48	* *	39	* *	48	* *
Douglas	435	8.3 (25)	8.5 (27)		441	7.7 (12)	400	5.3 (28)	438	79.5 (4)
Ford	564	5.0 (2)	6.0 (14)		580	7.1 (6)	523	2.9 (6)	575	77.4 (8)
Franklin	153	5.2 (3)	5.9 (13)		153	5.2 (2)	145	4.8 (26)	153	60.8 (27)
Geary	326	6.4 (8)	8.3 (25)		327	8.9 (21)	297	2.0 (3)	324	71.0 (18)
Graham	32	* * * *	* *		32	* *	26	* *	31	* *
Greeley	9	* * * *	* *		9	* *	7	* *	9	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8F
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Infant Health Indicators by Local Agency

Comparison of Infant Health Indicators ^(2, 3)										
Local Agency	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)							
	Number	% (Rank)	%	(Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Greenwood	39	* * * *	*	*	39	* *	37	* *	39	* *
Harvey	237	8.4 (26)	4.2	(3)	240	6.7 (4)	219	4.6 (21)	241	83.4 (2)
Jefferson	49	* * * *	*	*	49	* *	46	* *	48	* *
Johnson	1,601	7.1 (13)	6.7	(21)	1,690	7.3 (7)	1,476	3.5 (9)	1,683	81.8 (3)
Leavenworth	300	5.3 (4)	8.7	(29)	304	9.2 (24)	272	2.6 (5)	302	66.2 (23)
Lincoln	112	8.0 (21)	3.6	(2)	118	7.6 (11)	103	2.9 (6)	117	70.9 (19)
Linn	58	* * * *	*	*	62	* *	50	* *	62	* *
Lyon	285	7.7 (20)	6.0	(14)	283	7.4 (10)	262	4.6 (21)	285	75.8 (11)
Marion	205	5.9 (7)	8.3	(25)	207	9.7 (27)	181	1.7 (2)	211	77.7 (7)
Marshall	45	* * * *	*	*	48	* *	40	* *	47	* *
Meade	67	* * * *	*	*	83	* *	62	* *	83	* *
Miami	132	3.0 (1)	10.6	(31)	132	4.5 (1)	126	0.8 (1)	132	75.0 (13)
NEK	258	8.9 (29)	8.5	(27)	275	8.7 (20)	233	6.4 (30)	273	57.5 (30)
Nemaha	15	* * * *	*	*	33	* *	13	* *	29	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8F
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Infant Health Indicators by Local Agency

Comparison of Infant Health Indicators ^(2, 3)									
Local Agency	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Neosho	130	8.5 (28)	5.4 (9)	144	8.3 (16)	117	4.3 (18)	143	59.4 (28)
Osborne	318	7.2 (17)	5.3 (7)	323	9.0 (23)	287	3.1 (8)	322	74.8 (15)
Reno	493	8.1 (22)	6.5 (19)	502	8.4 (17)	451	4.4 (19)	500	71.8 (17)
Riley	824	5.6 (5)	9.3 (30)	841	6.8 (5)	767	2.5 (4)	834	78.5 (6)
Saline	471	6.4 (8)	5.5 (10)	476	8.0 (14)	432	4.2 (15)	475	65.9 (24)
Scott	36	* * *	* *	36	* *	31	* *	36	* *
Sedgwick	3,976	9.6 (30)	7.1 (22)	4,005	8.5 (19)	3,627	5.7 (29)	3,993	70.4 (20)
SEK	58	* * *	* *	69	* *	54	* *	68	* *
Seward	346	6.4 (8)	6.6 (20)	368	7.9 (13)	313	3.8 (13)	370	78.9 (5)
Shawnee	1,335	7.1 (13)	6.4 (18)	1,354	9.5 (25)	1,203	4.2 (15)	1,346	69.8 (22)
Sheridan	7	* * *	* *	7	* *	6	* *	7	* *
Sherman	123	8.1 (22)	4.9 (4)	123	7.3 (7)	114	4.4 (19)	123	85.4 (1)
SKCPH	212	7.1 (13)	7.5 (23)	232	7.3 (7)	193	3.6 (11)	235	77.0 (9)
Stevens	39	* * *	* *	59	* *	38	* *	58	* *

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.
- (4) <2500 g.
- (5) >4000 g.
- (6) <37 weeks gestation.
- (7) 37 or more weeks gestation and <2500 g.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8F
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Infant Health Indicators by Local Agency

Comparison of Infant Health Indicators ^(2, 3)										
Local Agency	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)							
	Number	% (Rank)	%	(Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Sumner	141	15.6 (31)	5.0	(5)	145	14.5 (31)	120	9.2 (31)	144	70.1 (21)
SW KS WIC	731	7.0 (12)	5.7	(11)	746	8.4 (17)	663	4.7 (24)	749	77.0 (9)
Wichita	15	* * *	*	*	15	* *	14	* *	15	* *
Wilson	71	* * *	*	*	71	* *	63	* *	71	* *
Wyandotte	1,715	8.2 (24)	5.7	(11)	1,796	9.6 (26)	1,541	4.6 (21)	1,794	71.9 (16)
Unknown Local Agencies	20	* * *	*	*	20	* *	20	* *	19	* *
Kansas	18,226	7.6	6.6		18,726	8.4	16,603	4.4	18,670	72.2
Nation (Prior Year)	1,018,145	8.2	6.7		1,017,333	11.1	848,957	3.5	997,136	66.7

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3F
2008 - 2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Racial and Ethnic Distribution by Local Agency
Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Local Agency	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Barton	1,370	71.1	2.0	25.6	0.4	0.5	0.4	0.0
Butler	853	90.3	1.2	6.1	0.5	0.2	1.8	0.0
Chautauqua	140	92.9	0.7	3.6	0.7	0.0	2.1	0.0
Cherokee	2,169	83.6	6.1	6.1	2.0	1.1	1.2	0.0
Cloud	608	91.6	1.0	4.8	1.3	0.5	0.8	0.0
Cowley	1,082	74.9	2.5	19.3	0.8	1.7	0.8	0.0
Crawford	1,788	86.7	3.3	7.2	0.6	0.8	1.3	0.0
Doniphan	133	88.0	2.3	6.8	0.8	0.0	2.3	0.0
Douglas	1,500	65.5	8.7	10.7	8.5	4.5	2.1	0.0
Ford	2,029	21.8	1.5	76.0	0.2	0.4	0.0	0.0
Franklin	538	91.4	0.9	4.5	1.5	0.2	1.5	0.0
Geary	1,154	53.9	22.7	15.3	0.6	3.2	4.3	0.0
Graham	126	89.7	2.4	7.1	0.0	0.0	0.8	0.0
Greeley	37	*	*	*	*	*	*	*
Greenwood	140	93.6	0.7	4.3	0.0	0.0	1.4	0.0
Harvey	818	70.3	2.4	23.7	1.2	0.7	1.6	0.0
Jefferson	191	94.2	0.5	2.1	2.6	0.0	0.5	0.0
Johnson	5,750	46.4	13.0	35.3	0.5	3.2	1.5	0.0
Leavenworth	1,079	69.0	14.6	10.6	0.6	1.7	3.7	0.0
Lincoln	403	92.1	0.2	6.9	0.0	0.0	0.7	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3F
2008 - 2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Racial and Ethnic Distribution by Local Agency
Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Local Agency	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Linn	191	95.8	0.0	2.1	1.0	0.0	1.0	0.0
Lyon	1,082	61.1	4.7	31.3	0.4	1.4	1.1	0.0
Marion	642	90.8	0.2	6.9	0.9	0.5	0.8	0.0
Marshall	163	92.0	0.0	2.5	3.1	0.0	2.5	0.0
Meade	328	69.8	0.3	29.0	0.0	0.0	0.9	0.0
Miami	433	91.0	2.8	3.5	0.2	0.9	1.6	0.0
NEK	946	81.9	4.0	3.4	8.7	0.0	2.0	0.0
Nemaha	118	89.0	0.0	8.5	0.0	0.8	1.7	0.0
Neosho	465	91.6	2.4	4.1	0.6	0.2	1.1	0.0
Osborne	1,093	88.7	1.6	7.6	0.9	0.8	0.5	0.0
Reno	1,860	81.4	2.1	14.0	0.6	0.6	1.2	0.0
Riley	2,903	72.3	10.9	12.1	0.5	3.6	0.5	0.0
Saline	1,570	71.5	4.2	20.0	0.2	2.2	1.9	0.0
Scott	160	66.9	0.6	31.3	1.3	0.0	0.0	0.0
Sedgwick	13,421	45.2	18.2	29.6	1.5	3.7	1.8	0.0
SEK	184	94.6	0.0	3.3	1.1	1.1	0.0	0.0
Seward	1,456	15.4	1.6	80.6	1.5	0.7	0.2	0.0
Shawnee	4,694	57.8	14.4	23.2	2.2	0.8	1.7	0.0
Sheridan	25	*	*	*	*	*	*	*
Sherman	445	80.7	0.7	16.6	0.2	0.9	0.9	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3F
2008 - 2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Racial and Ethnic Distribution by Local Agency
Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Local Agency	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
SKCPH	835	85.4	0.6	12.6	0.4	0.7	0.4	0.0
Stevens	225	45.3	0.0	52.9	1.3	0.4	0.0	0.0
Sumner	479	88.9	0.8	6.7	1.5	0.4	1.7	0.0
SW KS WIC	2,565	29.6	0.9	66.3	0.0	2.8	0.4	0.0
Wichita	47	*	*	*	*	*	*	*
Wilson	257	96.1	0.8	1.9	0.4	0.0	0.8	0.0
Wyandotte	6,498	26.7	30.1	39.4	0.3	2.8	0.6	0.0
Unknown Local Agencies	23	*	*	*	*	*	*	*
Kansas	65,016	56.9	11.3	27.2	1.2	2.1	1.3	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4F
2008 - 2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Age Distribution by Local Agency
Combined 3 Years

Comparison of Age Distribution ⁽²⁾								
Local Agency	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Barton	1,370	0.2	6.8	14.5	61.7	15.6	1.2	0.0
Butler	853	0.1	6.8	16.9	61.9	13.6	0.5	0.2
Chautauqua	140	0.0	5.0	20.7	60.0	12.9	0.7	0.7
Cherokee	2,169	0.2	7.0	18.8	61.3	11.7	0.8	0.2
Cloud	608	0.7	6.7	16.9	58.9	15.0	1.8	0.0
Cowley	1,082	0.6	7.9	16.4	61.6	13.0	0.5	0.0
Crawford	1,788	0.2	7.0	15.8	63.8	12.6	0.5	0.1
Doniphan	133	0.8	4.5	13.5	63.2	18.0	0.0	0.0
Douglas	1,500	0.0	4.1	12.5	63.7	18.4	1.4	0.0
Ford	2,029	0.3	7.0	13.0	54.2	23.5	1.9	0.0
Franklin	538	0.4	5.9	18.0	60.0	14.7	0.7	0.2
Geary	1,154	0.3	4.8	14.9	67.9	11.4	0.6	0.1
Graham	126	0.0	4.0	19.8	61.9	13.5	0.8	0.0
Greeley	37	*	*	*	*	*	*	*
Greenwood	140	0.0	6.4	17.1	62.9	13.6	0.0	0.0
Harvey	818	0.2	5.9	13.4	63.4	15.8	1.2	0.0
Jefferson	191	0.0	5.2	18.8	58.6	16.8	0.5	0.0
Johnson	5,750	0.2	3.8	10.9	60.4	23.1	1.6	0.1
Leavenworth	1,079	0.1	6.0	15.6	64.7	12.5	1.1	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4F
2008 - 2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Age Distribution by Local Agency
Combined 3 Years

Comparison of Age Distribution ⁽²⁾								
Local Agency	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Lincoln	403	0.0	3.0	15.1	60.8	19.9	1.2	0.0
Linn	191	0.0	6.3	15.7	61.8	15.7	0.5	0.0
Lyon	1,082	0.6	6.3	14.8	61.6	14.8	2.0	0.0
Marion	642	0.3	4.8	14.5	64.8	14.5	1.1	0.0
Marshall	163	0.0	6.1	12.9	68.1	12.3	0.6	0.0
Meade	328	0.0	7.0	9.5	61.3	20.7	1.5	0.0
Miami	433	0.5	5.3	15.7	64.0	13.9	0.7	0.0
NEK	946	0.1	6.8	15.8	63.0	13.6	0.7	0.0
Nemaha	118	0.0	3.4	12.7	65.3	13.6	4.2	0.8
Neosho	465	0.2	6.2	18.3	61.9	12.5	0.9	0.0
Osborne	1,093	0.3	4.1	14.6	64.3	15.1	1.4	0.2
Reno	1,860	0.2	6.2	14.9	64.1	13.9	0.7	0.0
Riley	2,903	0.0	1.9	11.9	73.4	12.3	0.5	0.0
Saline	1,570	0.1	6.7	15.9	62.4	14.0	1.0	0.1
Scott	160	0.6	3.1	11.3	66.9	15.6	2.5	0.0
Sedgwick	13,421	0.3	6.8	13.9	60.7	17.1	1.3	0.0
SEK	184	0.5	4.3	15.8	63.6	15.8	0.0	0.0
Seward	1,456	0.2	7.3	13.6	56.6	20.9	1.4	0.0
Shawnee	4,694	0.3	6.4	14.9	60.9	16.2	1.2	0.1

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4F
2008 - 2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Age Distribution by Local Agency
Combined 3 Years

Comparison of Age Distribution ⁽²⁾								
Local Agency	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Sheridan	25	*	*	*	*	*	*	*
Sherman	445	0.0	6.1	15.1	63.1	14.2	1.6	0.0
SKCPH	835	0.2	6.7	14.4	64.6	13.4	0.7	0.0
Stevens	225	0.0	7.1	18.2	57.3	16.4	0.9	0.0
Sumner	479	0.0	6.7	16.7	61.2	14.6	0.8	0.0
SW KS WIC	2,565	0.3	8.3	15.0	56.1	18.4	1.8	0.1
Wichita	47	*	*	*	*	*	*	*
Wilson	257	0.4	8.2	15.6	62.3	13.6	0.0	0.0
Wyandotte	6,498	0.5	8.4	13.5	57.2	19.0	1.4	0.0
Unknown Local Agencies	23	*	*	*	*	*	*	*
Kansas	65,016	0.3	6.2	14.2	61.3	16.7	1.2	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5F
2008 - 2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Health Indicators by Local Agency
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)														
Local Agency	Prepregnancy BMI ⁽⁴⁾			Weight Gain ⁽⁵⁾			Anemia (Low Hb/Hct) ⁽⁶⁾				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Barton	1,335	4.3 (18)	57.3 (34)	1,172	21.6 (29)	49.0 (10)	137	19.0 (2)	276	20.3 (8)	1,139	5.9 (27)	1,136	10.5(28)
Butler	826	5.9 (40)	51.0 (4)	755	19.7 (19)	53.4 (29)	114	43.9 (23)	188	29.3(26)	723	5.4 (21)	723	11.3(33)
Chautauqua	138	5.1 (32)	58.7 (37)	124	18.5 (12)	50.0 (16)	12	* *	24	* *	120	5.0 (16)	120	10.0(23)
Cherokee	2,105	6.2 (42)	53.8 (15)	1,882	22.4 (35)	49.9 (14)	348	30.7 (16)	488	25.0(18)	1,832	5.5 (22)	1,832	10.7(29)
Cloud	595	4.9 (28)	54.6 (21)	528	22.3 (34)	52.8 (26)	72	* *	150	25.3(20)	515	6.2 (33)	519	9.2(16)
Cowley	1,066	4.5 (20)	54.7 (22)	952	20.7 (22)	52.4 (24)	132	32.6 (19)	160	22.5(14)	917	5.9 (27)	919	9.9(21)
Crawford	1,758	5.2 (33)	55.2 (23)	1,543	20.9 (24)	52.1 (23)	248	20.6 (4)	323	20.7 (9)	1,510	7.1 (39)	1,510	11.9(36)
Doniphan	130	0.8 (1)	62.3 (43)	110	13.6 (1)	65.5 (42)	23	* *	25	* *	107	5.6 (23)	107	12.1(38)
Douglas	1,463	3.8 (14)	48.4 (1)	1,253	17.6 (8)	54.7 (35)	199	26.6 (10)	384	26.8(24)	1,218	6.0 (29)	1,217	7.8(11)
Ford	1,999	2.2 (4)	57.5 (35)	1,777	23.7 (38)	46.7 (6)	254	21.3 (5)	230	13.5 (1)	1,729	5.3 (20)	1,727	7.6(9)
Franklin	516	3.7 (11)	51.6 (7)	453	16.8 (5)	53.0 (27)	52	* *	136	14.0 (2)	386	4.1 (6)	386	9.8(19)
Geary	1,119	4.6 (25)	53.4 (12)	900	18.0 (11)	53.7 (30)	171	39.2 (21)	197	34.0(28)	902	4.1 (6)	901	10.7(29)
Graham	119	5.0 (31)	48.7 (2)	99	* *	* *	15	* *	20	* *	106	4.7 (13)	106	8.5(13)
Greeley	35	* *	* *	31	* *	* *	3	* *	5	* *	33	* *	33	* *
Greenwood	135	2.2 (4)	60.0 (40)	123	22.0 (31)	55.3 (39)	13	* *	21	* *	120	5.0 (16)	121	18.2(44)
Harvey	803	4.5 (20)	54.5 (20)	713	22.6 (36)	48.0 (8)	108	36.1 (20)	122	21.3(12)	689	4.6 (12)	686	10.3(26)
Jefferson	188	3.7 (11)	58.0 (36)	169	16.0 (3)	58.0 (41)	31	* *	51	* *	165	6.7 (35)	165	9.1(14)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5F
2008 - 2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Health Indicators by Local Agency
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)														
Local Agency	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight	Over- weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
							Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)
Johnson	5,547	4.2 (15)	54.0 (16)	4,944	18.8 (14)	50.9 (20)	902	24.2 (7)	1,454	17.4 (4)	4,815	6.8 (37)	4,806	6.2(5)
Leavenwort	1,046	4.2 (15)	55.4 (26)	911	19.5 (17)	55.3 (39)	167	26.9 (11)	266	21.8(13)	860	5.6 (23)	859	9.7(18)
Lincoln	391	2.8 (7)	61.4 (42)	366	25.7 (41)	54.1 (33)	53	* *	89	* *	355	9.0 (43)	354	11.6(34)
Linn	189	5.8 (38)	62.4 (44)	173	28.9 (42)	46.2 (4)	24	* *	29	* *	169	4.7 (13)	170	5.3(1)
Lyon	1,051	3.2 (9)	59.0 (38)	925	21.0 (26)	49.8 (13)	139	10.8 (1)	195	20.0 (6)	895	5.1 (18)	895	12.7(39)
Marion	622	4.2 (15)	54.2 (19)	540	21.5 (28)	49.3 (11)	81	* *	134	23.9(15)	536	4.5 (9)	533	10.1(24)
Marshall	159	5.7 (36)	53.5 (13)	144	20.8 (23)	50.0 (16)	21	* *	35	* *	137	5.1 (18)	136	9.6(17)
Meade	320	2.8 (7)	56.9 (33)	279	22.2 (32)	50.2 (18)	35	* *	72	* *	269	6.7 (35)	269	11.2(32)
Miami	419	6.0 (41)	55.6 (28)	375	16.0 (3)	55.2 (36)	84	* *	120	25.8(21)	367	7.1 (39)	367	10.4(27)
NEK	921	3.7 (11)	55.4 (26)	822	19.3 (15)	55.2 (36)	131	26.0 (9)	197	28.4(25)	789	6.1 (30)	790	13.5(41)
Nemaha	106	0.9 (2)	55.7 (29)	99	* *	* *	27	* *	32	* *	105	2.9 (1)	105	5.7(3)
Neosho	451	7.1 (43)	52.3 (9)	413	20.6 (21)	49.9 (14)	71	* *	71	* *	398	5.8 (26)	397	13.4(40)
Osborne	1,068	4.5 (20)	50.6 (3)	935	20.9 (24)	46.5 (5)	182	20.3 (3)	231	16.5 (3)	914	4.0 (5)	914	9.1(14)
Reno	1,831	4.8 (27)	54.1 (17)	1,646	21.0 (26)	49.5 (12)	244	28.7 (14)	334	21.0(10)	1,567	3.8 (4)	1,570	12.0(37)
Riley	2,786	4.5 (20)	51.0 (4)	2,279	17.6 (8)	53.8 (31)	430	28.4 (13)	678	26.7(23)	2,221	4.5 (9)	2,221	9.8(19)
Saline	1,543	4.9 (28)	55.3 (25)	1,397	17.0 (6)	54.3 (34)	294	24.8 (8)	308	20.1 (7)	1,366	6.2 (33)	1,369	7.4(7)
Scott	156	3.2 (9)	59.0 (38)	137	17.5 (7)	54.0 (32)	21	* *	25	* *	133	3.0 (2)	133	7.5(8)

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.
- (4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
- (5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
- (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5F
2008 - 2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Health Indicators by Local Agency
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)														
Local Agency	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Sedgwick	13,120	4.5 (20)	53.3 (11)	11,998	20.2 (20)	51.5 (21)	2,570	31.5 (18)	3,130	24.7 (17)	11,695	5.6 (23)	11,690	9.9 (21)
SEK	181	5.5 (34)	53.6 (14)	167	24.0 (39)	46.1 (3)	29	* *	37	* *	162	4.9 (15)	162	14.8 (42)
Seward	1,432	2.5 (6)	56.7 (32)	1,290	22.2 (32)	47.7 (7)	238	27.3 (12)	214	24.3 (16)	1,251	7.5 (41)	1,251	6.5 (6)
Shawnee	4,582	4.9 (28)	54.1 (17)	4,019	19.3 (15)	52.5 (25)	466	40.3 (22)	788	29.8 (27)	3,932	6.8 (37)	3,937	7.7 (10)
Sheridan	25	* *	* *	21	* *	* *	5	* *	4	* *	22	* *	22	* *
Sherman	433	5.8 (38)	55.2 (23)	377	18.6 (13)	55.2 (36)	51	* *	83	* *	369	4.1 (6)	369	11.1 (31)
SKCPH	820	4.6 (25)	51.5 (6)	701	21.8 (30)	48.9 (9)	100	23.0 (6)	162	21.0 (10)	674	3.7 (3)	673	11.6 (34)
Stevens	221	0.9 (2)	61.1 (41)	191	15.7 (2)	50.8 (19)	37	* *	60	* *	183	14.2 (44)	183	5.5 (2)
Sumner	466	5.6 (35)	51.7 (8)	426	22.8 (37)	43.9 (1)	68	* *	123	17.9 (5)	407	6.1 (30)	408	15.0 (43)
SW KS WIC	2,517	5.7 (36)	52.6 (10)	2,291	24.1 (40)	45.3 (2)	350	30.3 (15)	427	25.1 (19)	2,255	7.8 (42)	2,250	5.8 (4)
Wichita	45	* *	* *	41	* *	* *	6	* *	6	* *	42	* *	42	* *
Wilson	254	7.9 (44)	55.9 (31)	239	17.6 (8)	53.1 (28)	43	* *	38	* *	228	6.1 (30)	228	10.1 (24)
Wyandotte	6,341	4.3 (18)	55.7 (29)	5,681	19.6 (18)	51.9 (22)	981	30.9 (17)	1,141	26.0 (22)	5,503	4.5 (9)	5,503	8.3 (12)
Unknown Local Agencies	21	* *	* *	20	* *	* *	1	* *	3	* *	22	* *	22	* *
Kansas	63,394	4.5	54.2	56,431	20.2	51.1	9,783	29.3	13,286	23.5	54,852	5.7	54,836	9.2

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6F
2008 - 2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Behavioral Indicators by Local Agency
Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2, 3)										
Local Agency	Medical Care			WIC Enrollment			Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prenatal (4)		During Pregnancy (5)		
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	
Barton	1,188	83.7 (17)	1.4 (8)	1,270	50.2 (4)	1,183	20.7 (15)	978	54.8 (33)	
Butler	750	70.1 (38)	9.9 (39)	760	38.9 (27)	721	14.7 (27)	548	35.6 (41)	
Chautauqua	123	74.0 (33)	6.5 (33)	131	54.2 (1)	116	17.2 (21)	104	74.0 (2)	
Cherokee	1,556	83.2 (20)	5.1 (27)	2,004	35.2 (35)	1,847	13.7 (30)	1,547	60.3 (25)	
Cloud	543	86.9 (10)	3.7 (17)	567	47.1 (9)	518	21.4 (12)	453	69.1 (4)	
Cowley	894	69.2 (39)	6.3 (32)	987	44.2 (15)	920	12.6 (32)	782	56.0 (32)	
Crawford	1,487	88.0 (5)	2.5 (14)	1,680	42.3 (16)	1,540	12.3 (33)	1,334	58.6 (27)	
Doniphan	116	77.6 (27)	7.8 (37)	123	39.8 (23)	117	35.0 (3)	92	* *	
Douglas	1,274	74.0 (33)	5.7 (30)	1,397	37.6 (32)	1,295	19.3 (19)	1,054	60.4 (24)	
Ford	1,666	71.6 (37)	14.4 (42)	1,915	45.3 (12)	1,730	9.7 (37)	1,538	51.8 (37)	
Franklin	203	76.4 (29)	3.0 (15)	498	38.2 (30)	463	5.4 (42)	370	62.4 (17)	
Geary	876	72.1 (36)	1.3 (6)	1,097	35.3 (34)	1,009	19.5 (18)	846	56.9 (30)	
Graham	106	92.5 (2)	1.9 (10)	117	48.7 (5)	104	26.9 (6)	88	* *	
Greeley	32	* *	* *	34	* *	34	* *	27	* *	
Greenwood	119	56.3 (44)	36.1 (44)	132	52.3 (2)	116	14.7 (27)	104	70.2 (3)	
Harvey	657	74.0 (33)	12.9 (41)	764	47.8 (6)	705	16.6 (23)	601	66.2 (8)	
Jefferson	160	85.6 (12)	6.9 (34)	174	39.7 (24)	162	6.2 (41)	136	50.7 (38)	
Johnson	4,561	64.9 (42)	3.7 (17)	5,294	32.8 (41)	4,928	13.3 (31)	4,016	62.0 (19)	
Leavenworth	931	85.5 (13)	0.6 (2)	965	32.5 (42)	925	20.2 (16)	699	68.5 (5)	
Lincoln	376	88.0 (5)	2.1 (11)	366	38.8 (28)	347	29.7 (5)	293	61.4 (22)	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6F
2008 - 2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Behavioral Indicators by Local Agency
Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2, 3)										
Local Agency	Medical Care			WIC Enrollment		Multivitamin Consumption				
	1st Trimester		No Care	1st Trimester		Prenatal (4)		During Pregnancy (5)		
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	
Linn	175	80.0 (25)	7.4 (35)	180	47.8 (6)	167	21.0 (14)	148	61.5 (20)	
Lyon	946	84.8 (14)	0.7 (3)	988	41.0 (21)	916	16.0 (24)	756	62.2 (18)	
Marion	553	83.9 (16)	1.8 (9)	598	44.3 (14)	568	21.8 (10)	446	67.0 (7)	
Marshall	148	92.6 (1)	0.0 (1)	153	41.8 (17)	133	30.8 (4)	108	65.7 (9)	
Meade	280	80.7 (23)	5.0 (26)	292	51.0 (3)	271	42.8 (2)	237	65.4 (11)	
Miami	389	81.2 (22)	1.0 (5)	391	34.3 (38)	371	16.7 (22)	296	53.4 (34)	
NEK	713	83.3 (18)	5.6 (29)	898	39.6 (25)	832	10.3 (35)	740	65.0 (12)	
Nemaha	109	87.2 (7)	5.5 (28)	110	38.2 (30)	97	* *	86	* *	
Neosho	386	88.9 (3)	4.4 (22)	433	38.6 (29)	404	9.2 (39)	343	67.1 (6)	
Osborne	952	87.0 (9)	2.3 (13)	1,011	41.1 (20)	959	25.9 (7)	815	60.6 (23)	
Reno	1,635	84.4 (15)	2.1 (11)	1,703	39.5 (26)	1,586	21.5 (11)	1,337	63.2 (15)	
Riley	2,184	83.3 (18)	0.8 (4)	2,668	34.6 (37)	2,528	18.7 (20)	2,007	61.5 (20)	
Saline	1,234	74.9 (31)	4.1 (21)	1,451	34.1 (40)	1,348	5.0 (43)	1,151	52.2 (35)	
Scott	135	81.5 (21)	5.9 (31)	146	45.9 (11)	136	68.4 (1)	124	57.3 (29)	
Sedgwick	12,080	74.9 (31)	4.6 (24)	12,454	28.3 (44)	11,578	15.4 (25)	9,583	47.5 (39)	
SEK	153	88.9 (3)	3.9 (20)	173	44.5 (13)	170	20.0 (17)	145	64.1 (14)	
Seward	1,183	65.4 (41)	11.7 (40)	1,349	34.3 (38)	1,227	8.6 (40)	1,021	64.8 (13)	
Shawnee	3,720	77.1 (28)	3.7 (17)	4,360	47.2 (8)	4,082	10.0 (36)	3,484	52.2 (35)	
Sheridan	22	* *	* *	22	* *	20	* *	17	* *	
Sherman	248	87.1 (8)	4.4 (22)	414	41.3 (19)	378	25.9 (7)	324	74.7 (1)	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6F
2008 - 2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Behavioral Indicators by Local Agency
Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2, 3)									
Local Agency	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Pregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
SKCPH	702	79.8 (26)	7.4 (35)	769	46.4 (10)	738	21.1 (13)	611	65.6 (10)
Stevens	194	75.3 (30)	3.1 (16)	206	32.0 (43)	197	22.3 (9)	161	60.2 (26)
Sumner	435	80.2 (24)	4.6 (24)	440	40.5 (22)	405	15.1 (26)	325	62.5 (16)
SW KS WIC	2,154	60.6 (43)	23.4 (43)	2,380	34.7 (36)	2,219	11.0 (34)	1,852	39.6 (40)
Wichita	43	* *	* *	45	* *	42	* *	33	* *
Wilson	236	86.0 (11)	1.3 (6)	246	41.5 (18)	226	14.6 (29)	198	56.1 (31)
Wyandotte	5,629	65.5 (40)	7.8 (37)	6,006	37.2 (33)	5,557	9.3 (38)	4,735	57.5 (28)
Unknown Local Agencies	20	* *	* *	23	* *	23	* *	18	* *
Kansas	54,276	75.1	5.7	60,184	37.0	55,958	14.7	46,711	56.0

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7F
2008 - 2010 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by Local Agency
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
Local Agency	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Barton	1,338	38.2 (16)	1,156	24.4 (21)	354	28.2 (22)	1,096	13.6 (10)	1,161	8.4 (6)
Butler	818	49.4 (38)	685	30.7 (33)	252	32.1 (14)	618	13.4 (8)	690	8.4 (6)
Chautauqua	124	38.7 (18)	89	* *	29	* *	105	41.0 (37)	87	* *
Cherokee	2,086	39.5 (19)	1,786	27.4 (28)	551	19.1 (30)	1,709	32.7 (33)	1,804	31.6 (32)
Cloud	602	40.7 (21)	526	21.5 (13)	167	29.3 (20)	510	18.8 (20)	531	14.1 (18)
Cowley	1,041	42.4 (29)	819	28.4 (30)	274	26.3 (27)	864	20.9 (23)	822	20.3 (24)
Crawford	1,677	45.0 (33)	1,319	26.9 (26)	472	31.1 (18)	1,447	39.6 (36)	1,326	38.2 (35)
Doniphan	117	41.9 (25)	85	* *	24	* *	87	* *	88	* *
Douglas	1,468	40.9 (22)	1,234	19.8 (11)	385	42.1 (2)	1,179	10.2 (4)	1,236	6.4 (2)
Ford	1,956	9.0 (2)	1,639	4.5 (2)	105	31.4 (17)	1,704	8.9 (2)	1,637	8.4 (6)
Franklin	533	44.1 (31)	468	28.4 (30)	142	27.5 (23)	403	13.4 (8)	470	9.6 (9)
Geary	1,134	36.9 (14)	905	20.7 (12)	257	33.9 (9)	942	37.4 (35)	908	34.1 (33)
Graham	121	40.5 (20)	99	* *	34	* *	100	24.0 (25)	101	25.7 (30)
Greeley	37	* *	32	* *	7	* *	27	* *	32	* *
Greenwood	118	56.8 (43)	60	* *	27	* *	110	30.9 (31)	87	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7F
2008 - 2010 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by Local Agency
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
Local Agency	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Harvey	802	37.2 (15)	687	21.7 (14)	209	34.9 (6)	667	7.0 (1)	689	4.2 (1)
Jefferson	48	* *	21	* *	4	* *	48	* *	25	* *
Johnson	5,652	23.9 (7)	4,959	10.6 (5)	821	43.0 (1)	4,490	18.4 (19)	4,972	18.1 (23)
Leavenwort	1,059	46.8 (35)	892	25.8 (24)	288	34.7 (7)	788	41.5 (38)	895	40.1 (36)
Lincoln	392	42.3 (27)	334	25.4 (22)	120	33.3 (11)	324	15.1 (15)	350	13.1 (16)
Linn	188	56.4 (42)	170	37.6 (37)	84	* *	159	15.1 (15)	172	12.8 (15)
Lyon	1,065	31.9 (11)	942	17.5 (10)	226	35.0 (4)	854	13.7 (12)	943	10.9 (11)
Marion	631	38.5 (17)	548	24.3 (20)	168	32.1 (14)	495	10.7 (5)	550	7.5 (4)
Marshall	159	50.3 (39)	134	32.1 (34)	53	* *	132	53.8 (41)	136	51.5 (39)
Meade	303	19.1 (4)	254	12.6 (7)	40	* *	263	16.3 (18)	258	14.0 (17)
Miami	395	44.8 (32)	335	25.7 (23)	113	33.6 (10)	306	20.3 (22)	334	16.8 (20)
NEK	817	47.7 (36)	618	29.6 (32)	224	24.1 (29)	697	30.7 (30)	626	27.5 (31)
Nemaha	108	46.3 (34)	87	* *	37	* *	91	* *	88	* *
Neosho	395	41.8 (23)	237	23.6 (18)	77	* *	346	28.3 (29)	244	17.6 (22)
Osborne	1,049	41.8 (23)	871	23.5 (17)	301	32.9 (12)	848	23.0 (24)	885	20.7 (26)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7F
2008 - 2010 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by Local Agency
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
Local Agency	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Reno	1,827	42.3 (27)	1,595	23.6 (18)	554	35.0 (4)	1,496	14.6 (13)	1,602	12.4 (14)
Riley	2,795	28.4 (8)	2,044	15.6 (8)	404	30.0 (19)	2,220	15.0 (14)	2,137	12.0 (13)
Saline	1,479	42.0 (26)	1,233	26.4 (25)	430	26.5 (25)	1,235	12.6 (6)	1,237	9.9 (10)
Scott	130	28.5 (9)	97	* *	24	* *	115	32.2 (32)	100	21.0 (27)
Sedgwick	13,267	31.1 (10)	12,097	16.3 (9)	2,844	34.7 (7)	10,766	9.0 (3)	12,083	7.3 (3)
SEK	177	48.6 (37)	151	27.8 (29)	58	* *	153	24.2 (26)	154	22.1 (28)
Seward	1,418	6.8 (1)	1,264	3.2 (1)	66	* *	1,177	13.6 (10)	1,284	11.7 (12)
Shawnee	4,340	35.7 (13)	3,435	22.8 (16)	928	27.2 (24)	3,590	13.0 (7)	3,459	7.5 (4)
Sheridan	24	* *	21	* *	4	* *	20	* *	21	* *
Sherman	435	34.9 (12)	372	22.0 (15)	103	32.0 (16)	354	26.3 (28)	372	23.4 (29)
SKCPH	791	42.5 (30)	640	26.9 (26)	212	26.4 (26)	652	37.1 (34)	646	34.4 (34)
Stevens	223	19.3 (5)	189	8.5 (4)	27	* *	179	25.7 (27)	191	20.4 (25)
Sumner	466	53.9 (40)	414	33.8 (35)	175	29.1 (21)	367	47.7 (39)	416	49.0 (38)
SW KS WIC	2,526	11.4 (3)	2,280	5.9 (3)	197	37.6 (3)	2,092	15.3 (17)	2,280	14.3 (19)
Wichita	46	* *	37	* *	8	* *	36	* *	37	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7F
2008 - 2010 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by Local Agency
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
Local Agency	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Wilson	252	56.0 (41)	230	34.3 (36)	110	32.7 (13)	220	47.7 (39)	230	43.0 (37)
Wyandotte	6,335	21.1 (6)	5,508	12.4 (6)	870	26.3 (27)	5,304	19.6 (21)	5,600	17.3 (21)
Unknown Local Agencies	22	* *	22	* *	11	* *	18	* *	23	* *
Kansas	62,786	31.6	53,620	17.9	12,870	31.8	51,403	17.6	54,019	15.1

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8F
2008 - 2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Infant Health Indicators by Local Agency
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)									
Local Agency	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Barton	1,138	6.7 (14)	5.3 (8)	1,078	9.8 (28)	958	3.4 (13)	1,131	74.4 (14)
Butler	762	7.6 (28)	6.3 (20)	687	9.5 (22)	612	3.4 (13)	744	69.4 (23)
Chautauqua	122	5.7 (8)	4.1 (2)	118	7.6 (4)	106	2.8 (6)	122	72.1 (17)
Cherokee	1,884	7.2 (24)	5.6 (12)	1,745	9.5 (22)	1,571	3.5 (17)	1,838	58.9 (39)
Cloud	527	6.8 (16)	3.8 (1)	500	12.6 (40)	430	3.7 (21)	527	69.8 (21)
Cowley	928	7.5 (26)	7.5 (32)	871	11.4 (38)	748	4.8 (32)	938	58.5 (40)
Crawford	1,506	8.5 (36)	4.4 (4)	1,448	10.3 (33)	1,271	4.9 (35)	1,520	59.1 (38)
Doniphan	116	12.1 (43)	10.3 (42)	107	13.1 (43)	92	* *	114	61.4 (34)
Douglas	1,264	6.7 (14)	7.0 (26)	1,184	8.9 (17)	1,063	4.0 (27)	1,254	77.7 (4)
Ford	1,694	4.8 (3)	6.0 (16)	1,651	8.9 (17)	1,465	2.0 (2)	1,602	74.6 (13)
Franklin	462	5.6 (6)	5.4 (10)	426	8.2 (9)	389	3.9 (25)	457	61.5 (33)
Geary	927	6.9 (18)	6.3 (20)	874	8.9 (17)	794	3.3 (12)	896	70.5 (20)
Graham	100	11.0 (42)	14.0 (43)	96	* *	82	* *	99	* *
Greeley	32	* *	* *	29	* *	26	* *	32	* *

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.
- (4) <2500 g.
- (5) >4000 g.
- (6) <37 weeks gestation.
- (7) 37 or more weeks gestation and <2500 g.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8F
2008 - 2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Infant Health Indicators by Local Agency
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)										
Local Agency	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)							
	Number	% (Rank)	%	(Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Greenwood	119	5.0 (4)	5.0	(7)	114	8.8 (14)	104	4.8 (32)	117	60.7 (35)
Harvey	685	7.9 (29)	6.1	(18)	654	10.2 (32)	579	3.8 (23)	695	80.7 (1)
Jefferson	169	8.3 (35)	6.5	(24)	155	12.9 (41)	134	5.2 (38)	166	65.1 (30)
Johnson	4,730	6.8 (16)	7.1	(30)	4,622	7.6 (4)	4,020	3.4 (13)	4,877	80.2 (3)
Leavenwort	897	7.9 (29)	7.0	(26)	823	8.9 (17)	739	4.3 (30)	910	63.5 (32)
Lincoln	350	7.1 (22)	8.0	(34)	335	8.4 (11)	291	2.7 (3)	354	69.2 (24)
Linn	163	9.2 (40)	4.9	(6)	161	11.2 (37)	137	7.3 (41)	167	74.9 (12)
Lyon	926	7.0 (21)	6.0	(16)	847	8.4 (11)	774	3.6 (18)	910	73.4 (16)
Marion	523	6.1 (10)	9.0	(41)	500	8.8 (14)	444	2.9 (8)	533	75.0 (11)
Marshall	139	8.6 (37)	4.3	(3)	135	9.6 (25)	118	5.1 (37)	140	60.0 (37)
Meade	242	5.4 (5)	8.3	(37)	246	10.6 (34)	196	3.1 (9)	264	80.3 (2)
Miami	386	6.2 (11)	8.5	(39)	349	7.7 (7)	320	3.8 (23)	358	74.0 (15)
NEK	792	8.2 (32)	7.8	(33)	799	8.3 (10)	697	4.9 (35)	822	57.4 (43)
Nemaha	74	*	*	* *	101	9.9 (30)	58	* *	104	60.6 (36)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8F
2008 - 2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Infant Health Indicators by Local Agency
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)									
Local Agency	Birthweight			Preterm ⁽⁶⁾		Full Term LBW ⁽⁷⁾		Ever Breastfed	
	Low ⁽⁴⁾	High ⁽⁵⁾		Number	%	Number	%	Number	%
	Number	% (Rank)	% (Rank)						
Neosho	319	6.6 (13)	5.3 (8)	389	10.8 (35)	265	3.4 (13)	417	58.3 (41)
Osborne	901	7.5 (26)	5.5 (11)	863	9.7 (27)	749	3.6 (18)	908	75.4 (10)
Reno	1,614	6.9 (18)	5.9 (15)	1,496	8.8 (14)	1,348	3.7 (21)	1,604	67.0 (28)
Riley	2,347	5.6 (6)	8.0 (34)	2,145	6.9 (2)	1,976	2.7 (3)	2,339	77.6 (6)
Saline	1,384	7.9 (29)	6.3 (20)	1,307	9.6 (25)	1,159	4.2 (28)	1,395	63.9 (31)
Scott	129	4.7 (2)	7.0 (26)	123	7.3 (3)	112	2.7 (3)	129	68.2 (27)
Sedgwick	11,991	8.9 (39)	7.0 (26)	11,196	8.6 (13)	10,146	4.8 (32)	11,566	69.1 (25)
SEK	147	8.2 (32)	8.2 (36)	158	7.6 (4)	128	7.0 (40)	164	57.9 (42)
Seward	1,212	5.8 (9)	6.7 (25)	1,190	9.4 (21)	1,022	3.1 (9)	1,266	70.8 (19)
Shawnee	3,988	7.4 (25)	5.6 (12)	3,784	11.1 (36)	3,309	3.9 (25)	3,990	69.7 (22)
Sheridan	23	*	* * *	20	* *	16	* *	23	* *
Sherman	380	7.1 (22)	4.7 (5)	351	8.0 (8)	323	2.8 (6)	371	77.1 (7)
SKCPH	638	6.9 (18)	8.3 (37)	635	9.8 (28)	527	3.2 (11)	682	76.0 (9)
Stevens	117	3.4 (1)	8.5 (39)	172	5.2 (1)	101	1.0 (1)	189	76.7 (8)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8F
2008 - 2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Infant Health Indicators by Local Agency
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)										
Local Agency	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)							
	Number	% (Rank)	%	(Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Sumner	422	10.9 (41)	7.1	(30)	394	11.4 (38)	340	6.8 (39)	427	68.4 (26)
SW KS WIC	2,233	6.5 (12)	6.1	(18)	2,137	9.5 (22)	1,886	3.6 (18)	1,921	77.7 (4)
Wichita	41	* * *	*	*	39	* *	33	* *	41	* *
Wilson	240	8.8 (38)	5.8	(14)	232	12.9 (41)	200	4.5 (31)	226	66.4 (29)
Wyandotte	5,495	8.2 (32)	6.3	(20)	5,247	10.0 (31)	4,570	4.2 (28)	5,549	71.2 (18)
Unknown Local Agencies	23	* * *	*	*	23	* *	23	* *	22	* *
Kansas	55,301	7.5	6.5		52,556	9.2	46,451	4.0	54,920	70.7

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

