

Table 3F
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Racial and Ethnic Distribution by Local Agency

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Local Agency	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Barton	424	71.9	1.2	25.5	0.5	0.2	0.7	0.0
Butler	253	86.6	0.8	7.9	1.6	0.0	3.2	0.0
Chautauqua	44	*	*	*	*	*	*	*
Cherokee	648	80.7	7.3	8.2	1.7	1.1	1.1	0.0
Cloud	164	92.1	0.0	2.4	1.2	0.6	3.7	0.0
Cowley	361	76.7	1.4	18.0	0.8	1.9	1.1	0.0
Crawford	550	86.9	3.3	6.4	1.3	1.1	1.1	0.0
Doniphan	40	*	*	*	*	*	*	*
Douglas	525	64.8	8.0	10.7	8.6	5.9	2.1	0.0
Ford	674	19.6	3.0	76.0	0.3	0.9	0.3	0.0
Franklin	189	92.1	1.1	3.7	2.1	0.5	0.5	0.0
Geary	406	55.9	22.2	15.3	0.7	2.7	3.2	0.0
Graham	46	*	*	*	*	*	*	*
Greeley	14	*	*	*	*	*	*	*
Greenwood	35	*	*	*	*	*	*	*
Harvey	251	73.3	3.2	21.5	0.8	0.4	0.8	0.0
Jefferson	57	*	*	*	*	*	*	*
Johnson	1,898	47.5	14.6	33.6	0.3	2.8	1.2	0.0
Leavenworth	367	66.8	16.3	11.2	1.4	1.1	3.3	0.0
Lincoln	120	85.8	0.0	8.3	0.8	1.7	3.3	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3F
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Racial and Ethnic Distribution by Local Agency

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Local Agency	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Linn	69	*	*	*	*	*	*	*
Lyon	309	61.5	5.5	30.7	0.3	1.0	1.0	0.0
Marion	198	90.4	1.0	6.6	0.0	1.5	0.5	0.0
Marshall	60	*	*	*	*	*	*	*
Meade	106	67.9	0.0	31.1	0.9	0.0	0.0	0.0
Miami	158	91.1	1.3	5.7	0.0	0.6	1.3	0.0
NEK	314	76.4	6.4	4.5	9.9	0.0	2.9	0.0
Nemaha	42	*	*	*	*	*	*	*
Neosho	154	92.2	1.3	5.2	0.0	0.0	1.3	0.0
Osborne	384	85.4	1.8	10.7	0.8	0.3	1.0	0.0
Reno	600	80.8	2.8	13.5	0.2	0.3	2.3	0.0
Riley	1,039	68.5	11.1	12.4	1.9	4.4	1.6	0.0
Saline	486	71.8	5.6	19.3	0.2	0.6	2.5	0.0
Scott	64	*	*	*	*	*	*	*
Sedgwick	4,536	45.2	17.0	30.2	1.4	3.8	2.4	0.0
SEK	61	*	*	*	*	*	*	*
Seward	447	15.7	2.2	81.0	0.4	0.4	0.2	0.0
Shawnee	1,580	57.0	15.8	22.1	1.5	0.9	2.8	0.0
Sheridan	8	*	*	*	*	*	*	*
Sherman	152	73.0	0.7	23.7	0.7	2.0	0.0	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3F
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Racial and Ethnic Distribution by Local Agency

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Local Agency	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
SKCPH	287	82.2	1.4	13.9	0.0	1.0	1.4	0.0
Stevens	72	*	*	*	*	*	*	*
Sumner	136	83.8	1.5	8.8	2.9	0.0	2.9	0.0
SW KS WIC	824	27.5	1.2	67.0	0.0	4.1	0.1	0.0
Wichita	8	*	*	*	*	*	*	*
Wilson	84	*	*	*	*	*	*	*
Wyandotte	2,101	26.2	27.0	39.3	0.6	6.1	0.9	0.0
Unknown Local Agencies	25	*	*	*	*	*	*	*
Kansas	21,370	56.0	11.3	27.2	1.2	2.6	1.7	0.0
Nation (Prior Year)	1,162,961	41.6	22.7	29.2	1.1	3.4	1.3	0.7

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4F
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Age Distribution by Local Agency

Comparison of Age Distribution ⁽²⁾								
Local Agency	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Barton	424	0.0	6.8	14.9	64.2	13.0	1.2	0.0
Butler	253	0.0	3.6	13.0	68.0	14.2	1.2	0.0
Chautauqua	44	*	*	*	*	*	*	*
Cherokee	648	0.8	6.6	14.0	63.3	14.4	0.8	0.2
Cloud	164	0.0	4.3	18.3	56.7	18.9	1.8	0.0
Cowley	361	0.0	6.6	15.5	62.9	14.1	0.8	0.0
Crawford	550	0.5	3.5	16.2	62.9	15.8	0.9	0.2
Doniphan	40	*	*	*	*	*	*	*
Douglas	525	0.2	3.2	9.3	63.8	21.3	2.1	0.0
Ford	674	0.0	5.2	12.8	58.8	20.9	2.4	0.0
Franklin	189	0.0	4.8	13.8	61.4	19.0	1.1	0.0
Geary	406	0.2	3.7	16.0	66.3	12.8	1.0	0.0
Graham	46	*	*	*	*	*	*	*
Greeley	14	*	*	*	*	*	*	*
Greenwood	35	*	*	*	*	*	*	*
Harvey	251	0.4	6.8	13.9	54.2	22.7	2.0	0.0
Jefferson	57	*	*	*	*	*	*	*
Johnson	1,898	0.1	3.0	9.5	62.2	23.4	1.7	0.1
Leavenworth	367	0.3	3.5	15.8	66.8	12.8	0.8	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4F
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Age Distribution by Local Agency

Comparison of Age Distribution ⁽²⁾								
Local Agency	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Lincoln	120	0.0	3.3	10.8	65.0	20.0	0.8	0.0
Linn	69	*	*	*	*	*	*	*
Lyon	309	0.6	7.1	11.7	65.4	13.9	1.3	0.0
Marion	198	0.0	4.5	15.7	61.6	17.7	0.5	0.0
Marshall	60	*	*	*	*	*	*	*
Meade	106	0.0	5.7	14.2	58.5	19.8	1.9	0.0
Miami	158	0.0	3.8	17.1	70.9	8.2	0.0	0.0
NEK	314	0.0	8.0	14.3	65.3	11.8	0.6	0.0
Nemaha	42	*	*	*	*	*	*	*
Neosho	154	0.0	4.5	14.9	66.9	11.7	0.6	1.3
Osborne	384	0.5	4.7	14.3	62.8	16.9	0.8	0.0
Reno	600	0.5	4.5	15.7	64.8	13.8	0.7	0.0
Riley	1,039	0.0	1.3	10.7	74.4	13.5	0.2	0.0
Saline	486	0.6	7.2	13.8	61.3	15.6	1.4	0.0
Scott	64	*	*	*	*	*	*	*
Sedgwick	4,536	0.4	5.2	12.9	60.6	19.3	1.5	0.1
SEK	61	*	*	*	*	*	*	*
Seward	447	0.2	9.4	12.5	53.5	22.6	1.6	0.2
Shawnee	1,580	0.5	5.6	14.2	62.7	15.8	1.1	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4F
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Age Distribution by Local Agency

Comparison of Age Distribution ⁽²⁾								
Local Agency	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Sheridan	8	*	*	*	*	*	*	*
Sherman	152	0.0	6.6	16.4	61.2	12.5	3.3	0.0
SKCPH	287	0.3	4.9	13.2	66.6	14.3	0.7	0.0
Stevens	72	*	*	*	*	*	*	*
Sumner	136	0.0	3.7	13.2	64.7	16.9	1.5	0.0
SW KS WIC	824	0.4	5.0	16.0	56.7	20.6	1.3	0.0
Wichita	8	*	*	*	*	*	*	*
Wilson	84	*	*	*	*	*	*	*
Wyandotte	2,101	0.4	6.9	12.9	57.8	20.4	1.6	0.0
Unknown Local Agencies	25	*	*	*	*	*	*	*
Kansas	21,370	0.3	5.1	13.2	62.1	17.9	1.3	0.1
Nation (Prior Year)	1,162,961	0.3	5.3	11.5	59.4	21.7	1.7	0.2

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5F
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Health Indicators by Local Agency

Comparison of Maternal Health Indicators ^(2, 3)														
Local Agency	Prepregnancy BMI ⁽⁴⁾			Weight Gain ⁽⁵⁾			Anemia (Low Hb/Hct) ⁽⁶⁾				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Barton	419	3.3 (8)	62.5 (32)	365	28.2 (29)	46.3 (10)	47	* *	62	* *	374	7.2 (17)	374	12.6(23)
Butler	248	4.8 (26)	55.2 (15)	224	24.6 (22)	48.7 (17)	36	* *	38	* *	231	6.1 (13)	231	13.4(24)
Chautauqua	43	* *	* *	38	* *	* *	9	* *	5	* *	39	* *	39	* *
Cherokee	627	6.7 (30)	52.0 (4)	566	24.6 (22)	45.2 (6)	104	42.3 (6)	128	25.0 (6)	571	8.2 (23)	571	15.1(28)
Cloud	160	3.1 (4)	53.1 (7)	137	24.8 (24)	46.7 (11)	16	* *	33	* *	143	4.2 (6)	144	7.6(3)
Cowley	354	3.1 (4)	59.0 (28)	306	21.9 (9)	47.4 (15)	41	* *	47	* *	316	6.0 (12)	316	12.0(22)
Crawford	542	6.1 (27)	57.4 (23)	479	22.1 (14)	50.7 (24)	71	* *	92	* *	486	9.9 (29)	491	13.8(25)
Doniphan	38	* *	* *	26	* *	* *	5	* *	10	* *	29	* *	30	* *
Douglas	516	3.3 (8)	52.9 (5)	432	20.1 (5)	51.9 (29)	67	* *	124	29.0 (9)	426	9.9 (29)	426	7.7(4)
Ford	669	2.4 (2)	61.1 (31)	589	25.0 (25)	47.5 (16)	77	* *	70	* *	593	7.4 (20)	591	8.6(9)
Franklin	183	3.3 (8)	58.5 (25)	157	22.9 (17)	51.6 (28)	21	* *	44	* *	137	7.3 (18)	137	13.9(26)
Geary	387	3.6 (14)	51.4 (3)	310	21.9 (9)	51.0 (26)	52	* *	64	* *	323	6.2 (14)	322	8.7(10)
Graham	44	* *	* *	38	* *	* *	6	* *	6	* *	38	* *	40	* *
Greeley	14	* *	* *	12	* *	* *	2	* *	3	* *	12	* *	12	* *
Greenwood	35	* *	* *	33	* *	* *	4	* *	4	* *	33	* *	33	* *
Harvey	246	2.0 (1)	57.3 (22)	218	22.9 (17)	47.2 (13)	31	* *	43	* *	226	3.1 (1)	226	15.9(30)
Jefferson	56	* *	* *	50	* *	* *	9	* *	18	* *	51	* *	50	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5F
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Health Indicators by Local Agency

Comparison of Maternal Health Indicators ^(2, 3)														
Local Agency	Prepregnancy BMI ⁽⁴⁾			Weight Gain ⁽⁵⁾			Anemia (Low Hb/Hct) ⁽⁶⁾				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Johnson	1,836	3.1 (4)	54.7 (11)	1,616	22.1 (14)	47.0 (12)	263	21.3 (1)	446	16.6 (1)	1,665	10.9 (31)	1,664	7.8(5)
Leavenwort	350	4.3 (22)	52.9 (5)	290	17.9 (3)	49.3 (19)	54	* *	80	* *	302	5.6 (9)	301	11.0(17)
Lincoln	119	2.5 (3)	58.8 (26)	104	36.5 (31)	41.3 (3)	14	* *	13	* *	103	7.8 (21)	104	10.6(16)
Linn	69	* *	* *	60	* *	* *	4	* *	6	* *	61	* *	61	* *
Lyon	306	3.3 (8)	60.5 (30)	267	22.8 (16)	49.8 (22)	34	* *	55	* *	269	8.9 (28)	268	14.2(27)
Marion	195	3.6 (14)	55.9 (20)	167	26.3 (27)	45.5 (8)	35	* *	33	* *	171	3.5 (5)	172	10.5(15)
Marshall	57	* *	* *	51	* *	* *	9	* *	18	* *	55	* *	55	* *
Meade	106	3.8 (18)	53.8 (9)	96	* *	* *	11	* *	21	* *	96	* *	95	* *
Miami	155	3.2 (7)	54.8 (12)	141	12.1 (1)	60.3 (31)	28	* *	35	* *	148	3.4 (4)	148	11.5(18)
NEK	306	6.9 (31)	55.6 (18)	256	23.0 (19)	45.7 (9)	36	* *	55	* *	256	4.3 (7)	256	11.7(20)
Nemaha	42	* *	* *	36	* *	* *	4	* *	6	* *	36	* *	36	* *
Neosho	147	3.4 (12)	53.7 (8)	132	22.0 (12)	49.2 (18)	23	* *	24	* *	141	7.8 (21)	141	9.9(13)
Osborne	374	3.7 (16)	50.3 (1)	326	21.5 (8)	45.4 (7)	59	* *	81	* *	335	3.3 (3)	334	8.4(8)
Reno	587	4.3 (22)	55.4 (16)	511	19.6 (4)	50.7 (24)	68	* *	106	23.6 (5)	526	5.5 (8)	527	10.2(14)
Riley	993	3.7 (16)	50.7 (2)	825	21.0 (6)	51.4 (27)	151	41.7 (5)	231	26.8 (7)	861	8.8 (27)	866	9.5(11)
Saline	478	4.4 (24)	55.6 (18)	434	15.9 (2)	53.7 (30)	67	* *	65	* *	439	7.3 (18)	438	9.6(12)
Scott	63	* *	* *	58	* *	* *	12	* *	15	* *	57	* *	57	* *

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.
- (4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
- (5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
- (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5F
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Health Indicators by Local Agency

Comparison of Maternal Health Indicators ^(2, 3)														
Local Agency	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Sedgwick	4,402	3.9 (19)	55.9 (20)	3,986	21.9 (9)	49.5 (21)	836	29.3 (3)	1,016	22.3 (2)	4,112	8.5 (25)	4,113	11.5(18)
SEK	60	* *	* *	52	* *	* *	6	* *	14	* *	53	* *	53	* *
Seward	440	3.4 (12)	59.5 (29)	402	25.1 (26)	43.8 (5)	74	* *	55	* *	408	8.3 (24)	407	4.7(1)
Shawnee	1,549	4.4 (24)	54.6 (10)	1,365	21.0 (6)	50.4 (23)	160	43.8 (7)	240	28.3 (8)	1,398	6.4 (15)	1,398	8.1(7)
Sheridan	7	* *	* *	7	* *	* *	2	* *	0	* *	8	* *	8	* *
Sherman	147	4.1 (21)	55.1 (14)	123	30.9 (30)	38.2 (1)	19	* *	34	* *	128	3.1 (1)	128	15.6(29)
SKCPH	282	6.4 (29)	58.9 (27)	241	24.1 (20)	47.3 (14)	30	* *	61	* *	248	6.9 (16)	248	11.7(20)
Stevens	70	* *	* *	62	* *	* *	15	* *	18	* *	62	* *	63	* *
Sumner	132	6.1 (27)	57.6 (24)	119	24.4 (21)	42.9 (4)	20	* *	32	* *	123	5.7 (10)	123	19.5(31)
SW KS WIC	810	7.4 (32)	54.8 (12)	724	28.0 (28)	39.6 (2)	106	26.4 (2)	124	22.6 (4)	733	8.5 (25)	733	6.0(2)
Wichita	8	* *	* *	8	* *	* *	1	* *	0	* *	8	* *	8	* *
Wilson	81	* *	* *	80	* *	* *	12	* *	13	* *	82	* *	82	* *
Wyandotte	2,034	4.0 (20)	55.5 (17)	1,821	22.0 (12)	49.4 (20)	310	36.5 (4)	371	22.4 (3)	1,851	5.7 (10)	1,849	8.0(6)
Unknown Local Agencies	25	* *	* *	24	* *	* *	2	* *	3	* *	24	* *	24	* *
Kansas	20,811	4.1	55.5	18,364	22.5	48.6	3,063	30.9	4,062	22.1	18,787	7.6	18,793	10.1
Nation (Prior Year)	1,106,370	4.5	53.4	835,927	21.5	48.0	164,201	33.9	346,981	28.1	419,956	5.2	467,675	6.3

- (1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6F
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Behavioral Indicators by Local Agency

Comparison of Maternal Behavioral Indicators ^(2, 3)										
Local Agency	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	% (Rank)		Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Barton	370	85.4 (12)	0.8 (5)		420	45.2 (8)	424	18.4 (16)	338	62.1 (18)
Butler	231	86.1 (10)	0.9 (6)		252	36.9 (23)	253	19.0 (14)	198	67.2 (14)
Chautauqua	39	* *	* *		43	* *	44	* *	38	* *
Cherokee	424	88.4 (7)	2.4 (14)		644	34.6 (27)	648	12.3 (26)	537	69.3 (9)
Cloud	144	90.3 (4)	2.1 (12)		163	52.1 (1)	164	25.6 (6)	142	67.6 (10)
Cowley	270	72.2 (26)	5.9 (24)		357	42.3 (14)	361	16.1 (19)	292	59.6 (25)
Crawford	432	87.7 (9)	2.8 (17)		547	42.6 (12)	550	14.7 (22)	458	55.9 (27)
Doniphan	31	* *	* *		40	* *	40	* *	30	* *
Douglas	391	70.6 (27)	11.0 (28)		522	42.7 (11)	525	18.7 (15)	429	64.1 (17)
Ford	562	76.3 (24)	9.6 (27)		667	46.0 (6)	674	12.6 (25)	588	60.4 (23)
Franklin	127	83.5 (17)	7.1 (26)		188	44.1 (9)	189	13.8 (23)	150	65.3 (15)
Geary	309	79.0 (22)	1.0 (7)		405	42.0 (15)	406	20.4 (13)	334	59.9 (24)
Graham	41	* *	* *		45	* *	46	* *	37	* *
Greeley	7	* *	* *		14	* *	14	* *	11	* *
Greenwood	31	* *	* *		35	* *	35	* *	28	* *
Harvey	224	81.7 (19)	4.0 (19)		251	48.2 (3)	251	23.5 (8)	212	67.5 (11)
Jefferson	50	* *	* *		57	* *	57	* *	49	* *
Johnson	1,576	67.1 (28)	2.4 (14)		1,892	35.1 (26)	1,898	15.6 (20)	1,552	60.6 (21)
Leavenworth	302	88.1 (8)	1.3 (9)		365	31.0 (31)	367	21.5 (11)	292	74.3 (2)
Lincoln	106	91.5 (1)	1.9 (11)		120	48.3 (2)	120	29.2 (2)	106	61.3 (20)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6F
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Behavioral Indicators by Local Agency

Comparison of Maternal Behavioral Indicators ^(2, 3)										
Local Agency	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Prepregnancy ⁽⁴⁾		During Pregnancy ⁽⁵⁾	
	Number	% (Rank)	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Linn	61	* *	* *	* *	69	* *	69	* *	64	* *
Lyon	255	91.0 (2)	0.4 (3)		309	46.0 (6)	309	21.4 (12)	264	60.6 (21)
Marion	166	86.1 (10)	1.2 (8)		195	39.5 (19)	198	28.3 (3)	168	69.6 (8)
Marshall	54	* *	* *	* *	60	* *	60	* *	50	* *
Meade	93	* *	* *	* *	103	46.6 (4)	106	45.3 (1)	92	* *
Miami	146	82.2 (18)	0.0 (1)		158	34.2 (28)	158	12.0 (28)	120	64.2 (16)
NEK	214	88.8 (6)	4.7 (22)		312	46.5 (5)	314	15.0 (21)	260	73.5 (3)
Nemaha	36	* *	* *	* *	42	* *	42	* *	38	* *
Neosho	130	90.8 (3)	2.3 (13)		153	39.9 (18)	154	16.2 (18)	125	70.4 (6)
Osborne	327	84.4 (15)	3.1 (18)		379	38.8 (20)	384	24.7 (7)	322	52.8 (29)
Reno	520	83.8 (16)	2.7 (16)		597	41.2 (16)	600	22.2 (10)	472	62.1 (18)
Riley	801	84.8 (13)	0.7 (4)		1,038	35.9 (25)	1,039	26.8 (5)	839	67.5 (11)
Saline	305	79.7 (20)	5.2 (23)		486	34.2 (28)	486	8.6 (30)	406	67.5 (11)
Scott	59	* *	* *	* *	64	* *	64	* *	57	* *
Sedgwick	3,910	76.0 (25)	4.3 (20)		4,528	29.3 (32)	4,536	16.7 (17)	3,699	52.1 (30)
SEK	51	* *	* *	* *	61	* *	61	* *	56	* *
Seward	355	63.4 (29)	18.0 (29)		446	37.4 (22)	447	12.1 (27)	380	70.3 (7)
Shawnee	1,235	77.6 (23)	4.5 (21)		1,576	42.6 (12)	1,580	13.4 (24)	1,343	52.9 (28)
Sheridan	8	* *	* *	* *	8	* *	8	* *	4	* *
Sherman	125	84.8 (13)	1.6 (10)		152	40.1 (17)	152	27.6 (4)	127	76.4 (1)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6F
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Behavioral Indicators by Local Agency

Comparison of Maternal Behavioral Indicators ^(2, 3)													
Local Agency	Medical Care				WIC Enrollment			Multivitamin Consumption					
	1st Trimester		No Care		1st Trimester			Prepregnancy (4)		During Pregnancy (5)			
	Number	% (Rank)	%	% (Rank)	Number	%	% (Rank)	Number	%	% (Rank)			
SKCPH	246	79.7 (20)	6.1	(25)	286	44.1	(9)	287	23.0	(9)	237	70.9	(5)
Stevens	62	* *	* *	* *	71	* *	* *	72	* *	* *	56	* *	* *
Sumner	123	90.2 (5)	0.0	(1)	136	33.1	(30)	136	11.8	(29)	114	71.9	(4)
SW KS WIC	665	52.0 (31)	28.0	(31)	820	38.7	(21)	824	8.6	(30)	673	42.1	(31)
Wichita	8	* *	* *	* *	8	* *	* *	8	* *	* *	7	* *	* *
Wilson	75	* *	* *	* *	83	* *	* *	84	* *	* *	73	* *	* *
Wyandotte	1,452	57.9 (30)	19.5	(30)	2,093	36.8	(24)	2,101	5.9	(32)	1,782	57.5	(26)
Unknown Local Agencies	17	* *	* *	* *	25	* *	* *	25	* *	* *	22	* *	* *
Kansas	17,166	76.1	6.3		21,285	37.4		21,370	16.4		17,671	59.5	
Nation (Prior Year)	429,770	83.1	4.1		1,074,863	36.0		380,154	19.2		418,448	85.0	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7F
2011 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by Local Agency

Comparison of Smoking Indicators (2, 3)										
Local Agency	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Barton	421	39.9 (16)	366	25.4 (20)	111	30.6 (6)	332	14.8 (17)	364	8.8 (15)
Butler	251	53.4 (32)	229	33.6 (29)	102	28.4 (7)	196	7.1 (5)	229	2.6 (3)
Chautauqua	42	* *	37	* *	14	* *	37	* *	37	* *
Cherokee	627	40.0 (17)	556	28.2 (26)	182	20.9 (12)	526	33.3 (26)	555	34.6 (26)
Cloud	164	42.1 (23)	142	23.2 (15)	52	* *	142	22.5 (22)	144	16.0 (18)
Cowley	351	38.5 (14)	257	26.1 (22)	85	* *	284	19.4 (19)	239	18.8 (21)
Crawford	529	43.3 (27)	421	25.2 (19)	142	28.2 (8)	450	39.6 (30)	422	39.1 (30)
Doniphan	35	* *	26	* *	4	* *	26	* *	26	* *
Douglas	511	33.5 (12)	396	17.4 (10)	106	46.2 (2)	422	11.1 (13)	414	8.7 (13)
Ford	646	7.3 (2)	529	5.1 (3)	27	* *	565	7.4 (7)	529	5.5 (7)
Franklin	188	41.5 (21)	159	28.9 (27)	50	* *	150	10.0 (9)	161	6.8 (11)
Geary	399	40.6 (19)	307	19.5 (11)	95	* *	332	30.4 (25)	314	22.3 (24)
Graham	44	* *	38	* *	15	* *	36	* *	38	* *
Greeley	13	* *	11	* *	1	* *	11	* *	11	* *
Greenwood	33	* *	30	* *	10	* *	28	* *	31	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7F
2011 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by Local Agency

Comparison of Smoking Indicators (2, 3)										
Local Agency	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Harvey	249	37.8 (13)	215	22.8 (13)	64	* *	210	2.9 (1)	214	1.9 (2)
Jefferson	49	* *	28	* *	11	* *	43	* *	29	* *
Johnson	1,883	23.3 (5)	1,654	9.1 (4)	285	50.9 (1)	1,537	17.6 (18)	1,655	18.2 (20)
Leavenworth	356	44.4 (28)	284	24.3 (16)	94	* *	287	34.8 (28)	286	35.3 (27)
Lincoln	115	42.6 (25)	95	* *	35	* *	104	20.2 (20)	96	* *
Linn	68	* *	59	* *	23	* *	63	* *	60	* *
Lyon	302	33.4 (11)	259	15.4 (8)	74	* *	262	11.8 (15)	243	5.8 (8)
Marion	196	42.3 (24)	167	22.8 (13)	54	* *	167	4.8 (2)	168	4.8 (6)
Marshall	60	* *	53	* *	20	* *	50	* *	53	* *
Meade	104	24.0 (6)	92	* *	19	* *	91	* *	94	* *
Miami	157	46.5 (30)	143	32.2 (28)	48	* *	115	20.9 (21)	143	16.1 (19)
NEK	263	42.6 (25)	172	34.9 (30)	60	* *	225	33.3 (26)	176	30.1 (25)
Nemaha	17	* *	11	* *	4	* *	11	* *	13	* *
Neosho	145	41.4 (20)	103	26.2 (23)	34	* *	120	25.8 (24)	103	20.4 (22)
Osborne	370	41.9 (22)	294	26.2 (23)	95	* *	312	25.6 (23)	297	21.2 (23)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7F
2011 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by Local Agency

Comparison of Smoking Indicators (2, 3)										
Local Agency	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Reno	593	44.5 (29)	516	25.6 (21)	171	36.3 (4)	465	11.0 (12)	518	7.1 (12)
Riley	1,012	28.1 (8)	831	14.0 (6)	195	42.1 (3)	830	7.1 (5)	837	4.4 (4)
Saline	476	40.1 (18)	429	24.5 (17)	135	25.9 (11)	401	6.7 (4)	431	4.6 (5)
Scott	50	* *	33	* *	9	* *	45	* *	34	* *
Sedgwick	4,500	29.2 (9)	4,055	16.7 (9)	905	33.8 (5)	3,670	8.7 (8)	4,063	5.8 (8)
SEK	61	* *	50	* *	16	* *	55	* *	50	* *
Seward	442	4.3 (1)	398	1.8 (1)	12	* *	376	10.4 (10)	402	8.7 (13)
Shawnee	1,486	33.0 (10)	1,120	20.0 (12)	284	26.8 (10)	1,256	10.5 (11)	1,106	6.1 (10)
Sheridan	7	* *	5	* *	1	* *	4	* *	7	* *
Sherman	149	26.8 (7)	127	14.2 (7)	29	* *	123	4.9 (3)	128	1.6 (1)
SKCPH	283	39.6 (15)	235	26.4 (25)	79	* *	229	41.5 (31)	234	38.9 (29)
Stevens	71	* *	62	* *	15	* *	56	* *	63	* *
Sumner	135	49.6 (31)	121	24.8 (18)	49	* *	111	37.8 (29)	120	35.8 (28)
SW KS WIC	819	11.6 (3)	703	5.0 (2)	61	* *	671	11.5 (14)	701	11.4 (17)
Wichita	8	* *	6	* *	1	* *	7	* *	6	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7F
2011 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by Local Agency

Comparison of Smoking Indicators (2, 3)										
Local Agency	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Wilson	83	* *	81	* *	30	* *	72	* *	80	* *
Wyandotte	2,062	19.4 (4)	1,800	11.9 (5)	271	27.7 (9)	1,760	12.5 (16)	1,845	11.2 (16)
Unknown Local Agencies	25	* *	23	* *	8	* *	22	* *	22	* *
Kansas	20,850	30.4	17,728	17.5	4,187	32.7	17,287	15.1	17,791	12.6
Nation (Prior Year)	814,631	23.0	715,034	12.1	137,883	38.1	660,570	16.3	562,126	15.6

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8F
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Infant Health Indicators by Local Agency

Comparison of Infant Health Indicators ^(2, 3)										
Local Agency	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)							
	Number	% (Rank)	%	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Barton	353	5.1 (3)	5.4	(10)	354	7.9 (10)	321	3.4 (8)	356	74.4 (18)
Butler	220	7.7 (18)	5.0	(8)	223	7.6 (8)	202	5.4 (24)	223	76.7 (14)
Chautauqua	35	* * *	* *	*	36	* *	32	* *	37	* *
Cherokee	564	8.5 (25)	6.2	(16)	566	9.5 (21)	508	5.7 (26)	566	60.1 (30)
Cloud	133	7.5 (16)	9.0	(29)	135	11.1 (30)	117	2.6 (3)	136	84.6 (3)
Cowley	283	6.7 (8)	5.7	(12)	303	8.9 (15)	253	4.7 (20)	303	64.0 (29)
Crawford	460	6.3 (7)	4.6	(5)	477	9.0 (16)	414	2.9 (4)	480	65.4 (27)
Doniphan	28	* * *	* *	*	28	* *	23	* *	27	* *
Douglas	411	5.6 (4)	6.3	(18)	422	7.1 (7)	379	3.2 (7)	423	80.4 (7)
Ford	549	7.3 (15)	4.9	(6)	558	8.4 (13)	496	4.2 (17)	557	77.6 (12)
Franklin	156	7.1 (13)	4.5	(2)	158	12.7 (31)	135	3.0 (5)	159	64.2 (28)
Geary	314	8.0 (23)	9.6	(30)	316	6.0 (3)	294	5.4 (24)	315	75.6 (15)
Graham	36	* * *	* *	*	39	* *	32	* *	39	* *
Greeley	12	* * *	* *	*	12	* *	11	* *	12	* *

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.
- (4) <2500 g.
- (5) >4000 g.
- (6) <37 weeks gestation.
- (7) 37 or more weeks gestation and <2500 g.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8F
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Infant Health Indicators by Local Agency

Comparison of Infant Health Indicators ^(2, 3)										
Local Agency	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)							
	Number	% (Rank)	%	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Greenwood	31	* * * *	*	*	32	* *	30	* *	32	* *
Harvey	217	7.8 (20)	7.8	(26)	219	10.0 (25)	195	5.1 (23)	219	88.1 (1)
Jefferson	49	* * * *	*	*	49	* *	40	* *	49	* *
Johnson	1,566	6.7 (8)	7.2	(23)	1,630	6.3 (4)	1,459	3.9 (14)	1,627	82.7 (4)
Leavenworth	281	10.0 (29)	5.3	(9)	304	7.9 (10)	257	6.6 (30)	304	66.1 (26)
Lincoln	98	* * * *	*	*	104	5.8 (1)	92	* *	104	78.8 (11)
Linn	61	* * * *	*	*	61	* *	51	* *	61	* *
Lyon	265	7.9 (21)	7.5	(24)	265	8.3 (12)	243	3.7 (11)	264	79.5 (10)
Marion	165	7.9 (21)	6.7	(20)	163	11.0 (29)	144	6.3 (28)	166	81.9 (6)
Marshall	52	* * * *	*	*	52	* *	47	* *	52	* *
Meade	88	* * * *	*	*	89	* *	78	* *	90	* *
Miami	147	6.8 (11)	7.5	(24)	148	9.5 (21)	133	3.8 (12)	147	66.7 (25)
NEK	243	9.9 (28)	4.5	(2)	263	9.9 (24)	218	6.0 (27)	263	58.2 (31)
Nemaha	32	* * * *	*	*	36	* *	31	* *	36	* *

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.
- (4) <2500 g.
- (5) >4000 g.
- (6) <37 weeks gestation.
- (7) 37 or more weeks gestation and <2500 g.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8F
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Infant Health Indicators by Local Agency

Comparison of Infant Health Indicators ^(2, 3)										
Local Agency	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)		Number	%	Number	%	Number	%
	Number	% (Rank)	% (Rank)	(Rank)						
Neosho	130	4.6 (2)	6.2 (16)		139	5.8 (1)	121	1.7 (1)	139	72.7 (21)
Osborne	316	6.0 (5)	6.0 (15)		321	7.8 (9)	289	3.1 (6)	323	79.6 (9)
Reno	503	6.0 (5)	5.8 (13)		506	6.5 (5)	467	3.6 (10)	507	73.8 (19)
Riley	840	7.1 (13)	6.7 (20)		845	6.5 (5)	785	4.7 (20)	840	82.6 (5)
Saline	400	8.0 (23)	4.5 (2)		430	10.7 (27)	355	3.4 (8)	429	70.4 (23)
Scott	55	* * * *	* * * *		56	* *	48	* *	56	* *
Sedgwick	3,998	8.6 (27)	6.5 (19)		4,010	8.7 (14)	3,645	4.6 (19)	4,004	73.2 (20)
SEK	50	* * * *	* * * *		52	* *	46	* *	52	* *
Seward	368	4.1 (1)	8.4 (27)		381	9.2 (19)	333	2.1 (2)	382	80.1 (8)
Shawnee	1,347	8.5 (25)	5.5 (11)		1,361	9.2 (19)	1,219	4.7 (20)	1,358	70.3 (24)
Sheridan	8	* * * *	* * * *		8	* *	8	* *	8	* *
Sherman	121	12.4 (30)	3.3 (1)		121	9.1 (17)	110	6.4 (29)	121	86.8 (2)
SKCPH	227	7.0 (12)	8.8 (28)		238	10.1 (26)	204	3.9 (14)	233	76.8 (13)
Stevens	36	* * * *	* * * *		62	* *	32	* *	63	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8F
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Infant Health Indicators by Local Agency

Comparison of Infant Health Indicators ^(2, 3)									
Local Agency	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Sumner	119	7.6 (17)	6.7 (20)	122	10.7 (27)	106	3.8 (12)	122	75.4 (17)
SW KS WIC	716	6.7 (8)	4.9 (6)	721	9.8 (23)	644	3.9 (14)	718	75.6 (15)
Wichita	8	*	*	8	*	8	*	8	*
Wilson	75	*	*	77	*	68	*	77	*
Wyandotte	1,786	7.7 (18)	5.8 (13)	1,832	9.1 (17)	1,619	4.3 (18)	1,817	72.2 (22)
Unknown Local Agencies	24	*	*	24	*	24	*	24	*
Kansas	17,976	7.6	6.2	18,356	8.5	16,366	4.3	18,328	74.2
Nation (Prior Year)	880,933	8.0	6.7	948,286	10.7	749,107	3.4	871,811	68.2

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3F
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Racial and Ethnic Distribution by Local Agency
Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Local Agency	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Barton	1,342	70.9	1.7	25.9	0.4	0.6	0.5	0.0
Butler	820	88.3	1.1	7.3	1.0	0.2	2.1	0.0
Chautauqua	139	92.8	0.7	2.9	0.0	0.0	3.6	0.0
Cherokee	2,102	82.2	6.6	6.7	2.1	1.2	1.2	0.0
Cloud	575	91.5	1.0	3.8	1.4	0.5	1.7	0.0
Cowley	1,087	75.1	2.1	19.3	0.8	1.7	1.0	0.0
Crawford	1,709	87.0	3.0	6.8	1.0	0.9	1.3	0.0
Doniphan	130	90.8	2.3	4.6	0.8	0.0	1.5	0.0
Douglas	1,562	65.2	9.3	10.8	8.1	4.7	1.9	0.0
Ford	2,057	21.0	2.1	75.9	0.2	0.7	0.1	0.0
Franklin	561	92.2	0.5	4.3	1.6	0.4	1.1	0.0
Geary	1,220	54.7	22.3	15.3	0.8	3.1	3.8	0.0
Graham	137	89.8	1.5	7.3	0.0	0.0	1.5	0.0
Greeley	41	*	*	*	*	*	*	*
Greenwood	131	93.1	0.8	4.6	0.8	0.0	0.8	0.0
Harvey	818	71.5	2.9	21.9	1.3	0.9	1.5	0.0
Jefferson	183	94.0	0.5	2.2	2.7	0.0	0.5	0.0
Johnson	5,804	47.0	13.6	34.5	0.4	3.2	1.4	0.0
Leavenworth	1,090	69.2	14.2	11.4	0.8	1.2	3.2	0.0
Lincoln	394	90.1	0.3	7.6	0.3	0.5	1.3	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3F
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Racial and Ethnic Distribution by Local Agency
Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Local Agency	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Linn	206	95.6	0.0	2.4	1.5	0.0	0.5	0.0
Lyon	1,009	60.8	5.0	31.1	0.4	1.5	1.3	0.0
Marion	657	90.9	0.5	6.4	0.6	0.9	0.8	0.0
Marshall	160	89.4	0.6	3.1	1.9	0.0	5.0	0.0
Meade	312	70.8	0.3	27.9	0.3	0.0	0.6	0.0
Miami	455	91.2	2.4	4.0	0.2	0.7	1.5	0.0
NEK	962	80.1	5.0	3.7	9.1	0.0	2.0	0.0
Nemaha	123	90.2	0.0	7.3	0.8	0.8	0.8	0.0
Neosho	466	91.8	1.9	4.5	0.4	0.0	1.3	0.0
Osborne	1,154	87.7	1.4	8.6	1.0	0.6	0.8	0.0
Reno	1,821	80.7	2.4	14.3	0.4	0.5	1.7	0.0
Riley	3,076	71.2	10.7	12.5	0.9	3.8	1.0	0.0
Saline	1,563	70.9	4.7	20.3	0.2	1.7	2.2	0.0
Scott	175	68.0	0.0	32.0	0.0	0.0	0.0	0.0
Sedgwick	13,808	45.2	17.7	29.8	1.5	3.7	2.1	0.0
SEK	193	94.8	0.5	3.1	1.6	0.0	0.0	0.0
Seward	1,407	15.2	2.0	81.7	0.5	0.4	0.1	0.0
Shawnee	4,816	57.7	14.8	22.6	2.1	0.8	2.0	0.0
Sheridan	26	*	*	*	*	*	*	*
Sherman	453	78.4	0.7	19.2	0.4	1.1	0.2	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3F
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Racial and Ethnic Distribution by Local Agency
Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Local Agency	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
SKCPH	865	85.5	0.8	12.0	0.2	0.7	0.7	0.0
Stevens	231	46.8	0.4	51.5	1.3	0.0	0.0	0.0
Sumner	457	86.9	1.3	7.0	2.0	0.2	2.6	0.0
SW KS WIC	2,547	28.5	1.0	66.7	0.0	3.5	0.3	0.0
Wichita	44	*	*	*	*	*	*	*
Wilson	260	94.2	1.5	1.5	0.4	0.0	2.3	0.0
Wyandotte	6,487	26.4	28.8	39.3	0.5	4.2	0.8	0.0
Unknown Local Agencies	48	*	*	*	*	*	*	*
Kansas	65,683	56.6	11.2	27.2	1.2	2.3	1.5	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4F
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Age Distribution by Local Agency
Combined 3 Years

Comparison of Age Distribution ⁽²⁾								
Local Agency	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Barton	1,342	0.1	7.1	13.6	63.5	14.3	1.4	0.0
Butler	820	0.0	5.7	16.0	63.8	13.7	0.6	0.2
Chautauqua	139	0.0	5.0	18.7	61.9	13.7	0.0	0.7
Cherokee	2,102	0.4	6.8	17.4	61.8	12.7	0.9	0.1
Cloud	575	0.7	6.4	17.0	57.7	16.3	1.7	0.0
Cowley	1,087	0.5	7.1	16.4	62.2	13.2	0.6	0.0
Crawford	1,709	0.3	5.7	16.1	63.4	13.7	0.6	0.1
Doniphan	130	0.8	2.3	11.5	66.2	18.5	0.0	0.8
Douglas	1,562	0.1	3.8	11.1	63.8	19.8	1.5	0.0
Ford	2,057	0.2	6.4	12.9	55.5	22.5	2.4	0.0
Franklin	561	0.2	5.2	16.6	61.0	16.0	1.1	0.0
Geary	1,220	0.2	4.2	14.4	68.2	12.1	0.7	0.1
Graham	137	0.0	5.8	18.2	60.6	14.6	0.7	0.0
Greeley	41	*	*	*	*	*	*	*
Greenwood	131	0.0	4.6	16.0	62.6	16.0	0.0	0.8
Harvey	818	0.4	6.2	15.3	59.5	17.6	1.0	0.0
Jefferson	183	0.0	3.8	16.4	56.3	23.0	0.5	0.0
Johnson	5,804	0.1	3.3	10.1	60.8	23.8	1.8	0.1
Leavenworth	1,090	0.2	5.2	14.4	66.4	13.1	0.6	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4F
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Age Distribution by Local Agency
Combined 3 Years

Comparison of Age Distribution ⁽²⁾								
Local Agency	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Lincoln	394	0.0	3.3	13.2	62.2	19.8	1.5	0.0
Linn	206	0.0	6.3	13.1	62.1	18.0	0.5	0.0
Lyon	1,009	0.7	5.7	13.8	63.3	15.2	1.3	0.0
Marion	657	0.2	4.9	14.9	64.2	14.9	0.9	0.0
Marshall	160	0.0	5.0	17.5	63.8	13.1	0.6	0.0
Meade	312	0.0	6.7	11.5	60.3	20.5	1.0	0.0
Miami	455	0.4	4.2	17.6	65.3	11.9	0.7	0.0
NEK	962	0.1	7.0	15.1	63.4	13.8	0.6	0.0
Nemaha	123	0.0	3.3	11.4	65.9	13.0	5.7	0.8
Neosho	466	0.0	6.4	16.3	63.9	12.0	0.9	0.4
Osborne	1,154	0.3	4.2	14.4	63.9	15.9	1.0	0.2
Reno	1,821	0.3	5.9	15.0	63.9	14.2	0.6	0.0
Riley	3,076	0.0	1.7	11.2	73.8	12.8	0.3	0.0
Saline	1,563	0.3	6.6	15.0	62.3	14.7	1.2	0.1
Scott	175	0.0	2.9	11.4	64.6	19.4	1.7	0.0
Sedgwick	13,808	0.3	6.3	13.4	60.5	18.1	1.3	0.0
SEK	193	0.5	4.1	18.1	61.1	16.1	0.0	0.0
Seward	1,407	0.1	8.2	12.7	55.4	22.0	1.4	0.1
Shawnee	4,816	0.4	6.2	14.9	61.1	16.2	1.2	0.1

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4F
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Age Distribution by Local Agency
Combined 3 Years

Comparison of Age Distribution ⁽²⁾								
Local Agency	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Sheridan	26	*	*	*	*	*	*	*
Sherman	453	0.0	6.0	16.1	62.7	13.5	1.8	0.0
SKCPH	865	0.2	6.0	14.5	64.2	14.2	0.9	0.0
Stevens	231	0.0	6.9	15.2	56.7	19.5	1.7	0.0
Sumner	457	0.0	6.1	15.5	61.5	15.5	1.3	0.0
SW KS WIC	2,547	0.4	6.8	15.1	56.6	19.4	1.8	0.0
Wichita	44	*	*	*	*	*	*	*
Wilson	260	0.4	7.7	14.2	65.8	11.5	0.4	0.0
Wyandotte	6,487	0.4	7.8	13.3	57.1	19.8	1.6	0.0
Unknown Local Agencies	48	*	*	*	*	*	*	*
Kansas	65,683	0.3	5.8	13.8	61.5	17.4	1.2	0.1

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5F
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Health Indicators by Local Agency
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)														
Local Agency	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	%	%	Number	%	%	Number	%	Number	%	Number	%	Number	%
	(Rank)	(Rank)		(Rank)	(Rank)		(Rank)	(Rank)	(Rank)	(Rank)	(Rank)	(Rank)	(Rank)	
Barton	1,309	4.2 (19)	59.5 (36)	1,126	23.2 (33)	49.0 (15)	139	18.0 (2)	241	18.3 (7)	1,164	7.0 (26)	1,161	12.4(29)
Butler	791	5.9 (39)	53.6 (10)	711	21.2 (23)	51.9 (29)	118	44.1 (23)	137	30.7(27)	746	6.4 (17)	746	12.5(30)
Chautauqua	136	3.7 (15)	60.3 (39)	123	18.7 (5)	48.0 (10)	19	* *	22	* *	125	4.8 (10)	125	12.0(26)
Cherokee	2,021	6.4 (43)	53.5 (9)	1,804	22.8 (31)	48.3 (14)	350	35.7 (19)	447	25.3(21)	1,845	7.2 (30)	1,845	13.2(35)
Cloud	560	5.4 (36)	52.9 (7)	484	23.6 (34)	49.6 (17)	63	* *	144	23.6(17)	501	6.4 (17)	506	10.3(15)
Cowley	1,065	3.8 (16)	55.6 (23)	926	21.8 (25)	50.5 (20)	133	33.8 (17)	149	22.1(14)	947	6.9 (23)	949	11.3(20)
Crawford	1,678	5.8 (37)	55.6 (23)	1,457	21.8 (25)	51.7 (27)	224	22.8 (8)	277	18.1 (6)	1,490	8.7 (39)	1,496	13.7(38)
Doniphan	125	0.0 (1)	59.2 (34)	101	16.8 (3)	58.4 (44)	20	* *	29	* *	108	6.5 (19)	109	8.3(6)
Douglas	1,520	3.3 (10)	49.8 (1)	1,276	18.8 (7)	52.7 (34)	199	34.2 (18)	391	27.4(24)	1,297	8.0 (35)	1,297	8.8(9)
Ford	2,033	2.4 (4)	58.6 (33)	1,788	25.4 (41)	45.8 (5)	241	19.1 (3)	222	15.3 (1)	1,790	6.8 (22)	1,786	9.0(10)
Franklin	535	3.6 (13)	54.2 (13)	465	19.8 (12)	52.9 (36)	60	* *	148	15.5 (2)	404	5.7 (13)	404	11.6(23)
Geary	1,170	4.4 (23)	53.2 (8)	928	19.4 (10)	52.0 (31)	168	38.7 (21)	211	35.1(28)	969	4.5 (5)	967	10.2(14)
Graham	128	4.7 (30)	52.3 (5)	106	21.7 (24)	51.9 (29)	15	* *	23	* *	113	5.3 (11)	115	12.2(27)
Greeley	39	* *	* *	33	* *	* *	3	* *	6	* *	35	* *	35	* *
Greenwood	127	2.4 (4)	64.6 (44)	113	19.5 (11)	54.9 (41)	14	* *	18	* *	113	6.2 (14)	114	17.5(43)
Harvey	799	3.9 (17)	56.6 (31)	703	22.8 (31)	47.9 (9)	109	37.6 (20)	119	20.2(10)	723	4.6 (7)	722	13.0(34)
Jefferson	179	3.4 (11)	55.3 (20)	159	18.2 (4)	54.7 (40)	30	* *	51	* *	163	6.7 (20)	162	12.3(28)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5F
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Health Indicators by Local Agency
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)														
Local Agency	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Johnson	5,564	3.6 (13)	54.8 (17)	4,876	20.1 (13)	49.7 (18)	851	22.7 (7)	1,500	17.4 (4)	5,052	9.0 (40)	5,042	7.3(4)
Leavenwort	1,045	4.3 (21)	55.4 (21)	879	19.2 (9)	52.3 (33)	166	27.1 (11)	247	19.0 (8)	909	6.7 (20)	906	11.4(21)
Lincoln	381	2.6 (8)	61.4 (43)	343	30.0 (44)	49.9 (19)	51	* *	61	* *	355	10.1 (43)	355	13.2(35)
Linn	206	6.3 (42)	60.7 (41)	185	24.3 (39)	48.1 (12)	25	* *	25	* *	185	7.6 (33)	186	7.0(3)
Lyon	978	2.5 (6)	59.8 (37)	845	23.6 (34)	46.9 (7)	120	16.7 (1)	177	20.3(11)	870	7.4 (32)	869	13.8(39)
Marion	635	3.5 (12)	57.0 (32)	543	22.7 (30)	48.1 (12)	88	* *	131	21.4(13)	571	4.2 (4)	569	10.7(17)
Marshall	155	5.2 (35)	55.5 (22)	141	27.7 (43)	43.3 (3)	19	* *	40	* *	146	9.6 (41)	145	13.8(39)
Meade	305	3.0 (9)	56.1 (28)	268	20.1 (13)	53.7 (38)	30	* *	58	* *	272	9.9 (42)	271	11.1(18)
Miami	439	4.3 (21)	55.8 (27)	386	14.5 (1)	56.2 (43)	80	* *	113	26.5(22)	411	6.3 (16)	411	11.9(25)
NEK	932	4.7 (30)	55.6 (23)	808	20.3 (18)	52.7 (34)	120	26.7 (10)	188	27.1(23)	799	6.9 (23)	800	14.1(41)
Nemaha	112	0.9 (2)	60.7 (41)	100	19.0 (8)	56.0 (42)	22	* *	22	* *	110	4.5 (5)	110	9.1(11)
Neosho	451	5.8 (37)	52.5 (6)	407	22.4 (28)	51.1 (26)	71	* *	70	* *	422	7.6 (33)	421	14.3(42)
Osborne	1,121	4.4 (23)	51.0 (3)	970	20.9 (21)	45.9 (6)	188	21.3 (4)	234	15.8 (3)	1,007	4.0 (3)	1,006	9.3(13)
Reno	1,781	4.4 (23)	55.1 (19)	1,558	20.2 (15)	50.7 (22)	216	30.6 (14)	308	23.4(16)	1,597	4.6 (7)	1,600	12.9(33)
Riley	2,924	4.4 (23)	50.6 (2)	2,373	18.7 (5)	53.2 (37)	468	33.3 (16)	717	25.1(20)	2,485	6.2 (14)	2,490	10.6(16)
Saline	1,532	4.5 (27)	55.7 (26)	1,374	16.4 (2)	54.2 (39)	252	22.6 (6)	247	19.0 (8)	1,392	7.2 (30)	1,394	8.2(5)
Scott	172	4.7 (30)	56.4 (30)	152	21.1 (22)	48.0 (10)	24	* *	31	* *	152	2.6 (1)	152	11.2(19)

- (1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5F
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Health Indicators by Local Agency
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)														
Local Agency	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Sedgwick	13,402	4.1 (18)	54.6 (15)	12,083	20.8 (20)	50.8 (24)	2,584	30.1 (13)	2,990	23.7 (18)	12,460	7.1 (28)	12,457	11.4 (21)
SEK	189	4.8 (34)	51.9 (4)	170	25.3 (40)	47.6 (8)	26	* *	43	* *	175	6.9 (23)	175	12.6 (31)
Seward	1,381	2.5 (6)	59.2 (34)	1,244	24.1 (38)	45.0 (4)	230	27.4 (12)	188	28.7 (25)	1,266	8.5 (36)	1,267	5.8 (1)
Shawnee	4,690	4.6 (28)	54.1 (12)	4,078	20.2 (15)	51.7 (27)	474	43.9 (22)	767	29.5 (26)	4,184	7.1 (28)	4,186	8.4 (7)
Sheridan	25	* *	* *	23	* *	* *	4	* *	3	* *	25	* *	25	* *
Sherman	439	4.6 (28)	54.7 (16)	377	22.0 (27)	50.9 (25)	56	* *	82	* *	390	3.8 (2)	390	13.6 (37)
SKCPH	847	4.7 (30)	54.9 (18)	724	22.4 (28)	49.3 (16)	101	21.8 (5)	159	22.6 (15)	743	4.6 (7)	743	11.6 (23)
Stevens	225	2.2 (3)	60.4 (40)	198	20.2 (15)	50.5 (20)	44	* *	61	* *	201	15.9 (44)	202	8.4 (7)
Sumner	440	5.9 (39)	54.3 (14)	397	23.7 (36)	41.3 (1)	63	* *	112	17.9 (5)	413	7.0 (26)	414	17.9 (44)
SW KS WIC	2,495	6.2 (41)	53.9 (11)	2,240	25.7 (42)	42.9 (2)	333	26.4 (9)	407	20.9 (12)	2,292	8.6 (37)	2,287	6.7 (2)
Wichita	42	* *	* *	38	* *	* *	6	* *	2	* *	40	* *	40	* *
Wilson	256	7.0 (44)	59.8 (37)	242	24.0 (37)	52.1 (32)	42	* *	37	* *	244	8.6 (37)	244	12.7 (32)
Wyandotte	6,299	4.2 (19)	56.1 (28)	5,579	20.7 (19)	50.7 (22)	971	33.1 (15)	1,154	24.1 (19)	5,666	5.4 (12)	5,663	9.1 (11)
Unknown Local Agencies	46	* *	* *	44	* *	* *	3	* *	6	* *	46	* *	46	* *
Kansas	63,724	4.3	54.9	55,948	21.2	50.1	9,633	29.8	12,815	22.8	57,413	6.9	57,405	10.4

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6F
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Behavioral Indicators by Local Agency
Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2, 3)											
Local Agency	Medical Care				WIC Enrollment			Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester			Prepregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	% (Rank)		Number	% (Rank)		Number	% (Rank)	Number	% (Rank)
Barton	1,152	84.8 (16)	1.1 (7)		1,331	47.1 (6)		1,342	20.0 (21)	1,087	56.5 (36)
Butler	727	75.1 (33)	5.8 (31)		819	36.1 (31)		820	16.5 (25)	629	45.2 (41)
Chautauqua	124	72.6 (38)	8.9 (37)		137	48.9 (2)		139	20.1 (20)	121	77.7 (1)
Cherokee	1,421	84.8 (16)	3.8 (21)		2,092	32.5 (41)		2,102	13.4 (34)	1,727	64.6 (20)
Cloud	507	87.4 (8)	3.0 (15)		571	47.6 (4)		575	24.2 (11)	499	70.7 (5)
Cowley	850	70.2 (39)	6.1 (33)		1,075	42.0 (17)		1,087	14.1 (31)	902	57.9 (30)
Crawford	1,372	87.2 (9)	3.2 (16)		1,706	41.5 (18)		1,709	14.2 (30)	1,438	57.2 (34)
Doniphan	111	82.0 (24)	4.5 (25)		130	33.8 (39)		130	36.2 (4)	96	* *
Douglas	1,262	74.2 (34)	7.0 (35)		1,557	38.2 (27)		1,562	20.2 (18)	1,260	63.9 (21)
Ford	1,679	72.8 (37)	12.9 (41)		2,044	44.6 (11)		2,057	11.3 (38)	1,790	55.0 (37)
Franklin	235	79.1 (27)	6.0 (32)		557	39.5 (25)		561	8.9 (42)	429	60.6 (27)
Geary	911	74.2 (34)	1.3 (8)		1,218	35.1 (34)		1,220	20.4 (17)	997	59.7 (29)
Graham	115	92.2 (2)	1.7 (10)		136	47.1 (6)		137	28.5 (5)	111	63.1 (22)
Greeley	29	* *	* *		41	* *		41	* *	31	* *
Greenwood	110	60.9 (43)	30.9 (44)		131	43.5 (14)		131	16.8 (24)	111	74.8 (3)
Harvey	669	76.7 (30)	10.0 (39)		813	47.8 (3)		818	19.6 (22)	685	67.4 (13)
Jefferson	158	83.5 (21)	9.5 (38)		183	44.3 (13)		183	9.8 (41)	152	41.4 (43)
Johnson	4,612	66.4 (40)	3.4 (17)		5,780	33.3 (40)		5,804	14.4 (29)	4,677	61.8 (25)
Leavenworth	913	86.7 (10)	0.8 (3)		1,086	31.2 (42)		1,090	22.1 (14)	840	71.8 (4)
Lincoln	356	89.9 (3)	1.7 (10)		393	42.2 (16)		394	26.6 (7)	338	60.4 (28)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6F
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Behavioral Indicators by Local Agency
Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2, 3)										
Local Agency	Medical Care			WIC Enrollment			Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester			Prepregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	% (Rank)	Number	% (Rank)		Number	% (Rank)	Number	% (Rank)
Linn	188	80.3 (26)	8.0 (36)	206	47.6 (4)		206	23.3 (12)	181	65.2 (17)
Lyon	854	87.6 (7)	0.7 (2)	1,006	41.5 (18)		1,009	18.0 (23)	835	63.0 (23)
Marion	559	84.8 (16)	1.4 (9)	649	40.8 (21)		657	25.0 (10)	525	69.7 (10)
Marshall	146	93.2 (1)	0.0 (1)	160	44.4 (12)		160	45.6 (3)	133	69.9 (7)
Meade	265	85.3 (15)	3.8 (21)	307	50.8 (1)		312	46.5 (2)	275	69.8 (9)
Miami	406	82.3 (22)	1.0 (6)	455	34.3 (37)		455	14.9 (28)	354	57.9 (30)
NEK	685	85.5 (14)	5.1 (28)	957	40.1 (22)		962	12.8 (35)	804	69.3 (11)
Nemaha	109	89.0 (5)	3.7 (19)	123	35.8 (32)		123	13.8 (32)	109	65.1 (18)
Neosho	387	89.4 (4)	4.1 (24)	464	39.0 (26)		466	12.7 (36)	391	70.3 (6)
Osborne	988	86.2 (12)	2.8 (13)	1,145	40.1 (22)		1,154	25.7 (9)	961	56.7 (35)
Reno	1,567	84.1 (20)	2.4 (12)	1,812	40.0 (24)		1,821	21.6 (16)	1,495	62.8 (24)
Riley	2,317	84.2 (19)	0.9 (4)	3,074	34.1 (38)		3,076	22.0 (15)	2,457	65.7 (15)
Saline	1,126	76.4 (31)	4.7 (26)	1,559	35.0 (35)		1,563	6.0 (44)	1,315	57.6 (32)
Scott	152	78.9 (28)	5.3 (30)	172	44.8 (10)		175	74.9 (1)	154	53.2 (38)
Sedgwick	12,117	75.5 (32)	4.8 (27)	13,771	28.6 (43)		13,808	16.4 (27)	11,304	49.8 (40)
SEK	154	85.7 (13)	5.2 (29)	193	43.0 (15)		193	20.2 (18)	172	65.7 (15)
Seward	1,111	64.5 (41)	13.8 (42)	1,397	34.4 (36)		1,407	10.2 (40)	1,163	66.7 (14)
Shawnee	3,701	77.8 (29)	3.7 (19)	4,792	45.0 (8)		4,816	11.7 (37)	4,044	51.7 (39)
Sheridan	25	* *	* *	26	* *		26	* *	20	* *
Sherman	278	86.3 (11)	2.9 (14)	453	38.2 (27)		453	27.2 (6)	373	75.9 (2)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6F
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Behavioral Indicators by Local Agency
Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2, 3)										
Local Agency	Medical Care			WIC Enrollment			Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester			Pregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	% (Rank)	Number	% (Rank)	%	Number	% (Rank)	Number	% (Rank)
SKCPH	734	81.1 (25)	6.1 (33)	855	44.9 (9)		865	22.3 (13)	710	69.9 (7)
Stevens	201	73.6 (36)	4.0 (23)	228	28.1 (44)		231	26.0 (8)	188	69.1 (12)
Sumner	413	82.1 (23)	3.4 (17)	457	37.2 (29)		457	13.6 (33)	371	64.7 (19)
SW KS WIC	2,111	55.7 (44)	26.7 (43)	2,525	35.4 (33)		2,547	11.2 (39)	2,094	42.2 (42)
Wichita	40	* *	* *	44	* *		44	* *	35	* *
Wilson	235	88.9 (6)	0.9 (4)	258	41.5 (18)		260	16.5 (25)	224	61.6 (26)
Wyandotte	5,159	63.0 (42)	11.4 (40)	6,442	36.2 (30)		6,487	7.7 (43)	5,470	57.4 (33)
Unknown Local Agencies	37	* *	* *	48	* *		48	* *	40	* *
Kansas	53,378	75.5	6.0	65,378	36.4		65,683	15.8	54,117	57.7

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7F
2009 - 2011 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by Local Agency
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
Local Agency	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Barton	1,317	37.9 (16)	1,127	24.2 (19)	337	29.7 (17)	1,069	14.2 (15)	1,126	8.3 (8)
Butler	807	49.6 (38)	716	30.6 (36)	276	31.5 (15)	623	11.1 (6)	723	6.8 (4)
Chautauqua	123	43.1 (29)	93	* *	35	* *	108	47.2 (41)	92	* *
Cherokee	2,033	40.1 (20)	1,767	27.9 (32)	566	20.8 (29)	1,695	33.8 (33)	1,770	33.7 (33)
Cloud	574	41.6 (24)	495	22.4 (17)	165	28.5 (20)	496	19.8 (22)	500	13.4 (18)
Cowley	1,053	40.4 (22)	784	27.6 (30)	259	26.6 (25)	878	20.4 (23)	766	19.6 (25)
Crawford	1,606	45.2 (34)	1,253	26.9 (28)	445	29.2 (18)	1,380	40.4 (36)	1,261	38.7 (36)
Doniphan	121	38.0 (18)	89	* *	19	* *	90	* *	90	* *
Douglas	1,532	37.5 (15)	1,249	18.7 (13)	373	42.4 (2)	1,243	10.5 (5)	1,268	6.9 (5)
Ford	1,990	8.8 (2)	1,635	5.0 (2)	103	29.1 (19)	1,747	8.8 (3)	1,639	8.1 (7)
Franklin	558	43.2 (30)	480	29.6 (34)	145	21.4 (28)	427	12.9 (11)	484	8.7 (9)
Geary	1,200	36.9 (14)	942	20.2 (14)	254	35.0 (9)	990	34.3 (34)	949	28.9 (32)
Graham	132	40.2 (21)	108	25.0 (22)	38	* *	108	29.6 (29)	109	28.4 (30)
Greeley	40	* *	33	* *	5	* *	31	* *	33	* *
Greenwood	110	53.6 (42)	57	* *	23	* *	102	31.4 (31)	83	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7F
2009 - 2011 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by Local Agency
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
Local Agency	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Harvey	804	37.9 (16)	692	21.1 (15)	212	37.7 (6)	674	6.1 (1)	692	2.9 (1)
Jefferson	81	* *	40	* *	14	* *	75	* *	43	* *
Johnson	5,734	23.5 (7)	5,016	9.8 (4)	829	46.4 (1)	4,627	18.0 (20)	5,021	17.7 (24)
Leavenworth	1,068	46.0 (37)	879	24.5 (21)	287	36.2 (8)	832	37.7 (35)	880	38.4 (35)
Lincoln	386	43.5 (32)	316	26.3 (25)	118	32.2 (14)	336	17.9 (19)	331	14.8 (21)
Linn	203	52.7 (41)	182	32.4 (39)	86	* *	179	14.5 (16)	183	10.4 (15)
Lyon	992	32.2 (12)	862	16.8 (11)	222	38.7 (4)	830	12.9 (11)	845	8.8 (11)
Marion	650	39.4 (19)	561	24.4 (20)	173	32.4 (12)	518	8.5 (2)	562	6.6 (3)
Marshall	156	44.9 (33)	135	26.7 (27)	49	* *	129	46.5 (40)	136	41.2 (37)
Meade	295	19.0 (4)	253	14.2 (7)	45	* *	267	14.6 (17)	259	13.1 (17)
Miami	438	42.9 (28)	380	26.6 (26)	122	27.0 (23)	336	20.8 (24)	380	16.1 (23)
NEK	832	45.6 (36)	603	31.0 (38)	211	23.7 (27)	702	31.8 (32)	614	28.5 (31)
Nemaha	88	* *	64	* *	24	* *	73	* *	67	* *
Neosho	396	43.4 (31)	231	27.7 (31)	82	* *	346	30.9 (30)	237	21.9 (27)
Osborne	1,120	42.1 (26)	920	25.1 (23)	314	30.9 (16)	926	25.1 (26)	932	21.7 (26)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7F
2009 - 2011 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by Local Agency
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
Local Agency	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Reno	1,799	42.2 (27)	1,562	23.4 (18)	527	37.8 (5)	1,475	13.1 (13)	1,568	9.9 (13)
Riley	2,984	28.2 (8)	2,264	14.3 (8)	481	36.8 (7)	2,425	12.3 (10)	2,292	8.7 (9)
Saline	1,522	40.9 (23)	1,305	25.4 (24)	435	26.9 (24)	1,289	11.8 (8)	1,308	9.1 (12)
Scott	143	29.4 (9)	101	14.9 (9)	26	* *	127	23.6 (25)	105	15.2 (22)
Sedgwick	13,698	29.8 (10)	12,370	16.4 (10)	2,816	34.2 (10)	11,239	9.4 (4)	12,371	6.9 (5)
SEK	188	45.2 (34)	156	26.9 (28)	57	* *	168	25.6 (27)	159	22.6 (28)
Seward	1,378	5.3 (1)	1,232	2.7 (1)	46	* *	1,153	12.0 (9)	1,247	9.9 (13)
Shawnee	4,492	34.7 (13)	3,473	22.0 (16)	924	27.8 (21)	3,745	11.7 (7)	3,483	6.2 (2)
Sheridan	24	* *	21	* *	4	* *	20	* *	23	* *
Sherman	445	30.1 (11)	387	17.1 (12)	97	* *	364	18.7 (21)	389	14.1 (20)
SKCPH	834	41.8 (25)	689	27.9 (32)	234	25.2 (26)	683	40.4 (36)	683	36.7 (34)
Stevens	228	21.5 (6)	199	11.1 (5)	32	* *	186	26.9 (28)	202	24.3 (29)
Sumner	451	52.5 (40)	400	30.8 (37)	167	32.3 (13)	363	43.0 (38)	402	43.5 (39)
SW KS WIC	2,521	10.9 (3)	2,216	5.2 (3)	176	39.2 (3)	2,084	13.3 (14)	2,215	12.7 (16)
Wichita	43	* *	34	* *	7	* *	34	* *	34	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7F
2009 - 2011 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by Local Agency
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
Local Agency	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Wilson	256	51.2 (39)	238	30.3 (35)	105	33.3 (11)	222	45.9 (39)	237	42.2 (38)
Wyandotte	6,373	19.9 (5)	5,554	11.6 (6)	841	27.8 (21)	5,418	16.7 (18)	5,656	13.9 (19)
Unknown Local Agencies	47	* *	45	* *	19	* *	40	* *	45	* *
Kansas	63,865	30.9	54,208	17.6	12,825	32.5	52,872	16.8	54,494	14.0

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8F
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Infant Health Indicators by Local Agency
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)										
Local Agency	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)		Number	%	Number	%	Number	%
	Number	% (Rank)	% (Rank)	Number						
Barton	1,101	5.8 (4)	4.6 (8)	1,106	8.1 (13)	1,000	3.7 (10)	1,110	74.1 (16)	
Butler	718	6.8 (16)	6.1 (20)	730	7.7 (10)	661	4.1 (17)	727	70.4 (28)	
Chautauqua	119	8.4 (37)	4.2 (5)	123	6.5 (5)	110	5.5 (37)	125	70.4 (28)	
Cherokee	1,817	7.4 (23)	5.6 (13)	1,822	8.6 (19)	1,655	4.3 (23)	1,819	58.8 (43)	
Cloud	482	7.3 (22)	6.0 (17)	488	10.0 (35)	430	3.0 (3)	489	74.4 (15)	
Cowley	891	7.5 (25)	7.1 (29)	929	10.4 (39)	786	4.7 (32)	930	61.4 (41)	
Crawford	1,418	8.1 (32)	3.9 (2)	1,461	10.2 (37)	1,266	4.6 (29)	1,460	61.4 (41)	
Doniphan	107	12.1 (43)	10.3 (43)	108	13.9 (43)	92	* *	107	64.5 (37)	
Douglas	1,268	6.8 (16)	6.7 (25)	1,293	7.5 (8)	1,169	4.1 (17)	1,288	79.2 (4)	
Ford	1,681	5.6 (3)	5.7 (15)	1,726	8.5 (17)	1,529	2.9 (2)	1,719	75.6 (14)	
Franklin	473	5.9 (6)	5.1 (10)	474	9.1 (26)	426	3.8 (13)	477	64.6 (36)	
Geary	954	7.8 (28)	7.7 (34)	956	8.3 (14)	873	4.2 (22)	950	72.6 (18)	
Graham	107	7.5 (25)	8.4 (39)	111	9.9 (33)	96	* *	108	71.3 (23)	
Greeley	34	* *	* *	34	* *	30	* *	34	* *	

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8F
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Infant Health Indicators by Local Agency
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)									
Local Agency	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Greenwood	110	3.6 (1)	4.5 (7)	111	4.5 (1)	105	3.8 (13)	111	69.4 (31)
Harvey	688	7.4 (23)	6.8 (26)	693	8.4 (16)	625	4.5 (27)	695	83.7 (1)
Jefferson	159	6.3 (8)	4.4 (6)	160	15.0 (44)	135	3.7 (10)	158	67.1 (32)
Johnson	4,728	6.8 (16)	7.2 (30)	4,982	6.9 (7)	4,379	3.8 (13)	4,973	81.5 (3)
Leavenworth	879	8.1 (32)	7.5 (32)	911	8.6 (19)	801	5.1 (34)	910	64.8 (34)
Lincoln	326	8.0 (30)	7.7 (34)	349	6.6 (6)	304	4.3 (23)	347	71.8 (20)
Linn	178	6.7 (14)	3.9 (2)	183	10.9 (41)	158	4.4 (26)	183	76.5 (12)
Lyon	855	7.8 (28)	6.5 (23)	852	8.3 (14)	781	4.5 (27)	850	75.9 (13)
Marion	533	6.8 (16)	8.6 (40)	537	9.3 (30)	475	3.8 (13)	543	77.3 (8)
Marshall	137	8.8 (38)	2.2 (1)	141	9.9 (33)	123	6.5 (40)	140	64.3 (38)
Meade	239	6.7 (14)	9.6 (42)	261	9.2 (28)	214	5.1 (34)	261	77.8 (7)
Miami	407	6.6 (12)	8.8 (41)	410	7.6 (9)	377	3.7 (10)	407	70.0 (30)
NEK	782	8.8 (38)	6.6 (24)	830	8.6 (19)	711	5.6 (38)	828	58.1 (44)
Nemaha	73	*	*	109	6.4 (4)	68	*	104	61.5 (40)

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.
- (4) <2500 g.
- (5) >4000 g.
- (6) <37 weeks gestation.
- (7) 37 or more weeks gestation and <2500 g.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8F
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Infant Health Indicators by Local Agency
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)									
Local Agency	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Neosho	330	6.4 (10)	5.2 (11)	416	8.9 (23)	299	3.0 (3)	414	64.7 (35)
Osborne	942	6.6 (12)	6.4 (22)	967	8.9 (23)	851	3.2 (5)	967	76.7 (10)
Reno	1,546	7.2 (21)	5.4 (12)	1,558	8.0 (11)	1,416	4.1 (17)	1,561	70.8 (25)
Riley	2,454	6.1 (7)	7.9 (36)	2,479	6.2 (3)	2,300	3.5 (9)	2,467	79.2 (4)
Saline	1,337	7.7 (27)	6.2 (21)	1,384	9.5 (32)	1,205	4.3 (23)	1,380	66.4 (33)
Scott	146	8.2 (35)	4.8 (9)	146	10.3 (38)	128	4.7 (32)	148	72.3 (19)
Sedgwick	12,156	9.0 (40)	6.8 (26)	12,213	8.5 (17)	11,100	5.1 (34)	12,187	70.8 (25)
SEK	149	10.1 (41)	6.0 (17)	172	8.7 (22)	137	5.8 (39)	171	62.0 (39)
Seward	1,156	5.4 (2)	7.9 (36)	1,215	9.1 (26)	1,040	3.4 (8)	1,216	73.6 (17)
Shawnee	4,064	8.0 (30)	5.7 (15)	4,108	10.1 (36)	3,632	4.6 (29)	4,100	70.6 (27)
Sheridan	25	*	*	25	*	22	*	25	*
Sherman	375	8.3 (36)	4.0 (4)	375	8.0 (11)	345	4.1 (17)	375	81.6 (2)
SKCPH	664	6.5 (11)	8.0 (38)	706	8.9 (23)	594	3.2 (5)	709	76.7 (10)
Stevens	121	5.8 (4)	7.4 (31)	196	5.1 (2)	112	2.7 (1)	197	78.7 (6)

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.
- (4) <2500 g.
- (5) >4000 g.
- (6) <37 weeks gestation.
- (7) 37 or more weeks gestation and <2500 g.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8F
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Infant Health Indicators by Local Agency
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)										
Local Agency	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)							
	Number	% (Rank)	%	(Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Sumner	395	11.6 (42)	6.8	(26)	407	12.5 (42)	344	6.7 (41)	406	71.7 (21)
SW KS WIC	2,193	6.9 (20)	5.6	(13)	2,231	9.3 (30)	1,974	4.1 (17)	2,228	76.9 (9)
Wichita	38	* * *	* *		38	* *	34	* *	38	* *
Wilson	239	6.3 (8)	7.5	(32)	242	10.7 (40)	212	3.3 (7)	241	71.0 (24)
Wyandotte	5,426	8.1 (32)	6.0	(17)	5,594	9.2 (28)	4,902	4.6 (29)	5,581	71.4 (22)
Unknown Local Agencies	47	* * *	* *		47	* *	47	* *	46	* *
Kansas	55,067	7.6	6.4		56,409	8.6	50,073	4.4	56,310	72.2

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

