

Table 3A
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Clinic	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaska Native	Asian/Pacific Islander	Multiple Races	All Other/Unknown
		%	%	%	%	%	%	%
Allen	112	92.0	5.4	1.8	0.9	0.0	0.0	0.0
Anderson	50	*	*	*	*	*	*	*
Atchison	118	89.8	5.9	1.7	0.0	0.0	2.5	0.0
Barber-Kio	2	*	*	*	*	*	*	*
Barber-Med	37	*	*	*	*	*	*	*
Barton	279	70.6	2.5	26.2	0.4	0.0	0.4	0.0
Bourbon	135	91.9	3.0	2.2	0.0	0.7	2.2	0.0
Brown	110	76.4	2.7	3.6	12.7	0.0	4.5	0.0
Butler	286	92.7	1.0	4.2	0.0	0.0	2.1	0.0
Chase	15	*	*	*	*	*	*	*
Chautauqua	32	*	*	*	*	*	*	*
Cherokee	219	90.9	1.4	4.6	1.4	0.5	1.4	0.0
Clark	4	*	*	*	*	*	*	*
Clay	67	*	*	*	*	*	*	*
Cloud	72	*	*	*	*	*	*	*
Coffey	24	*	*	*	*	*	*	*
Comanche	6	*	*	*	*	*	*	*
Cowley-Ark	201	66.7	3.0	28.9	0.5	0.5	0.5	0.0
Cowley-Win	155	88.4	1.9	3.9	1.3	3.9	0.6	0.0
Crawford-G	1	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3A
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Clinic	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaska Native	Asian/Pacific Islander	Multiple Races	All Other/Unknown
		%	%	%	%	%	%	%
Crawford-P	381	82.7	3.9	11.0	0.0	1.3	1.0	0.0
Decatur	9	*	*	*	*	*	*	*
DickinsonA	94	*	*	*	*	*	*	*
DickinsonH	11	*	*	*	*	*	*	*
Doniphan	43	*	*	*	*	*	*	*
Douglas	463	65.7	6.0	10.2	10.2	5.2	2.8	0.0
Edwards	27	*	*	*	*	*	*	*
Elk	13	*	*	*	*	*	*	*
Ellis	143	83.9	2.1	14.0	0.0	0.0	0.0	0.0
Ellsworth	28	*	*	*	*	*	*	*
Finney	688	30.5	0.9	65.7	0.0	2.5	0.4	0.0
Ford	646	22.1	1.2	76.3	0.2	0.2	0.0	0.0
Franklin	166	89.8	2.4	4.2	1.8	0.0	1.8	0.0
Geary	340	53.5	23.5	15.0	0.0	2.9	5.0	0.0
Gove	7	*	*	*	*	*	*	*
Graham	14	*	*	*	*	*	*	*
Grant	68	*	*	*	*	*	*	*
Gray	57	*	*	*	*	*	*	*
Greeley	10	*	*	*	*	*	*	*
Greenwood	44	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3A
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Clinic	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaska Native	Asian/Pacific Islander	Multiple Races	All Other/Unknown
		%	%	%	%	%	%	%
Hamilton	18	*	*	*	*	*	*	*
Harper	51	*	*	*	*	*	*	*
Harvey	251	69.3	1.6	27.5	0.4	0.0	1.2	0.0
Haskell	24	*	*	*	*	*	*	*
Hodgeman	12	*	*	*	*	*	*	*
Jackson	70	*	*	*	*	*	*	*
Jefferson	65	*	*	*	*	*	*	*
Jewell	14	*	*	*	*	*	*	*
Johnson-M	862	45.2	14.5	35.2	0.7	2.6	1.9	0.0
Johnson-O	982	46.0	11.2	37.4	0.5	3.3	1.6	0.0
Kearny	21	*	*	*	*	*	*	*
Kingman	57	*	*	*	*	*	*	*
Kiowa	9	*	*	*	*	*	*	*
Labette	166	88.0	7.2	4.2	0.6	0.0	0.0	0.0
Lane	13	*	*	*	*	*	*	*
Leavenworth	356	66.0	17.4	8.7	0.6	2.5	4.8	0.0
Lincoln	22	*	*	*	*	*	*	*
Linn	54	*	*	*	*	*	*	*
Logan	13	*	*	*	*	*	*	*
Lyon	343	58.9	5.0	34.7	0.0	0.9	0.6	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3A
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Clinic	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaska Native	Asian/Pacific Islander	Multiple Races	All Other/Unknown
		%	%	%	%	%	%	%
Marion	46	*	*	*	*	*	*	*
Marshall	63	*	*	*	*	*	*	*
Mc Pherson	145	89.7	2.1	7.6	0.7	0.0	0.0	0.0
Meade	49	*	*	*	*	*	*	*
Miami	136	90.4	2.2	4.4	0.0	1.5	1.5	0.0
Mitchell	43	*	*	*	*	*	*	*
Montgome-C	152	74.3	9.2	11.8	1.3	1.3	2.0	0.0
Montgome-I	178	84.8	6.7	5.1	2.2	1.1	0.0	0.0
Morris	32	*	*	*	*	*	*	*
Nemaha	24	*	*	*	*	*	*	*
Neosho-Ch	143	92.3	2.8	2.8	0.7	0.7	0.7	0.0
Neosho-Er	10	*	*	*	*	*	*	*
Ness	8	*	*	*	*	*	*	*
Norton	26	*	*	*	*	*	*	*
Osage	89	*	*	*	*	*	*	*
Osborne	19	*	*	*	*	*	*	*
Ottawa	22	*	*	*	*	*	*	*
Pawnee	43	*	*	*	*	*	*	*
Phillips	33	*	*	*	*	*	*	*
Pottawat-S	12	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3A
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Clinic	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaska Native	Asian/Pacific Islander	Multiple Races	All Other/Unknown
		%	%	%	%	%	%	%
Pottawat-W	28	*	*	*	*	*	*	*
Pratt	68	*	*	*	*	*	*	*
Rawlins	10	*	*	*	*	*	*	*
Reno	494	81.0	1.8	14.4	0.8	0.8	1.2	0.0
Republic	33	*	*	*	*	*	*	*
Rice	70	*	*	*	*	*	*	*
Riley-Ft	428	69.4	13.1	13.6	0.7	3.3	0.0	0.0
Riley-Manh	398	73.1	11.8	8.5	1.3	5.0	0.3	0.0
Rooks	36	*	*	*	*	*	*	*
Rush	12	*	*	*	*	*	*	*
Russell	40	*	*	*	*	*	*	*
Saline	493	73.8	4.1	18.3	0.2	2.2	1.4	0.0
Scott	36	*	*	*	*	*	*	*
Sedgwick-C	1,078	41.9	13.4	34.9	1.0	6.8	2.0	0.0
Sedgwick-M	1,940	33.4	28.3	33.5	1.0	2.8	1.0	0.0
Sedgwick-S	1,131	67.7	7.3	17.5	2.8	2.7	1.9	0.0
Seneca	13	*	*	*	*	*	*	*
Seward	496	16.1	1.0	77.8	3.4	1.2	0.4	0.0
ShawneeM&I	168	35.1	14.3	48.2	1.8	0.6	0.0	0.0
ShawneeMa	1,183	57.1	15.6	22.6	1.8	0.8	2.1	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3A
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Sheridan	7	*	*	*	*	*	*	*
Sherman	70	*	*	*	*	*	*	*
Smith	17	*	*	*	*	*	*	*
Stafford	40	*	*	*	*	*	*	*
Stanton	23	*	*	*	*	*	*	*
Stevens	66	*	*	*	*	*	*	*
Sumner	158	90.5	0.0	7.6	1.3	0.6	0.0	0.0
Thomas	50	*	*	*	*	*	*	*
Trego	14	*	*	*	*	*	*	*
Wabaunsee	18	*	*	*	*	*	*	*
Wallace	1	*	*	*	*	*	*	*
Washington	25	*	*	*	*	*	*	*
Wichita	11	*	*	*	*	*	*	*
Wilson	81	*	*	*	*	*	*	*
Woodson	2	*	*	*	*	*	*	*
Wyandotte	2,112	27.3	31.0	39.3	0.2	1.8	0.4	0.0
Kansas	20,703	56.9	11.3	27.3	1.2	2.0	1.2	0.0
Nation (Prior Year)	1,252,728	41.7	23.3	27.9	1.1	3.2	1.0	1.8

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Age Distribution by Clinic

Comparison of Age Distribution ⁽²⁾								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Allen	112	0.0	6.3	15.2	63.4	13.4	0.9	0.9
Anderson	50	*	*	*	*	*	*	*
Atchison	118	0.0	6.8	16.9	62.7	11.9	1.7	0.0
Barber-Kio	2	*	*	*	*	*	*	*
Barber-Med	37	*	*	*	*	*	*	*
Barton	279	0.0	6.5	19.0	57.3	16.8	0.4	0.0
Bourbon	135	0.0	4.4	20.0	65.9	9.6	0.0	0.0
Brown	110	0.0	7.3	16.4	65.5	10.0	0.9	0.0
Butler	286	0.3	7.0	16.1	61.9	14.0	0.7	0.0
Chase	15	*	*	*	*	*	*	*
Chautauqua	32	*	*	*	*	*	*	*
Cherokee	219	0.5	9.1	17.8	56.6	14.6	0.9	0.5
Clark	4	*	*	*	*	*	*	*
Clay	67	*	*	*	*	*	*	*
Cloud	72	*	*	*	*	*	*	*
Coffey	24	*	*	*	*	*	*	*
Comanche	6	*	*	*	*	*	*	*
Cowley-Ark	201	0.5	10.4	14.9	60.7	12.9	0.5	0.0
Cowley-Win	155	0.6	7.1	16.1	61.9	14.2	0.0	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Age Distribution by Clinic

Comparison of Age Distribution ⁽²⁾								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Crawford-G	1	*	*	*	*	*	*	*
Crawford-P	381	0.3	8.9	13.6	63.5	13.1	0.5	0.0
Decatur	9	*	*	*	*	*	*	*
DickinsonA	94	*	*	*	*	*	*	*
DickinsonH	11	*	*	*	*	*	*	*
Doniphan	43	*	*	*	*	*	*	*
Douglas	463	0.0	4.1	13.6	63.5	17.1	1.7	0.0
Edwards	27	*	*	*	*	*	*	*
Elk	13	*	*	*	*	*	*	*
Ellis	143	0.0	4.2	11.2	69.9	11.9	2.8	0.0
Ellsworth	28	*	*	*	*	*	*	*
Finney	688	0.1	10.5	16.0	53.8	18.2	1.2	0.3
Ford	646	0.3	7.3	12.8	54.8	24.0	0.8	0.0
Franklin	166	0.6	7.2	18.1	58.4	15.1	0.0	0.6
Geary	340	0.6	5.6	17.9	65.0	10.3	0.6	0.0
Gove	7	*	*	*	*	*	*	*
Graham	14	*	*	*	*	*	*	*
Grant	68	*	*	*	*	*	*	*
Gray	57	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Age Distribution by Clinic

Comparison of Age Distribution ⁽²⁾								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Greeley	10	*	*	*	*	*	*	*
Greenwood	44	*	*	*	*	*	*	*
Hamilton	18	*	*	*	*	*	*	*
Harper	51	*	*	*	*	*	*	*
Harvey	251	0.0	5.6	8.0	66.9	16.7	2.8	0.0
Haskell	24	*	*	*	*	*	*	*
Hodgeman	12	*	*	*	*	*	*	*
Jackson	70	*	*	*	*	*	*	*
Jefferson	65	*	*	*	*	*	*	*
Jewell	14	*	*	*	*	*	*	*
Johnson-M	862	0.2	4.1	13.6	61.0	20.2	0.9	0.0
Johnson-O	982	0.3	5.1	10.1	60.5	22.5	1.4	0.1
Kearny	21	*	*	*	*	*	*	*
Kingman	57	*	*	*	*	*	*	*
Kiowa	9	*	*	*	*	*	*	*
Labette	166	0.0	4.8	21.1	65.7	7.8	0.6	0.0
Lane	13	*	*	*	*	*	*	*
Leavenworth	356	0.0	5.9	19.4	61.5	11.0	2.2	0.0
Lincoln	22	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Age Distribution by Clinic

Comparison of Age Distribution ⁽²⁾								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Linn	54	*	*	*	*	*	*	*
Logan	13	*	*	*	*	*	*	*
Lyon	343	0.3	9.0	13.7	59.8	13.7	3.5	0.0
Marion	46	*	*	*	*	*	*	*
Marshall	63	*	*	*	*	*	*	*
Mc Pherson	145	0.7	3.4	15.9	62.1	17.2	0.7	0.0
Meade	49	*	*	*	*	*	*	*
Miami	136	0.0	7.4	11.0	67.6	14.0	0.0	0.0
Mitchell	43	*	*	*	*	*	*	*
Montgome-C	152	0.0	6.6	19.7	60.5	11.8	0.7	0.7
Montgome-I	178	0.0	7.9	16.9	65.2	9.6	0.6	0.0
Morris	32	*	*	*	*	*	*	*
Nemaha	24	*	*	*	*	*	*	*
Neosho-Ch	143	0.7	4.2	21.0	61.5	12.6	0.0	0.0
Neosho-Er	10	*	*	*	*	*	*	*
Ness	8	*	*	*	*	*	*	*
Norton	26	*	*	*	*	*	*	*
Osage	89	*	*	*	*	*	*	*
Osborne	19	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Age Distribution by Clinic

Comparison of Age Distribution ⁽²⁾								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Ottawa	22	*	*	*	*	*	*	*
Pawnee	43	*	*	*	*	*	*	*
Phillips	33	*	*	*	*	*	*	*
Pottawat-S	12	*	*	*	*	*	*	*
Pottawat-W	28	*	*	*	*	*	*	*
Pratt	68	*	*	*	*	*	*	*
Rawlins	10	*	*	*	*	*	*	*
Reno	494	0.2	5.9	15.2	66.2	11.5	1.0	0.0
Republic	33	*	*	*	*	*	*	*
Rice	70	*	*	*	*	*	*	*
Riley-Ft	428	0.0	0.5	12.9	75.7	10.5	0.5	0.0
Riley-Manh	398	0.0	2.5	13.1	71.9	11.8	0.8	0.0
Rooks	36	*	*	*	*	*	*	*
Rush	12	*	*	*	*	*	*	*
Russell	40	*	*	*	*	*	*	*
Saline	493	0.0	7.5	16.6	61.5	13.6	0.8	0.0
Scott	36	*	*	*	*	*	*	*
Sedgwick-C	1,078	0.2	6.5	12.8	62.2	17.3	1.0	0.1
Sedgwick-M	1,940	0.4	6.9	14.8	60.5	15.9	1.5	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Age Distribution by Clinic

Comparison of Age Distribution ⁽²⁾								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Sedgwick-S	1,131	0.2	6.7	14.9	62.2	14.4	1.7	0.0
Seneca	13	*	*	*	*	*	*	*
Seward	496	0.4	6.7	15.1	57.3	19.2	1.4	0.0
ShawneeM&I	168	0.6	9.5	11.9	56.5	19.0	2.4	0.0
ShawneeMa	1,183	0.2	6.3	14.0	62.5	16.1	1.0	0.0
Sheridan	7	*	*	*	*	*	*	*
Sherman	70	*	*	*	*	*	*	*
Smith	17	*	*	*	*	*	*	*
Stafford	40	*	*	*	*	*	*	*
Stanton	23	*	*	*	*	*	*	*
Stevens	66	*	*	*	*	*	*	*
Sumner	158	0.0	5.7	17.1	63.3	13.9	0.0	0.0
Thomas	50	*	*	*	*	*	*	*
Trego	14	*	*	*	*	*	*	*
Wabaunsee	18	*	*	*	*	*	*	*
Wallace	1	*	*	*	*	*	*	*
Washington	25	*	*	*	*	*	*	*
Wichita	11	*	*	*	*	*	*	*
Wilson	81	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Age Distribution by Clinic

Comparison of Age Distribution ⁽²⁾								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Woodson	2	*	*	*	*	*	*	*
Wyandotte	2,112	0.7	8.6	13.4	58.0	18.1	1.2	0.0
Kansas	20,703	0.3	6.5	14.6	61.6	15.9	1.2	0.0
Nation (Prior Year)	1,252,728	0.4	6.5	12.7	59.9	19.0	1.4	0.1

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators ^(2, 3)														
Clinic	Prepregnancy BMI ⁽⁴⁾			Weight Gain ⁽⁵⁾			Anemia (Low Hb/Hct) ⁽⁶⁾				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Allen	112	9.8 (10)	49.1 (29)	108	21.3 (11)	49.1 (31)	20	* *	35	* *	97	* *	97	* *
Anderson	50	* *	* *	47	* *	* *	9	* *	8	* *	38	* *	38	* *
Atchison	116	13.8 (34)	44.0 (8)	106	27.4 (35)	43.4 (13)	20	* *	30	* *	90	* *	90	* *
Barber-Kio	2	* *	* *	1	* *	* *	0	* *	0	* *	1	* *	1	* *
Barber-Med	37	* *	* *	32	* *	* *	2	* *	12	* *	27	* *	27	* *
Barton	274	10.6 (15)	45.6 (16)	260	27.3 (34)	41.5 (7)	27	* *	58	* *	219	4.1 (24)	219	7.3(25)
Bourbon	135	9.6 (8)	50.4 (36)	122	26.2 (31)	39.3 (4)	8	* *	31	* *	108	4.6 (28)	108	8.3(29)
Brown	110	6.4 (2)	52.7 (39)	99	* *	* *	19	* *	22	* *	95	* *	95	* *
Butler	283	12.7 (30)	41.3 (3)	269	21.6 (13)	46.8 (26)	32	* *	89	* *	208	2.4 (9)	208	9.6(33)
Chase	14	* *	* *	11	* *	* *	3	* *	2	* *	12	* *	12	* *
Chautauqua	32	* *	* *	27	* *	* *	2	* *	5	* *	25	* *	25	* *
Cherokee	218	12.8 (31)	46.3 (19)	196	22.4 (18)	43.9 (15)	24	* *	39	* *	168	4.2 (26)	168	5.4(10)
Clark	3	* *	* *	3	* *	* *	0	* *	2	* *	4	* *	4	* *
Clay	67	* *	* *	61	* *	* *	13	* *	13	* *	52	* *	52	* *
Cloud	72	* *	* *	65	* *	* *	5	* *	16	* *	55	* *	56	* *
Coffey	24	* *	* *	24	* *	* *	2	* *	7	* *	16	* *	16	* *
Comanche	6	* *	* *	5	* *	* *	1	* *	3	* *	3	* *	3	* *

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
- (4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).
- (5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).
- (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators ^(2, 3)														
Clinic	Prepregnancy BMI ⁽⁴⁾			Weight Gain ⁽⁵⁾			Anemia (Low Hb/Hct) ⁽⁶⁾				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Cowley-Ark	201	8.5 (3)	50.2 (35)	185	22.7 (20)	42.2 (10)	26	* *	34	* *	158	3.8 (21)	158	8.9(31)
Cowley-Win	154	11.7 (27)	49.4 (30)	147	20.4 (6)	50.3 (34)	14	* *	24	* *	128	1.6 (3)	128	6.3(20)
Crawford-G	1	* *	* *	1	* *	* *	1	* *	0	* *	1	* *	1	* *
Crawford-P	377	9.8 (10)	49.9 (34)	336	23.5 (26)	46.7 (25)	66	* *	72	* *	300	4.7 (29)	299	5.7(13)
Decatur	9	* *	* *	9	* *	* *	0	* *	3	* *	7	* *	7	* *
DickinsonA	93	* *	* *	86	* *	* *	15	* *	20	* *	70	* *	70	* *
DickinsonH	11	* *	* *	10	* *	* *	2	* *	4	* *	10	* *	10	* *
Doniphan	43	* *	* *	35	* *	* *	8	* *	6	* *	28	* *	28	* *
Douglas	459	13.1 (32)	41.8 (4)	409	20.3 (4)	53.1 (36)	67	* *	117	27.4 (7)	347	3.2 (17)	346	4.0(5)
Edwards	27	* *	* *	21	* *	* *	4	* *	7	* *	15	* *	15	* *
Elk	13	* *	* *	12	* *	* *	0	* *	2	* *	9	* *	9	* *
Ellis	142	10.6 (15)	46.5 (20)	127	22.8 (22)	42.5 (11)	33	* *	41	* *	101	3.0 (14)	101	6.9(24)
Ellsworth	28	* *	* *	28	* *	* *	3	* *	14	* *	20	* *	20	* *
Finney	682	14.8 (38)	40.6 (1)	640	28.3 (37)	40.0 (5)	99	* *	98	* *	570	6.7 (36)	570	3.5(4)
Ford	637	5.7 (1)	49.5 (33)	580	23.4 (25)	45.2 (21)	90	* *	78	* *	532	2.6 (11)	532	4.1(6)
Franklin	164	11.0 (19)	45.1 (13)	145	16.6 (1)	42.8 (12)	13	* *	32	* *	119	2.5 (10)	119	8.4(30)
Geary	336	11.0 (19)	44.3 (10)	283	21.2 (10)	48.1 (29)	55	* *	50	* *	256	5.1 (31)	256	9.8(34)

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
- (4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).
- (5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).
- (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators ^(2, 3)														
Clinic	Prepregnancy BMI ⁽⁴⁾			Weight Gain ⁽⁵⁾			Anemia (Low Hb/Hct) ⁽⁶⁾				Gestational Diabetes During Pregnancy	Hypertension During Pregnancy		
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)			Number	% (Rank)
Gove	7	* *	* *	7	* *	* *	2	* *	0	* *	7	* *	7	* *
Graham	14	* *	* *	11	* *	* *	4	* *	1	* *	12	* *	12	* *
Grant	65	* *	* *	61	* *	* *	10	* *	20	* *	53	* *	53	* *
Gray	57	* *	* *	48	* *	* *	7	* *	19	* *	41	* *	41	* *
Greeley	10	* *	* *	10	* *	* *	2	* *	2	* *	10	* *	10	* *
Greenwood	44	* *	* *	44	* *	* *	3	* *	7	* *	40	* *	40	* *
Hamilton	18	* *	* *	14	* *	* *	2	* *	5	* *	14	* *	14	* *
Harper	51	* *	* *	44	* *	* *	5	* *	8	* *	38	* *	37	* *
Harvey	250	10.8 (18)	44.4 (11)	228	27.2 (33)	41.7 (9)	30	* *	46	* *	192	3.1 (16)	190	6.8(22)
Haskell	24	* *	* *	22	* *	* *	5	* *	8	* *	19	* *	19	* *
Hodgeman	12	* *	* *	12	* *	* *	2	* *	2	* *	9	* *	9	* *
Jackson	70	* *	* *	67	* *	* *	8	* *	12	* *	61	* *	61	* *
Jefferson	65	* *	* *	60	* *	* *	10	* *	18	* *	53	* *	53	* *
Jewell	14	* *	* *	14	* *	* *	1	* *	2	* *	13	* *	13	* *
Johnson-M	851	10.2 (12)	45.1 (13)	789	20.5 (7)	44.7 (20)	138	22.5 (1)	195	18.5 (2)	670	4.0 (22)	671	3.4(1)
Johnson-O	968	11.4 (23)	44.2 (9)	895	22.5 (19)	44.0 (16)	176	28.4 (3)	205	14.6 (1)	758	4.0 (22)	757	5.2(8)
Kearny	21	* *	* *	19	* *	* *	4	* *	7	* *	19	* *	19	* *

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
- (4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).
- (5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).
- (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators ^(2, 3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Kingman	57	* *	* *	51	* *	* *	7	* *	12	* *	41	* *	41	* *
Kiowa	9	* *	* *	7	* *	* *	2	* *	1	* *	8	* *	8	* *
Labette	165	9.1 (5)	50.9 (37)	150	22.7 (20)	53.3 (37)	26	* *	36	* *	132	3.0 (14)	132	11.4(36)
Lane	12	* *	* *	10	* *	* *	4	* *	1	* *	8	* *	8	* *
Leavenwort	352	10.5 (13)	48.3 (26)	323	21.7 (15)	49.8 (33)	55	* *	99	* *	253	1.6 (3)	254	5.1(7)
Lincoln	22	* *	* *	22	* *	* *	1	* *	5	* *	16	* *	16	* *
Linn	52	* *	* *	49	* *	* *	3	* *	10	* *	45	* *	45	* *
Logan	13	* *	* *	10	* *	* *	2	* *	3	* *	9	* *	9	* *
Lyon	341	11.4 (23)	52.5 (38)	312	22.8 (22)	52.2 (35)	48	* *	64	* *	266	2.3 (8)	266	9.8(34)
Marion	46	* *	* *	40	* *	* *	6	* *	6	* *	31	* *	31	* *
Marshall	62	* *	* *	55	* *	* *	11	* *	13	* *	46	* *	46	* *
Mc Pherson	145	14.5 (37)	48.3 (26)	138	23.9 (29)	37.7 (2)	30	* *	30	* *	120	0.8 (1)	120	5.8(16)
Meade	49	* *	* *	44	* *	* *	7	* *	12	* *	39	* *	39	* *
Miami	135	14.1 (35)	48.1 (24)	130	19.2 (2)	53.8 (38)	32	* *	42	* *	104	4.8 (30)	104	5.8(16)
Mitchell	43	* *	* *	43	* *	* *	8	* *	13	* *	38	* *	38	* *
Montgome-C	152	13.2 (33)	47.4 (22)	138	32.6 (38)	34.8 (1)	24	* *	45	* *	117	1.7 (5)	117	3.4(1)
Montgome-I	176	19.9 (39)	43.8 (6)	160	28.1 (36)	43.8 (14)	28	* *	49	* *	141	2.1 (7)	141	7.8(28)

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
- (4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).
- (5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).
- (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators ^(2, 3)														
Clinic	Prepregnancy BMI ⁽⁴⁾			Weight Gain ⁽⁵⁾			Anemia (Low Hb/Hct) ⁽⁶⁾				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Morris	32	* *	* *	28	* *	* *	5	* *	6	* *	25	* *	25	* *
Nemaha	24	* *	* *	23	* *	* *	7	* *	10	* *	19	* *	19	* *
Neosho-Ch	141	14.2 (36)	48.2 (25)	132	19.7 (3)	40.9 (6)	20	* *	23	* *	108	0.9 (2)	108	5.6(12)
Neosho-Er	10	* *	* *	10	* *	* *	3	* *	2	* *	9	* *	9	* *
Ness	8	* *	* *	7	* *	* *	0	* *	5	* *	4	* *	4	* *
Norton	26	* *	* *	22	* *	* *	2	* *	7	* *	17	* *	17	* *
Osage	88	* *	* *	75	* *	* *	13	* *	22	* *	68	* *	68	* *
Osborne	19	* *	* *	18	* *	* *	1	* *	3	* *	17	* *	17	* *
Ottawa	22	* *	* *	20	* *	* *	3	* *	7	* *	16	* *	16	* *
Pawnee	43	* *	* *	42	* *	* *	4	* *	11	* *	36	* *	36	* *
Phillips	33	* *	* *	31	* *	* *	1	* *	5	* *	24	* *	24	* *
Pottawat-S	12	* *	* *	11	* *	* *	3	* *	6	* *	10	* *	10	* *
Pottawat-W	28	* *	* *	24	* *	* *	4	* *	11	* *	21	* *	21	* *
Pratt	68	* *	* *	59	* *	* *	8	* *	21	* *	46	* *	46	* *
Rawlins	10	* *	* *	9	* *	* *	1	* *	4	* *	8	* *	8	* *
Reno	493	12.0 (28)	45.2 (15)	462	26.0 (30)	41.6 (8)	66	* *	102	18.6 (3)	376	3.5 (19)	377	7.7(26)
Republic	33	* *	* *	33	* *	* *	5	* *	5	* *	30	* *	30	* *

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
- (4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).
- (5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).
- (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators ^(2, 3)														
Clinic	Prepregnancy BMI ⁽⁴⁾			Weight Gain ⁽⁵⁾			Anemia (Low Hb/Hct) ⁽⁶⁾				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Rice	70	* *	* *	60	* *	* *	9	* *	15	* *	53	* *	53	* *
Riley-Ft	426	9.2 (6)	41.1 (2)	352	21.6 (13)	49.7 (32)	56	* *	78	* *	273	1.8 (6)	273	6.2(19)
Riley-Manh	394	9.6 (8)	43.9 (7)	349	20.9 (8)	47.9 (28)	50	* *	97	* *	293	5.5 (34)	293	5.5(11)
Rooks	36	* *	* *	35	* *	* *	8	* *	6	* *	30	* *	30	* *
Rush	12	* *	* *	11	* *	* *	2	* *	2	* *	10	* *	10	* *
Russell	40	* *	* *	37	* *	* *	5	* *	5	* *	31	* *	31	* *
Saline	490	11.4 (23)	47.6 (23)	458	21.0 (9)	44.3 (18)	109	25.7 (2)	126	25.4 (5)	413	4.1 (24)	413	6.8(22)
Scott	36	* *	* *	34	* *	* *	5	* *	8	* *	30	* *	30	* *
Sedgwick-C	1,069	11.0 (19)	47.1 (21)	1,021	20.3 (4)	44.1 (17)	196	38.8 (8)	283	26.9 (6)	899	4.4 (27)	898	5.7(13)
Sedgwick-M	1,929	10.5 (13)	45.9 (18)	1,823	23.6 (27)	45.9 (22)	389	31.4 (5)	537	22.7 (4)	1,534	3.4 (18)	1,533	5.8(16)
Sedgwick-S	1,123	12.6 (29)	44.6 (12)	1,063	23.3 (24)	46.4 (24)	237	33.8 (6)	336	27.7 (8)	914	3.6 (20)	915	7.7(26)
Seneca	12	* *	* *	13	* *	* *	2	* *	6	* *	12	* *	12	* *
Seward	493	8.7 (4)	45.6 (16)	450	22.0 (17)	44.4 (19)	82	* *	81	* *	393	5.3 (33)	391	6.6(21)
ShawneeM&I	168	10.7 (17)	49.4 (30)	136	26.5 (32)	38.2 (3)	7	* *	8	* *	119	5.9 (35)	119	3.4(1)
ShawneeMa	1,169	11.0 (19)	49.4 (30)	1,081	21.3 (11)	48.8 (30)	132	35.6 (7)	225	28.4 (9)	944	5.1 (31)	947	5.7(13)
Sheridan	7	* *	* *	5	* *	* *	3	* *	1	* *	5	* *	5	* *
Sherman	70	* *	* *	63	* *	* *	6	* *	17	* *	53	* *	53	* *

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
- (4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).
- (5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).
- (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators ^(2, 3)															
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy		
	Under- weight	Over- weight	% (Rank)	% (Rank)	< Ideal	> Ideal	3rd Trimester		Postpartum						
							Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	
Smith	17	* *	* *	* *	14	* *	* *	3	* *	8	* *	15	* *	15	* *
Stafford	39	* *	* *	* *	33	* *	* *	3	* *	6	* *	27	* *	27	* *
Stanton	23	* *	* *	* *	21	* *	* *	3	* *	6	* *	21	* *	21	* *
Stevens	66	* *	* *	* *	55	* *	* *	8	* *	17	* *	44	* *	44	* *
Sumner	158	11.4 (23)	43.7 (5)		148	23.6 (27)	45.9 (22)	25	* *	43	* *	117	2.6 (11)	117	9.4(32)
Thomas	48	* *	* *	* *	41	* *	* *	5	* *	10	* *	37	* *	37	* *
Trego	14	* *	* *	* *	13	* *	* *	0	* *	2	* *	12	* *	12	* *
Wabaunsee	18	* *	* *	* *	18	* *	* *	0	* *	6	* *	15	* *	15	* *
Wallace	1	* *	* *	* *	1	* *	* *	0	* *	1	* *	0	* *	0	* *
Washington	25	* *	* *	* *	24	* *	* *	2	* *	5	* *	20	* *	19	* *
Wichita	11	* *	* *	* *	11	* *	* *	1	* *	4	* *	10	* *	10	* *
Wilson	79	* *	* *	* *	77	* *	* *	13	* *	14	* *	66	* *	66	* *
Woodson	2	* *	* *	* *	2	* *	* *	0	* *	0	* *	2	* *	2	* *
Wyandotte	2,077	9.5 (7)	48.9 (28)		1,924	21.9 (16)	46.9 (27)	320	29.7 (4)	358	28.5(10)	1,688	2.8 (13)	1,689	5.2(8)
Kansas	20,516	10.9	46.6		18,896	22.9	45.6	3,213	29.3	4,533	24.0	16,226	3.7	16,224	6.2
Nation (Prior Year)	1,175,574	10.9	44.5		922,658	25.0	42.8	180,176	33.5	414,735	29.8	230,547	3.9	280,574	4.3

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
- (4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).
- (5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).
- (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6A
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators ^(2, 3)											
Clinic	Medical Care				WIC Enrollment			Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester			Prepregnancy ⁽⁴⁾		During Pregnancy ⁽⁵⁾	
	Number	% (Rank)	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)
Allen	102	95.1 (1)	0.0 (1)		96	* *		69	* *	64	* *
Anderson	48	* *	* *		40	* *		38	* *	29	* *
Atchison	104	87.5 (7)	2.9 (17)		95	* *		81	* *	71	* *
Barber-Kio	1	* *	* *		2	* *		2	* *	2	* *
Barber-Med	32	* *	* *		30	* *		22	* *	19	* *
Barton	254	81.1 (15)	1.2 (9)		217	56.7 (3)		157	19.7 (2)	132	63.6 (4)
Bourbon	115	91.3 (3)	0.9 (7)		113	48.7 (6)		77	* *	78	* *
Brown	96	* *	* *		95	* *		64	* *	83	* *
Butler	254	70.5 (29)	13.4 (35)		193	48.2 (7)		154	12.3 (10)	117	37.6 (24)
Chase	11	* *	* *		12	* *		9	* *	8	* *
Chautauqua	27	* *	* *		28	* *		16	* *	17	* *
Cherokee	177	76.3 (20)	9.6 (32)		167	50.3 (4)		115	9.6 (18)	108	48.1 (17)
Clark	3	* *	* *		2	* *		0	* *	1	* *
Clay	59	* *	* *		55	* *		35	* *	32	* *
Cloud	64	* *	* *		58	* *		44	* *	36	* *
Coffey	24	* *	* *		14	* *		14	* *	8	* *
Comanche	5	* *	* *		3	* *		3	* *	2	* *
Cowley-Ark	168	60.7 (37)	8.9 (30)		151	42.4 (16)		116	6.9 (20)	103	51.5 (14)
Cowley-Win	146	78.8 (18)	3.4 (18)		118	61.0 (1)		78	* *	69	* *
Crawford-G	1	* *	* *		1	* *		0	* *	0	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6A
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators ^(2, 3)										
Clinic	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Prepregnancy ⁽⁴⁾		During Pregnancy ⁽⁵⁾	
	Number	% (Rank)	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Crawford-P	329	87.5 (7)	1.2 (9)		311	44.7 (11)	235	8.1 (19)	212	61.8 (6)
Decatur	7	*	*	*	6	*	5	*	4	*
DickinsonA	84	*	*	*	74	*	62	*	49	*
DickinsonH	10	*	*	*	10	*	6	*	6	*
Doniphan	36	*	*	*	33	*	27	*	26	*
Douglas	403	70.0 (30)	6.9 (28)		362	42.3 (17)	258	12.4 (8)	223	48.0 (18)
Edwards	18	*	*	*	20	*	19	*	13	*
Elk	11	*	*	*	9	*	5	*	4	*
Ellis	128	85.9 (9)	1.6 (13)		112	33.9 (30)	95	*	84	*
Ellsworth	28	*	*	*	16	*	16	*	10	*
Finney	578	66.1 (32)	19.0 (37)		551	37.4 (25)	413	5.8 (23)	357	29.7 (25)
Ford	549	72.7 (27)	14.2 (36)		538	49.1 (5)	347	5.5 (25)	336	49.4 (15)
Franklin	95	*	*	*	129	41.1 (19)	91	*	91	*
Geary	274	73.0 (26)	0.7 (5)		284	45.8 (10)	195	15.9 (4)	183	47.0 (19)
Gove	7	*	*	*	5	*	3	*	3	*
Graham	12	*	*	*	12	*	4	*	5	*
Grant	54	*	*	*	51	*	36	*	27	*
Gray	49	*	*	*	43	*	32	*	24	*
Greeley	10	*	*	*	7	*	7	*	7	*
Greenwood	40	*	*	*	36	*	20	*	21	*

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6A
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators ^(2, 3)										
Clinic	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	% (Rank)		Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Hamilton	15	* *	* *		14	* *	10	* *	11	* *
Harper	41	* *	* *		43	* *	28	* *	29	* *
Harvey	212	73.6 (25)	12.7 (34)		202	48.0 (8)	138	11.6 (13)	128	61.7 (7)
Haskell	21	* *	* *		21	* *	16	* *	15	* *
Hodgeman	12	* *	* *		8	* *	7	* *	4	* *
Jackson	42	* *	* *		63	* *	39	* *	42	* *
Jefferson	52	* *	* *		48	* *	36	* *	33	* *
Jewell	14	* *	* *		12	* *	8	* *	10	* *
Johnson-M	729	64.2 (34)	1.2 (9)		640	35.0 (29)	454	11.7 (12)	402	64.2 (2)
Johnson-O	796	61.4 (36)	5.3 (27)		766	33.4 (31)	568	11.4 (15)	489	57.7 (11)
Kearny	19	* *	* *		17	* *	10	* *	9	* *
Kingman	49	* *	* *		47	* *	39	* *	36	* *
Kiowa	8	* *	* *		9	* *	6	* *	8	* *
Labette	128	85.9 (9)	2.3 (15)		132	40.2 (23)	92	* *	89	* *
Lane	9	* *	* *		10	* *	7	* *	9	* *
Leavenworth	320	84.4 (11)	0.9 (7)		244	36.1 (28)	202	12.4 (8)	151	61.6 (8)
Lincoln	22	* *	* *		15	* *	14	* *	9	* *
Linn	48	* *	* *		43	* *	30	* *	31	* *
Logan	6	* *	* *		10	* *	8	* *	7	* *
Lyon	312	81.4 (14)	0.6 (4)		265	44.2 (14)	193	14.0 (6)	169	58.0 (9)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6A
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators ^(2, 3)										
Clinic	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Prepregnancy ⁽⁴⁾		During Pregnancy ⁽⁵⁾	
	Number	% (Rank)	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Marion	40	* *	* *	* *	34	* *	27	* *	21	* *
Marshall	56	* *	* *	* *	53	* *	33	* *	25	* *
Mc Pherson	138	88.4 (6)	0.7 (5)		114	37.7 (24)	74	* *	67	* *
Meade	44	* *	* *	* *	35	* *	26	* *	25	* *
Miami	129	79.1 (17)	0.0 (1)		94	* *	74	* *	62	* *
Mitchell	43	* *	* *	* *	34	* *	24	* *	22	* *
Montgome-C	118	79.7 (16)	9.3 (31)		118	43.2 (15)	84	* *	72	* *
Montgome-I	136	91.9 (2)	3.7 (22)		139	44.6 (12)	102	13.7 (7)	88	* *
Morris	26	* *	* *	* *	26	* *	14	* *	13	* *
Nemaha	23	* *	* *	* *	19	* *	13	* *	12	* *
Neosho-Ch	119	89.9 (4)	3.4 (18)		113	40.7 (20)	87	* *	73	* *
Neosho-Er	10	* *	* *	* *	9	* *	5	* *	4	* *
Ness	8	* *	* *	* *	3	* *	5	* *	1	* *
Norton	23	* *	* *	* *	15	* *	14	* *	8	* *
Osage	76	* *	* *	* *	65	* *	51	* *	37	* *
Osborne	18	* *	* *	* *	14	* *	6	* *	10	* *
Ottawa	19	* *	* *	* *	16	* *	11	* *	10	* *
Pawnee	41	* *	* *	* *	36	* *	28	* *	25	* *
Phillips	31	* *	* *	* *	25	* *	20	* *	21	* *
Pottawat-S	10	* *	* *	* *	10	* *	4	* *	5	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6A
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators ^(2, 3)										
Clinic	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Prepregnancy ⁽⁴⁾		During Pregnancy ⁽⁵⁾	
	Number	% (Rank)	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Pottawat-W	18	* *	* *	* *	23	* *	15	* *	16	* *
Pratt	60	* *	* *	* *	46	* *	41	* *	29	* *
Rawlins	5	* *	* *	* *	9	* *	6	* *	7	* *
Reno	450	83.6 (12)	2.0 (14)		374	40.4 (22)	291	23.7 (1)	247	64.0 (3)
Republic	33	* *	* *	* *	26	* *	13	* *	14	* *
Rice	59	* *	* *	* *	59	* *	42	* *	41	* *
Riley-Ft	308	89.6 (5)	0.0 (1)		301	36.5 (27)	253	11.5 (14)	193	40.9 (22)
Riley-Manh	332	75.9 (21)	1.2 (9)		298	41.6 (18)	219	18.3 (3)	175	55.4 (13)
Rooks	35	* *	* *	* *	30	* *	17	* *	18	* *
Rush	11	* *	* *	* *	11	* *	8	* *	7	* *
Russell	37	* *	* *	* *	30	* *	23	* *	21	* *
Saline	413	74.3 (24)	3.4 (18)		378	30.4 (32)	271	5.9 (22)	242	48.8 (16)
Scott	33	* *	* *	* *	28	* *	18	* *	18	* *
Sedgwick-C	998	75.5 (22)	4.2 (25)		855	30.1 (33)	615	15.6 (5)	572	46.3 (20)
Sedgwick-M	1,820	70.9 (28)	3.7 (22)		1,457	27.5 (35)	1,082	10.9 (16)	863	40.3 (23)
Sedgwick-S	1,055	77.8 (19)	3.6 (21)		899	28.7 (34)	609	10.0 (17)	543	42.5 (21)
Seneca	13	* *	* *	* *	10	* *	3	* *	3	* *
Seward	427	66.0 (33)	11.5 (33)		398	37.4 (25)	267	6.0 (21)	238	64.3 (1)
ShawneeM&I	132	63.6 (35)	2.3 (15)		148	60.1 (2)	102	0.0 (26)	113	61.9 (5)
ShawneeMa	1,030	75.5 (22)	5.0 (26)		921	47.9 (9)	683	5.7 (24)	626	57.0 (12)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6A
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators ^(2, 3)										
Clinic	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Sheridan	5	* *	* *	* *	4	* *	2	* *	1	* *
Sherman	47	* *	* *	* *	58	* *	39	* *	38	* *
Smith	12	* *	* *	* *	13	* *	9	* *	10	* *
Stafford	33	* *	* *	* *	33	* *	25	* *	23	* *
Stanton	21	* *	* *	* *	21	* *	11	* *	12	* *
Stevens	55	* *	* *	* *	49	* *	38	* *	29	* *
Sumner	145	83.4 (13)	4.1 (24)		119	44.5 (13)	84	* *	68	* *
Thomas	36	* *	* *	* *	36	* *	23	* *	26	* *
Trego	13	* *	* *	* *	9	* *	6	* *	6	* *
Wabaunsee	16	* *	* *	* *	10	* *	10	* *	7	* *
Wallace	1	* *	* *	* *	0	* *	1	* *	0	* *
Washington	24	* *	* *	* *	20	* *	15	* *	14	* *
Wichita	11	* *	* *	* *	9	* *	6	* *	5	* *
Wilson	76	* *	* *	* *	71	* *	50	* *	47	* *
Woodson	2	* *	* *	* *	1	* *	0	* *	0	* *
Wyandotte	1,922	66.7 (31)	7.0 (29)		1,657	40.5 (21)	1,171	12.3 (10)	1,047	57.8 (10)
Kansas	18,064	74.9	5.2		16,091	39.8	11,645	12.0	10,265	53.0
Nation (Prior Year)	612,462	79.4	4.7		1,034,046	31.3	98,170	15.7	238,509	75.9

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2008 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Allen	111	46.8 (33)	100	32.0 (34)	38	* *	96	* *	98	* *
Anderson	48	* *	43	* *	17	* *	39	* *	43	* *
Atchison	74	* *	38	* *	20	* *	67	* *	38	* *
Barber-Kio	2	* *	1	* *	0	* *	2	* *	1	* *
Barber-Med	37	* *	32	* *	14	* *	31	* *	32	* *
Barton	272	39.7 (21)	249	25.3 (24)	80	* *	218	10.1 (11)	252	6.7 (8)
Bourbon	128	43.8 (27)	102	25.5 (25)	34	* *	111	31.5 (30)	103	33.0 (31)
Brown	105	44.8 (31)	85	* *	26	* *	91	* *	85	* *
Butler	262	52.7 (36)	198	34.3 (35)	78	* *	191	14.7 (16)	196	7.7 (11)
Chase	14	* *	12	* *	7	* *	12	* *	12	* *
Chautauqua	30	* *	22	* *	6	* *	25	* *	21	* *
Cherokee	201	39.3 (19)	163	28.8 (30)	50	* *	160	30.0 (28)	169	29.6 (29)
Clark	1	* *	0	* *	0	* *	1	* *	0	* *
Clay	65	* *	57	* *	25	* *	53	* *	59	* *
Cloud	70	* *	61	* *	11	* *	57	* *	61	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2008 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Coffey	23	* *	23	* *	8	* *	14	* *	23	* *
Comanche	5	* *	4	* *	0	* *	3	* *	4	* *
Cowley-Ark	186	39.2 (18)	151	25.8 (26)	43	* *	150	21.3 (24)	153	20.9 (23)
Cowley-Win	153	51.0 (35)	141	31.9 (33)	57	* *	120	20.8 (23)	142	21.1 (24)
Crawford-G	1	* *	1	* *	0	* *	1	* *	1	* *
Crawford-P	360	41.7 (25)	284	23.2 (19)	97	* *	309	40.1 (33)	285	38.9 (33)
Decatur	9	* *	7	* *	1	* *	6	* *	7	* *
DickinsonA	89	* *	77	* *	27	* *	74	* *	79	* *
DickinsonH	11	* *	10	* *	5	* *	10	* *	10	* *
Doniphan	31	* *	22	* *	9	* *	23	* *	24	* *
Douglas	447	44.1 (29)	381	20.7 (15)	118	44.9 (1)	358	10.1 (11)	382	7.3 (9)
Edwards	27	* *	20	* *	5	* *	20	* *	21	* *
Elk	13	* *	11	* *	2	* *	9	* *	11	* *
Ellis	140	29.3 (10)	121	9.1 (4)	34	* *	111	6.3 (3)	121	8.3 (13)
Ellsworth	27	* *	27	* *	10	* *	16	* *	28	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2008 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Finney	674	13.4 (3)	627	7.0 (3)	68	* *	556	17.3 (18)	626	15.7 (19)
Ford	612	7.7 (1)	533	3.8 (2)	29	* *	522	7.5 (7)	527	6.6 (6)
Franklin	163	44.2 (30)	147	25.2 (23)	47	* *	126	11.1 (13)	147	9.5 (14)
Geary	333	41.4 (24)	270	21.1 (17)	98	* *	284	39.8 (32)	273	38.8 (32)
Gove	7	* *	7	* *	0	* *	5	* *	7	* *
Graham	14	* *	12	* *	7	* *	12	* *	12	* *
Grant	65	* *	62	* *	5	* *	50	* *	62	* *
Gray	55	* *	47	* *	3	* *	43	* *	47	* *
Greeley	10	* *	10	* *	3	* *	7	* *	10	* *
Greenwood	41	* *	33	* *	14	* *	36	* *	35	* *
Hamilton	18	* *	15	* *	2	* *	14	* *	15	* *
Harper	45	* *	25	* *	10	* *	41	* *	26	* *
Harvey	247	35.2 (15)	210	24.8 (22)	61	* *	203	5.9 (1)	211	6.2 (4)
Haskell	24	* *	23	* *	3	* *	21	* *	23	* *
Hodgeman	12	* *	9	* *	3	* *	8	* *	9	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2008 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Jackson	69	* *	64	* *	27	* *	62	* *	65	* *
Jefferson	16	* *	9	* *	1	* *	16	* *	11	* *
Jewell	14	* *	13	* *	3	* *	12	* *	13	* *
Johnson-M	839	25.5 (6)	742	11.7 (7)	130	37.7 (4)	634	20.2 (22)	748	20.6 (22)
Johnson-O	962	23.8 (5)	855	11.2 (6)	147	43.5 (2)	766	17.9 (19)	858	18.8 (21)
Kearny	21	* *	20	* *	2	* *	17	* *	20	* *
Kingman	50	* *	42	* *	9	* *	47	* *	48	* *
Kiowa	8	* *	4	* *	2	* *	8	* *	5	* *
Labette	161	33.5 (14)	139	20.9 (16)	33	* *	131	30.5 (29)	143	29.4 (28)
Lane	3	* *	0	* *	0	* *	4	* *	0	* *
Leavenwort	347	47.0 (34)	297	28.3 (29)	95	* *	243	46.5 (34)	301	40.5 (34)
Lincoln	20	* *	17	* *	6	* *	15	* *	17	* *
Linn	53	* *	47	* *	21	* *	43	* *	49	* *
Logan	13	* *	10	* *	4	* *	11	* *	10	* *
Lyon	338	30.2 (12)	304	16.1 (11)	63	* *	260	12.7 (14)	306	10.5 (16)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2008 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Marion	46	* *	40	* *	6	* *	34	* *	40	* *
Marshall	63	* *	52	* *	24	* *	53	* *	53	* *
Mc Pherson	143	40.6 (23)	132	30.3 (32)	47	* *	114	7.0 (4)	133	4.5 (2)
Meade	44	* *	37	* *	8	* *	35	* *	37	* *
Miami	114	54.4 (38)	98	* *	39	* *	85	* *	97	* *
Mitchell	42	* *	41	* *	14	* *	34	* *	42	* *
Montgome-C	147	39.5 (20)	125	27.2 (27)	38	* *	118	36.4 (31)	128	30.5 (30)
Montgome-I	171	39.8 (22)	148	29.1 (31)	46	* *	131	22.1 (26)	149	22.8 (27)
Morris	31	* *	27	* *	11	* *	26	* *	27	* *
Nemaha	24	* *	21	* *	9	* *	19	* *	21	* *
Neosho-Ch	134	37.3 (17)	101	17.8 (13)	28	* *	111	18.9 (21)	102	11.8 (17)
Neosho-Er	10	* *	8	* *	1	* *	9	* *	8	* *
Ness	7	* *	7	* *	2	* *	3	* *	7	* *
Norton	26	* *	23	* *	6	* *	15	* *	23	* *
Osage	85	* *	76	* *	24	* *	64	* *	76	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2008 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Osborne	18	* *	18	* *	7	* *	13	* *	18	* *
Ottawa	18	* *	15	* *	4	* *	15	* *	15	* *
Pawnee	42	* *	41	* *	15	* *	36	* *	41	* *
Phillips	17	* *	3	* *	3	* *	17	* *	3	* *
Pottawat-S	12	* *	11	* *	2	* *	10	* *	11	* *
Pottawat-W	28	* *	22	* *	7	* *	23	* *	23	* *
Pratt	66	* *	58	* *	17	* *	46	* *	60	* *
Rawlins	10	* *	8	* *	1	* *	9	* *	8	* *
Reno	478	45.6 (32)	417	24.7 (21)	151	31.1 (6)	372	18.3 (20)	419	17.4 (20)
Republic	32	* *	31	* *	13	* *	26	* *	32	* *
Rice	70	* *	58	* *	18	* *	59	* *	58	* *
Riley-Ft	413	28.6 (8)	296	16.6 (12)	50	* *	297	22.9 (27)	329	21.9 (26)
Riley-Manh	370	29.2 (9)	282	19.1 (14)	59	* *	295	7.1 (5)	319	6.6 (6)
Rooks	35	* *	26	* *	17	* *	30	* *	29	* *
Rush	12	* *	10	* *	4	* *	11	* *	10	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2008 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Russell	40	* *	36	* *	11	* *	30	* *	37	* *
Saline	433	43.9 (28)	357	27.5 (28)	130	24.6 (8)	347	8.6 (8)	360	6.4 (5)
Scott	34	* *	29	* *	7	* *	29	* *	29	* *
Sedgwick-C	1,054	29.8 (11)	986	14.7 (10)	216	35.6 (5)	853	7.3 (6)	978	7.8 (12)
Sedgwick-M	1,902	30.3 (13)	1,760	14.3 (9)	378	39.2 (3)	1,451	6.0 (2)	1,761	5.9 (3)
Sedgwick-S	1,113	42.0 (26)	1,036	21.8 (18)	339	31.0 (7)	893	8.8 (9)	1,036	7.6 (10)
Seneca	13	* *	13	* *	8	* *	10	* *	13	* *
Seward	482	8.9 (2)	430	3.3 (1)	32	* *	400	15.3 (17)	439	14.1 (18)
ShawneeM&I	156	26.3 (7)	117	11.1 (5)	22	* *	146	8.9 (10)	117	1.7 (1)
ShawneeMa	1,075	37.1 (16)	872	23.4 (20)	237	22.8 (10)	881	14.1 (15)	872	9.5 (14)
Sheridan	7	* *	5	* *	1	* *	4	* *	5	* *
Sherman	69	* *	58	* *	20	* *	57	* *	57	* *
Smith	14	* *	11	* *	3	* *	12	* *	12	* *
Stafford	39	* *	30	* *	9	* *	32	* *	31	* *
Stanton	22	* *	20	* *	2	* *	21	* *	20	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2008 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Stevens	66	* *	52	* *	10	* *	49	* *	52	* *
Sumner	150	54.0 (37)	135	34.8 (36)	57	* *	115	53.0 (35)	134	53.7 (35)
Thomas	47	* *	36	* *	10	* *	36	* *	36	* *
Trego	12	* *	10	* *	4	* *	11	* *	11	* *
Wabaunsee	18	* *	17	* *	5	* *	10	* *	17	* *
Wallace	0	* *	0	* *	0	* *	0	* *	0	* *
Washington	25	* *	24	* *	5	* *	20	* *	23	* *
Wichita	11	* *	9	* *	2	* *	9	* *	9	* *
Wilson	79	* *	73	* *	35	* *	70	* *	73	* *
Woodson	2	* *	2	* *	0	* *	1	* *	2	* *
Wyandotte	2,024	23.2 (4)	1,754	14.2 (8)	300	23.3 (9)	1,646	21.6 (25)	1,789	21.7 (25)
Kansas	19,771	32.9	17,140	18.5	4,232	30.8	15,818	17.6	17,316	16.1
Nation (Prior Year)	895,227	26.6	716,629	15.8	161,864	35.1	572,120	20.9	486,636	22.6

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators ^(2, 3)									
Clinic	Birthweight			Preterm ⁽⁶⁾		Full Term LBW ⁽⁷⁾		Ever Breastfed	
	Low ⁽⁴⁾	High ⁽⁵⁾		Number	%	Number	%	Number	%
	Number	% (Rank)	% (Rank)						
Allen	107	6.5 (22)	5.6 (11)	91	* *	85	* *	106	48.1 (37)
Anderson	46	* *	* *	37	* *	36	* *	43	* *
Atchison	101	5.9 (13)	5.9 (17)	84	* *	79	* *	105	53.3 (34)
Barber-Kio	1	* *	* *	1	* *	1	* *	1	* *
Barber-Med	30	* *	* *	25	* *	19	* *	31	* *
Barton	243	7.0 (24)	5.8 (15)	202	13.9 (31)	172	2.9 (18)	237	75.9 (9)
Bourbon	117	6.0 (15)	6.0 (20)	101	11.9 (23)	88	* *	116	64.7 (22)
Brown	86	* *	* *	88	* *	74	* *	90	* *
Butler	264	9.8 (36)	5.7 (13)	180	14.4 (32)	153	3.3 (23)	240	72.9 (15)
Chase	12	* *	* *	10	* *	10	* *	11	* *
Chautauqua	26	* *	* *	23	* *	20	* *	23	* *
Cherokee	194	7.7 (26)	5.7 (13)	147	12.9 (30)	128	3.1 (22)	182	61.0 (29)
Clark	4	* *	* *	2	* *	2	* *	3	* *
Clay	60	* *	* *	49	* *	41	* *	59	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators ^(2, 3)										
Clinic	Birthweight				Preterm ⁽⁶⁾		Full Term LBW ⁽⁷⁾		Ever Breastfed	
	Low ⁽⁴⁾		High ⁽⁵⁾		Number	%	Number	%	Number	%
	Number	% (Rank)	% (Rank)	(Rank)						
Cloud	63	* * * *	* * * *	52	* *	42	* *	59	* *	
Coffey	23	* * * *	* * * *	14	* *	12	* *	24	* *	
Comanche	5	* * * *	* * * *	2	* *	2	* *	5	* *	
Cowley-Ark	180	6.1 (17)	7.2 (31)	135	11.1 (21)	119	4.2 (28)	172	48.3 (36)	
Cowley-Win	140	7.9 (27)	7.1 (28)	110	12.7 (29)	96	* *	139	64.0 (23)	
Crawford-G	1	* * * *	* * * *	1	* *	1	* *	1	* *	
Crawford-P	323	8.7 (32)	5.9 (17)	271	9.2 (9)	245	3.7 (24)	317	59.6 (30)	
Decatur	7	* * * *	* * * *	6	* *	6	* *	6	* *	
DickinsonA	78	* * * *	* * * *	66	* *	61	* *	80	* *	
DickinsonH	9	* * * *	* * * *	9	* *	9	* *	9	* *	
Doniphan	37	* * * *	* * * *	27	* *	23	* *	34	* *	
Douglas	407	5.4 (7)	7.1 (28)	313	12.1 (25)	273	2.6 (16)	389	75.6 (10)	
Edwards	20	* * * *	* * * *	14	* *	9	* *	19	* *	
Elk	12	* * * *	* * * *	8	* *	8	* *	11	* *	

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
- (4) <2500 g.
- (5) >4000 g.
- (6) <37 weeks gestation.
- (7) 37 or more weeks gestation and <2500 g.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators ^(2, 3)									
Clinic	Birthweight			Preterm ⁽⁶⁾		Full Term LBW ⁽⁷⁾		Ever Breastfed	
	Low ⁽⁴⁾	High ⁽⁵⁾		Number	%	Number	%	Number	%
	Number	% (Rank)	% (Rank)						
Ellis	118	8.5 (31)	0.8 (1)	96	* *	80	* *	115	81.7 (2)
Ellsworth	28	* *	* *	16	* *	14	* *	27	* *
Finney	618	5.7 (11)	6.5 (23)	514	10.1 (14)	457	2.0 (7)	341	78.6 (4)
Ford	562	4.8 (3)	5.9 (17)	483	9.7 (12)	432	1.4 (4)	440	74.3 (11)
Franklin	145	6.2 (20)	5.5 (10)	110	10.9 (20)	98	* *	139	54.0 (33)
Geary	287	5.2 (4)	5.2 (7)	234	7.7 (3)	215	2.3 (11)	261	69.0 (18)
Gove	7	* *	* *	5	* *	4	* *	7	* *
Graham	11	* *	* *	10	* *	8	* *	12	* *
Grant	62	* *	* *	47	* *	41	* *	35	* *
Gray	37	* *	* *	34	* *	26	* *	44	* *
Greeley	10	* *	* *	7	* *	7	* *	10	* *
Greenwood	40	* *	* *	35	* *	29	* *	38	* *
Hamilton	13	* *	* *	11	* *	10	* *	5	* *
Harper	37	* *	* *	37	* *	28	* *	42	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators ^(2, 3)										
Clinic	Birthweight				Preterm ⁽⁶⁾		Full Term LBW ⁽⁷⁾		Ever Breastfed	
	Low ⁽⁴⁾		High ⁽⁵⁾		Number	%	Number	%	Number	%
	Number	% (Rank)	Number	% (Rank)						
Harvey	214	9.3 (34)	5.6 (11)		180	17.2 (34)	149	2.7 (17)	219	78.5 (5)
Haskell	23	* *	* *		20	* *	19	* *	12	* *
Hodgeman	12	* *	* *		8	* *	7	* *	10	* *
Jackson	66	* *	* *		60	* *	51	* *	62	* *
Jefferson	59	* *	* *		44	* *	39	* *	57	* *
Jewell	13	* *	* *		12	* *	8	* *	12	* *
Johnson-M	716	8.4 (30)	7.1 (28)		579	9.3 (11)	486	2.5 (15)	730	77.9 (7)
Johnson-O	852	5.3 (6)	6.9 (26)		691	8.4 (5)	614	2.4 (14)	801	79.7 (3)
Kearny	19	* *	* *		16	* *	13	* *	9	* *
Kingman	43	* *	* *		41	* *	33	* *	48	* *
Kiowa	8	* *	* *		8	* *	7	* *	8	* *
Labette	149	4.7 (2)	10.1 (38)		117	12.0 (24)	103	1.0 (2)	133	55.6 (32)
Lane	10	* *	* *		7	* *	7	* *	10	* *
Leavenworth	299	9.4 (35)	4.0 (3)		216	8.8 (6)	195	4.1 (26)	304	62.2 (27)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators ^(2, 3)										
Clinic	Birthweight				Preterm ⁽⁶⁾		Full Term LBW ⁽⁷⁾		Ever Breastfed	
	Low ⁽⁴⁾		High ⁽⁵⁾		Number	%	Number	%	Number	%
	Number	% (Rank)	% (Rank)	(Rank)						
Lincoln	22	* * * *	* * * *	15	* *	14	* *	20	* *	
Linn	46	* * * *	* * * *	39	* *	30	* *	45	* *	
Logan	10	* * * *	* * * *	8	* *	7	* *	9	* *	
Lyon	301	6.0 (15)	6.0 (20)	236	8.9 (7)	214	0.9 (1)	289	73.0 (14)	
Marion	39	* * * *	* * * *	28	* *	24	* *	39	* *	
Marshall	54	* * * *	* * * *	46	* *	42	* *	52	* *	
Mc Pherson	133	3.8 (1)	5.3 (9)	106	4.7 (1)	100	2.0 (7)	132	74.2 (12)	
Meade	38	* * * *	* * * *	30	* *	25	* *	36	* *	
Miami	126	5.6 (9)	6.3 (22)	87	* *	76	* *	98	* *	
Mitchell	40	* * * *	* * * *	33	* *	30	* *	37	* *	
Montgome-C	134	8.2 (29)	5.2 (7)	105	16.2 (33)	86	* *	123	62.6 (25)	
Montgome-I	154	10.4 (38)	3.2 (2)	120	10.8 (19)	107	4.7 (29)	147	61.9 (28)	
Morris	29	* * * *	* * * *	23	* *	19	* *	28	* *	
Nemaha	21	* * * *	* * * *	18	* *	13	* *	23	* *	

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
- (4) <2500 g.
- (5) >4000 g.
- (6) <37 weeks gestation.
- (7) 37 or more weeks gestation and <2500 g.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators ^(2, 3)									
Clinic	Birthweight			Preterm ⁽⁶⁾		Full Term LBW ⁽⁷⁾		Ever Breastfed	
	Low ⁽⁴⁾	High ⁽⁵⁾		Number	%	Number	%	Number	%
	Number	% ^(Rank)	% ^(Rank)						
Neosho-Ch	109	5.5 (8)	7.3 (32)	103	12.6 (28)	78	* *	132	52.3 (35)
Neosho-Er	10	* *	* *	9	* *	9	* *	10	* *
Ness	8	* *	* *	3	* *	2	* *	8	* *
Norton	20	* *	* *	12	* *	10	* *	19	* *
Osage	83	* *	* *	64	* *	49	* *	74	* *
Osborne	17	* *	* *	13	* *	13	* *	16	* *
Ottawa	19	* *	* *	14	* *	13	* *	15	* *
Pawnee	39	* *	* *	35	* *	32	* *	38	* *
Phillips	29	* *	* *	23	* *	21	* *	28	* *
Pottawat-S	10	* *	* *	8	* *	7	* *	10	* *
Pottawat-W	24	* *	* *	19	* *	18	* *	23	* *
Pratt	57	* *	* *	39	* *	38	* *	52	* *
Rawlins	9	* *	* *	8	* *	8	* *	9	* *
Reno	438	5.9 (13)	7.5 (34)	338	10.4 (16)	299	2.3 (11)	418	58.9 (31)

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
- (4) <2500 g.
- (5) >4000 g.
- (6) <37 weeks gestation.
- (7) 37 or more weeks gestation and <2500 g.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators ^(2, 3)										
Clinic	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)		Number	%	Number	%	Number	%
	Number	% (Rank)	% (Rank)	(Rank)						
Republic	32	* * * *	* * * *	26	* *	17	* *	33	* *	
Rice	59	* * * *	* * * *	49	* *	41	* *	55	* *	
Riley-Ft	353	5.7 (11)	7.4 (33)	230	10.4 (16)	206	1.9 (5)	344	77.9 (7)	
Riley-Manh	346	6.1 (17)	5.8 (15)	254	9.1 (8)	230	2.2 (10)	335	78.5 (5)	
Rooks	33	* * * *	* * * *	29	* *	26	* *	32	* *	
Rush	11	* * * *	* * * *	10	* *	9	* *	11	* *	
Russell	36	* * * *	* * * *	27	* *	22	* *	33	* *	
Saline	447	8.7 (32)	4.9 (4)	353	11.6 (22)	309	2.9 (18)	444	62.4 (26)	
Scott	28	* * * *	* * * *	26	* *	25	* *	27	* *	
Sedgwick-C	1,002	7.0 (24)	7.9 (36)	803	9.8 (13)	717	2.9 (18)	942	74.0 (13)	
Sedgwick-M	1,778	9.8 (36)	6.8 (24)	1,350	10.1 (14)	1,206	4.1 (26)	1,476	67.1 (19)	
Sedgwick-S	1,053	6.1 (17)	7.0 (27)	840	8.2 (4)	768	2.0 (7)	965	62.7 (24)	
Seneca	12	* * * *	* * * *	10	* *	8	* *	13	* *	
Seward	424	5.2 (4)	5.0 (5)	356	10.4 (16)	315	1.3 (3)	432	71.1 (17)	

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators ^(2, 3)										
Clinic	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)		Number	%	Number	%	Number	%
	Number	% (Rank)	% (Rank)	(Rank)						
ShawneeM&I	124	5.6 (9)	8.9 (37)		119	9.2 (9)	105	1.9 (5)	119	83.2 (1)
ShawneeMa	1,047	6.7 (23)	5.0 (5)		844	12.1 (25)	734	2.3 (11)	1,039	65.7 (20)
Sheridan	6	* * * *	* * * *		3	* *	2	* *	6	* *
Sherman	62	* * * *	* * * *		51	* *	45	* *	58	* *
Smith	15	* * * *	* * * *		11	* *	9	* *	15	* *
Stafford	30	* * * *	* * * *		27	* *	23	* *	28	* *
Stanton	21	* * * *	* * * *		19	* *	16	* *	9	* *
Stevens	32	* * * *	* * * *		38	* *	21	* *	55	* *
Sumner	146	6.2 (20)	7.5 (34)		109	6.4 (2)	102	3.9 (25)	143	65.0 (21)
Thomas	44	* * * *	* * * *		30	* *	28	* *	40	* *
Trego	11	* * * *	* * * *		9	* *	6	* *	11	* *
Wabaunsee	17	* * * *	* * * *		10	* *	8	* *	16	* *
Wallace	1	* * * *	* * * *		0	* *	0	* *	1	* *
Washington	23	* * * *	* * * *		20	* *	17	* *	23	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators ^(2, 3)										
Clinic	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)		Number	%	Number	%	Number	%
	Number	% (Rank)	% (Rank)	(Rank)						
Wichita	11	* * * *	* * * *	9	* *	7	* *	11	* *	
Wilson	76	* * * *	* * * *	67	* *	56	* *	62	* *	
Woodson	2	* * * *	* * * *	1	* *	1	* *	2	* *	
Wyandotte	1,855	8.0 (28)	6.8 (24)	1,485	12.1 (25)	1,287	2.9 (18)	1,785	71.8 (16)	
Kansas	18,210	7.2	6.5	14,503	10.7	12,744	2.8	16,938	69.4	
Nation (Prior Year)	1,039,036	8.2	6.7	947,705	11.9	856,467	3.7	936,194	64.6	

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3A
2006 - 2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Racial and Ethnic Distribution by Clinic
Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Clinic	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaska Native	Asian/Pacific Islander	Multiple Races	All Other/Unknown
		%	%	%	%	%	%	%
Allen	300	92.0	4.3	2.0	0.7	0.7	0.3	0.0
Anderson	159	93.1	0.6	4.4	0.6	1.3	0.0	0.0
Atchison	396	87.1	10.1	1.0	0.0	0.5	1.3	0.0
Barber-Kio	8	*	*	*	*	*	*	*
Barber-Med	95	*	*	*	*	*	*	*
Barton	845	73.4	1.4	24.9	0.1	0.0	0.2	0.0
Bourbon	371	92.5	3.5	2.7	0.0	0.5	0.8	0.0
Brown	288	77.4	2.4	3.1	14.9	0.0	2.1	0.0
Butler	802	91.4	2.0	4.5	0.4	0.1	1.6	0.0
Chase	41	*	*	*	*	*	*	*
Chautauqua	86	*	*	*	*	*	*	*
Cherokee	672	92.1	1.5	3.7	1.2	0.7	0.7	0.0
Cheyenne	5	*	*	*	*	*	*	*
Clark	13	*	*	*	*	*	*	*
Clay	206	96.6	0.5	1.5	1.0	0.5	0.0	0.0
Cloud	235	94.0	1.3	3.4	0.4	0.9	0.0	0.0
Coffey	75	*	*	*	*	*	*	*
Comanche	14	*	*	*	*	*	*	*
Cowley-Ark	543	74.4	2.2	22.1	0.6	0.4	0.4	0.0
Cowley-Win	465	87.7	1.5	4.9	1.5	3.7	0.6	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3A
2006 - 2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Racial and Ethnic Distribution by Clinic
Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Clinic	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaska Native	Asian/Pacific Islander	Multiple Races	All Other/Unknown
		%	%	%	%	%	%	%
Crawford-G	11	*	*	*	*	*	*	*
Crawford-P	1,200	86.3	3.3	7.8	0.2	1.8	0.8	0.0
Decatur	31	*	*	*	*	*	*	*
DickinsonA	295	93.6	0.7	4.4	1.0	0.0	0.3	0.0
DickinsonH	45	*	*	*	*	*	*	*
Doniphan	123	86.2	4.1	7.3	1.6	0.0	0.8	0.0
Douglas	1,433	64.3	8.4	10.7	9.6	4.7	2.2	0.0
Edwards	76	*	*	*	*	*	*	*
Elk	56	*	*	*	*	*	*	*
Ellis	439	85.4	2.3	11.2	0.2	0.9	0.0	0.0
Ellsworth	88	*	*	*	*	*	*	*
Finney	2,082	34.0	1.1	62.8	0.0	2.0	0.1	0.0
Ford	1,872	23.2	0.8	75.1	0.2	0.7	0.1	0.0
Franklin	507	90.1	2.4	4.3	1.2	0.4	1.6	0.0
Ft. Riley	474	81.9	8.9	6.5	0.6	2.1	0.0	0.0
Geary	948	54.0	24.3	13.8	0.5	3.7	3.7	0.0
Gove	35	*	*	*	*	*	*	*
Graham	50	*	*	*	*	*	*	*
Grant	246	27.2	0.0	72.8	0.0	0.0	0.0	0.0
Gray	170	68.8	0.0	30.6	0.0	0.0	0.6	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3A
2006 - 2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Racial and Ethnic Distribution by Clinic
Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Clinic	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaska Native	Asian/Pacific Islander	Multiple Races	All Other/Unknown
		%	%	%	%	%	%	%
Greeley	23	*	*	*	*	*	*	*
Greenwood	130	93.8	0.8	3.8	0.0	0.8	0.8	0.0
Hamilton	55	*	*	*	*	*	*	*
Harper	154	81.8	0.0	16.2	0.6	0.0	1.3	0.0
Harvey	772	69.4	2.1	26.2	0.6	0.0	1.7	0.0
Haskell	65	*	*	*	*	*	*	*
Hodgeman	30	*	*	*	*	*	*	*
Jackson	242	79.3	0.0	5.4	14.5	0.0	0.8	0.0
Jefferson	199	95.5	0.0	1.5	2.0	0.5	0.5	0.0
Jewell	41	*	*	*	*	*	*	*
Johnson-M	2,576	45.7	15.4	34.4	0.5	2.8	1.2	0.0
Johnson-O	2,888	47.6	10.0	37.7	0.5	3.4	0.8	0.0
Kearny	54	*	*	*	*	*	*	*
Kingman	164	93.3	0.0	4.3	1.8	0.6	0.0	0.0
Kiowa	40	*	*	*	*	*	*	*
Labette	528	88.3	6.4	3.8	0.6	0.2	0.8	0.0
Lane	39	*	*	*	*	*	*	*
Leavenwort	1,055	70.0	16.0	8.9	0.7	1.7	2.7	0.0
Lincoln	67	*	*	*	*	*	*	*
Linn	179	96.1	0.0	2.8	0.0	0.0	1.1	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3A
2006 - 2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Racial and Ethnic Distribution by Clinic
Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Clinic	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaska Native	Asian/Pacific Islander	Multiple Races	All Other/Unknown
		%	%	%	%	%	%	%
Logan	28	*	*	*	*	*	*	*
Lyon	1,091	54.5	5.1	38.7	0.1	1.1	0.5	0.0
Marion	149	92.6	0.7	6.0	0.7	0.0	0.0	0.0
Marshall	172	93.0	0.6	2.9	2.3	0.0	1.2	0.0
Mc Pherson	435	88.7	1.4	8.0	1.4	0.2	0.2	0.0
Meade	123	60.2	0.0	39.8	0.0	0.0	0.0	0.0
Miami	389	93.1	2.3	3.1	0.0	0.5	1.0	0.0
Mitchell	105	95.2	0.0	4.8	0.0	0.0	0.0	0.0
Montgome-C	467	71.5	13.3	10.1	3.0	0.9	1.3	0.0
Montgome-I	534	82.4	7.7	6.6	2.2	0.9	0.2	0.0
Morris	87	*	*	*	*	*	*	*
Nemaha	75	*	*	*	*	*	*	*
Neosho-Ch	407	91.9	2.5	4.2	0.5	0.7	0.2	0.0
Neosho-Er	45	*	*	*	*	*	*	*
Ness	23	*	*	*	*	*	*	*
Norton	81	*	*	*	*	*	*	*
Osage	241	97.9	0.0	1.2	0.4	0.0	0.4	0.0
Osborne	55	*	*	*	*	*	*	*
Ottawa	76	*	*	*	*	*	*	*
Pawnee	148	83.1	2.0	13.5	0.0	0.7	0.7	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3A
2006 - 2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Racial and Ethnic Distribution by Clinic
Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Clinic	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaska Native	Asian/Pacific Islander	Multiple Races	All Other/Unknown
		%	%	%	%	%	%	%
Phillips	103	93.2	1.0	4.9	0.0	1.0	0.0	0.0
Pottawat-S	59	*	*	*	*	*	*	*
Pottawat-W	70	*	*	*	*	*	*	*
Pratt	234	86.3	0.0	13.2	0.0	0.4	0.0	0.0
Rawlins	23	*	*	*	*	*	*	*
Reno	1,446	83.2	2.9	11.8	0.6	0.9	0.6	0.0
Republic	91	*	*	*	*	*	*	*
Rice	211	79.6	1.9	18.5	0.0	0.0	0.0	0.0
Riley-Ft	1,031	72.6	12.6	10.7	0.7	3.2	0.2	0.0
Riley-Manh	1,176	75.3	10.4	7.3	0.6	6.1	0.3	0.0
Rooks	119	95.0	0.8	1.7	0.8	0.8	0.8	0.0
Rush	47	*	*	*	*	*	*	*
Russell	118	93.2	4.2	1.7	0.8	0.0	0.0	0.0
Saline	1,351	73.1	4.3	19.5	0.3	1.8	1.1	0.0
Scott	142	66.2	0.7	33.1	0.0	0.0	0.0	0.0
Sedgwick-C	3,163	45.3	13.6	32.2	1.1	6.4	1.2	0.0
Sedgwick-E	622	44.1	18.8	32.0	1.3	2.6	1.3	0.0
Sedgwick-M	5,347	36.0	26.3	32.6	1.5	2.7	1.1	0.0
Sedgwick-S	3,247	69.2	7.9	16.4	2.3	2.5	1.7	0.0
Seneca	32	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3A
2006 - 2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Racial and Ethnic Distribution by Clinic
Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Clinic	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaska Native	Asian/Pacific Islander	Multiple Races	All Other/Unknown
		%	%	%	%	%	%	%
Seward	445	19.6	1.8	78.0	0.2	0.2	0.2	0.0
Seward	997	18.5	1.4	76.3	2.4	1.1	0.3	0.0
ShawneeM&I	643	35.6	15.6	43.9	2.6	1.2	1.1	0.0
ShawneeMa	3,635	58.7	17.2	19.7	2.1	0.7	1.6	0.0
Sheridan	29	*	*	*	*	*	*	*
Sherman	232	77.6	0.0	20.7	0.4	0.4	0.9	0.0
Smith	62	*	*	*	*	*	*	*
Stafford	121	71.1	0.0	28.1	0.0	0.8	0.0	0.0
Stanton	58	*	*	*	*	*	*	*
Stevens	36	*	*	*	*	*	*	*
Stevens	169	45.6	0.0	53.8	0.0	0.6	0.0	0.0
Sumner	427	94.4	0.5	4.2	0.5	0.2	0.2	0.0
Thomas	158	86.1	1.3	9.5	0.6	1.3	1.3	0.0
Trego	48	*	*	*	*	*	*	*
Wabaunsee	55	*	*	*	*	*	*	*
Wallace	8	*	*	*	*	*	*	*
Washington	81	*	*	*	*	*	*	*
Wichita	57	*	*	*	*	*	*	*
Wilson	294	95.9	0.3	2.7	0.0	0.3	0.7	0.0
Woodson	28	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3A
2006 - 2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Racial and Ethnic Distribution by Clinic
Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Wyandotte	6,483	25.8	31.6	39.8	0.2	2.3	0.3	0.0
Kansas	62,530	58.3	11.4	26.2	1.2	2.0	0.9	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2006 - 2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Age Distribution by Clinic
Combined 3 Years

Comparison of Age Distribution ⁽²⁾								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Allen	300	0.7	9.0	15.7	63.7	10.0	0.3	0.7
Anderson	159	0.0	8.2	12.6	66.7	11.9	0.6	0.0
Atchison	396	0.0	7.6	18.7	61.1	11.6	0.8	0.3
Barber-Kio	8	*	*	*	*	*	*	*
Barber-Med	95	*	*	*	*	*	*	*
Barton	845	0.1	6.3	17.6	61.7	14.0	0.4	0.0
Bourbon	371	0.3	5.9	18.9	65.5	8.9	0.5	0.0
Brown	288	0.0	6.9	17.4	64.6	9.7	1.0	0.3
Butler	802	0.1	6.4	16.5	62.0	14.2	0.9	0.0
Chase	41	*	*	*	*	*	*	*
Chautauqua	86	*	*	*	*	*	*	*
Cherokee	672	0.1	7.9	18.0	60.1	12.9	0.6	0.3
Cheyenne	5	*	*	*	*	*	*	*
Clark	13	*	*	*	*	*	*	*
Clay	206	0.0	4.4	12.6	67.0	15.0	1.0	0.0
Cloud	235	0.4	7.2	15.3	62.1	13.2	1.7	0.0
Coffey	75	*	*	*	*	*	*	*
Comanche	14	*	*	*	*	*	*	*
Cowley-Ark	543	0.7	9.6	16.4	57.8	14.5	0.9	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2006 - 2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Age Distribution by Clinic
Combined 3 Years

Comparison of Age Distribution ⁽²⁾								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Cowley-Win	465	0.2	6.9	18.3	61.3	13.3	0.0	0.0
Crawford-G	11	*	*	*	*	*	*	*
Crawford-P	1,200	0.2	7.0	15.9	63.3	12.7	0.8	0.2
Decatur	31	*	*	*	*	*	*	*
DickinsonA	295	0.3	6.4	13.2	62.0	17.6	0.3	0.0
DickinsonH	45	*	*	*	*	*	*	*
Doniphan	123	0.0	4.9	17.1	65.0	13.0	0.0	0.0
Douglas	1,433	0.1	4.0	12.9	64.2	17.3	1.4	0.0
Edwards	76	*	*	*	*	*	*	*
Elk	56	*	*	*	*	*	*	*
Ellis	439	0.2	5.2	12.5	68.8	11.2	1.8	0.2
Ellsworth	88	*	*	*	*	*	*	*
Finney	2,082	0.5	10.3	13.6	54.9	19.3	1.2	0.1
Ford	1,872	0.4	6.9	12.0	58.1	21.4	1.0	0.2
Franklin	507	1.0	7.7	16.8	61.5	12.4	0.4	0.2
Ft. Riley	474	0.2	1.3	11.6	77.6	8.9	0.4	0.0
Geary	948	0.4	5.8	16.2	66.2	10.8	0.5	0.0
Gove	35	*	*	*	*	*	*	*
Graham	50	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2006 - 2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Age Distribution by Clinic
Combined 3 Years

Comparison of Age Distribution ⁽²⁾								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Grant	246	0.4	6.5	14.2	56.9	19.9	2.0	0.0
Gray	170	0.0	5.9	9.4	51.8	30.0	2.4	0.6
Greeley	23	*	*	*	*	*	*	*
Greenwood	130	0.0	6.9	16.9	63.1	13.1	0.0	0.0
Hamilton	55	*	*	*	*	*	*	*
Harper	154	0.0	7.1	11.7	67.5	13.6	0.0	0.0
Harvey	772	0.1	6.0	11.4	65.7	14.9	1.9	0.0
Haskell	65	*	*	*	*	*	*	*
Hodgeman	30	*	*	*	*	*	*	*
Jackson	242	0.0	7.0	13.2	69.8	9.9	0.0	0.0
Jefferson	199	0.0	10.6	21.6	56.3	10.6	1.0	0.0
Jewell	41	*	*	*	*	*	*	*
Johnson-M	2,576	0.3	4.0	12.2	62.3	20.1	1.0	0.0
Johnson-O	2,888	0.1	4.6	11.3	62.6	20.0	1.3	0.1
Kearny	54	*	*	*	*	*	*	*
Kingman	164	0.0	4.9	13.4	64.0	17.7	0.0	0.0
Kiowa	40	*	*	*	*	*	*	*
Labette	528	0.0	6.8	17.4	65.7	9.5	0.6	0.0
Lane	39	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2006 - 2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Age Distribution by Clinic
Combined 3 Years

Comparison of Age Distribution ⁽²⁾								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Leavenwort	1,055	0.3	5.8	18.3	63.9	10.6	1.1	0.0
Lincoln	67	*	*	*	*	*	*	*
Linn	179	0.0	6.1	16.8	63.7	11.7	1.7	0.0
Logan	28	*	*	*	*	*	*	*
Lyon	1,091	0.5	6.6	13.3	61.9	15.8	2.0	0.0
Marion	149	0.0	3.4	8.1	68.5	17.4	2.7	0.0
Marshall	172	0.0	5.2	10.5	69.8	14.5	0.0	0.0
Mc Pherson	435	0.2	5.3	13.6	63.9	16.1	0.9	0.0
Meade	123	0.0	5.7	11.4	64.2	17.9	0.8	0.0
Miami	389	0.0	7.7	15.4	65.8	10.5	0.5	0.0
Mitchell	105	0.0	4.8	19.0	54.3	21.9	0.0	0.0
Montgome-C	467	1.3	8.6	18.2	60.2	11.3	0.2	0.2
Montgome-I	534	0.2	6.9	16.3	65.7	10.3	0.6	0.0
Morris	87	*	*	*	*	*	*	*
Nemaha	75	*	*	*	*	*	*	*
Neosho-Ch	407	0.7	5.9	21.4	61.7	10.1	0.2	0.0
Neosho-Er	45	*	*	*	*	*	*	*
Ness	23	*	*	*	*	*	*	*
Norton	81	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2006 - 2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Age Distribution by Clinic
Combined 3 Years

Comparison of Age Distribution ⁽²⁾								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Osage	241	0.4	3.3	17.8	63.9	14.1	0.0	0.4
Osborne	55	*	*	*	*	*	*	*
Ottawa	76	*	*	*	*	*	*	*
Pawnee	148	0.7	4.1	16.9	66.9	10.1	1.4	0.0
Phillips	103	1.0	1.9	15.5	60.2	21.4	0.0	0.0
Pottawat-S	59	*	*	*	*	*	*	*
Pottawat-W	70	*	*	*	*	*	*	*
Pratt	234	0.4	8.5	16.2	64.1	10.3	0.4	0.0
Rawlins	23	*	*	*	*	*	*	*
Reno	1,446	0.3	4.8	15.5	66.7	12.0	0.6	0.1
Republic	91	*	*	*	*	*	*	*
Rice	211	0.5	7.1	14.7	57.8	18.0	1.9	0.0
Riley-Ft	1,031	0.1	0.6	11.0	78.3	9.8	0.3	0.0
Riley-Manh	1,176	0.0	3.6	13.5	69.3	13.0	0.5	0.1
Rooks	119	0.0	4.2	13.4	63.0	16.8	2.5	0.0
Rush	47	*	*	*	*	*	*	*
Russell	118	0.0	7.6	15.3	68.6	8.5	0.0	0.0
Saline	1,351	0.2	8.0	16.7	61.4	13.0	0.7	0.1
Scott	142	0.7	7.0	6.3	71.8	12.0	1.4	0.7

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2006 - 2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Age Distribution by Clinic
Combined 3 Years

Comparison of Age Distribution ⁽²⁾								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Sedgwick-C	3,163	0.3	6.6	12.9	62.7	16.3	1.2	0.1
Sedgwick-E	622	0.5	5.5	13.5	61.6	17.7	1.3	0.0
Sedgwick-M	5,347	0.4	7.2	14.4	59.8	16.7	1.4	0.1
Sedgwick-S	3,247	0.2	6.8	14.2	63.1	14.4	1.2	0.1
Seneca	32	*	*	*	*	*	*	*
Seward	445	0.4	7.9	13.3	56.6	20.4	0.9	0.4
Seward	997	0.5	7.4	13.5	56.6	20.3	1.7	0.0
ShawneeM&I	643	0.9	8.9	14.2	58.6	16.2	1.2	0.0
ShawneeMa	3,635	0.4	6.1	14.1	64.0	14.5	0.7	0.1
Sheridan	29	*	*	*	*	*	*	*
Sherman	232	0.0	5.2	20.7	58.2	13.4	2.6	0.0
Smith	62	*	*	*	*	*	*	*
Stafford	121	0.0	4.1	11.6	56.2	26.4	1.7	0.0
Stanton	58	*	*	*	*	*	*	*
Stevens	36	*	*	*	*	*	*	*
Stevens	169	0.6	5.9	14.8	62.7	15.4	0.6	0.0
Sumner	427	0.2	6.1	16.9	63.5	13.1	0.2	0.0
Thomas	158	0.0	4.4	12.0	66.5	15.8	0.6	0.6
Trego	48	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2006 - 2008 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Age Distribution by Clinic
Combined 3 Years

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Wabaunsee	55	*	*	*	*	*	*	*
Wallace	8	*	*	*	*	*	*	*
Washington	81	*	*	*	*	*	*	*
Wichita	57	*	*	*	*	*	*	*
Wilson	294	0.0	11.6	12.9	63.3	12.2	0.0	0.0
Woodson	28	*	*	*	*	*	*	*
Wyandotte	6,483	0.7	8.2	14.3	58.5	17.1	1.2	0.1
Kansas	62,530	0.3	6.4	14.3	62.2	15.6	1.1	0.1

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2006 - 2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Health Indicators by Clinic
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Allen	297	11.8 (46)	48.5 (59)	270	21.9 (35)	47.0 (41)	49	* *	86	* *	271	4.1 (64)	271	7.4(68)
Anderson	151	13.9 (60)	41.1 (4)	142	24.6 (55)	43.0 (16)	22	* *	28	* *	144	0.7 (9)	144	6.9(66)
Atchison	383	13.3 (57)	47.5 (54)	283	26.9 (63)	43.1 (17)	66	* *	66	* *	345	2.3 (50)	345	7.0(67)
Barber-Kio	7	* *	* *	6	* *	* *	2	* *	1	* *	7	* *	7	* *
Barber-Med	93	* *	* *	84	* *	* *	9	* *	21	* *	82	* *	82	* *
Barton	818	10.0 (22)	45.5 (33)	746	25.2 (56)	43.6 (19)	96	* *	145	22.8(12)	731	2.1 (47)	731	3.7(47)
Bourbon	366	11.5 (41)	50.8 (65)	330	23.3 (45)	42.7 (14)	47	* *	73	* *	333	2.4 (52)	333	3.3(41)
Brown	276	8.3 (8)	46.0 (44)	228	18.0 (8)	56.6 (68)	51	* *	47	* *	252	0.0 (1)	252	4.8(59)
Butler	772	13.5 (59)	45.5 (33)	716	20.5 (21)	48.7 (55)	90	* *	212	28.8(25)	686	1.2 (25)	686	5.2(63)
Chase	37	* *	* *	30	* *	* *	5	* *	5	* *	37	* *	37	* *
Chautauqua	85	* *	* *	64	* *	* *	4	* *	10	* *	70	* *	70	* *
Cherokee	624	10.6 (29)	45.5 (33)	503	19.3 (13)	49.5 (60)	72	* *	146	24.0(16)	571	1.8 (40)	571	1.9(15)
Cheyenne	5	* *	* *	5	* *	* *	0	* *	1	* *	5	* *	5	* *
Clark	11	* *	* *	5	* *	* *	0	* *	3	* *	9	* *	9	* *
Clay	198	8.1 (7)	49.5 (61)	181	12.7 (1)	52.5 (66)	25	* *	43	* *	181	3.9 (63)	181	2.2(22)
Cloud	228	13.2 (54)	43.0 (12)	202	24.3 (52)	49.0 (56)	27	* *	49	* *	199	1.5 (31)	200	2.5(27)
Coffey	75	* *	* *	73	* *	* *	6	* *	21	* *	66	* *	66	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).
(5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2006 - 2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Health Indicators by Clinic
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Comanche	14	* *	* *	12	* *	* *	2	* *	5	* *	10	* *	10	* *
Cowley-Ark	530	11.7 (44)	44.5 (24)	481	25.2 (56)	41.6 (12)	86	* *	68	* *	471	1.9 (42)	471	4.0(52)
Cowley-Win	446	11.9 (49)	45.5 (33)	404	20.8 (25)	47.5 (43)	52	* *	56	* *	404	0.7 (9)	404	2.2(22)
Crawford-G	11	* *	* *	10	* *	* *	1	* *	0	* *	11	* *	11	* *
Crawford-P	1,133	13.2 (54)	44.6 (25)	953	21.0 (29)	48.5 (53)	162	13.6 (1)	189	16.9 (4)	1,020	1.9 (42)	1,019	2.5(27)
Decatur	31	* *	* *	27	* *	* *	6	* *	5	* *	27	* *	27	* *
DickinsonA	280	10.4 (27)	40.7 (2)	251	23.9 (50)	45.8 (30)	34	* *	59	* *	254	2.4 (52)	254	3.5(45)
DickinsonH	45	* *	* *	42	* *	* *	10	* *	12	* *	42	* *	42	* *
Doniphan	118	5.1 (1)	56.8 (70)	90	* *	* *	25	* *	24	* *	98	* *	98	* *
Douglas	1,378	11.7 (44)	43.1 (15)	1,157	18.8 (10)	51.4 (65)	192	31.8 (15)	351	29.3(27)	1,168	1.7 (37)	1,167	2.0(19)
Edwards	74	* *	* *	61	* *	* *	9	* *	17	* *	59	* *	59	* *
Elk	53	* *	* *	47	* *	* *	2	* *	8	* *	50	* *	50	* *
Ellis	421	14.0 (62)	37.8 (1)	381	19.2 (12)	43.8 (20)	89	* *	103	13.6 (1)	370	2.2 (49)	370	4.6(54)
Ellsworth	85	* *	* *	76	* *	* *	9	* *	22	* *	76	* *	76	* *
Finney	2,025	11.6 (43)	44.7 (26)	1,863	25.6 (59)	39.9 (8)	257	29.2 (12)	265	23.8(15)	1,860	2.6 (57)	1,860	1.4(7)
Ford	1,824	7.0 (5)	45.3 (29)	1,652	24.3 (52)	43.2 (18)	258	22.5 (3)	187	19.8 (8)	1,661	1.2 (25)	1,661	1.9(15)
Franklin	484	11.4 (39)	46.3 (48)	423	17.5 (6)	48.2 (49)	48	* *	115	17.4 (5)	425	0.7 (9)	425	4.7(57)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).
(5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2006 - 2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Health Indicators by Clinic
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight	Over- weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
							Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)
Ft. Riley	399	11.8 (46)	41.9 (7)	334	17.7 (7)	50.3 (64)	79	* *	17	* *	411	0.2 (4)	411	0.2(3)
Geary	916	10.9 (34)	45.4 (31)	771	20.4 (19)	48.5 (53)	150	38.0 (19)	137	28.5(24)	775	2.5 (55)	775	4.6(54)
Gove	33	* *	* *	29	* *	* *	6	* *	8	* *	32	* *	32	* *
Graham	50	* *	* *	42	* *	* *	6	* *	9	* *	46	* *	46	* *
Grant	242	9.1 (12)	53.7 (69)	233	20.2 (16)	40.8 (11)	41	* *	67	* *	226	1.3 (28)	226	0.4(4)
Gray	164	5.5 (3)	45.7 (41)	144	31.3 (65)	34.7 (2)	22	* *	42	* *	143	0.7 (9)	143	4.9(60)
Greeley	23	* *	* *	22	* *	* *	3	* *	6	* *	22	* *	22	* *
Greenwood	127	7.9 (6)	52.0 (67)	105	21.9 (35)	46.7 (37)	15	* *	27	* *	118	0.8 (14)	118	3.4(42)
Hamilton	55	* *	* *	46	* *	* *	9	* *	19	* *	47	* *	47	* *
Harper	146	17.8 (70)	42.5 (10)	118	33.1 (68)	39.0 (6)	17	* *	17	* *	130	1.5 (31)	129	4.7(57)
Harvey	755	10.7 (30)	47.5 (54)	687	22.6 (43)	46.7 (37)	108	28.7 (11)	126	26.2(19)	678	2.1 (47)	676	2.4(24)
Haskell	64	* *	* *	58	* *	* *	14	* *	25	* *	59	* *	59	* *
Hodgeman	30	* *	* *	29	* *	* *	6	* *	7	* *	26	* *	26	* *
Jackson	234	10.7 (30)	46.6 (49)	202	23.8 (49)	44.1 (23)	37	* *	48	* *	221	0.5 (5)	221	1.4(7)
Jefferson	191	14.7 (64)	48.2 (57)	161	18.0 (8)	48.4 (51)	26	* *	51	* *	173	2.9 (60)	173	1.2(6)
Jewell	39	* *	* *	39	* *	* *	4	* *	6	* *	40	* *	40	* *
Johnson-M	2,466	10.2 (25)	44.3 (21)	2,089	20.6 (24)	46.6 (36)	421	22.6 (4)	596	18.1 (6)	2,209	3.3 (62)	2,210	1.6(11)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).
(5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2006 - 2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Health Indicators by Clinic
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight	Over- weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
							Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)
Johnson-O	2,766	10.8 (32)	43.9 (18)	2,445	20.9 (26)	45.1 (26)	475	24.8 (7)	827	15.0 (3)	2,477	4.4 (66)	2,476	2.7(30)
Kearny	53	* *	* *	50	* *	* *	15	* *	19	* *	52	* *	52	* *
Kingman	158	13.9 (60)	44.3 (21)	120	22.5 (42)	47.5 (43)	21	* *	40	* *	136	4.4 (66)	136	5.1(61)
Kiowa	38	* *	* *	29	* *	* *	3	* *	5	* *	34	* *	34	* *
Labette	517	10.1 (23)	48.4 (58)	431	23.0 (44)	48.3 (50)	86	* *	114	18.4 (7)	471	1.3 (28)	471	4.2(53)
Lane	37	* *	* *	30	* *	* *	9	* *	4	* *	31	* *	31	* *
Leavenwort	1,006	11.0 (35)	46.0 (44)	884	20.9 (26)	50.1 (63)	161	41.0 (20)	210	22.4(11)	867	0.9 (17)	868	2.0(19)
Lincoln	65	* *	* *	54	* *	* *	9	* *	12	* *	57	* *	57	* *
Linn	173	11.0 (35)	49.1 (60)	160	28.8 (64)	43.8 (20)	23	* *	33	* *	162	0.6 (8)	162	3.1(37)
Logan	27	* *	* *	22	* *	* *	5	* *	6	* *	22	* *	22	* *
Lyon	1,058	8.8 (11)	50.1 (64)	970	23.5 (47)	47.7 (47)	157	17.2 (2)	183	14.2 (2)	958	1.5 (31)	958	5.2(63)
Marion	144	10.4 (27)	44.4 (23)	132	20.5 (21)	34.8 (3)	16	* *	29	* *	128	0.8 (14)	128	3.9(50)
Marshall	166	9.6 (15)	45.2 (28)	124	26.6 (60)	42.7 (14)	27	* *	28	* *	136	0.0 (1)	136	1.5(10)
Mc Pherson	418	13.4 (58)	47.1 (53)	379	23.7 (48)	42.2 (13)	69	* *	83	* *	380	1.1 (20)	380	3.4(42)
Meade	122	6.6 (4)	50.0 (63)	106	24.5 (54)	49.1 (57)	18	* *	21	* *	106	1.9 (42)	106	1.9(15)
Miami	368	14.7 (64)	45.4 (31)	343	20.1 (15)	49.6 (61)	78	* *	89	* *	334	1.8 (40)	334	1.8(14)
Mitchell	103	9.7 (16)	45.6 (39)	94	* *	* *	18	* *	32	* *	91	* *	91	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).
(5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2006 - 2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Health Indicators by Clinic
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Montgome-C	463	11.9 (49)	43.0 (12)	391	26.6 (60)	40.2 (9)	73	* *	123	29.3(27)	408	0.7 (9)	408	2.9(33)
Montgome-I	529	18.5 (71)	42.0 (8)	436	26.6 (60)	46.8 (39)	96	* *	137	24.8(18)	467	0.9 (17)	467	3.4(42)
Morris	84	* *	* *	68	* *	* *	11	* *	15	* *	75	* *	75	* *
Nemaha	74	* *	* *	67	* *	* *	20	* *	24	* *	66	* *	66	* *
Neosho-Ch	397	15.1 (68)	46.1 (46)	363	17.4 (5)	47.1 (42)	65	* *	60	* *	358	1.1 (20)	358	2.0(19)
Neosho-Er	42	* *	* *	38	* *	* *	12	* *	8	* *	42	* *	42	* *
Ness	22	* *	* *	19	* *	* *	4	* *	11	* *	17	* *	17	* *
Norton	80	* *	* *	61	* *	* *	7	* *	16	* *	69	* *	69	* *
Osage	235	11.1 (37)	45.5 (33)	189	23.3 (45)	47.6 (46)	28	* *	48	* *	209	2.4 (52)	209	2.9(33)
Osborne	54	* *	* *	49	* *	* *	8	* *	7	* *	51	* *	51	* *
Ottawa	75	* *	* *	66	* *	* *	9	* *	20	* *	65	* *	65	* *
Pawnee	145	15.2 (69)	53.1 (68)	128	32.0 (66)	40.6 (10)	11	* *	22	* *	128	0.8 (14)	128	3.1(37)
Phillips	98	* *	* *	82	* *	* *	5	* *	13	* *	86	* *	86	* *
Pottawat-S	59	* *	* *	56	* *	* *	5	* *	29	* *	56	* *	56	* *
Pottawat-W	69	* *	* *	59	* *	* *	10	* *	24	* *	60	* *	60	* *
Pratt	223	9.9 (18)	43.0 (12)	193	20.2 (16)	46.1 (33)	29	* *	56	* *	191	0.5 (5)	191	1.6(11)
Rawlins	23	* *	* *	21	* *	* *	3	* *	10	* *	21	* *	21	* *

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
- (4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).
- (5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).
- (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2006 - 2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Health Indicators by Clinic
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Reno	1,419	11.8 (46)	46.7 (50)	1,282	21.8 (34)	46.3 (35)	213	28.6 (10)	346	22.8(12)	1,243	1.7 (37)	1,244	2.4(24)
Republic	87	* *	* *	81	* *	* *	12	* *	12	* *	83	* *	83	* *
Rice	204	9.3 (13)	49.5 (61)	181	21.5 (33)	44.8 (25)	23	* *	34	* *	184	1.1 (20)	184	1.6(11)
Riley-Ft	1,021	9.5 (14)	41.5 (6)	806	20.0 (14)	49.3 (58)	142	26.8 (8)	180	38.9(29)	740	1.2 (25)	740	3.0(35)
Riley-Manh	1,133	10.2 (25)	41.0 (3)	934	17.2 (4)	48.4 (51)	141	24.1 (6)	236	21.6(10)	969	2.5 (55)	969	3.1(37)
Rooks	116	12.1 (53)	43.1 (15)	102	25.5 (58)	38.2 (4)	17	* *	25	* *	110	0.9 (17)	110	3.6(46)
Rush	46	* *	* *	41	* *	* *	8	* *	7	* *	41	* *	41	* *
Russell	116	14.7 (64)	41.4 (5)	102	32.4 (67)	30.4 (1)	18	* *	34	* *	103	1.9 (42)	103	1.9(15)
Saline	1,328	11.4 (39)	44.8 (27)	1,216	21.2 (30)	45.4 (27)	248	23.8 (5)	297	20.9 (9)	1,201	1.5 (31)	1,201	3.2(40)
Scott	140	14.3 (63)	42.9 (11)	123	17.1 (3)	55.3 (67)	19	* *	31	* *	127	0.0 (1)	127	0.0(1)
Sedgwick-C	3,110	11.5 (41)	45.6 (39)	2,886	20.2 (16)	45.5 (28)	594	37.5 (18)	731	27.9(23)	2,842	2.3 (50)	2,841	2.4(24)
Sedgwick-E	545	9.9 (18)	42.4 (9)	448	18.8 (10)	46.9 (40)	104	26.9 (9)	102	27.5(22)	542	1.1 (20)	542	0.0(1)
Sedgwick-M	5,166	9.9 (18)	45.9 (42)	4,758	21.2 (30)	45.9 (31)	1,077	29.6 (14)	1,501	23.4(14)	4,686	1.5 (31)	4,685	3.0(35)
Sedgwick-S	3,171	13.2 (54)	44.2 (20)	2,962	20.4 (19)	47.5 (43)	692	33.1 (16)	896	26.5(21)	2,896	1.9 (42)	2,897	3.8(49)
Seneca	30	* *	* *	31	* *	* *	5	* *	9	* *	31	* *	31	* *
Seward	407	10.1 (23)	46.7 (50)	368	16.0 (2)	44.6 (24)	81	* *	32	* *	417	0.5 (5)	417	0.5(5)
Seward	986	8.7 (10)	45.5 (33)	901	22.1 (37)	45.6 (29)	161	29.2 (12)	171	24.0(16)	854	3.0 (61)	852	4.6(54)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).
(5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2006 - 2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Health Indicators by Clinic
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)														
Clinic	Prepregnancy BMI ⁽⁴⁾			Weight Gain ⁽⁵⁾			Anemia (Low Hb/Hct) ⁽⁶⁾				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
ShawneeM&I	640	9.8 (17)	45.3 (29)	511	23.9 (50)	39.5 (7)	10	* *	44	* *	507	4.5 (68)	507	1.4 (7)
ShawneeMa	3,513	10.8 (32)	45.9 (42)	3,160	20.9 (26)	49.3 (58)	415	41.7 (21)	667	28.8 (25)	3,211	2.6 (57)	3,214	2.7 (30)
Sheridan	28	* *	* *	22	* *	* *	3	* *	5	* *	24	* *	24	* *
Sherman	217	12.0 (51)	47.9 (56)	185	22.2 (40)	45.9 (31)	16	* *	33	* *	195	1.5 (31)	195	5.1 (61)
Smith	61	* *	* *	42	* *	* *	11	* *	22	* *	54	* *	54	* *
Stafford	118	8.5 (9)	56.8 (70)	98	* *	* *	15	* *	19	* *	99	* *	99	* *
Stanton	58	* *	* *	54	* *	* *	7	* *	21	* *	54	* *	54	* *
Stevens	24	* *	* *	22	* *	* *	3	* *	5	* *	36	* *	36	* *
Stevens	169	5.3 (2)	51.5 (66)	151	20.5 (21)	49.7 (62)	26	* *	43	* *	142	4.2 (65)	142	5.6 (65)
Sumner	415	12.0 (51)	43.1 (15)	285	21.4 (32)	47.7 (47)	87	* *	93	* *	357	1.7 (37)	357	3.9 (50)
Thomas	154	14.9 (67)	46.1 (46)	140	22.1 (37)	38.6 (5)	20	* *	28	* *	141	2.8 (59)	141	2.8 (32)
Trego	44	* *	* *	37	* *	* *	5	* *	9	* *	45	* *	45	* *
Wabaunsee	54	* *	* *	48	* *	* *	5	* *	19	* *	50	* *	50	* *
Wallace	7	* *	* *	7	* *	* *	1	* *	1	* *	7	* *	7	* *
Washington	76	* *	* *	69	* *	* *	3	* *	15	* *	71	* *	70	* *
Wichita	57	* *	* *	54	* *	* *	7	* *	10	* *	53	* *	53	* *
Wilson	284	11.3 (38)	44.0 (19)	232	22.4 (41)	44.0 (22)	48	* *	45	* *	267	1.1 (20)	267	3.7 (47)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).
(5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2006 - 2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Health Indicators by Clinic
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Woodson	26	* *	* *	22	* *	* *	4	* *	3	* *	25	* *	25	* *
Wyandotte	6,225	9.9 (18)	46.8 (52)	5,748	22.1 (37)	46.1 (33)	1,103	36.0 (17)	1,132	26.4 (20)	5,747	1.4 (30)	5,748	2.5 (27)
Kansas	60,398	10.9	45.5	53,675	21.7	46.1	9,707	30.3	12,867	23.4	54,316	1.9	54,314	2.9

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).
(5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6A
2006 - 2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Behavioral Indicators by Clinic
Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2, 3)										
Clinic	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	% (Rank)		Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Allen	272	90.1 (5)	1.1 (17)		253	41.5 (42)	69	* *	151	84.1 (21)
Anderson	154	87.7 (15)	1.3 (21)		121	49.6 (12)	38	* *	66	* *
Atchison	352	90.1 (5)	1.7 (31)		329	37.1 (53)	81	* *	158	74.7 (44)
Barber-Kio	7	* *	* *		7	* *	2	* *	3	* *
Barber-Med	87	* *	* *		74	* *	22	* *	41	* *
Barton	747	80.3 (36)	0.9 (12)		671	57.8 (5)	157	19.7 (2)	367	86.9 (6)
Bourbon	332	93.7 (1)	0.9 (12)		298	45.0 (28)	77	* *	156	78.8 (38)
Brown	245	82.9 (31)	6.9 (63)		231	45.5 (25)	64	* *	165	81.2 (28)
Butler	715	76.8 (49)	6.7 (62)		557	44.2 (31)	154	12.3 (10)	317	77.0 (42)
Chase	36	* *	* *		30	* *	9	* *	18	* *
Chautauqua	70	* *	* *		74	* *	16	* *	41	* *
Cherokee	513	80.3 (36)	4.9 (54)		470	47.4 (19)	115	9.6 (18)	277	79.8 (33)
Cheyenne	5	* *	* *		4	* *	0	* *	2	* *
Clark	8	* *	* *		11	* *	0	* *	8	* *
Clay	187	82.9 (31)	3.7 (48)		154	48.1 (16)	35	* *	55	* *
Cloud	206	85.9 (19)	1.9 (32)		181	54.7 (6)	44	* *	41	* *
Coffey	74	* *	* *		52	* *	14	* *	28	* *
Comanche	12	* *	* *		10	* *	3	* *	6	* *
Cowley-Ark	466	67.2 (68)	5.2 (57)		430	40.5 (45)	116	6.9 (20)	241	79.3 (36)
Cowley-Win	416	79.8 (40)	1.9 (32)		367	49.3 (15)	78	* *	175	81.7 (27)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6A
2006 - 2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Behavioral Indicators by Clinic
Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2, 3)										
Clinic	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Crawford-G	11	* *	* *	* *	10	* *	0	* *	8	* *
Crawford-P	1,025	88.9 (8)	0.8 (9)		937	46.3 (23)	235	8.1 (19)	563	85.6 (15)
Decatur	27	* *	* *	* *	28	* *	5	* *	18	* *
DickinsonA	262	85.1 (21)	3.4 (45)		214	50.0 (11)	62	* *	115	80.0 (32)
DickinsonH	42	* *	* *	* *	38	* *	6	* *	21	* *
Doniphan	105	80.0 (39)	4.8 (53)		98	* *	27	* *	66	* *
Douglas	1,199	74.1 (58)	5.8 (60)		1,122	44.0 (32)	258	12.4 (8)	609	81.0 (30)
Edwards	62	* *	* *	* *	62	* *	19	* *	29	* *
Elk	51	* *	* *	* *	39	* *	5	* *	17	* *
Ellis	396	86.1 (18)	1.0 (15)		349	37.0 (54)	95	* *	215	85.6 (15)
Ellsworth	83	* *	* *	* *	61	* *	16	* *	30	* *
Finney	1,836	70.6 (62)	9.3 (67)		1,682	41.7 (39)	413	5.8 (23)	904	72.2 (48)
Ford	1,654	75.0 (56)	9.4 (68)		1,562	47.4 (19)	347	5.5 (25)	774	78.0 (39)
Franklin	386	81.3 (34)	1.6 (28)		346	39.9 (47)	91	* *	161	86.3 (11)
Ft. Riley	272	86.8 (16)	0.0 (1)		323	37.8 (52)	0	* *	265	100.0 (1)
Geary	752	73.7 (60)	0.4 (4)		769	45.6 (24)	195	15.9 (4)	431	77.5 (41)
Gove	32	* *	* *	* *	24	* *	3	* *	14	* *
Graham	46	* *	* *	* *	41	* *	4	* *	21	* *
Grant	226	85.0 (24)	4.9 (54)		195	36.9 (55)	36	* *	109	86.2 (13)
Gray	149	83.2 (30)	2.0 (34)		137	51.1 (10)	32	* *	69	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6A
2006 - 2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Behavioral Indicators by Clinic
Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2, 3)										
Clinic	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Prenatal (4)		During Pregnancy (5)	
	Number	% (Rank)	%	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Greeley	22	* *	* *	* *	16	* *	7	* *	10	* *
Greenwood	105	73.3 (61)	18.1	(70)	108	51.9 (8)	20	* *	56	* *
Hamilton	45	* *	* *	* *	41	* *	10	* *	21	* *
Harper	127	76.4 (53)	7.1	(64)	123	49.6 (12)	28	* *	63	* *
Harvey	665	74.3 (57)	9.8	(69)	633	47.9 (17)	138	11.6 (13)	312	84.3 (20)
Haskell	59	* *	* *	* *	57	* *	16	* *	29	* *
Hodgeman	28	* *	* *	* *	20	* *	7	* *	10	* *
Jackson	188	84.6 (26)	3.7	(48)	188	42.6 (37)	39	* *	101	90.1 (4)
Jefferson	164	87.8 (13)	1.2	(20)	138	31.9 (64)	36	* *	87	* *
Jewell	40	* *	* *	* *	29	* *	8	* *	17	* *
Johnson-M	2,199	67.8 (67)	1.1	(17)	1,945	32.6 (61)	454	11.7 (12)	1,078	86.6 (8)
Johnson-O	2,419	65.0 (69)	2.9	(42)	2,252	31.5 (65)	568	11.4 (15)	1,262	83.6 (24)
Kearny	51	* *	* *	* *	45	* *	10	* *	20	* *
Kingman	140	79.3 (44)	7.1	(64)	128	44.5 (30)	39	* *	69	* *
Kiowa	33	* *	* *	* *	36	* *	6	* *	23	* *
Labette	444	87.8 (13)	1.4	(22)	443	43.8 (33)	92	* *	262	86.6 (8)
Lane	32	* *	* *	* *	31	* *	7	* *	17	* *
Leavenworth	923	85.5 (20)	0.7	(5)	738	39.0 (50)	202	12.4 (8)	429	86.5 (10)
Lincoln	61	* *	* *	* *	47	* *	14	* *	23	* *
Linn	162	79.6 (42)	3.7	(48)	146	43.8 (33)	30	* *	82	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6A
2006 - 2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Behavioral Indicators by Clinic
Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2, 3)										
Clinic	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Logan	18	* *	* *	* *	24	* *	8	* *	17	* *
Lyon	992	81.7 (33)	0.8 (9)		823	43.3 (35)	193	14.0 (6)	458	84.5 (19)
Marion	136	80.1 (38)	1.5 (23)		106	58.5 (4)	27	* *	60	* *
Marshall	146	91.1 (4)	0.7 (5)		131	40.5 (45)	33	* *	50	* *
Mc Pherson	395	88.4 (10)	1.0 (15)		327	44.6 (29)	74	* *	190	86.3 (11)
Meade	109	75.2 (55)	5.5 (58)		101	46.5 (22)	26	* *	49	* *
Miami	353	76.8 (49)	0.3 (2)		263	35.4 (57)	74	* *	138	76.8 (43)
Mitchell	96	* *	* *	* *	77	* *	24	* *	43	* *
Montgome-C	383	84.1 (27)	3.4 (45)		386	42.7 (36)	84	* *	219	84.9 (18)
Montgome-I	439	88.4 (10)	2.5 (37)		445	41.6 (41)	102	13.7 (7)	258	83.3 (25)
Morris	73	* *	* *	* *	65	* *	14	* *	30	* *
Nemaha	70	* *	* *	* *	54	* *	13	* *	26	* *
Neosho-Ch	356	91.3 (2)	2.2 (35)		332	47.3 (21)	87	* *	162	80.9 (31)
Neosho-Er	41	* *	* *	* *	37	* *	5	* *	19	* *
Ness	21	* *	* *	* *	16	* *	5	* *	7	* *
Norton	74	* *	* *	* *	59	* *	14	* *	32	* *
Osage	205	91.2 (3)	1.5 (23)		183	51.9 (8)	51	* *	61	* *
Osborne	50	* *	* *	* *	41	* *	6	* *	23	* *
Ottawa	68	* *	* *	* *	56	* *	11	* *	29	* *
Pawnee	131	84.0 (28)	1.5 (23)		126	63.5 (2)	28	* *	62	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6A
2006 - 2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Behavioral Indicators by Clinic
Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2, 3)										
Clinic	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Phillips	92	* *	* *	* *	74	* *	20	* *	58	* *
Pottawat-S	52	* *	* *	* *	42	* *	4	* *	20	* *
Pottawat-W	52	* *	* *	* *	58	* *	15	* *	32	* *
Pratt	202	70.3 (63)	6.4 (61)		178	47.8 (18)	41	* *	104	85.6 (15)
Rawlins	16	* *	* *	* *	19	* *	6	* *	12	* *
Reno	1,303	85.1 (21)	1.5 (23)		1,123	42.0 (38)	291	23.7 (1)	678	86.9 (6)
Republic	86	* *	* *	* *	73	* *	13	* *	18	* *
Rice	188	76.6 (51)	1.6 (28)		171	40.9 (43)	42	* *	61	* *
Riley-Ft	646	87.9 (12)	0.3 (2)		766	35.0 (58)	253	11.5 (14)	414	72.5 (47)
Riley-Manh	918	77.3 (47)	1.1 (17)		877	45.5 (25)	219	18.3 (3)	481	83.8 (23)
Rooks	112	84.8 (25)	2.7 (40)		97	* *	17	* *	59	* *
Rush	41	* *	* *	* *	41	* *	8	* *	22	* *
Russell	109	89.0 (7)	0.9 (12)		95	* *	23	* *	57	* *
Saline	1,190	76.1 (54)	3.2 (44)		1,050	34.4 (60)	271	5.9 (22)	612	79.7 (34)
Scott	128	79.7 (41)	5.5 (58)		113	39.8 (48)	18	* *	66	* *
Sedgwick-C	2,895	76.6 (51)	3.5 (47)		2,604	32.3 (62)	615	15.6 (5)	1,494	79.5 (35)
Sedgwick-E	536	77.2 (48)	1.5 (23)		405	32.1 (63)	0	* *	311	100.0 (1)
Sedgwick-M	4,901	74.0 (59)	2.8 (41)		3,980	26.3 (67)	1,082	10.9 (16)	1,895	72.8 (46)
Sedgwick-S	2,986	81.0 (35)	1.6 (28)		2,649	28.9 (66)	609	10.0 (17)	1,399	77.7 (40)
Seneca	32	* *	* *	* *	26	* *	3	* *	9	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6A
2006 - 2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Behavioral Indicators by Clinic
Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2, 3)										
Clinic	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Prenatal (4)		During Pregnancy (5)	
	Number	% (Rank)	% (Rank)		Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Seward	409	69.9 (64)	0.7 (5)		350	34.9 (59)	0	* *	268	100.0 (1)
Seward	878	68.0 (66)	7.2 (66)		801	38.5 (51)	267	6.0 (21)	328	74.1 (45)
ShawneeM&I	516	61.4 (70)	2.5 (37)		605	62.6 (3)	102	0.0 (26)	270	84.1 (21)
ShawneeMa	3,217	78.0 (45)	3.1 (43)		2,794	45.1 (27)	683	5.7 (24)	1,421	81.1 (29)
Sheridan	23	* *	* *		21	* *	2	* *	14	* *
Sherman	174	85.1 (21)	4.0 (51)		176	54.5 (7)	39	* *	106	88.7 (5)
Smith	39	* *	* *		51	* *	9	* *	31	* *
Stafford	103	79.6 (42)	4.9 (54)		101	64.4 (1)	25	* *	61	* *
Stanton	54	* *	* *		48	* *	11	* *	25	* *
Stevens	35	* *	* *		16	* *	0	* *	5	* *
Stevens	153	77.8 (46)	2.6 (39)		130	39.2 (49)	38	* *	47	* *
Sumner	348	86.2 (17)	2.3 (36)		320	40.6 (44)	84	* *	167	86.2 (13)
Thomas	132	83.3 (29)	0.8 (9)		119	49.6 (12)	23	* *	87	* *
Trego	46	* *	* *		37	* *	6	* *	17	* *
Wabaunsee	51	* *	* *		33	* *	10	* *	19	* *
Wallace	8	* *	* *		5	* *	1	* *	4	* *
Washington	75	* *	* *		63	* *	15	* *	35	* *
Wichita	54	* *	* *		46	* *	6	* *	30	* *
Wilson	271	88.6 (9)	0.7 (5)		252	41.7 (39)	50	* *	139	79.1 (37)
Woodson	24	* *	* *		21	* *	0	* *	11	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6A
2006 - 2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Behavioral Indicators by Clinic
Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2, 3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Pregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Wyandotte	5,883	69.3 (65)	4.2 (52)	5,137	35.7 (56)	1,171	12.3 (10)	2,640	83.3 (25)
Kansas	54,561	76.9	3.2	48,744	39.3	11,645	12.0	26,186	81.6

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2006 - 2008 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by Clinic
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Allen	294	45.2 (48)	263	32.7 (56)	96	* *	247	36.0 (58)	261	32.6 (57)
Anderson	153	57.5 (68)	140	33.6 (57)	59	* *	119	33.6 (55)	141	20.6 (36)
Atchison	321	44.9 (46)	235	21.3 (24)	74	* *	290	32.8 (52)	239	28.5 (53)
Barber-Kio	7	* *	6	* *	3	* *	6	* *	6	* *
Barber-Med	95	* *	85	* *	30	* *	75	* *	86	* *
Barton	826	39.8 (29)	745	24.4 (32)	229	29.7 (25)	662	18.4 (23)	747	7.4 (9)
Bourbon	359	47.6 (56)	322	28.0 (47)	111	26.1 (31)	289	28.4 (46)	323	30.7 (56)
Brown	270	42.2 (39)	218	25.7 (36)	65	* *	221	41.6 (61)	220	38.6 (61)
Butler	762	53.8 (63)	646	33.6 (57)	248	30.6 (24)	539	19.5 (25)	645	9.5 (15)
Chase	39	* *	35	* *	18	* *	30	* *	35	* *
Chautauqua	84	* *	61	* *	18	* *	70	* *	61	* *
Cherokee	623	41.7 (38)	519	25.6 (34)	133	21.8 (34)	450	32.9 (54)	537	27.6 (52)
Cheyenne	5	* *	5	* *	0	* *	4	* *	5	* *
Clark	8	* *	5	* *	0	* *	9	* *	5	* *
Clay	202	52.0 (61)	179	30.2 (51)	74	* *	149	30.2 (49)	179	25.1 (49)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2006 - 2008 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by Clinic
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Cloud	229	41.0 (33)	195	21.5 (26)	60	* *	179	6.7 (1)	194	2.6 (1)
Coffey	74	* *	72	* *	24	* *	51	* *	73	* *
Comanche	12	* *	11	* *	4	* *	10	* *	11	* *
Cowley-Ark	519	41.6 (37)	448	28.1 (48)	151	23.2 (32)	423	24.6 (36)	453	21.4 (39)
Cowley-Win	457	53.4 (62)	409	34.7 (61)	174	27.6 (28)	357	17.9 (21)	410	13.2 (20)
Crawford-G	11	* *	11	* *	7	* *	11	* *	11	* *
Crawford-P	1,145	44.6 (45)	952	26.5 (38)	330	33.9 (16)	913	38.4 (59)	952	34.0 (59)
Decatur	31	* *	27	* *	8	* *	28	* *	27	* *
DickinsonA	283	43.8 (43)	254	25.6 (34)	85	* *	211	13.3 (11)	257	7.4 (9)
DickinsonH	45	* *	39	* *	22	* *	37	* *	40	* *
Doniphan	99	* *	68	* *	29	* *	79	* *	70	* *
Douglas	1,390	44.0 (44)	1,151	20.7 (23)	362	44.2 (6)	1,096	12.6 (10)	1,157	7.1 (7)
Edwards	76	* *	64	* *	21	* *	61	* *	65	* *
Elk	56	* *	52	* *	13	* *	37	* *	52	* *
Ellis	431	37.8 (24)	377	14.6 (16)	122	53.3 (1)	344	12.5 (9)	375	8.8 (14)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2006 - 2008 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by Clinic
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Ellsworth	85	* *	77	* *	30	* *	61	* *	79	* *
Finney	2,041	15.7 (6)	1,866	7.1 (5)	227	48.9 (2)	1,676	19.7 (26)	1,871	17.7 (33)
Ford	1,808	10.7 (2)	1,519	4.9 (3)	119	35.3 (11)	1,544	9.2 (4)	1,528	7.3 (8)
Franklin	499	46.7 (52)	434	27.2 (44)	128	35.2 (12)	335	20.9 (31)	447	14.1 (23)
Ft. Riley	391	31.7 (16)	374	12.3 (11)	79	* *	314	31.2 (51)	386	30.1 (54)
Geary	914	43.5 (42)	752	21.3 (24)	268	38.8 (8)	756	39.9 (60)	755	41.1 (62)
Gove	34	* *	31	* *	9	* *	23	* *	31	* *
Graham	49	* *	44	* *	21	* *	40	* *	44	* *
Grant	243	13.2 (4)	235	5.5 (4)	22	* *	194	17.5 (20)	235	14.0 (22)
Gray	167	13.8 (5)	146	8.9 (6)	16	* *	137	15.3 (15)	148	12.2 (19)
Greeley	23	* *	22	* *	4	* *	16	* *	22	* *
Greenwood	121	47.1 (55)	73	* *	30	* *	107	33.6 (55)	107	22.4 (43)
Hamilton	54	* *	47	* *	3	* *	40	* *	47	* *
Harper	141	40.4 (31)	105	21.9 (27)	32	* *	122	27.9 (42)	109	15.6 (26)
Harvey	759	35.2 (23)	660	23.5 (30)	188	27.1 (29)	621	6.8 (2)	663	4.8 (4)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2006 - 2008 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by Clinic
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Haskell	63	* *	61	* *	8	* *	54	* *	61	* *
Hodgeman	28	* *	24	* *	5	* *	20	* *	24	* *
Jackson	230	47.0 (54)	207	26.6 (41)	74	* *	183	27.9 (42)	212	19.3 (34)
Jefferson	127	54.3 (64)	87	* *	33	* *	102	44.1 (64)	96	* *
Jewell	41	* *	39	* *	8	* *	30	* *	39	* *
Johnson-M	2,491	27.3 (12)	2,184	11.2 (9)	414	47.1 (4)	1,880	21.9 (32)	2,195	20.9 (37)
Johnson-O	2,820	25.3 (10)	2,500	11.0 (8)	423	47.3 (3)	2,185	18.8 (24)	2,502	16.7 (30)
Kearny	53	* *	52	* *	9	* *	45	* *	52	* *
Kingman	139	32.4 (19)	121	12.4 (12)	19	* *	125	32.8 (52)	134	32.8 (58)
Kiowa	37	* *	28	* *	6	* *	33	* *	29	* *
Labette	513	40.9 (32)	447	24.6 (33)	134	20.9 (35)	437	30.2 (49)	467	26.1 (50)
Lane	29	* *	21	* *	3	* *	22	* *	22	* *
Leavenwort	1,028	48.7 (59)	867	27.8 (46)	283	28.6 (26)	738	42.4 (62)	878	42.6 (63)
Lincoln	62	* *	53	* *	21	* *	45	* *	53	* *
Linn	177	59.3 (70)	159	44.0 (64)	77	* *	144	20.8 (29)	161	14.9 (24)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2006 - 2008 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by Clinic
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Logan	27	* *	22	* *	7	* *	24	* *	22	* *
Lyon	1,070	30.7 (14)	974	15.7 (18)	210	36.7 (9)	808	16.5 (18)	977	13.4 (21)
Marion	148	33.8 (22)	134	20.1 (22)	33	* *	105	11.4 (7)	135	5.2 (5)
Marshall	170	58.8 (69)	135	43.0 (63)	61	* *	131	54.2 (66)	140	47.9 (65)
Mc Pherson	427	41.0 (33)	387	26.6 (41)	123	26.8 (30)	320	10.6 (6)	390	4.6 (3)
Meade	115	33.0 (20)	98	* *	23	* *	98	* *	99	* *
Miami	344	54.9 (65)	282	32.3 (54)	106	35.8 (10)	245	27.8 (41)	280	20.0 (35)
Mitchell	104	39.4 (27)	94	* *	31	* *	77	* *	95	* *
Montgome-C	451	40.1 (30)	392	26.5 (38)	121	19.0 (36)	380	33.9 (57)	402	26.6 (51)
Montgome-I	510	41.2 (35)	437	26.5 (38)	142	23.2 (32)	433	28.6 (47)	449	24.3 (47)
Morris	81	* *	70	* *	27	* *	63	* *	72	* *
Nemaha	74	* *	67	* *	21	* *	54	* *	66	* *
Neosho-Ch	396	45.7 (50)	341	25.8 (37)	125	35.2 (12)	330	27.9 (42)	342	21.6 (40)
Neosho-Er	45	* *	41	* *	11	* *	36	* *	41	* *
Ness	22	* *	20	* *	6	* *	16	* *	20	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2006 - 2008 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by Clinic
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Norton	80	* *	71	* *	24	* *	59	* *	70	* *
Osage	231	46.8 (53)	200	30.5 (52)	70	* *	180	28.3 (45)	200	25.0 (48)
Osborne	53	* *	50	* *	15	* *	40	* *	50	* *
Ottawa	72	* *	63	* *	17	* *	55	* *	63	* *
Pawnee	146	48.6 (58)	128	29.7 (50)	47	* *	124	24.2 (35)	128	22.7 (44)
Phillips	83	* *	58	* *	22	* *	64	* *	58	* *
Pottawat-S	57	* *	55	* *	7	* *	41	* *	55	* *
Pottawat-W	68	* *	57	* *	22	* *	55	* *	58	* *
Pratt	223	38.1 (26)	126	32.5 (55)	51	* *	153	26.8 (40)	158	23.4 (45)
Rawlins	22	* *	20	* *	5	* *	19	* *	20	* *
Reno	1,421	46.1 (51)	1,261	23.9 (31)	443	34.5 (15)	1,109	22.5 (34)	1,267	17.4 (32)
Republic	89	* *	83	* *	30	* *	72	* *	84	* *
Rice	208	48.1 (57)	181	34.3 (60)	64	* *	173	30.1 (48)	182	22.0 (41)
Riley-Ft	992	25.5 (11)	687	14.7 (17)	116	31.0 (20)	740	24.6 (36)	776	24.2 (46)
Riley-Manh	1,126	33.0 (20)	888	18.9 (21)	193	35.2 (12)	851	8.8 (3)	940	6.1 (6)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2006 - 2008 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by Clinic
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Rooks	117	55.6 (66)	99	* *	47	* *	95	* *	105	30.5 (55)
Rush	45	* *	40	* *	19	* *	41	* *	40	* *
Russell	117	45.3 (49)	105	36.2 (62)	41	* *	95	* *	108	44.4 (64)
Saline	1,239	44.9 (46)	1,020	27.0 (43)	365	30.7 (23)	991	14.8 (13)	1,023	7.9 (11)
Scott	134	24.6 (9)	116	17.2 (19)	25	* *	111	26.1 (38)	118	22.0 (41)
Sedgwick-C	3,103	32.0 (18)	2,855	17.7 (20)	716	31.6 (19)	2,568	13.7 (12)	2,858	11.8 (18)
Sedgwick-E	620	31.8 (17)	533	12.2 (10)	76	* *	402	12.4 (8)	539	8.2 (12)
Sedgwick-M	5,216	29.3 (13)	4,711	13.0 (13)	932	44.5 (5)	3,902	9.4 (5)	4,724	8.7 (13)
Sedgwick-S	3,176	43.1 (41)	2,896	23.3 (29)	986	33.6 (17)	2,588	15.5 (16)	2,899	11.7 (17)
Seneca	32	* *	32	* *	15	* *	26	* *	32	* *
Seward	439	11.2 (3)	408	3.9 (2)	36	* *	341	20.8 (29)	406	17.0 (31)
Seward	965	8.3 (1)	859	3.7 (1)	53	* *	796	16.2 (17)	867	15.8 (27)
ShawneeM&I	607	31.0 (15)	471	13.2 (14)	109	42.2 (7)	594	18.0 (22)	467	3.2 (2)
ShawneeMa	3,420	38.0 (25)	2,874	22.8 (28)	774	27.8 (27)	2,672	17.2 (19)	2,873	11.3 (16)
Sheridan	25	* *	18	* *	3	* *	20	* *	18	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2006 - 2008 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by Clinic
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Sherman	221	43.0 (40)	187	28.3 (49)	60	* *	167	20.4 (28)	186	15.1 (25)
Smith	56	* *	40	* *	13	* *	49	* *	42	* *
Stafford	119	41.2 (35)	97	* *	34	* *	100	15.0 (14)	99	* *
Stanton	57	* *	52	* *	3	* *	47	* *	53	* *
Stevens	35	* *	35	* *	6	* *	17	* *	35	* *
Stevens	168	20.8 (7)	145	9.0 (7)	23	* *	128	20.3 (27)	146	15.8 (27)
Sumner	388	51.8 (60)	297	31.3 (53)	116	31.0 (20)	306	47.7 (65)	299	49.5 (66)
Thomas	152	39.5 (28)	135	27.4 (45)	37	* *	117	26.5 (39)	136	16.2 (29)
Trego	46	* *	35	* *	14	* *	37	* *	37	* *
Wabaunsee	54	* *	51	* *	14	* *	32	* *	51	* *
Wallace	6	* *	6	* *	3	* *	5	* *	6	* *
Washington	81	* *	74	* *	18	* *	63	* *	73	* *
Wichita	57	* *	52	* *	7	* *	47	* *	52	* *
Wilson	289	57.1 (67)	261	33.7 (59)	129	31.0 (20)	246	42.7 (63)	265	38.5 (60)
Woodson	27	* *	21	* *	8	* *	21	* *	22	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2006 - 2008 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by Clinic
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Wyandotte	6,269	24.2 (8)	5,632	13.3 (15)	1,006	32.4 (18)	5,036	22.4 (33)	5,688	21.0 (38)
Kansas	60,208	34.1	52,408	18.6	13,124	34.0	47,541	20.5	52,940	17.1

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2006 - 2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Infant Health Indicators by Clinic
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)									
Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)	High (5)							
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Allen	273	7.3 (31)	4.4 (11)	232	12.9 (46)	200	3.5 (35)	275	50.5 (64)
Anderson	149	6.7 (23)	4.7 (15)	115	11.3 (33)	102	5.9 (55)	144	50.7 (63)
Atchison	342	7.6 (36)	5.6 (23)	294	9.5 (15)	261	3.1 (29)	337	49.9 (66)
Barber-Kio	7	* * *	* *	6	* *	6	* *	7	* *
Barber-Med	82	* * *	* *	66	* *	51	* *	82	* *
Barton	735	7.6 (36)	4.8 (16)	596	12.2 (40)	519	3.5 (35)	710	72.3 (25)
Bourbon	335	6.9 (24)	6.0 (28)	275	12.4 (41)	236	3.4 (33)	332	58.1 (54)
Brown	167	7.2 (28)	7.2 (46)	203	10.3 (22)	140	2.9 (25)	214	53.3 (61)
Butler	721	9.2 (55)	5.8 (25)	504	14.9 (60)	426	3.5 (35)	692	73.4 (21)
Chase	35	* * *	* *	27	* *	25	* *	35	* *
Chautauqua	63	* * *	* *	59	* *	45	* *	63	* *
Cherokee	586	7.7 (39)	7.3 (47)	400	13.0 (49)	347	2.3 (16)	561	57.9 (55)
Cheyenne	5	* * *	* *	4	* *	4	* *	5	* *
Clark	7	* * *	* *	7	* *	5	* *	8	* *

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
- (4) <2500 g.
- (5) >4000 g.
- (6) <37 weeks gestation.
- (7) 37 or more weeks gestation and <2500 g.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2006 - 2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Infant Health Indicators by Clinic
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)									
Clinic	Birthweight			Preterm ⁽⁶⁾		Full Term LBW ⁽⁷⁾		Ever Breastfed	
	Low ⁽⁴⁾	High ⁽⁵⁾		Number	%	Number	%	Number	%
	Number	% ^(Rank)	% ^(Rank)						
Clay	186	7.5 (33)	3.8 (7)	136	14.0 (56)	117	3.4 (33)	182	63.2 (43)
Cloud	200	8.0 (44)	3.5 (4)	154	16.2 (62)	128	3.1 (29)	193	67.4 (36)
Coffey	72	* * *	* *	51	* *	47	* *	73	* *
Comanche	12	* * *	* *	8	* *	7	* *	12	* *
Cowley-Ark	482	7.3 (31)	8.3 (56)	385	9.4 (13)	341	3.8 (42)	462	50.4 (65)
Cowley-Win	412	8.3 (47)	6.6 (33)	325	12.9 (46)	281	5.3 (54)	407	62.2 (46)
Crawford-G	10	* * *	* *	9	* *	8	* *	10	* *
Crawford-P	992	9.3 (58)	4.9 (17)	794	12.6 (43)	687	4.4 (47)	999	59.8 (52)
Decatur	27	* * *	* *	26	* *	26	* *	26	* *
DickinsonA	257	6.2 (13)	7.8 (53)	188	9.0 (9)	169	2.4 (20)	253	75.5 (14)
DickinsonH	40	* * *	* *	35	* *	32	* *	41	* *
Doniphan	89	* * *	* *	82	* *	57	* *	101	46.5 (67)
Douglas	1,200	6.5 (19)	6.9 (40)	918	11.3 (33)	802	2.6 (22)	1,145	77.6 (8)
Edwards	61	* * *	* *	51	* *	42	* *	62	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2006 - 2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Infant Health Indicators by Clinic
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)										
Clinic	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)							
	Number	% (Rank)	%	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Elk	48	* * * *	*	*	36	* *	28	* *	50	* *
Ellis	380	5.8 (8)	4.5 (12)	4.5 (12)	306	8.5 (6)	274	2.2 (13)	375	77.9 (7)
Ellsworth	79	* * * *	*	*	56	* *	47	* *	78	* *
Finney	1,867	6.2 (13)	6.3 (30)	6.3 (30)	1,535	10.7 (25)	1,357	2.4 (20)	1,416	74.4 (20)
Ford	1,629	5.5 (6)	5.8 (25)	5.8 (25)	1,398	10.3 (22)	1,232	2.2 (13)	1,423	72.7 (22)
Franklin	435	6.9 (24)	5.3 (20)	5.3 (20)	287	13.2 (50)	246	2.8 (24)	419	53.9 (60)
Ft. Riley	398	5.3 (4)	12.6 (63)	12.6 (63)	260	11.2 (31)	230	3.9 (45)	393	74.6 (18)
Geary	797	6.6 (22)	7.0 (44)	7.0 (44)	628	8.3 (5)	571	2.3 (16)	734	68.5 (35)
Gove	18	* * * *	*	*	21	* *	9	* *	30	* *
Graham	40	* * * *	*	*	34	* *	30	* *	43	* *
Grant	228	9.2 (55)	7.5 (50)	7.5 (50)	186	11.3 (33)	161	3.7 (39)	183	75.4 (15)
Gray	121	5.8 (8)	6.6 (33)	6.6 (33)	116	12.1 (38)	90	* *	136	83.8 (1)
Greeley	15	* * * *	*	*	15	* *	11	* *	21	* *
Greenwood	115	8.7 (50)	3.5 (4)	3.5 (4)	99	* *	83	* *	109	62.4 (45)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2006 - 2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Infant Health Indicators by Clinic
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)										
Clinic	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)		Number	%	Number	%	Number	%
	Number	% (Rank)	% (Rank)	(Rank)						
Hamilton	44	* * * *	* * * *	34	* *	32	* *	32	* *	
Harper	92	* * * *	* * * *	106	11.3 (33)	67	* *	129	76.0 (12)	
Harvey	592	9.1 (54)	5.9 (27)	572	13.6 (53)	419	3.6 (38)	681	78.3 (5)	
Haskell	62	* * * *	* * * *	54	* *	51	* *	44	* *	
Hodgeman	29	* * * *	* * * *	19	* *	18	* *	25	* *	
Jackson	195	8.2 (46)	8.2 (55)	173	14.5 (59)	131	2.3 (16)	214	76.2 (11)	
Jefferson	169	6.5 (19)	8.3 (56)	121	13.2 (50)	104	3.8 (42)	172	57.0 (57)	
Jewell	40	* * * *	* * * *	29	* *	25	* *	36	* *	
Johnson-M	2,160	7.6 (36)	7.4 (49)	1,699	9.8 (19)	1,466	3.3 (32)	2,167	78.2 (6)	
Johnson-O	2,486	5.3 (4)	8.1 (54)	1,976	8.2 (4)	1,767	1.9 (6)	2,373	80.5 (3)	
Kearny	51	* * * *	* * * *	44	* *	36	* *	36	* *	
Kingman	93	* * * *	* * * *	107	15.9 (61)	67	* *	126	75.4 (15)	
Kiowa	31	* * * *	* * * *	30	* *	25	* *	32	* *	
Labette	474	7.8 (41)	7.0 (44)	400	10.8 (26)	355	3.7 (39)	456	54.6 (59)	

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
- (4) <2500 g.
- (5) >4000 g.
- (6) <37 weeks gestation.
- (7) 37 or more weeks gestation and <2500 g.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2006 - 2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Infant Health Indicators by Clinic
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)									
Clinic	Birthweight			Preterm ⁽⁶⁾		Full Term LBW ⁽⁷⁾		Ever Breastfed	
	Low ⁽⁴⁾	High ⁽⁵⁾							
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Lane	30	* * * *		25	* *	20	* *	29	* *
Leavenwort	865	8.3 (47)	4.5 (12)	625	9.3 (11)	553	2.9 (25)	893	61.4 (49)
Lincoln	54	* * * *		43	* *	34	* *	59	* *
Linn	160	6.9 (24)	5.0 (18)	132	13.6 (53)	113	4.4 (47)	156	60.3 (50)
Logan	22	* * * *		20	* *	19	* *	21	* *
Lyon	983	6.2 (13)	6.7 (37)	734	9.9 (20)	660	2.1 (11)	941	72.5 (24)
Marion	133	7.5 (33)	3.0 (1)	93	* *	81	* *	131	82.4 (2)
Marshall	129	11.6 (62)	3.9 (8)	105	10.5 (24)	87	* *	140	57.9 (55)
Mc Pherson	388	6.4 (16)	6.7 (37)	288	8.7 (8)	261	2.3 (16)	383	77.3 (9)
Meade	99	* * * *		88	* *	71	* *	94	* *
Miami	352	6.5 (19)	5.7 (24)	233	12.4 (41)	202	4.5 (49)	305	67.2 (37)
Mitchell	88	* * * *		66	* *	61	* *	85	* *
Montgome-C	417	9.4 (59)	3.1 (2)	347	14.1 (57)	294	5.1 (53)	402	55.2 (58)
Montgome-I	473	7.2 (28)	3.4 (3)	392	11.0 (28)	347	2.0 (8)	452	60.0 (51)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2006 - 2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Infant Health Indicators by Clinic
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)									
Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
	Number	% (Rank)	% (Rank)						
Morris	66	* * * *	*	57	* *	40	* *	70	* *
Nemaha	57	* * * *	*	49	* *	33	* *	69	* *
Neosho-Ch	297	8.4 (49)	7.7 (52)	306	13.7 (55)	212	3.8 (42)	376	50.8 (62)
Neosho-Er	39	* * * *	*	35	* *	28	* *	43	* *
Ness	21	* * * *	*	14	* *	13	* *	21	* *
Norton	72	* * * *	*	53	* *	44	* *	68	* *
Osage	199	5.5 (6)	6.0 (28)	170	12.9 (46)	135	1.5 (3)	200	59.0 (53)
Osborne	39	* * * *	*	38	* *	25	* *	50	* *
Ottawa	56	* * * *	*	49	* *	37	* *	61	* *
Pawnee	125	8.8 (51)	4.0 (10)	111	14.4 (58)	95	* *	122	68.9 (34)
Phillips	88	* * * *	*	63	* *	53	* *	87	* *
Pottawat-S	52	* * * *	*	39	* *	37	* *	53	* *
Pottawat-W	63	* * * *	*	51	* *	46	* *	60	* *
Pratt	195	9.2 (55)	8.7 (60)	149	9.4 (13)	131	4.6 (50)	180	71.1 (27)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2006 - 2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Infant Health Indicators by Clinic
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)									
Clinic	Birthweight			Preterm ⁽⁶⁾		Full Term LBW ⁽⁷⁾		Ever Breastfed	
	Low ⁽⁴⁾	High ⁽⁵⁾		Number	%	Number	%	Number	%
	Number	% (Rank)	% (Rank)						
Rawlins	21	* * * *		17	* *	15	* *	20	* *
Reno	1,271	6.4 (16)	8.6 (58)	994	11.0 (28)	872	2.1 (11)	1,239	63.5 (42)
Republic	58	* * * *		66	* *	38	* *	83	* *
Rice	186	7.5 (33)	6.5 (31)	151	12.6 (43)	131	4.6 (50)	180	65.6 (38)
Riley-Ft	811	5.1 (3)	8.6 (58)	560	7.3 (3)	516	1.4 (2)	786	76.8 (10)
Riley-Manh	1,008	5.8 (8)	6.9 (40)	727	11.0 (28)	644	2.0 (8)	973	78.6 (4)
Rooks	101	5.9 (11)	5.0 (18)	90	* *	77	* *	104	70.2 (29)
Rush	40	* * * *		35	* *	32	* *	39	* *
Russell	103	9.7 (61)	3.9 (8)	86	* *	73	* *	98	* *
Saline	1,211	9.4 (59)	5.5 (22)	949	11.9 (37)	830	4.0 (46)	1,206	61.8 (48)
Scott	98	* * * *		101	6.9 (2)	81	* *	117	70.1 (31)
Sedgwick-C	2,876	7.2 (28)	6.6 (33)	2,394	9.5 (15)	2,148	2.9 (25)	2,760	69.1 (32)
Sedgwick-E	528	8.1 (45)	6.6 (33)	322	11.2 (31)	283	1.8 (4)	531	69.1 (32)
Sedgwick-M	4,831	8.9 (52)	6.5 (31)	3,605	9.1 (10)	3,247	3.7 (39)	4,171	64.7 (39)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2006 - 2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Infant Health Indicators by Clinic
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)									
Clinic	Birthweight			Preterm ⁽⁶⁾		Full Term LBW ⁽⁷⁾		Ever Breastfed	
	Low ⁽⁴⁾		High ⁽⁵⁾						
	Number	% ^(Rank)	% ^(Rank)	Number	% ^(Rank)	Number	% ^(Rank)	Number	% ^(Rank)
Sedgwick-S	2,976	7.0 (27)	7.6 (51)	2,444	9.3 (11)	2,201	2.6 (22)	2,812	63.8 (41)
Seneca	24	* *	* *	26	* *	16	* *	32	* *
Seward	389	6.4 (16)	9.3 (62)	321	10.0 (21)	277	1.8 (4)	411	70.3 (28)
Seward	864	4.9 (2)	6.7 (37)	719	9.5 (15)	637	1.9 (6)	872	72.6 (23)
ShawneeM&I	492	7.9 (42)	7.3 (47)	484	16.7 (63)	394	2.0 (8)	484	72.3 (25)
ShawneeMa	3,118	7.7 (39)	5.4 (21)	2,519	12.1 (38)	2,123	3.1 (29)	3,180	64.4 (40)
Sheridan	24	* *	* *	17	* *	15	* *	23	* *
Sherman	200	15.0 (63)	3.5 (4)	149	13.4 (52)	129	10.1 (56)	192	74.5 (19)
Smith	53	* *	* *	43	* *	37	* *	53	* *
Stafford	94	* *	* *	83	* *	75	* *	89	* *
Stanton	52	* *	* *	45	* *	40	* *	35	* *
Stevens	0	* *	* *	15	* *	0	* *	31	* *
Stevens	99	* *	* *	114	9.6 (18)	64	* *	148	75.7 (13)
Sumner	334	6.0 (12)	6.9 (40)	280	8.6 (7)	224	2.2 (13)	350	62.9 (44)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2006 - 2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Infant Health Indicators by Clinic
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)										
Clinic	Birthweight				Preterm ⁽⁶⁾		Full Term LBW ⁽⁷⁾		Ever Breastfed	
	Low ⁽⁴⁾		High ⁽⁵⁾		Number	%	Number	%	Number	%
	Number	% (Rank)	Number	% (Rank)						
Thomas	146	3.4 (1)	8.9 (61)		109	6.4 (1)	102	1.0 (1)	140	75.0 (17)
Trego	35	* * *	* *		34	* *	24	* *	41	* *
Wabaunsee	51	* * *	* *		31	* *	28	* *	47	* *
Wallace	8	* * *	* *		5	* *	5	* *	8	* *
Washington	45	* * *	* *		58	* *	37	* *	72	* *
Wichita	37	* * *	* *		43	* *	22	* *	53	* *
Wilson	267	9.0 (53)	4.5 (12)		234	12.8 (45)	201	5.0 (52)	244	61.9 (47)
Woodson	23	* * *	* *		18	* *	14	* *	22	* *
Wyandotte	5,745	7.9 (42)	6.9 (40)		4,637	10.9 (27)	4,080	3.0 (28)	5,600	70.2 (29)
Kansas	54,016	7.3	6.6		43,191	10.7	37,507	2.9	52,161	68.6

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

