

Table 3A
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Allen	91	*	*	*	*	*	*	*
Anderson	50	*	*	*	*	*	*	*
Atchison	155	88.4	7.1	1.3	1.9	0.0	1.3	0.0
Barber-Med	46	*	*	*	*	*	*	*
Barton	257	65.0	1.6	31.1	0.4	0.8	1.2	0.0
Bourbon	117	92.3	1.7	3.4	0.9	0.0	1.7	0.0
Brown	99	*	*	*	*	*	*	*
Butler	282	88.3	1.1	7.1	0.7	0.7	2.1	0.0
Chase	16	*	*	*	*	*	*	*
Chautauqua	32	*	*	*	*	*	*	*
Cherokee	212	88.7	0.9	3.8	1.9	3.8	0.9	0.0
Cheyenne	3	*	*	*	*	*	*	*
Clark	4	*	*	*	*	*	*	*
Clay	59	*	*	*	*	*	*	*
Cloud	89	*	*	*	*	*	*	*
Coffey	25	*	*	*	*	*	*	*
Comanche	11	*	*	*	*	*	*	*
Cowley-Ark	204	62.7	3.4	31.9	1.0	0.0	1.0	0.0
Cowley-Win	159	86.2	1.9	6.9	0.6	3.1	1.3	0.0
Crawford-G	1	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3A
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Crawford-P	400	84.3	2.8	10.0	0.0	1.5	1.5	0.0
Decatur	10	*	*	*	*	*	*	*
DickinsonA	122	91.8	0.0	5.7	0.8	0.8	0.8	0.0
DickinsonH	12	*	*	*	*	*	*	*
Doniphan	37	*	*	*	*	*	*	*
Douglas	507	63.1	10.5	11.6	8.1	5.3	1.4	0.0
Edwards	25	*	*	*	*	*	*	*
Elk	20	*	*	*	*	*	*	*
Ellis	180	84.4	0.0	13.9	1.1	0.6	0.0	0.0
Ellsworth	33	*	*	*	*	*	*	*
Finney	718	30.8	0.6	65.3	0.1	2.9	0.3	0.0
Ford	686	21.6	1.0	76.7	0.1	0.6	0.0	0.0
Franklin	187	90.4	0.5	7.0	0.5	0.0	1.6	0.0
Geary	393	49.9	24.9	15.5	0.8	3.6	5.3	0.0
Gove	12	*	*	*	*	*	*	*
Graham	17	*	*	*	*	*	*	*
Grant	71	*	*	*	*	*	*	*
Gray	45	*	*	*	*	*	*	*
Greeley	15	*	*	*	*	*	*	*
Greenwood	49	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3A
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaska Native	Asian/Pacific Islander	Multiple Races	All Other/Unknown
		%	%	%	%	%	%	%
Hamilton	31	*	*	*	*	*	*	*
Harper	55	*	*	*	*	*	*	*
Harvey	284	71.5	2.5	22.2	1.8	0.7	1.4	0.0
Haskell	20	*	*	*	*	*	*	*
Hodgeman	16	*	*	*	*	*	*	*
Jackson	75	*	*	*	*	*	*	*
Jefferson	68	*	*	*	*	*	*	*
Jewell	9	*	*	*	*	*	*	*
Johnson-M	850	47.9	14.9	31.2	0.7	3.2	2.1	0.0
Johnson-O	1,066	43.6	12.0	39.5	0.5	3.6	0.8	0.0
Kearny	19	*	*	*	*	*	*	*
Kingman	59	*	*	*	*	*	*	*
Kiowa	16	*	*	*	*	*	*	*
Labette	194	86.6	7.7	1.5	2.1	0.5	1.5	0.0
Lane	13	*	*	*	*	*	*	*
Leavenworth	359	71.6	13.1	10.6	0.3	0.8	3.6	0.0
Lincoln	21	*	*	*	*	*	*	*
Linn	66	*	*	*	*	*	*	*
Logan	15	*	*	*	*	*	*	*
Lyon	323	54.8	4.3	36.5	0.6	2.5	1.2	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3A
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Marion	47	*	*	*	*	*	*	*
Marshall	48	*	*	*	*	*	*	*
Mc Pherson	144	89.6	1.4	6.9	0.7	0.0	1.4	0.0
Meade	44	*	*	*	*	*	*	*
Miami	146	92.5	3.4	2.1	0.0	0.7	1.4	0.0
Mitchell	54	*	*	*	*	*	*	*
Montgome-C	157	70.1	10.2	12.1	5.7	0.0	1.9	0.0
Montgome-I	187	82.4	5.9	9.1	2.1	0.5	0.0	0.0
Morris	23	*	*	*	*	*	*	*
Nemaha	34	*	*	*	*	*	*	*
Neosho-Ch	135	91.1	3.7	3.7	0.0	0.0	1.5	0.0
Neosho-Er	11	*	*	*	*	*	*	*
Ness	5	*	*	*	*	*	*	*
Norton	32	*	*	*	*	*	*	*
Osage	95	*	*	*	*	*	*	*
Osborne	14	*	*	*	*	*	*	*
Ottawa	28	*	*	*	*	*	*	*
Pawnee	62	*	*	*	*	*	*	*
Phillips	36	*	*	*	*	*	*	*
Pottawat-S	18	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3A
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Pottawat-W	20	*	*	*	*	*	*	*
Pratt	85	*	*	*	*	*	*	*
Rawlins	12	*	*	*	*	*	*	*
Reno	499	80.4	2.6	15.4	0.2	0.4	1.0	0.0
Republic	43	*	*	*	*	*	*	*
Rice	75	*	*	*	*	*	*	*
Riley-Ft	541	72.8	10.9	13.1	0.4	2.8	0.0	0.0
Riley-Manh	412	75.5	10.4	9.2	0.2	3.4	1.2	0.0
Rooks	50	*	*	*	*	*	*	*
Rush	9	*	*	*	*	*	*	*
Russell	42	*	*	*	*	*	*	*
Saline	531	68.4	4.9	21.3	0.4	2.6	2.4	0.0
Scott	51	*	*	*	*	*	*	*
Sedgwick-C	1,230	40.7	13.3	34.6	1.2	8.1	2.2	0.0
Sedgwick-M	2,059	32.3	27.6	34.2	1.7	2.7	1.4	0.0
Sedgwick-S	1,432	65.6	8.3	19.4	2.3	2.2	2.2	0.0
Seneca	9	*	*	*	*	*	*	*
Seward	532	15.0	1.7	82.1	0.6	0.4	0.2	0.0
ShawneeM&I	262	37.4	14.5	44.7	1.5	1.1	0.8	0.0
ShawneeMa	1,248	59.9	15.9	19.9	2.2	0.6	1.4	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3A
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaska Native	Asian/Pacific Islander	Multiple Races	All Other/Unknown
		%	%	%	%	%	%	%
Sheridan	10	*	*	*	*	*	*	*
Sherman	60	*	*	*	*	*	*	*
Smith	20	*	*	*	*	*	*	*
Stafford	43	*	*	*	*	*	*	*
Stanton	14	*	*	*	*	*	*	*
Stevens	89	*	*	*	*	*	*	*
Sumner	157	88.5	1.3	4.5	1.9	0.0	3.8	0.0
Thomas	60	*	*	*	*	*	*	*
Trego	18	*	*	*	*	*	*	*
Wabaunsee	25	*	*	*	*	*	*	*
Washington	27	*	*	*	*	*	*	*
Wichita	18	*	*	*	*	*	*	*
Wilson	101	95.0	1.0	3.0	1.0	0.0	0.0	0.0
Woodson	8	*	*	*	*	*	*	*
Wyandotte	2,286	26.2	29.9	40.3	0.3	2.8	0.4	0.0
Unknown Clinics	3	*	*	*	*	*	*	*
Kansas	22,411	56.4	11.3	27.6	1.3	2.2	1.3	0.0
Nation (Prior Year)	1,307,887	42.0	24.2	28.1	1.0	3.1	1.1	0.5

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Age Distribution by Clinic

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Allen	91	*	*	*	*	*	*	*
Anderson	50	*	*	*	*	*	*	*
Atchison	155	0.0	7.1	16.8	62.6	13.5	0.0	0.0
Barber-Med	46	*	*	*	*	*	*	*
Barton	257	0.0	8.6	14.4	61.9	14.0	1.2	0.0
Bourbon	117	0.0	7.7	18.8	59.0	13.7	0.9	0.0
Brown	99	*	*	*	*	*	*	*
Butler	282	0.0	7.8	19.9	59.9	11.7	0.4	0.4
Chase	16	*	*	*	*	*	*	*
Chautauqua	32	*	*	*	*	*	*	*
Cherokee	212	0.0	8.0	19.8	59.9	10.8	0.9	0.5
Cheyenne	3	*	*	*	*	*	*	*
Clark	4	*	*	*	*	*	*	*
Clay	59	*	*	*	*	*	*	*
Cloud	89	*	*	*	*	*	*	*
Coffey	25	*	*	*	*	*	*	*
Comanche	11	*	*	*	*	*	*	*
Cowley-Ark	204	1.0	7.4	19.1	58.3	12.7	1.5	0.0
Cowley-Win	159	0.6	8.8	13.2	64.8	12.6	0.0	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Age Distribution by Clinic

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Crawford-G	1	*	*	*	*	*	*	*
Crawford-P	400	0.3	7.0	16.5	63.0	12.5	0.5	0.3
Decatur	10	*	*	*	*	*	*	*
DickinsonA	122	0.0	9.8	13.1	61.5	14.8	0.8	0.0
DickinsonH	12	*	*	*	*	*	*	*
Doniphan	37	*	*	*	*	*	*	*
Douglas	507	0.0	3.9	11.4	63.1	20.1	1.4	0.0
Edwards	25	*	*	*	*	*	*	*
Elk	20	*	*	*	*	*	*	*
Ellis	180	0.0	3.3	10.6	73.9	10.6	1.7	0.0
Ellsworth	33	*	*	*	*	*	*	*
Finney	718	0.6	9.1	15.7	54.0	18.8	1.8	0.0
Ford	686	0.4	7.9	14.0	53.9	21.7	1.9	0.1
Franklin	187	0.0	5.3	21.9	56.7	14.4	1.6	0.0
Geary	393	0.3	4.8	14.5	67.9	12.2	0.3	0.0
Gove	12	*	*	*	*	*	*	*
Graham	17	*	*	*	*	*	*	*
Grant	71	*	*	*	*	*	*	*
Gray	45	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Age Distribution by Clinic

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Greeley	15	*	*	*	*	*	*	*
Greenwood	49	*	*	*	*	*	*	*
Hamilton	31	*	*	*	*	*	*	*
Harper	55	*	*	*	*	*	*	*
Harvey	284	0.7	7.4	18.3	59.9	13.4	0.4	0.0
Haskell	20	*	*	*	*	*	*	*
Hodgeman	16	*	*	*	*	*	*	*
Jackson	75	*	*	*	*	*	*	*
Jefferson	68	*	*	*	*	*	*	*
Jewell	9	*	*	*	*	*	*	*
Johnson-M	850	0.5	3.9	11.2	58.9	23.8	1.8	0.0
Johnson-O	1,066	0.0	3.8	10.7	59.9	24.2	1.3	0.1
Kearny	19	*	*	*	*	*	*	*
Kingman	59	*	*	*	*	*	*	*
Kiowa	16	*	*	*	*	*	*	*
Labette	194	0.0	10.3	19.1	59.3	10.3	1.0	0.0
Lane	13	*	*	*	*	*	*	*
Leavenwort	359	0.0	6.7	17.3	62.7	12.8	0.6	0.0
Lincoln	21	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Age Distribution by Clinic

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Linn	66	*	*	*	*	*	*	*
Logan	15	*	*	*	*	*	*	*
Lyon	323	0.6	3.7	13.3	66.3	14.6	1.5	0.0
Marion	47	*	*	*	*	*	*	*
Marshall	48	*	*	*	*	*	*	*
Mc Pherson	144	0.0	4.9	9.7	68.8	16.7	0.0	0.0
Meade	44	*	*	*	*	*	*	*
Miami	146	0.7	4.8	17.1	63.0	13.7	0.7	0.0
Mitchell	54	*	*	*	*	*	*	*
Montgome-C	157	0.6	1.9	17.8	66.2	13.4	0.0	0.0
Montgome-I	187	0.0	2.1	15.5	69.5	12.3	0.0	0.5
Morris	23	*	*	*	*	*	*	*
Nemaha	34	*	*	*	*	*	*	*
Neosho-Ch	135	0.0	9.6	18.5	60.7	10.4	0.7	0.0
Neosho-Er	11	*	*	*	*	*	*	*
Ness	5	*	*	*	*	*	*	*
Norton	32	*	*	*	*	*	*	*
Osage	95	*	*	*	*	*	*	*
Osborne	14	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Age Distribution by Clinic

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Ottawa	28	*	*	*	*	*	*	*
Pawnee	62	*	*	*	*	*	*	*
Phillips	36	*	*	*	*	*	*	*
Pottawat-S	18	*	*	*	*	*	*	*
Pottawat-W	20	*	*	*	*	*	*	*
Pratt	85	*	*	*	*	*	*	*
Rawlins	12	*	*	*	*	*	*	*
Reno	499	0.0	7.0	17.0	62.5	13.0	0.4	0.0
Republic	43	*	*	*	*	*	*	*
Rice	75	*	*	*	*	*	*	*
Riley-Ft	541	0.0	0.7	10.7	78.0	10.2	0.4	0.0
Riley-Manh	412	0.0	3.4	14.6	69.7	12.1	0.2	0.0
Rooks	50	*	*	*	*	*	*	*
Rush	9	*	*	*	*	*	*	*
Russell	42	*	*	*	*	*	*	*
Saline	531	0.2	7.2	14.5	64.6	12.4	1.1	0.0
Scott	51	*	*	*	*	*	*	*
Sedgwick-C	1,230	0.2	7.6	12.6	58.5	19.8	1.1	0.1
Sedgwick-M	2,059	0.3	7.2	13.1	61.1	16.9	1.3	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Age Distribution by Clinic

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Sedgwick-S	1,432	0.4	5.9	15.4	61.7	15.4	1.3	0.0
Seneca	9	*	*	*	*	*	*	*
Seward	532	0.0	7.7	14.3	54.7	21.8	1.5	0.0
ShawneeM&I	262	0.8	6.5	17.6	55.0	19.1	1.1	0.0
ShawneeMa	1,248	0.2	6.8	15.5	61.4	14.7	1.3	0.0
Sheridan	10	*	*	*	*	*	*	*
Sherman	60	*	*	*	*	*	*	*
Smith	20	*	*	*	*	*	*	*
Stafford	43	*	*	*	*	*	*	*
Stanton	14	*	*	*	*	*	*	*
Stevens	89	*	*	*	*	*	*	*
Sumner	157	0.0	5.7	19.7	60.5	12.7	1.3	0.0
Thomas	60	*	*	*	*	*	*	*
Trego	18	*	*	*	*	*	*	*
Wabaunsee	25	*	*	*	*	*	*	*
Washington	27	*	*	*	*	*	*	*
Wichita	18	*	*	*	*	*	*	*
Wilson	101	0.0	6.9	19.8	61.4	11.9	0.0	0.0
Woodson	8	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Age Distribution by Clinic

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Wyandotte	2,286	0.4	7.6	13.7	57.9	18.9	1.4	0.0
Unknown Clinics	3	*	*	*	*	*	*	*
Kansas	22,411	0.3	6.3	14.6	61.1	16.6	1.1	0.1
Nation (Prior Year)	1,307,887	0.3	6.3	12.6	59.7	19.4	1.4	0.3

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2, 3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Allen	89	* *	* *	80	* *	* *	19	* *	24	* *	82	* *	82	* *
Anderson	49	* *	* *	43	* *	* *	8	* *	6	* *	45	* *	45	* *
Atchison	147	6.8 (31)	59.9 (37)	133	15.0 (2)	56.4 (34)	23	* *	29	* *	134	7.5 (27)	133	15.8(36)
Barber-Med	44	* *	* *	33	* *	* *	7	* *	10	* *	37	* *	37	* *
Barton	252	4.8 (21)	55.6 (26)	213	15.5 (5)	52.6 (24)	28	* *	40	* *	217	3.7 (5)	215	13.5(31)
Bourbon	114	3.5 (6)	58.8 (36)	99	* *	* *	17	* *	21	* *	101	5.9 (16)	101	11.9(25)
Brown	97	* *	* *	87	* *	* *	8	* *	26	* *	87	* *	87	* *
Butler	263	6.8 (31)	51.7 (9)	234	17.9 (10)	49.1 (12)	41	* *	55	* *	255	8.6 (34)	255	12.9(29)
Chase	15	* *	* *	13	* *	* *	1	* *	0	* *	14	* *	14	* *
Chautauqua	32	* *	* *	32	* *	* *	4	* *	4	* *	32	* *	32	* *
Cherokee	192	7.8 (36)	54.7 (23)	169	13.0 (1)	58.0 (36)	24	* *	45	* *	184	4.9 (10)	185	13.5(31)
Cheyenne	3	* *	* *	3	* *	* *	0	* *	1	* *	3	* *	3	* *
Clark	4	* *	* *	3	* *	* *	1	* *	0	* *	3	* *	3	* *
Clay	59	* *	* *	51	* *	* *	4	* *	16	* *	53	* *	53	* *
Cloud	89	* *	* *	80	* *	* *	13	* *	27	* *	78	* *	81	* *
Coffey	23	* *	* *	19	* *	* *	1	* *	9	* *	20	* *	20	* *
Comanche	11	* *	* *	11	* *	* *	4	* *	2	* *	11	* *	11	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2, 3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Cowley-Ark	198	4.0 (11)	54.0 (18)	172	21.5 (28)	53.5 (29)	31	* * *	19	* * *	175	8.0 (31)	175	12.6(28)
Cowley-Win	158	5.7 (29)	51.3 (8)	141	21.3 (27)	53.2 (27)	21	* * *	26	* * *	141	2.8 (3)	142	11.3(22)
Crawford-G	1	* * *	* * *	0	* * *	* * *	0	* * *	0	* * *	0	* * *	0	* * *
Crawford-P	388	7.2 (34)	53.1 (14)	336	19.3 (16)	56.3 (33)	54	* * *	54	* * *	348	7.2 (23)	348	13.8(33)
Decatur	10	* * *	* * *	9	* * *	* * *	1	* * *	1	* * *	9	* * *	9	* * *
DickinsonA	116	5.2 (24)	57.8 (34)	104	15.4 (4)	54.8 (31)	12	* * *	30	* * *	111	7.2 (23)	107	10.3(17)
DickinsonH	12	* * *	* * *	10	* * *	* * *	1	* * *	3	* * *	10	* * *	10	* * *
Doniphan	35	* * *	* * *	31	* * *	* * *	5	* * *	9	* * *	33	* * *	33	* * *
Douglas	491	3.7 (8)	48.1 (2)	421	16.9 (7)	55.1 (32)	70	* * *	137	21.2 (6)	432	6.7 (21)	432	9.0(10)
Edwards	25	* * *	* * *	25	* * *	* * *	3	* * *	4	* * *	25	* * *	25	* * *
Elk	19	* * *	* * *	18	* * *	* * *	1	* * *	4	* * *	19	* * *	19	* * *
Ellis	166	3.0 (5)	51.8 (10)	140	19.3 (16)	47.9 (11)	42	* * *	43	* * *	157	1.9 (1)	157	5.7(2)
Ellsworth	33	* * *	* * *	32	* * *	* * *	4	* * *	5	* * *	31	* * *	31	* * *
Finney	701	7.6 (35)	50.6 (6)	630	23.3 (31)	46.5 (10)	99	* * *	104	15.4 (2)	645	7.4 (26)	643	7.2(6)
Ford	677	2.5 (4)	55.5 (25)	604	22.8 (30)	45.5 (6)	80	* * *	61	* * *	601	6.3 (18)	600	9.8(14)
Franklin	180	2.2 (3)	55.6 (26)	160	18.1 (13)	56.9 (35)	19	* * *	56	* * *	128	4.7 (9)	128	10.9(20)
Geary	378	5.3 (25)	51.1 (7)	301	15.9 (6)	52.5 (23)	55	* * *	83	* * *	314	3.5 (4)	314	10.5(18)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2, 3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy	Hypertension During Pregnancy		
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)			Number	% (Rank)
Gove	9	* *	* *	8	* *	* *	0	* *	4	* *	11	* *	11	* *
Graham	17	* *	* *	14	* *	* *	2	* *	2	* *	14	* *	14	* *
Grant	69	* *	* *	63	* *	* *	7	* *	22	* *	66	* *	64	* *
Gray	45	* *	* *	40	* *	* *	2	* *	7	* *	40	* *	40	* *
Greeley	13	* *	* *	12	* *	* *	1	* *	2	* *	14	* *	14	* *
Greenwood	47	* *	* *	43	* *	* *	5	* *	5	* *	44	* *	45	* *
Hamilton	28	* *	* *	25	* *	* *	3	* *	10	* *	27	* *	27	* *
Harper	53	* *	* *	44	* *	* *	7	* *	9	* *	43	* *	43	* *
Harvey	277	4.7 (20)	53.1 (14)	241	26.6 (36)	45.2 (5)	40	* *	37	* *	246	6.1 (17)	246	11.8(24)
Haskell	20	* *	* *	17	* *	* *	3	* *	7	* *	17	* *	17	* *
Hodgeman	15	* *	* *	15	* *	* *	3	* *	4	* *	16	* *	16	* *
Jackson	70	* *	* *	62	* *	* *	15	* *	19	* *	64	* *	64	* *
Jefferson	67	* *	* *	61	* *	* *	11	* *	20	* *	62	* *	62	* *
Jewell	9	* *	* *	9	* *	* *	1	* *	1	* *	9	* *	9	* *
Johnson-M	804	5.1 (23)	55.2 (24)	700	19.3 (16)	50.3 (15)	134	26.1 (2)	228	16.2 (3)	733	8.5 (33)	731	8.1(7)
Johnson-O	1,008	4.0 (11)	54.5 (22)	909	19.5 (19)	51.6 (20)	173	22.5 (1)	364	19.0 (5)	953	8.2 (32)	952	6.7(5)
Kearny	19	* *	* *	17	* *	* *	2	* *	4	* *	17	* *	17	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2, 3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Kingman	59	* *	* *	51	* *	* *	7	* *	6	* *	51	* *	51	* *
Kiowa	15	* *	* *	12	* *	* *	1	* *	4	* *	13	* *	13	* *
Labette	190	7.9 (37)	52.6 (13)	178	25.8 (35)	42.7 (2)	31	* *	43	* *	179	5.6 (13)	179	11.2(21)
Lane	13	* *	* *	12	* *	* *	0	* *	2	* *	12	* *	12	* *
Leavenwort	347	4.6 (17)	57.3 (32)	297	19.2 (15)	51.5 (19)	61	* *	75	* *	303	8.9 (36)	302	9.9(15)
Lincoln	19	* *	* *	17	* *	* *	4	* *	7	* *	19	* *	19	* *
Linn	66	* *	* *	62	* *	* *	10	* *	10	* *	61	* *	61	* *
Logan	15	* *	* *	13	* *	* *	2	* *	2	* *	13	* *	13	* *
Lyon	306	2.0 (2)	63.1 (39)	267	23.6 (32)	43.8 (3)	45	* *	51	* *	282	7.1 (22)	282	14.2(35)
Marion	44	* *	* *	38	* *	* *	5	* *	14	* *	42	* *	42	* *
Marshall	47	* *	* *	41	* *	* *	5	* *	10	* *	42	* *	42	* *
Mc Pherson	142	5.6 (27)	50.0 (4)	126	19.8 (22)	46.0 (7)	18	* *	35	* *	128	5.5 (12)	129	10.1(16)
Meade	44	* *	* *	37	* *	* *	4	* *	6	* *	37	* *	37	* *
Miami	137	6.6 (30)	57.7 (33)	120	18.3 (14)	49.2 (13)	22	* *	44	* *	131	9.2 (37)	131	13.0(30)
Mitchell	52	* *	* *	49	* *	* *	4	* *	12	* *	50	* *	50	* *
Montgome-C	153	3.9 (9)	61.4 (38)	139	18.0 (11)	51.8 (21)	37	* *	39	* *	140	7.9 (29)	140	9.3(11)
Montgome-I	184	8.2 (38)	57.1 (31)	155	23.9 (33)	44.5 (4)	24	* *	48	* *	158	3.8 (6)	158	10.8(19)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2, 3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy	Hypertension During Pregnancy		
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)			Number	% (Rank)
Morris	22	* *	* *	20	* *	* *	3	* *	8	* *	21	* *	21	* *
Nemaha	30	* *	* *	28	* *	* *	7	* *	9	* *	32	* *	32	* *
Neosho-Ch	131	9.2 (39)	50.4 (5)	120	22.5 (29)	53.3 (28)	20	* *	17	* *	123	9.8 (38)	123	13.8(33)
Neosho-Er	9	* *	* *	8	* *	* *	2	* *	2	* *	10	* *	10	* *
Ness	4	* *	* *	4	* *	* *	0	* *	3	* *	5	* *	5	* *
Norton	32	* *	* *	29	* *	* *	4	* *	7	* *	29	* *	29	* *
Osage	94	* *	* *	82	* *	* *	9	* *	18	* *	86	* *	86	* *
Osborne	14	* *	* *	10	* *	* *	2	* *	5	* *	10	* *	10	* *
Ottawa	26	* *	* *	23	* *	* *	3	* *	4	* *	25	* *	25	* *
Pawnee	60	* *	* *	46	* *	* *	5	* *	12	* *	52	* *	52	* *
Phillips	34	* *	* *	30	* *	* *	2	* *	9	* *	32	* *	32	* *
Pottawat-S	18	* *	* *	17	* *	* *	1	* *	11	* *	17	* *	17	* *
Pottawat-W	19	* *	* *	17	* *	* *	0	* *	7	* *	18	* *	18	* *
Pratt	81	* *	* *	67	* *	* *	10	* *	17	* *	71	* *	71	* *
Rawlins	12	* *	* *	11	* *	* *	1	* *	3	* *	11	* *	11	* *
Reno	489	4.9 (22)	55.8 (29)	426	19.7 (21)	50.9 (16)	54	* *	64	* *	438	2.5 (2)	437	16.5(37)
Republic	41	* *	* *	32	* *	* *	5	* *	11	* *	34	* *	34	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2, 3)																
Clinic	Prepregnancy BMI (4)				Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy		
	Under-weight		Over-weight		< Ideal		> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)		Number	% (Rank)	% (Rank)		Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Rice	72	* *	* *		63	* *	* *		7	* *	18	* *	68	* *	67	* *
Riley-Ft	504	5.4 (26)	47.8 (1)		383	17.2 (9)	59.3 (37)		96	* *	114	18.4 (4)	415	4.1 (8)	414	12.1(26)
Riley-Manh	394	4.6 (17)	48.2 (3)		333	15.0 (2)	52.0 (22)		64	* *	94	* *	350	4.0 (7)	350	8.9(9)
Rooks	50	* *	* *		45	* *	* *		5	* *	7	* *	45	* *	45	* *
Rush	9	* *	* *		9	* *	* *		1	* *	3	* *	9	* *	9	* *
Russell	42	* *	* *		37	* *	* *		6	* *	9	* *	37	* *	37	* *
Saline	516	4.1 (13)	53.9 (17)		473	16.9 (7)	53.1 (25)		89	* *	105	14.3 (1)	482	5.6 (13)	486	6.2(3)
Scott	50	* *	* *		45	* *	* *		8	* *	6	* *	45	* *	45	* *
Sedgwick-C	1,190	4.5 (16)	53.8 (16)		1,074	19.8 (22)	49.8 (14)		225	31.6 (4)	227	27.3 (9)	1,112	6.6 (20)	1,112	9.3(11)
Sedgwick-M	1,990	3.5 (6)	54.4 (20)		1,802	20.1 (24)	50.9 (16)		420	27.9 (3)	440	22.3 (7)	1,857	5.8 (15)	1,858	11.7(23)
Sedgwick-S	1,394	5.6 (27)	52.1 (11)		1,274	20.3 (26)	54.0 (30)		269	36.8 (6)	321	28.0 (10)	1,309	6.3 (18)	1,311	12.2(27)
Seneca	7	* *	* *		6	* *	* *		1	* *	1	* *	8	* *	8	* *
Seward	518	1.9 (1)	58.7 (35)		468	24.1 (34)	46.2 (9)		93	* *	64	* *	478	8.8 (35)	482	6.4(4)
ShawneeM&I	259	3.9 (9)	52.1 (11)		187	29.4 (37)	46.0 (7)		4	* *	28	* *	189	7.9 (29)	189	5.3(1)
ShawneeMa	1,202	4.6 (17)	54.2 (19)		1,086	18.0 (11)	53.1 (25)		148	43.9 (7)	208	30.8 (11)	1,114	7.2 (23)	1,116	8.2(8)
Sheridan	10	* *	* *		10	* *	* *		2	* *	2	* *	10	* *	10	* *
Sherman	59	* *	* *		51	* *	* *		5	* *	5	* *	52	* *	52	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2, 3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy	Hypertension During Pregnancy		
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)			Number	% (Rank)
Smith	20	* *	* *	17	* *	* *	4	* *	4	* *	16	* *	16	* *
Stafford	43	* *	* *	38	* *	* *	4	* *	5	* *	38	* *	38	* *
Stanton	14	* *	* *	14	* *	* *	1	* *	7	* *	14	* *	14	* *
Stevens	86	* *	* *	76	* *	* *	14	* *	26	* *	79	* *	79	* *
Sumner	147	4.1 (13)	54.4 (20)	134	20.1 (24)	41.0 (1)	20	* *	37	* *	144	7.6 (28)	144	16.7(38)
Thomas	57	* *	* *	52	* *	* *	11	* *	11	* *	54	* *	54	* *
Trego	16	* *	* *	14	* *	* *	3	* *	3	* *	16	* *	16	* *
Wabaunsee	24	* *	* *	24	* *	* *	3	* *	10	* *	24	* *	24	* *
Washington	25	* *	* *	21	* *	* *	1	* *	8	* *	23	* *	23	* *
Wichita	18	* *	* *	15	* *	* *	3	* *	2	* *	15	* *	15	* *
Wilson	101	6.9 (33)	56.4 (30)	93	* *	* *	17	* *	14	* *	93	* *	93	* *
Woodson	8	* *	* *	8	* *	* *	1	* *	4	* *	8	* *	8	* *
Wyandotte	2,228	4.2 (15)	55.7 (28)	1,973	19.5 (19)	51.2 (18)	351	31.9 (5)	393	22.4 (8)	2,004	5.2 (11)	2,005	9.5(13)
Unknown Clinics	3	* *	* *	3	* *	* *	0	* *	1	* *	3	* *	3	* *
Kansas	21,646	4.6	54.3	19,129	19.9	51.1	3,394	29.5	4,491	22.7	19,697	6.4	19,699	10.3
Nation (Prior Year)	1,224,766	4.7	52.1	963,624	21.4	48.2	188,530	33.8	416,710	29.6	346,704	4.4	395,960	5.2

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6A
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2, 3)										
Clinic	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	% (Rank)		Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Allen	77	* *	* *		91	* *	91	* *	83	* *
Anderson	36	* *	* *		50	* *	50	* *	45	* *
Atchison	136	89.0 (2)	2.2 (11)		155	31.0 (30)	155	9.7 (33)	127	66.9 (7)
Barber-Med	37	* *	* *		45	* *	46	* *	40	* *
Barton	214	85.0 (10)	1.4 (7)		257	51.0 (2)	257	19.8 (5)	215	49.8 (31)
Bourbon	97	* *	* *		117	47.0 (3)	117	12.8 (26)	104	59.6 (18)
Brown	86	* *	* *		98	* *	99	* *	85	* *
Butler	241	64.7 (32)	13.3 (33)		282	38.3 (16)	282	16.7 (13)	207	31.9 (37)
Chase	14	* *	* *		16	* *	16	* *	12	* *
Chautauqua	32	* *	* *		31	* *	32	* *	30	* *
Cherokee	143	79.0 (18)	5.6 (27)		210	28.6 (37)	212	15.1 (20)	151	55.6 (23)
Cheyenne	1	* *	* *		3	* *	3	* *	2	* *
Clark	3	* *	* *		4	* *	4	* *	3	* *
Clay	53	* *	* *		59	* *	59	* *	56	* *
Cloud	81	* *	* *		88	* *	89	* *	77	* *
Coffey	20	* *	* *		25	* *	25	* *	17	* *
Comanche	11	* *	* *		11	* *	11	* *	9	* *
Cowley-Ark	152	57.9 (36)	7.9 (30)		202	40.6 (12)	204	12.7 (27)	171	53.2 (27)
Cowley-Win	131	75.6 (23)	3.8 (18)		156	44.2 (7)	159	13.2 (24)	136	64.0 (9)
Crawford-G	0	* *	* *		1	* *	1	* *	1	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6A
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2, 3)										
Clinic	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	% (Rank)		Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Crawford-P	329	83.3 (13)	3.6 (17)		400	42.3 (11)	400	13.3 (23)	349	59.9 (17)
Decatur	4	* *	* *		10	* *	10	* *	7	* *
DickinsonA	112	85.7 (8)	0.9 (6)		121	46.3 (6)	122	19.7 (7)	91	* *
DickinsonH	10	* *	* *		12	* *	12	* *	11	* *
Doniphan	33	* *	* *		37	* *	37	* *	25	* *
Douglas	428	75.7 (22)	4.7 (22)		507	33.1 (27)	507	20.5 (4)	397	63.0 (12)
Edwards	23	* *	* *		24	* *	25	* *	21	* *
Elk	18	* *	* *		20	* *	20	* *	16	* *
Ellis	154	84.4 (11)	2.6 (14)		179	30.2 (33)	180	26.7 (1)	139	74.8 (1)
Ellsworth	32	* *	* *		33	* *	33	* *	27	* *
Finney	581	58.9 (35)	25.6 (36)		712	34.0 (25)	718	13.0 (25)	593	35.1 (36)
Ford	584	70.0 (27)	16.4 (35)		682	43.0 (9)	686	7.4 (36)	611	52.7 (28)
Franklin	32	* *	* *		184	42.9 (10)	187	4.3 (39)	152	53.3 (26)
Geary	285	67.0 (30)	1.4 (7)		392	33.2 (26)	393	18.6 (8)	321	60.7 (15)
Gove	10	* *	* *		12	* *	12	* *	8	* *
Graham	14	* *	* *		17	* *	17	* *	14	* *
Grant	60	* *	* *		70	* *	71	* *	56	* *
Gray	36	* *	* *		44	* *	45	* *	39	* *
Greeley	14	* *	* *		15	* *	15	* *	10	* *
Greenwood	42	* *	* *		49	* *	49	* *	40	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6A
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2, 3)										
Clinic	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	% (Rank)		Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Hamilton	27	* *	* *		31	* *	31	* *	24	* *
Harper	44	* *	* *		53	* *	55	* *	45	* *
Harvey	225	74.7 (24)	14.7 (34)		281	47.0 (3)	284	16.2 (15)	238	68.9 (4)
Haskell	15	* *	* *		20	* *	20	* *	17	* *
Hodgeman	16	* *	* *		16	* *	16	* *	13	* *
Jackson	38	* *	* *		74	* *	75	* *	62	* *
Jefferson	60	* *	* *		68	* *	68	* *	54	* *
Jewell	9	* *	* *		9	* *	9	* *	9	* *
Johnson-M	660	69.7 (28)	2.9 (15)		846	31.4 (29)	850	15.9 (17)	660	70.2 (3)
Johnson-O	858	64.0 (33)	4.5 (20)		1,059	29.8 (34)	1,066	12.6 (28)	867	59.6 (18)
Kearny	15	* *	* *		19	* *	19	* *	15	* *
Kingman	49	* *	* *		58	* *	59	* *	53	* *
Kiowa	13	* *	* *		16	* *	16	* *	12	* *
Labette	151	86.8 (6)	5.3 (25)		193	39.4 (14)	194	16.0 (16)	166	55.4 (24)
Lane	12	* *	* *		13	* *	13	* *	10	* *
Leavenwort	304	88.5 (3)	0.0 (1)		357	34.7 (24)	359	22.6 (3)	275	72.4 (2)
Lincoln	19	* *	* *		21	* *	21	* *	16	* *
Linn	62	* *	* *		66	* *	66	* *	57	* *
Logan	2	* *	* *		15	* *	15	* *	12	* *
Lyon	279	86.7 (7)	0.7 (4)		322	38.8 (15)	323	15.5 (19)	265	63.4 (11)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6A
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2, 3)										
Clinic	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	% (Rank)		Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Marion	40	* *	* *		47	* *	47	* *	31	* *
Marshall	42	* *	* *		48	* *	48	* *	40	* *
Mc Pherson	126	88.1 (4)	0.8 (5)		144	38.2 (18)	144	22.9 (2)	121	65.3 (8)
Meade	37	* *	* *		43	* *	44	* *	36	* *
Miami	129	85.3 (9)	1.6 (9)		146	30.8 (31)	146	14.4 (21)	107	54.2 (25)
Mitchell	48	* *	* *		53	* *	54	* *	44	* *
Montgome-C	109	81.7 (15)	5.5 (26)		156	29.5 (35)	157	17.2 (9)	132	68.2 (6)
Montgome-I	132	83.3 (13)	4.5 (20)		186	36.6 (19)	187	11.8 (30)	160	61.3 (13)
Morris	18	* *	* *		23	* *	23	* *	19	* *
Nemaha	32	* *	* *		34	* *	34	* *	30	* *
Neosho-Ch	117	88.0 (5)	6.8 (29)		135	35.6 (22)	135	9.6 (34)	118	68.6 (5)
Neosho-Er	7	* *	* *		11	* *	11	* *	9	* *
Ness	5	* *	* *		5	* *	5	* *	3	* *
Norton	29	* *	* *		32	* *	32	* *	29	* *
Osage	73	* *	* *		90	* *	95	* *	89	* *
Osborne	10	* *	* *		14	* *	14	* *	12	* *
Ottawa	25	* *	* *		28	* *	28	* *	24	* *
Pawnee	50	* *	* *		61	* *	62	* *	50	* *
Phillips	32	* *	* *		36	* *	36	* *	32	* *
Pottawat-S	17	* *	* *		18	* *	18	* *	13	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6A
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2, 3)										
Clinic	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Pottawat-W	17	* *	* *	* *	20	* *	20	* *	17	* *
Pratt	70	* *	* *	* *	84	* *	85	* *	63	* *
Rawlins	0	* *	* *	* *	12	* *	12	* *	11	* *
Reno	424	84.0 (12)	2.1 (10)		495	40.0 (13)	499	19.8 (5)	414	59.4 (20)
Republic	34	* *	* *	* *	43	* *	43	* *	35	* *
Rice	67	* *	* *	* *	73	* *	75	* *	61	* *
Riley-Ft	383	93.2 (1)	0.5 (2)		540	28.5 (38)	541	16.3 (14)	424	59.2 (21)
Riley-Manh	329	76.6 (21)	0.6 (3)		412	38.3 (16)	412	17.2 (9)	339	61.1 (14)
Rooks	44	* *	* *	* *	50	* *	50	* *	40	* *
Rush	9	* *	* *	* *	8	* *	9	* *	7	* *
Russell	37	* *	* *	* *	42	* *	42	* *	34	* *
Saline	419	74.0 (25)	5.7 (28)		529	32.9 (28)	531	4.9 (38)	431	51.5 (30)
Scott	44	* *	* *	* *	49	* *	51	* *	45	* *
Sedgwick-C	1,097	77.3 (20)	4.4 (19)		1,225	29.1 (36)	1,230	12.3 (29)	1,023	46.8 (34)
Sedgwick-M	1,847	70.9 (26)	5.1 (24)		2,052	25.8 (39)	2,059	13.9 (22)	1,658	52.6 (29)
Sedgwick-S	1,281	80.2 (16)	2.3 (12)		1,428	30.3 (32)	1,432	17.0 (11)	1,166	39.5 (35)
Seneca	8	* *	* *	* *	9	* *	9	* *	9	* *
Seward	420	64.0 (33)	10.2 (32)		528	34.8 (23)	532	10.2 (32)	438	63.5 (10)
ShawneeM&I	173	68.8 (29)	2.3 (12)		259	66.4 (1)	262	8.4 (35)	246	49.2 (33)
ShawneeMa	994	80.2 (16)	3.3 (16)		1,243	43.2 (8)	1,248	7.2 (37)	1,018	49.6 (32)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6A
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2, 3)										
Clinic	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	% (Rank)		Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Sheridan	10	* *	* *		10	* *	10	* *	8	* *
Sherman	16	* *	* *		60	* *	60	* *	52	* *
Smith	15	* *	* *		19	* *	20	* *	18	* *
Stafford	38	* *	* *		43	* *	43	* *	38	* *
Stanton	14	* *	* *		14	* *	14	* *	7	* *
Stevens	79	* *	* *		88	* *	89	* *	72	* *
Sumner	143	79.0 (18)	4.9 (23)		157	35.7 (21)	157	15.9 (17)	121	60.3 (16)
Thomas	20	* *	* *		60	* *	60	* *	49	* *
Trego	15	* *	* *		18	* *	18	* *	15	* *
Wabaunsee	21	* *	* *		25	* *	25	* *	20	* *
Washington	23	* *	* *		27	* *	27	* *	20	* *
Wichita	15	* *	* *		18	* *	18	* *	15	* *
Wilson	92	* *	* *		101	46.5 (5)	101	16.8 (12)	88	* *
Woodson	8	* *	* *		8	* *	8	* *	6	* *
Wyandotte	1,981	65.6 (31)	8.0 (31)		2,267	35.9 (20)	2,286	10.4 (31)	1,910	56.9 (22)
Unknown Clinics	3	* *	* *		3	* *	3	* *	1	* *
Kansas	18,492	75.4	5.7		22,287	35.7	22,411	14.6	18,386	55.7
Nation (Prior Year)	580,485	80.1	3.1		1,187,387	31.7	266,567	19.3	269,548	80.6

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Allen	86	* *	68	* *	22	* *	80	* *	68	* *
Anderson	48	* *	37	* *	18	* *	44	* *	38	* *
Atchison	109	48.6 (34)	81	* *	27	* *	89	* *	84	* *
Barber-Med	46	* *	37	* *	18	* *	40	* *	37	* *
Barton	255	36.9 (16)	215	23.3 (18)	68	* *	215	14.0 (13)	214	4.7 (1)
Bourbon	110	52.7 (37)	82	* *	34	* *	97	* *	82	* *
Brown	95	* *	70	* *	25	* *	82	* *	70	* *
Butler	275	44.0 (30)	234	26.5 (26)	77	* *	204	12.7 (11)	236	7.6 (11)
Chase	16	* *	14	* *	6	* *	12	* *	14	* *
Chautauqua	23	* *	13	* *	5	* *	22	* *	13	* *
Cherokee	204	37.7 (19)	182	24.7 (24)	53	* *	148	33.1 (29)	183	32.8 (31)
Cheyenne	3	* *	3	* *	0	* *	2	* *	3	* *
Clark	0	* *	1	* *	0	* *	2	* *	3	* *
Clay	59	* *	53	* *	25	* *	56	* *	53	* *
Cloud	89	* *	79	* *	29	* *	77	* *	81	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Coffey	22	* *	20	* *	8	* *	16	* *	20	* *
Comanche	11	* *	11	* *	5	* *	9	* *	11	* *
Cowley-Ark	194	33.5 (13)	133	23.3 (18)	35	* *	165	19.4 (22)	134	18.7 (24)
Cowley-Win	153	52.3 (36)	116	34.5 (35)	51	* *	132	25.8 (26)	115	26.1 (27)
Crawford-G	1	* *	0	* *	0	* *	1	* *	0	* *
Crawford-P	363	41.3 (25)	282	23.8 (22)	93	* *	327	40.4 (33)	283	42.4 (34)
Decatur	10	* *	9	* *	2	* *	7	* *	9	* *
DickinsonA	121	43.8 (29)	110	30.9 (31)	36	* *	90	* *	110	6.4 (9)
DickinsonH	12	* *	10	* *	3	* *	11	* *	10	* *
Doniphan	34	* *	26	* *	7	* *	24	* *	27	* *
Douglas	497	44.3 (31)	422	21.6 (16)	150	41.3 (2)	392	10.2 (7)	425	5.6 (7)
Edwards	25	* *	24	* *	5	* *	20	* *	24	* *
Elk	19	* *	16	* *	5	* *	15	* *	15	* *
Ellis	179	36.9 (16)	151	13.2 (9)	38	* *	139	7.2 (1)	151	5.3 (5)
Ellsworth	33	* *	30	* *	7	* *	27	* *	30	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Finney	706	11.9 (3)	618	5.8 (3)	54	* *	588	17.2 (18)	619	16.0 (21)
Ford	673	11.1 (2)	568	5.5 (2)	44	* *	601	10.5 (8)	570	10.5 (15)
Franklin	186	47.8 (33)	165	30.9 (31)	61	* *	151	13.9 (12)	166	8.4 (13)
Geary	386	35.0 (14)	307	19.2 (13)	82	* *	320	37.5 (31)	307	29.6 (30)
Gove	11	* *	11	* *	1	* *	7	* *	11	* *
Graham	17	* *	13	* *	6	* *	14	* *	14	* *
Grant	71	* *	67	* *	1	* *	56	* *	67	* *
Gray	45	* *	37	* *	1	* *	38	* *	38	* *
Greeley	15	* *	14	* *	3	* *	10	* *	14	* *
Greenwood	38	* *	14	* *	6	* *	35	* *	31	* *
Hamilton	30	* *	27	* *	1	* *	24	* *	27	* *
Harper	47	* *	33	* *	13	* *	35	* *	28	* *
Harvey	275	36.0 (15)	234	19.7 (14)	70	* *	231	9.5 (6)	236	5.1 (4)
Haskell	20	* *	17	* *	3	* *	17	* *	17	* *
Hodgeman	16	* *	16	* *	7	* *	12	* *	15	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Jackson	75	* *	63	* *	20	* *	61	* *	63	* *
Jefferson	17	* *	1	* *	0	* *	19	* *	2	* *
Jewell	9	* *	9	* *	6	* *	9	* *	9	* *
Johnson-M	836	27.0 (7)	726	10.1 (5)	131	45.8 (1)	655	21.8 (24)	726	20.1 (26)
Johnson-O	1,052	20.2 (5)	939	9.7 (4)	135	40.0 (4)	862	15.8 (16)	940	15.5 (20)
Kearny	19	* *	17	* *	3	* *	15	* *	17	* *
Kingman	56	* *	43	* *	17	* *	52	* *	45	* *
Kiowa	15	* *	10	* *	1	* *	11	* *	11	* *
Labette	189	37.0 (18)	169	26.6 (27)	52	* *	165	30.9 (28)	170	28.8 (29)
Lane	3	* *	1	* *	0	* *	3	* *	1	* *
Leavenworth	351	45.9 (32)	294	25.2 (25)	95	* *	274	40.1 (32)	295	41.0 (33)
Lincoln	21	* *	15	* *	5	* *	16	* *	15	* *
Linn	66	* *	60	* *	26	* *	57	* *	60	* *
Logan	15	* *	13	* *	3	* *	12	* *	13	* *
Lyon	319	27.9 (9)	279	15.4 (11)	59	* *	265	12.1 (10)	278	8.6 (14)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Marion	47	* *	41	* *	8	* *	31	* *	41	* *
Marshall	45	* *	37	* *	8	* *	38	* *	38	* *
Mc Pherson	143	38.5 (20)	129	23.3 (18)	43	* *	120	9.2 (4)	129	5.4 (6)
Meade	38	* *	32	* *	9	* *	33	* *	33	* *
Miami	138	42.0 (26)	116	26.7 (29)	38	* *	102	16.7 (17)	116	14.7 (18)
Mitchell	53	* *	48	* *	14	* *	44	* *	49	* *
Montgome-C	156	41.0 (24)	134	30.6 (30)	43	* *	130	26.9 (27)	134	26.9 (28)
Montgome-I	181	40.3 (23)	141	31.2 (33)	44	* *	155	36.8 (30)	142	37.3 (32)
Morris	23	* *	18	* *	4	* *	17	* *	18	* *
Nemaha	32	* *	27	* *	13	* *	26	* *	27	* *
Neosho-Ch	103	49.5 (35)	52	* *	21	* *	92	* *	56	* *
Neosho-Er	9	* *	2	* *	0	* *	8	* *	2	* *
Ness	5	* *	4	* *	1	* *	3	* *	4	* *
Norton	31	* *	26	* *	12	* *	28	* *	28	* *
Osage	91	* *	74	* *	29	* *	83	* *	75	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Osborne	14	* *	10	* *	5	* *	12	* *	10	* *
Ottawa	27	* *	19	* *	9	* *	24	* *	21	* *
Pawnee	60	* *	46	* *	13	* *	49	* *	46	* *
Phillips	30	* *	24	* *	3	* *	19	* *	24	* *
Pottawat-S	18	* *	15	* *	3	* *	13	* *	16	* *
Pottawat-W	20	* *	17	* *	3	* *	17	* *	18	* *
Pratt	83	* *	69	* *	25	* *	62	* *	68	* *
Rawlins	12	* *	11	* *	4	* *	11	* *	11	* *
Reno	490	43.3 (28)	415	23.6 (21)	143	37.8 (6)	407	18.2 (20)	419	16.0 (21)
Republic	42	* *	31	* *	10	* *	33	* *	32	* *
Rice	75	* *	67	* *	23	* *	61	* *	68	* *
Riley-Ft	531	23.9 (6)	387	11.6 (6)	62	* *	422	19.9 (23)	394	17.3 (23)
Riley-Manh	406	32.3 (12)	318	18.6 (12)	70	* *	337	8.6 (3)	328	6.7 (10)
Rooks	49	* *	42	* *	19	* *	40	* *	43	* *
Rush	9	* *	9	* *	5	* *	7	* *	9	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Russell	42	* *	36	* *	16	* *	34	* *	36	* *
Saline	503	43.1 (27)	406	26.6 (27)	135	24.4 (9)	410	17.8 (19)	408	15.4 (19)
Scott	51	* *	44	* *	9	* *	44	* *	46	* *
Sedgwick-C	1,220	28.1 (10)	1,104	13.9 (10)	246	39.4 (5)	1,017	15.2 (15)	1,100	13.5 (17)
Sedgwick-M	2,038	27.4 (8)	1,849	12.9 (8)	360	41.1 (3)	1,652	9.3 (5)	1,852	7.7 (12)
Sedgwick-S	1,422	38.8 (22)	1,297	23.0 (17)	394	27.4 (8)	1,158	7.3 (2)	1,301	4.7 (1)
Seneca	8	* *	6	* *	2	* *	8	* *	6	* *
Seward	514	5.8 (1)	457	3.1 (1)	17	* *	435	14.7 (14)	463	11.7 (16)
ShawneeM&I	234	30.3 (11)	158	20.3 (15)	42	* *	214	18.7 (21)	162	4.9 (3)
ShawneeMa	1,177	38.5 (20)	945	24.4 (23)	273	28.6 (7)	977	10.8 (9)	951	6.0 (8)
Sheridan	9	* *	10	* *	1	* *	8	* *	10	* *
Sherman	59	* *	52	* *	10	* *	50	* *	52	* *
Smith	19	* *	14	* *	4	* *	17	* *	15	* *
Stafford	41	* *	36	* *	15	* *	35	* *	36	* *
Stanton	13	* *	12	* *	1	* *	7	* *	12	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Stevens	87	* *	77	* *	13	* *	70	* *	79	* *
Sumner	153	52.9 (38)	135	32.6 (34)	55	* *	117	46.2 (34)	138	49.3 (35)
Thomas	59	* *	53	* *	15	* *	48	* *	53	* *
Trego	16	* *	12	* *	5	* *	14	* *	12	* *
Wabaunsee	25	* *	24	* *	7	* *	20	* *	24	* *
Washington	27	* *	22	* *	5	* *	20	* *	22	* *
Wichita	18	* *	12	* *	2	* *	15	* *	12	* *
Wilson	99	* *	88	* *	40	* *	87	* *	88	* *
Woodson	8	* *	8	* *	2	* *	6	* *	8	* *
Wyandotte	2,255	19.8 (4)	2,000	12.1 (7)	299	23.7 (10)	1,897	22.5 (25)	2,004	19.4 (25)
Unknown Clinics	3	* *	3	* *	1	* *	1	* *	3	* *
Kansas	21,732	31.5	18,530	17.8	4,411	31.7	17,945	18.8	18,640	15.9
Nation (Prior Year)	999,199	23.1	856,953	14.2	146,933	35.5	655,095	18.9	578,852	19.9

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2, 3)										
Clinic	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)		Number	%	Number	%	Number	%
	Number	% (Rank)	% (Rank)	(Rank)						
Allen	81	* * * *	* *		82	* *	71	* *	82	* *
Anderson	33	* * * *	* *		43	* *	30	* *	43	* *
Atchison	136	6.6 (16)	5.9 (11)		139	6.5 (5)	127	3.1 (5)	139	54.7 (36)
Barber-Med	35	* * * *	* *		34	* *	29	* *	35	* *
Barton	204	4.4 (1)	2.9 (1)		206	9.2 (20)	185	3.2 (6)	204	72.1 (9)
Bourbon	98	* * * *	* *		99	* *	88	* *	99	* *
Brown	80	* * * *	* *		88	* *	76	* *	88	* *
Butler	246	5.7 (9)	7.3 (23)		250	5.2 (3)	233	3.4 (10)	248	70.2 (15)
Chase	13	* * * *	* *		13	* *	12	* *	13	* *
Chautauqua	30	* * * *	* *		31	* *	27	* *	32	* *
Cherokee	185	5.4 (7)	8.1 (25)		183	4.9 (2)	174	2.9 (4)	183	59.0 (33)
Cheyenne	2	* * * *	* *		2	* *	2	* *	2	* *
Clark	3	* * * *	* *		3	* *	2	* *	3	* *
Clay	52	* * * *	* *		52	* *	46	* *	52	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2, 3)										
Clinic	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)		Number	%	Number	%	Number	%
	Number	% (Rank)	% (Rank)	Number						
Cloud	78	* * * *	* * * *	77	* *	69	* *	78	* *	
Coffey	19	* * * *	* * * *	19	* *	18	* *	19	* *	
Comanche	11	* * * *	* * * *	11	* *	9	* *	11	* *	
Cowley-Ark	171	5.8 (10)	6.4 (18)	173	10.4 (30)	151	3.3 (7)	172	58.7 (34)	
Cowley-Win	135	13.3 (36)	8.9 (32)	136	13.2 (35)	115	8.7 (35)	138	69.6 (20)	
Crawford-G	0	* * * *	* * * *	0	* *	0	* *	0	* *	
Crawford-P	330	9.4 (31)	5.2 (8)	342	11.1 (32)	292	5.1 (25)	340	55.6 (35)	
Decatur	8	* * * *	* * * *	8	* *	7	* *	8	* *	
DickinsonA	106	6.6 (16)	10.4 (36)	105	9.5 (22)	95	* *	106	69.8 (19)	
DickinsonH	10	* * * *	* * * *	10	* *	9	* *	10	* *	
Doniphan	31	* * * *	* * * *	32	* *	30	* *	32	* *	
Douglas	422	6.4 (14)	5.2 (8)	430	7.7 (11)	390	3.8 (14)	427	77.8 (7)	
Edwards	23	* * * *	* * * *	23	* *	19	* *	23	* *	
Elk	19	* * * *	* * * *	19	* *	17	* *	19	* *	

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2, 3)										
Clinic	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)							
	Number	% (Rank)	% (Rank)		Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Ellis	136	4.4 (1)	10.3 (35)		150	7.3 (9)	124	1.6 (2)	150	72.0 (11)
Ellsworth	29	*	*	*	29	*	26	*	29	*
Finney	611	7.2 (20)	6.7 (21)		625	10.1 (29)	544	4.0 (17)	621	77.9 (6)
Ford	568	4.6 (3)	6.0 (13)		588	10.0 (28)	510	1.6 (2)	587	72.1 (9)
Franklin	164	5.5 (8)	4.9 (5)		163	9.2 (20)	146	3.4 (10)	165	68.5 (23)
Geary	314	8.9 (28)	5.1 (6)		313	9.9 (26)	282	5.3 (27)	311	71.4 (12)
Gove	11	*	*	*	11	*	10	*	9	*
Graham	12	*	*	*	13	*	12	*	13	*
Grant	65	*	*	*	66	*	59	*	67	*
Gray	36	*	*	*	38	*	32	*	37	*
Greeley	13	*	*	*	13	*	12	*	13	*
Greenwood	40	*	*	*	40	*	38	*	40	*
Hamilton	26	*	*	*	26	*	23	*	26	*
Harper	43	*	*	*	43	*	37	*	45	*

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2, 3)									
Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)	High (5)		Number	%	Number	%	Number	%
	Number	% (Rank)	% (Rank)						
Harvey	234	6.0 (12)	8.5 (29)	234	8.5 (13)	211	3.8 (14)	235	80.0 (4)
Haskell	15	* *	* *	16	* *	14	* *	16	* *
Hodgeman	15	* *	* *	15	* *	13	* *	15	* *
Jackson	65	* *	* *	65	* *	57	* *	65	* *
Jefferson	61	* *	* *	62	* *	49	* *	61	* *
Jewell	9	* *	* *	9	* *	8	* *	9	* *
Johnson-M	677	6.9 (19)	6.2 (16)	729	6.4 (4)	631	4.1 (18)	728	78.0 (5)
Johnson-O	884	6.3 (13)	8.8 (31)	933	7.5 (10)	813	3.7 (12)	935	81.4 (3)
Kearny	17	* *	* *	17	* *	15	* *	17	* *
Kingman	34	* *	* *	47	* *	32	* *	48	* *
Kiowa	13	* *	* *	13	* *	10	* *	13	* *
Labette	167	4.8 (5)	6.0 (13)	168	8.9 (17)	153	3.3 (7)	166	51.2 (37)
Lane	10	* *	* *	11	* *	9	* *	11	* *
Leavenwort	298	9.1 (29)	8.4 (26)	303	8.6 (15)	272	6.3 (32)	304	62.2 (31)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2, 3)										
Clinic	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)		Number	%	Number	%	Number	%
	Number	% (Rank)	% (Rank)	Number						
Lincoln	17	* * * *	* * * *	17	* *	17	* *	16	* *	
Linn	59	* * * *	* * * *	60	* *	57	* *	60	* *	
Logan	12	* * * *	* * * *	12	* *	11	* *	12	* *	
Lyon	273	8.4 (26)	5.9 (11)	272	9.6 (23)	246	5.7 (31)	269	72.9 (8)	
Marion	32	* * * *	* * * *	34	* *	31	* *	33	* *	
Marshall	40	* * * *	* * * *	41	* *	36	* *	41	* *	
Mc Pherson	122	4.9 (6)	6.6 (20)	123	6.5 (5)	114	0.9 (1)	123	69.1 (21)	
Meade	30	* * * *	* * * *	33	* *	27	* *	33	* *	
Miami	128	10.2 (33)	8.6 (30)	130	8.5 (13)	118	6.8 (33)	128	68.8 (22)	
Mitchell	36	* * * *	* * * *	47	* *	34	* *	47	* *	
Montgome-C	140	6.4 (14)	3.6 (4)	139	8.6 (15)	127	3.9 (16)	140	62.9 (29)	
Montgome-I	153	9.2 (30)	3.3 (2)	153	11.1 (32)	135	4.4 (20)	153	59.5 (32)	
Morris	15	* * * *	* * * *	18	* *	15	* *	17	* *	
Nemaha	21	* * * *	* * * *	32	* *	19	* *	31	* *	

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2, 3)										
Clinic	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)		Number	%	Number	%	Number	%
	Number	% (Rank)	% (Rank)	Number						
Neosho-Ch	66	* * * *	* *	123	13.8 (36)	57	* *	122	62.3 (30)	
Neosho-Er	4	* * * *	* *	10	* *	4	* *	10	* *	
Ness	4	* * * *	* *	4	* *	3	* *	4	* *	
Norton	28	* * * *	* *	28	* *	22	* *	27	* *	
Osage	85	* * * *	* *	82	* *	67	* *	87	* *	
Osborne	9	* * * *	* *	10	* *	9	* *	9	* *	
Ottawa	25	* * * *	* *	25	* *	24	* *	25	* *	
Pawnee	48	* * * *	* *	47	* *	44	* *	48	* *	
Phillips	32	* * * *	* *	32	* *	25	* *	32	* *	
Pottawat-S	16	* * * *	* *	16	* *	13	* *	16	* *	
Pottawat-W	18	* * * *	* *	18	* *	17	* *	18	* *	
Pratt	66	* * * *	* *	65	* *	61	* *	66	* *	
Rawlins	11	* * * *	* *	11	* *	10	* *	11	* *	
Reno	428	8.2 (22)	3.5 (3)	427	9.6 (23)	384	5.2 (26)	431	66.6 (26)	

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2, 3)										
Clinic	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)		Number	%	Number	%	Number	%
	Number	% (Rank)	% (Rank)	Number						
Republic	32	* * * *	* * * *	33	* *	32	* *	33	* *	
Rice	65	* * * *	* * * *	63	* *	54	* *	65	* *	
Riley-Ft	411	4.6 (3)	9.5 (34)	412	3.4 (1)	396	3.3 (7)	412	70.6 (14)	
Riley-Manh	345	6.7 (18)	6.1 (15)	347	6.6 (7)	322	3.7 (12)	347	83.6 (1)	
Rooks	45	* * * *	* * * *	45	* *	44	* *	45	* *	
Rush	9	* * * *	* * * *	8	* *	6	* *	9	* *	
Russell	35	* * * *	* * * *	35	* *	32	* *	35	* *	
Saline	466	8.8 (27)	8.4 (26)	478	9.8 (25)	418	5.3 (27)	476	63.2 (28)	
Scott	45	* * * *	* * * *	43	* *	40	* *	45	* *	
Sedgwick-C	1,089	7.9 (21)	7.7 (24)	1,089	6.9 (8)	1,009	4.5 (21)	1,090	67.9 (25)	
Sedgwick-M	1,815	10.1 (32)	6.2 (16)	1,829	8.9 (17)	1,650	5.5 (29)	1,822	70.0 (17)	
Sedgwick-S	1,278	8.3 (23)	6.8 (22)	1,280	8.3 (12)	1,169	5.0 (23)	1,278	68.3 (24)	
Seneca	5	* * * *	* * * *	8	* *	5	* *	8	* *	
Seward	442	5.9 (11)	8.4 (26)	466	9.9 (26)	394	4.1 (18)	464	64.0 (27)	

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2, 3)										
Clinic	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)		Number	%	Number	%	Number	%
	Number	% (Rank)	% (Rank)	Number						
ShawneeM&I	176	10.8 (34)	5.1 (6)		176	15.9 (37)	145	5.5 (29)	178	81.5 (2)
ShawneeMa	1,097	8.3 (23)	5.6 (10)		1,111	10.8 (31)	974	5.0 (23)	1,107	70.8 (13)
Sheridan	10	* * * *	* * * *		10	* *	8	* *	10	* *
Sherman	52	* * * *	* * * *		52	* *	51	* *	52	* *
Smith	15	* * * *	* * * *		15	* *	12	* *	16	* *
Stafford	35	* * * *	* * * *		37	* *	31	* *	34	* *
Stanton	12	* * * *	* * * *		14	* *	12	* *	14	* *
Stevens	46	* * * *	* * * *		75	* *	42	* *	76	* *
Sumner	135	11.1 (35)	8.9 (32)		140	12.1 (34)	118	6.8 (33)	140	70.0 (17)
Thomas	54	* * * *	* * * *		54	* *	47	* *	54	* *
Trego	16	* * * *	* * * *		16	* *	16	* *	16	* *
Wabaunsee	24	* * * *	* * * *		24	* *	24	* *	24	* *
Washington	21	* * * *	* * * *		22	* *	16	* *	22	* *
Wichita	15	* * * *	* * * *		15	* *	12	* *	15	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2, 3)														
Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed						
	Low (4)	High (5)		Number	%	Number	%	Number	%					
	Number	% (Rank)	% (Rank)							(Rank)	(Rank)	(Rank)		
Wilson	93	*	*	*	*	94	*	*	81	*	*	93	*	*
Woodson	8	*	*	*	*	8	*	*	7	*	*	8	*	*
Wyandotte	1,925	8.3 (23)	6.4 (18)	1,966	8.9 (17)	1,742	4.8 (22)	1,970	70.2 (15)					
Unknown Clinics	3	*	*	*	*	3	*	*	3	*	*	3	*	*
Kansas	18,865	7.7	6.5	19,327	8.8	17,104	4.4	19,312	70.3					
Nation (Prior Year)	1,085,134	8.2	6.6	1,025,278	11.5	890,981	3.6	952,980	65.4					

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3A
2007 - 2009 Pregnancy Nutrition Surveillance
Kansas
Comparison of Racial and Ethnic Distribution by Clinic
Combined 3 Years

(1)

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaska Native	Asian/Pacific Islander	Multiple Races	All Other/Unknown
		%	%	%	%	%	%	%
Allen	298	92.3	3.4	1.3	1.3	0.7	1.0	0.0
Anderson	155	92.9	0.6	4.5	0.6	1.3	0.0	0.0
Atchison	413	88.1	8.2	1.5	0.7	0.0	1.5	0.0
Barber-Kio	5	*	*	*	*	*	*	*
Barber-Med	110	95.5	2.7	0.9	0.0	0.9	0.0	0.0
Barton	815	69.6	1.5	28.0	0.2	0.2	0.5	0.0
Bourbon	390	93.3	2.8	2.1	0.3	0.3	1.3	0.0
Brown	293	76.1	2.7	4.1	14.7	0.0	2.4	0.0
Butler	815	90.8	1.3	5.4	0.4	0.4	1.7	0.0
Chase	40	*	*	*	*	*	*	*
Chautauqua	89	*	*	*	*	*	*	*
Cherokee	626	90.7	1.3	3.8	1.8	1.6	0.8	0.0
Cheyenne	6	*	*	*	*	*	*	*
Clark	10	*	*	*	*	*	*	*
Clay	185	95.7	0.5	1.6	1.1	0.5	0.5	0.0
Cloud	240	93.8	1.3	2.9	0.8	0.8	0.4	0.0
Coffey	69	*	*	*	*	*	*	*
Comanche	24	*	*	*	*	*	*	*
Cowley-Ark	558	67.9	2.9	27.4	0.7	0.4	0.7	0.0
Cowley-Win	462	86.8	1.7	6.3	1.3	3.0	0.9	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3A
2007 - 2009 Pregnancy Nutrition Surveillance
Kansas
Comparison of Racial and Ethnic Distribution by Clinic
Combined 3 Years

(1)

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Crawford-G	2	*	*	*	*	*	*	*
Crawford-P	1,155	84.1	3.5	9.7	0.1	1.5	1.2	0.0
Decatur	30	*	*	*	*	*	*	*
DickinsonA	304	92.8	0.0	5.6	1.0	0.3	0.3	0.0
DickinsonH	44	*	*	*	*	*	*	*
Doniphan	122	86.1	4.1	7.4	0.8	0.0	1.6	0.0
Douglas	1,442	63.9	9.1	10.9	9.2	5.2	1.8	0.0
Edwards	69	*	*	*	*	*	*	*
Elk	48	*	*	*	*	*	*	*
Ellis	471	84.7	1.5	13.0	0.4	0.4	0.0	0.0
Ellsworth	94	*	*	*	*	*	*	*
Finney	2,024	30.0	0.9	66.2	0.0	2.6	0.2	0.0
Ford	1,930	22.8	0.9	75.4	0.2	0.7	0.1	0.0
Franklin	509	90.6	1.8	4.7	1.2	0.4	1.4	0.0
Geary	1,056	53.1	24.4	15.2	0.6	2.8	3.9	0.0
Gove	31	*	*	*	*	*	*	*
Graham	51	*	*	*	*	*	*	*
Grant	210	19.5	0.0	80.0	0.0	0.0	0.5	0.0
Gray	157	67.5	0.0	31.8	0.0	0.0	0.6	0.0
Greeley	31	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3A
2007 - 2009 Pregnancy Nutrition Surveillance
Kansas
Comparison of Racial and Ethnic Distribution by Clinic
Combined 3 Years

(1)

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Greenwood	140	92.9	0.7	5.0	0.0	0.7	0.7	0.0
Hamilton	74	*	*	*	*	*	*	*
Harper	147	81.0	0.0	16.3	1.4	0.0	1.4	0.0
Harvey	790	69.9	2.5	24.7	1.1	0.3	1.5	0.0
Haskell	63	*	*	*	*	*	*	*
Hodgeman	39	*	*	*	*	*	*	*
Jackson	220	76.4	0.0	5.9	15.5	0.0	2.3	0.0
Jefferson	193	96.4	0.5	1.0	2.1	0.0	0.0	0.0
Jewell	39	*	*	*	*	*	*	*
Johnson-M	2,537	46.5	15.1	33.5	0.6	2.9	1.5	0.0
Johnson-O	2,977	46.3	11.0	38.0	0.5	3.3	0.9	0.0
Kearny	60	*	*	*	*	*	*	*
Kingman	173	91.9	0.0	5.2	2.3	0.6	0.0	0.0
Kiowa	33	*	*	*	*	*	*	*
Labette	536	87.7	7.3	2.6	1.3	0.4	0.7	0.0
Lane	38	*	*	*	*	*	*	*
Leavenwort	1,045	69.4	15.6	9.8	0.7	1.4	3.2	0.0
Lincoln	60	*	*	*	*	*	*	*
Linn	186	96.2	0.0	2.7	0.0	0.0	1.1	0.0
Logan	36	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3A
2007 - 2009 Pregnancy Nutrition Surveillance
Kansas
Comparison of Racial and Ethnic Distribution by Clinic
Combined 3 Years

(1)

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaska Native	Asian/Pacific Islander	Multiple Races	All Other/Unknown
		%	%	%	%	%	%	%
Lyon	1,027	54.2	5.0	38.6	0.2	1.4	0.7	0.0
Marion	136	92.6	0.0	5.9	0.7	0.7	0.0	0.0
Marshall	171	91.8	0.0	3.5	2.9	0.0	1.8	0.0
Mc Pherson	406	89.2	1.5	7.6	1.2	0.0	0.5	0.0
Meade	137	62.0	0.0	37.2	0.0	0.0	0.7	0.0
Miami	409	92.7	2.4	2.9	0.0	0.7	1.2	0.0
Mitchell	129	96.1	0.0	3.9	0.0	0.0	0.0	0.0
Montgome-C	448	71.2	12.3	10.0	4.2	0.7	1.6	0.0
Montgome-I	562	81.9	7.1	7.8	2.1	0.9	0.2	0.0
Morris	81	*	*	*	*	*	*	*
Nemaha	88	*	*	*	*	*	*	*
Neosho-Ch	402	91.3	2.5	4.5	0.5	0.5	0.7	0.0
Neosho-Er	36	*	*	*	*	*	*	*
Ness	18	*	*	*	*	*	*	*
Norton	85	*	*	*	*	*	*	*
Osage	264	96.6	0.0	1.9	0.8	0.0	0.8	0.0
Osborne	53	*	*	*	*	*	*	*
Ottawa	76	*	*	*	*	*	*	*
Pawnee	164	81.7	2.4	14.6	0.0	0.0	1.2	0.0
Phillips	107	93.5	0.0	5.6	0.0	0.9	0.0	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3A
2007 - 2009 Pregnancy Nutrition Surveillance
Kansas
Comparison of Racial and Ethnic Distribution by Clinic
Combined 3 Years

(1)

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Pottawat-S	60	*	*	*	*	*	*	*
Pottawat-W	70	*	*	*	*	*	*	*
Pratt	227	87.2	0.0	12.3	0.0	0.4	0.0	0.0
Rawlins	31	*	*	*	*	*	*	*
Reno	1,434	81.8	2.9	13.2	0.5	0.8	0.9	0.0
Republic	103	91.3	1.9	4.9	0.0	0.0	1.9	0.0
Rice	209	77.0	1.4	21.5	0.0	0.0	0.0	0.0
Riley-Ft	1,473	72.0	12.4	11.7	0.5	3.3	0.1	0.0
Riley-Manh	1,166	75.1	10.3	8.5	0.7	4.8	0.6	0.0
Rooks	130	95.4	0.8	0.8	1.5	1.5	0.0	0.0
Rush	35	*	*	*	*	*	*	*
Russell	126	92.1	4.8	1.6	1.6	0.0	0.0	0.0
Saline	1,437	71.7	4.4	19.7	0.4	2.3	1.5	0.0
Scott	143	59.4	0.7	39.9	0.0	0.0	0.0	0.0
Sedgwick-C	3,345	42.9	13.5	33.5	1.3	7.2	1.6	0.0
Sedgwick-M	5,872	33.9	27.4	33.4	1.6	2.6	1.1	0.0
Sedgwick-S	3,538	67.6	7.9	17.9	2.4	2.4	1.8	0.0
Seneca	28	*	*	*	*	*	*	*
Seward	1,478	17.4	1.4	78.3	1.8	0.7	0.3	0.0
ShawneeM&I	579	38.7	15.0	43.2	1.9	0.9	0.3	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3A
2007 - 2009 Pregnancy Nutrition Surveillance
Kansas
Comparison of Racial and Ethnic Distribution by Clinic
Combined 3 Years

(1)

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
ShawneeMa	3,724	58.2	15.9	21.5	2.3	0.7	1.5	0.0
Sheridan	27	*	*	*	*	*	*	*
Sherman	190	75.3	0.0	22.6	0.5	0.5	1.1	0.0
Smith	61	*	*	*	*	*	*	*
Stafford	118	66.9	0.0	32.2	0.0	0.8	0.0	0.0
Stanton	56	*	*	*	*	*	*	*
Stevens	250	45.2	0.0	53.6	0.8	0.4	0.0	0.0
Sumner	443	91.2	0.9	5.0	1.1	0.2	1.6	0.0
Thomas	160	86.9	1.9	10.0	0.0	0.0	1.3	0.0
Trego	50	*	*	*	*	*	*	*
Wabaunsee	62	*	*	*	*	*	*	*
Wallace	4	*	*	*	*	*	*	*
Washington	83	*	*	*	*	*	*	*
Wichita	46	*	*	*	*	*	*	*
Wilson	281	96.4	0.7	2.1	0.4	0.0	0.4	0.0
Woodson	25	*	*	*	*	*	*	*
Wyandotte	6,467	26.0	31.1	39.9	0.2	2.5	0.3	0.0
Unknown Clinics	3	*	*	*	*	*	*	*
Kansas	63,102	57.2	11.4	27.0	1.3	2.1	1.0	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Age Distribution by Clinic
Combined 3 Years

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Allen	298	0.3	8.1	15.4	63.8	11.4	0.3	0.7
Anderson	155	0.6	8.4	12.3	67.1	11.0	0.6	0.0
Atchison	413	0.0	6.5	17.4	61.5	13.8	0.7	0.0
Barber-Kio	5	*	*	*	*	*	*	*
Barber-Med	110	0.0	4.5	22.7	60.9	11.8	0.0	0.0
Barton	815	0.1	6.6	17.8	59.9	15.0	0.6	0.0
Bourbon	390	0.0	6.9	19.7	61.8	11.0	0.5	0.0
Brown	293	0.0	6.5	15.4	64.8	12.3	1.0	0.0
Butler	815	0.1	7.6	17.2	61.7	12.8	0.5	0.1
Chase	40	*	*	*	*	*	*	*
Chautauqua	89	*	*	*	*	*	*	*
Cherokee	626	0.2	8.6	18.7	58.9	12.5	0.8	0.3
Cheyenne	6	*	*	*	*	*	*	*
Clark	10	*	*	*	*	*	*	*
Clay	185	0.0	7.6	17.8	60.0	14.1	0.5	0.0
Cloud	240	0.0	5.4	17.5	65.4	10.4	1.3	0.0
Coffey	69	*	*	*	*	*	*	*
Comanche	24	*	*	*	*	*	*	*
Cowley-Ark	558	1.1	8.6	17.2	57.9	14.5	0.7	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Age Distribution by Clinic
Combined 3 Years

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Cowley-Win	462	0.4	7.6	15.4	64.5	12.1	0.0	0.0
Crawford-G	2	*	*	*	*	*	*	*
Crawford-P	1,155	0.2	7.2	16.3	63.1	12.5	0.6	0.2
Decatur	30	*	*	*	*	*	*	*
DickinsonA	304	0.3	7.2	14.5	61.8	15.8	0.3	0.0
DickinsonH	44	*	*	*	*	*	*	*
Doniphan	122	0.8	4.1	14.8	63.9	16.4	0.0	0.0
Douglas	1,442	0.1	3.8	13.2	63.9	17.7	1.4	0.0
Edwards	69	*	*	*	*	*	*	*
Elk	48	*	*	*	*	*	*	*
Ellis	471	0.0	3.8	11.7	70.9	11.5	1.9	0.2
Ellsworth	94	*	*	*	*	*	*	*
Finney	2,024	0.4	9.9	14.8	54.2	19.2	1.4	0.1
Ford	1,930	0.4	7.6	12.5	56.1	22.1	1.3	0.1
Franklin	509	0.8	6.3	19.8	58.2	14.1	0.6	0.2
Geary	1,056	0.5	4.9	15.4	67.2	11.6	0.4	0.0
Gove	31	*	*	*	*	*	*	*
Graham	51	*	*	*	*	*	*	*
Grant	210	0.0	4.8	15.7	60.0	17.6	1.9	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Age Distribution by Clinic
Combined 3 Years

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Gray	157	0.0	5.7	10.8	53.5	28.0	1.9	0.0
Greeley	31	*	*	*	*	*	*	*
Greenwood	140	0.0	5.7	17.1	62.1	15.0	0.0	0.0
Hamilton	74	*	*	*	*	*	*	*
Harper	147	0.0	7.5	12.2	70.1	10.2	0.0	0.0
Harvey	790	0.3	6.5	13.4	63.3	15.3	1.3	0.0
Haskell	63	*	*	*	*	*	*	*
Hodgeman	39	*	*	*	*	*	*	*
Jackson	220	0.0	7.3	15.9	66.8	10.0	0.0	0.0
Jefferson	193	0.0	8.3	17.6	60.1	13.5	0.5	0.0
Jewell	39	*	*	*	*	*	*	*
Johnson-M	2,537	0.2	4.3	12.1	60.8	21.4	1.2	0.0
Johnson-O	2,977	0.1	4.5	11.1	60.8	22.1	1.3	0.1
Kearny	60	*	*	*	*	*	*	*
Kingman	173	0.0	5.8	14.5	64.2	15.0	0.6	0.0
Kiowa	33	*	*	*	*	*	*	*
Labette	536	0.0	7.5	19.4	62.1	10.1	0.9	0.0
Lane	38	*	*	*	*	*	*	*
Leavenwort	1,045	0.1	5.9	18.4	63.3	11.3	1.1	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Age Distribution by Clinic
Combined 3 Years

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Lincoln	60	*	*	*	*	*	*	*
Linn	186	0.0	5.4	17.2	62.9	13.4	1.1	0.0
Logan	36	*	*	*	*	*	*	*
Lyon	1,027	0.4	6.2	13.7	62.7	14.7	2.2	0.0
Marion	136	0.0	2.2	12.5	66.9	14.7	3.7	0.0
Marshall	171	0.0	5.3	9.9	69.0	15.8	0.0	0.0
Mc Pherson	406	0.2	4.9	13.1	65.8	15.5	0.5	0.0
Meade	137	0.0	7.3	10.9	60.6	19.7	1.5	0.0
Miami	409	0.2	7.3	14.4	65.5	12.0	0.5	0.0
Mitchell	129	0.0	2.3	16.3	61.2	20.2	0.0	0.0
Montgome-C	448	0.7	6.0	17.6	62.7	12.5	0.2	0.2
Montgome-I	562	0.0	5.5	16.7	67.3	10.0	0.4	0.2
Morris	81	*	*	*	*	*	*	*
Nemaha	88	*	*	*	*	*	*	*
Neosho-Ch	402	0.2	7.0	19.7	62.4	10.4	0.2	0.0
Neosho-Er	36	*	*	*	*	*	*	*
Ness	18	*	*	*	*	*	*	*
Norton	85	*	*	*	*	*	*	*
Osage	264	0.0	5.7	18.2	65.5	9.8	0.0	0.8

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Age Distribution by Clinic
Combined 3 Years

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Osborne	53	*	*	*	*	*	*	*
Ottawa	76	*	*	*	*	*	*	*
Pawnee	164	1.2	6.7	14.6	68.3	7.9	1.2	0.0
Phillips	107	0.0	2.8	15.0	63.6	18.7	0.0	0.0
Pottawat-S	60	*	*	*	*	*	*	*
Pottawat-W	70	*	*	*	*	*	*	*
Pratt	227	0.4	9.3	17.6	58.1	13.7	0.9	0.0
Rawlins	31	*	*	*	*	*	*	*
Reno	1,434	0.1	5.6	16.5	65.8	11.4	0.6	0.0
Republic	103	0.0	3.9	22.3	55.3	16.5	1.9	0.0
Rice	209	0.5	6.2	13.4	58.9	20.1	1.0	0.0
Riley-Ft	1,473	0.0	0.7	10.8	78.1	10.1	0.3	0.0
Riley-Manh	1,166	0.0	3.3	13.6	70.3	12.2	0.4	0.1
Rooks	130	0.0	3.8	13.8	63.8	16.9	1.5	0.0
Rush	35	*	*	*	*	*	*	*
Russell	126	0.0	6.3	16.7	65.1	11.9	0.0	0.0
Saline	1,437	0.2	7.7	15.1	63.1	12.9	1.0	0.1
Scott	143	0.7	5.6	10.5	66.4	15.4	1.4	0.0
Sedgwick-C	3,345	0.2	6.6	13.0	60.9	18.2	1.0	0.1

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Age Distribution by Clinic
Combined 3 Years

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Sedgwick-M	5,872	0.4	7.3	13.8	60.1	16.9	1.4	0.1
Sedgwick-S	3,538	0.3	6.6	14.8	61.9	15.0	1.4	0.0
Seneca	28	*	*	*	*	*	*	*
Seward	1,478	0.3	7.7	13.9	55.8	20.6	1.6	0.0
ShawneeM&I	579	0.7	8.5	14.0	56.3	18.8	1.7	0.0
ShawneeMa	3,724	0.3	6.6	14.8	62.5	14.7	1.0	0.0
Sheridan	27	*	*	*	*	*	*	*
Sherman	190	0.0	4.7	18.4	57.4	16.3	3.2	0.0
Smith	61	*	*	*	*	*	*	*
Stafford	118	0.0	4.2	12.7	59.3	21.2	2.5	0.0
Stanton	56	*	*	*	*	*	*	*
Stevens	250	0.0	6.8	14.8	59.2	18.8	0.4	0.0
Sumner	443	0.0	5.0	18.7	62.8	12.9	0.7	0.0
Thomas	160	0.0	7.5	15.6	65.0	11.9	0.0	0.0
Trego	50	*	*	*	*	*	*	*
Wabaunsee	62	*	*	*	*	*	*	*
Wallace	4	*	*	*	*	*	*	*
Washington	83	*	*	*	*	*	*	*
Wichita	46	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Age Distribution by Clinic
Combined 3 Years

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Wilson	281	0.0	8.9	16.0	62.6	12.5	0.0	0.0
Woodson	25	*	*	*	*	*	*	*
Wyandotte	6,467	0.6	8.2	13.9	58.0	18.0	1.3	0.1
Unknown Clinics	3	*	*	*	*	*	*	*
Kansas	63,102	0.3	6.5	14.5	61.6	16.1	1.1	0.1

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Maternal Health Indicators by Clinic
Combined 3 Years

Comparison of Maternal Health Indicators (2, 3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Allen	296	5.1 (46)	54.4 (43)	273	18.3 (21)	56.4 (64)	51	* *	92	* *	267	6.7 (65)	267	12.4(66)
Anderson	154	4.5 (33)	50.6 (14)	143	22.4 (53)	47.6 (12)	26	* *	27	* *	137	2.2 (8)	137	10.9(64)
Atchison	400	5.8 (50)	56.5 (55)	349	19.5 (36)	53.9 (49)	61	* *	87	* *	354	4.0 (38)	353	12.2(65)
Barber-Kio	5	* *	* *	4	* *	* *	2	* *	1	* *	4	* *	4	* *
Barber-Med	108	3.7 (15)	57.4 (60)	90	* *	* *	13	* *	26	* *	89	* *	89	* *
Barton	803	4.5 (33)	52.6 (26)	724	20.9 (44)	48.8 (18)	97	* *	156	25.0 (19)	692	3.2 (21)	690	7.7(46)
Bourbon	384	3.1 (10)	56.8 (58)	343	19.5 (36)	52.8 (41)	47	* *	78	* *	339	3.2 (21)	339	6.5(28)
Brown	290	3.1 (10)	55.2 (49)	256	16.8 (6)	56.3 (63)	45	* *	65	* *	257	1.6 (4)	257	9.7(62)
Butler	791	6.3 (56)	50.2 (12)	728	18.7 (27)	52.5 (40)	103	38.8 (20)	216	28.7 (26)	692	4.2 (44)	692	8.8(58)
Chase	38	* *	* *	33	* *	* *	4	* *	3	* *	35	* *	35	* *
Chautauqua	89	* *	* *	79	* *	* *	7	* *	12	* *	79	* *	79	* *
Cherokee	604	5.0 (45)	51.8 (21)	526	16.2 (4)	54.8 (53)	74	* *	139	27.3 (25)	525	3.0 (18)	526	6.7(34)
Cheyenne	6	* *	* *	6	* *	* *	0	* *	1	* *	6	* *	6	* *
Clark	8	* *	* *	6	* *	* *	1	* *	2	* *	7	* *	7	* *
Clay	183	3.8 (18)	56.3 (54)	163	13.5 (1)	61.3 (67)	21	* *	47	* *	158	5.7 (62)	158	7.0(39)
Cloud	240	4.6 (35)	51.3 (19)	214	21.5 (48)	53.3 (47)	28	* *	62	* *	202	4.0 (38)	206	8.7(57)
Coffey	67	* *	* *	62	* *	* *	6	* *	25	* *	56	* *	56	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Maternal Health Indicators by Clinic
Combined 3 Years

Comparison of Maternal Health Indicators (2, 3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Comanche	24	* *	* *	22	* *	* *	6	* *	7	* *	20	* *	20	* *
Cowley-Ark	551	3.8 (18)	53.2 (33)	494	22.1 (51)	48.6 (17)	87	* *	69	* *	473	4.7 (50)	473	8.5(55)
Cowley-Win	458	6.6 (59)	52.6 (26)	421	19.2 (33)	56.1 (60)	58	* *	60	* *	404	1.7 (5)	405	6.2(25)
Crawford-G	2	* *	* *	1	* *	* *	1	* *	0	* *	1	* *	1	* *
Crawford-P	1,127	5.9 (51)	53.3 (35)	977	18.8 (29)	55.8 (59)	175	16.6 (2)	190	18.4 (8)	976	4.4 (48)	975	7.3(44)
Decatur	30	* *	* *	28	* *	* *	4	* *	5	* *	27	* *	27	* *
DickinsonA	296	4.7 (39)	51.0 (16)	272	19.9 (39)	52.2 (36)	36	* *	70	* *	264	5.3 (59)	260	7.7(46)
DickinsonH	44	* *	* *	40	* *	* *	9	* *	14	* *	40	* *	40	* *
Doniphan	119	2.5 (5)	63.9 (70)	97	* *	* *	25	* *	24	* *	98	* *	98	* *
Douglas	1,421	4.4 (29)	49.3 (7)	1,246	17.3 (11)	56.1 (60)	202	31.2 (16)	403	24.8 (15)	1,199	3.9 (36)	1,198	5.2(13)
Edwards	69	* *	* *	59	* *	* *	8	* *	16	* *	55	* *	55	* *
Elk	47	* *	* *	44	* *	* *	1	* *	9	* *	42	* *	42	* *
Ellis	453	4.4 (29)	49.0 (6)	397	18.6 (25)	49.4 (21)	106	23.6 (4)	126	16.7 (4)	394	2.5 (14)	394	5.6(18)
Ellsworth	94	* *	* *	91	* *	* *	9	* *	23	* *	84	* *	84	* *
Finney	1,987	6.7 (60)	50.1 (11)	1,839	23.3 (58)	45.7 (7)	270	28.1 (12)	289	24.9 (18)	1,805	5.0 (53)	1,803	4.0(5)
Ford	1,906	2.4 (4)	54.7 (47)	1,727	21.1 (46)	48.2 (14)	252	21.8 (3)	200	16.0 (2)	1,690	3.3 (24)	1,689	5.1(11)
Franklin	499	3.2 (12)	54.3 (42)	437	16.7 (5)	55.4 (58)	49	* *	133	16.5 (3)	381	2.4 (12)	381	8.4(52)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Maternal Health Indicators by Clinic
Combined 3 Years

Comparison of Maternal Health Indicators (2, 3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy	Hypertension During Pregnancy		
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)			Number	% (Rank)
Geary	1,031	4.6 (35)	51.6 (20)	860	17.6 (13)	54.3 (52)	165	38.8 (20)	193	29.5 (29)	854	3.4 (27)	854	7.8(48)
Gove	28	* *	* *	26	* *	* *	3	* *	10	* *	29	* *	29	* *
Graham	51	* *	* *	44	* *	* *	8	* *	8	* *	46	* *	46	* *
Grant	205	3.9 (20)	66.3 (71)	194	19.6 (38)	53.1 (45)	27	* *	60	* *	189	3.2 (21)	187	3.2(1)
Gray	157	2.5 (5)	57.3 (59)	135	24.4 (60)	49.6 (22)	17	* *	42	* *	129	2.3 (9)	129	7.0(39)
Greeley	29	* *	* *	27	* *	* *	4	* *	6	* *	29	* *	29	* *
Greenwood	137	2.9 (7)	60.6 (66)	128	22.7 (54)	52.3 (37)	16	* *	25	* *	126	4.0 (38)	127	9.4(60)
Hamilton	71	* *	* *	60	* *	* *	9	* *	25	* *	63	* *	63	* *
Harper	144	7.6 (65)	49.3 (7)	123	22.8 (55)	47.2 (11)	16	* *	22	* *	118	3.4 (27)	117	6.0(23)
Harvey	779	4.2 (24)	54.0 (39)	707	22.8 (55)	48.8 (18)	114	30.7 (15)	133	25.6 (21)	679	3.5 (30)	677	6.6(31)
Haskell	61	* *	* *	53	* *	* *	12	* *	22	* *	54	* *	54	* *
Hodgeman	38	* *	* *	38	* *	* *	7	* *	10	* *	36	* *	36	* *
Jackson	214	5.1 (46)	48.6 (5)	193	17.6 (13)	54.9 (54)	34	* *	46	* *	192	2.1 (6)	192	6.8(35)
Jefferson	192	7.3 (63)	55.2 (49)	172	16.9 (8)	55.2 (56)	29	* *	58	* *	169	4.7 (50)	169	6.5(28)
Jewell	39	* *	* *	39	* *	* *	5	* *	5	* *	38	* *	38	* *
Johnson-M	2,470	4.3 (26)	54.7 (47)	2,211	17.5 (12)	51.8 (33)	412	24.3 (6)	673	17.2 (6)	2,155	5.1 (55)	2,154	4.2(6)
Johnson-O	2,889	4.2 (24)	53.2 (33)	2,617	18.8 (29)	50.9 (26)	508	25.4 (8)	959	18.0 (7)	2,547	5.5 (60)	2,545	4.9(8)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Maternal Health Indicators by Clinic
Combined 3 Years

Comparison of Maternal Health Indicators (2, 3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Kearny	60	* *	* *	56	* *	* *	15	* *	19	* *	56	* *	56	* *
Kingman	173	4.6 (35)	45.7 (2)	148	17.6 (13)	54.1 (51)	19	* *	37	* *	140	5.0 (53)	140	7.1(43)
Kiowa	31	* *	* *	26	* *	* *	4	* *	6	* *	28	* *	28	* *
Labette	530	6.8 (61)	55.8 (53)	487	23.4 (59)	51.1 (31)	90	* *	125	23.2 (14)	478	3.3 (24)	478	8.4(52)
Lane	35	* *	* *	30	* *	* *	7	* *	4	* *	30	* *	30	* *
Leavenwort	1,019	4.9 (43)	53.5 (36)	905	18.6 (25)	55.0 (55)	171	29.2 (13)	246	19.5 (10)	857	3.7 (35)	857	5.5(17)
Lincoln	57	* *	* *	54	* *	* *	9	* *	16	* *	51	* *	51	* *
Linn	184	5.4 (49)	58.2 (61)	171	29.8 (67)	43.9 (3)	24	* *	30	* *	168	3.0 (18)	168	4.2(6)
Logan	36	* *	* *	29	* *	* *	6	* *	7	* *	28	* *	28	* *
Lyon	1,004	3.5 (14)	59.2 (63)	913	20.8 (43)	51.0 (29)	143	13.3 (1)	194	17.0 (5)	887	3.3 (24)	887	9.6(61)
Marion	133	9.0 (71)	42.9 (1)	119	18.5 (24)	44.5 (5)	14	* *	33	* *	114	0.9 (3)	114	7.0(39)
Marshall	167	7.2 (62)	50.3 (13)	140	22.9 (57)	47.9 (13)	23	* *	35	* *	137	0.7 (2)	137	5.1(11)
Mc Pherson	399	7.5 (64)	51.9 (22)	367	19.1 (32)	46.6 (10)	62	* *	92	* *	357	2.5 (14)	358	7.0(39)
Meade	136	3.7 (15)	59.6 (64)	116	21.6 (49)	50.9 (26)	18	* *	23	* *	115	5.2 (57)	115	5.2(13)
Miami	395	8.1 (67)	54.2 (40)	362	18.2 (20)	53.3 (47)	78	* *	116	25.0 (19)	352	5.1 (55)	352	6.5(28)
Mitchell	127	3.9 (20)	58.3 (62)	122	28.7 (66)	45.9 (8)	18	* *	37	* *	118	5.9 (63)	118	8.5(55)
Montgome-C	442	4.1 (22)	55.4 (51)	398	22.1 (51)	48.5 (16)	79	* *	121	26.4 (23)	385	3.6 (32)	385	6.2(25)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Maternal Health Indicators by Clinic
Combined 3 Years

Comparison of Maternal Health Indicators (2, 3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Montgome-I	557	8.1 (67)	52.1 (23)	484	21.9 (50)	50.4 (24)	90	* *	157	21.7 (12)	481	2.1 (6)	481	6.9(38)
Morris	79	* *	* *	72	* *	* *	12	* *	19	* *	71	* *	71	* *
Nemaha	84	* *	* *	80	* *	* *	21	* *	29	* *	80	* *	80	* *
Neosho-Ch	390	8.7 (70)	51.0 (16)	367	17.7 (17)	51.0 (29)	63	* *	63	* *	353	4.0 (38)	353	6.8(35)
Neosho-Er	34	* *	* *	33	* *	* *	9	* *	8	* *	34	* *	34	* *
Ness	17	* *	* *	15	* *	* *	1	* *	10	* *	13	* *	13	* *
Norton	85	* *	* *	75	* *	* *	10	* *	19	* *	73	* *	73	* *
Osage	262	6.1 (53)	53.1 (31)	229	17.9 (18)	52.8 (41)	29	* *	59	* *	228	3.9 (36)	228	5.7(20)
Osborne	53	* *	* *	47	* *	* *	6	* *	11	* *	47	* *	47	* *
Ottawa	74	* *	* *	67	* *	* *	9	* *	17	* *	66	* *	66	* *
Pawnee	162	8.0 (66)	55.6 (52)	141	27.0 (64)	46.1 (9)	11	* *	26	* *	142	2.8 (16)	142	4.9(8)
Phillips	105	2.9 (7)	61.0 (67)	93	* *	* *	6	* *	21	* *	90	* *	90	* *
Pottawat-S	60	* *	* *	58	* *	* *	5	* *	36	* *	57	* *	57	* *
Pottawat-W	68	* *	* *	62	* *	* *	7	* *	25	* *	60	* *	60	* *
Pratt	222	3.2 (12)	52.3 (24)	188	17.0 (9)	56.4 (64)	29	* *	61	* *	181	0.6 (1)	181	6.1(24)
Rawlins	30	* *	* *	28	* *	* *	4	* *	11	* *	28	* *	28	* *
Reno	1,419	4.9 (43)	54.2 (40)	1,280	20.1 (41)	51.9 (34)	180	26.7 (9)	314	24.8 (15)	1,216	2.3 (9)	1,216	8.3(51)

- (1) Reporting period is January 1 through December 31.
 - (2) Excludes records with unknown data or errors.
 - (3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
 - (4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
 - (5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
 - (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Maternal Health Indicators by Clinic
Combined 3 Years

Comparison of Maternal Health Indicators (2, 3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Republic	101	5.9 (51)	52.5 (25)	88	* *	* *	15	* *	19	* *	88	* *	88	* *
Rice	205	2.0 (2)	56.6 (56)	181	21.0 (45)	50.8 (25)	24	* *	45	* *	181	5.5 (60)	180	5.6(18)
Riley-Ft	1,426	4.3 (26)	50.0 (10)	1,131	17.0 (9)	56.1 (60)	229	24.5 (7)	281	31.7 (30)	1,097	2.3 (9)	1,096	6.6(31)
Riley-Manh	1,139	4.6 (35)	48.2 (4)	988	15.9 (3)	52.0 (35)	157	24.2 (5)	285	20.4 (11)	957	3.6 (32)	957	6.4(27)
Rooks	129	6.2 (55)	49.6 (9)	119	26.1 (61)	42.9 (2)	17	* *	27	* *	118	4.2 (44)	118	7.6(45)
Rush	35	* *	* *	32	* *	* *	5	* *	8	* *	31	* *	31	* *
Russell	125	6.4 (58)	48.0 (3)	114	28.1 (65)	39.5 (1)	17	* *	32	* *	110	3.6 (32)	110	3.6(4)
Saline	1,415	4.7 (39)	53.6 (38)	1,307	18.4 (23)	52.3 (37)	264	27.7 (10)	340	19.4 (9)	1,277	3.5 (30)	1,281	5.0(10)
Scott	141	4.3 (26)	54.6 (46)	125	17.6 (13)	55.2 (56)	19	* *	27	* *	126	2.4 (12)	126	3.2(1)
Sedgwick-C	3,287	4.7 (39)	52.8 (29)	3,042	19.0 (31)	50.9 (26)	619	35.1 (19)	816	28.9 (28)	2,977	4.2 (44)	2,976	5.7(20)
Sedgwick-M	5,769	4.1 (22)	53.1 (31)	5,320	19.2 (33)	51.5 (32)	1,206	29.9 (14)	1,603	23.0 (13)	5,130	3.4 (27)	5,130	6.8(35)
Sedgwick-S	3,482	6.3 (56)	50.9 (15)	3,231	19.9 (39)	53.0 (44)	728	33.8 (17)	1,002	26.4 (23)	3,134	4.1 (43)	3,137	8.4(52)
Seneca	25	* *	* *	24	* *	* *	4	* *	8	* *	26	* *	26	* *
Seward	1,450	2.9 (7)	54.4 (43)	1,328	21.4 (47)	50.1 (23)	249	27.7 (10)	218	24.8 (15)	1,294	5.2 (57)	1,296	5.3(15)
ShawneeM&I	574	4.4 (29)	52.6 (26)	432	26.4 (62)	44.4 (4)	12	* *	52	* *	424	6.6 (64)	424	3.3(3)
ShawneeMa	3,646	4.8 (42)	52.8 (29)	3,331	18.3 (21)	53.1 (45)	436	41.5 (22)	712	28.8 (27)	3,264	4.7 (50)	3,269	5.4(16)
Sheridan	26	* *	* *	23	* *	* *	5	* *	3	* *	24	* *	24	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Maternal Health Indicators by Clinic
Combined 3 Years

Comparison of Maternal Health Indicators (2, 3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Sherman	187	3.7 (15)	61.5 (68)	164	20.1 (41)	59.1 (66)	14	* *	27	* *	157	4.5 (49)	157	8.9(59)
Smith	61	* *	* *	51	* *	* *	11	* *	20	* *	52	* *	52	* *
Stafford	117	0.9 (1)	63.2 (69)	102	26.5 (63)	45.1 (6)	13	* *	15	* *	97	* *	97	* *
Stanton	56	* *	* *	54	* *	* *	7	* *	21	* *	54	* *	54	* *
Stevens	247	2.0 (2)	60.3 (65)	220	16.8 (6)	49.1 (20)	38	* *	63	* *	213	8.9 (66)	213	6.6(31)
Sumner	433	5.3 (48)	51.0 (16)	385	19.2 (33)	48.3 (15)	77	* *	113	12.4 (1)	378	4.0 (38)	378	9.8(63)
Thomas	155	8.4 (69)	53.5 (36)	141	15.6 (2)	53.9 (49)	23	* *	37	* *	139	2.9 (17)	139	7.9(50)
Trego	47	* *	* *	41	* *	* *	6	* *	9	* *	45	* *	45	* *
Wabaunsee	61	* *	* *	58	* *	* *	5	* *	25	* *	57	* *	57	* *
Wallace	4	* *	* *	4	* *	* *	1	* *	1	* *	3	* *	3	* *
Washington	81	* *	* *	71	* *	* *	4	* *	19	* *	70	* *	69	* *
Wichita	46	* *	* *	42	* *	* *	7	* *	10	* *	41	* *	41	* *
Wilson	279	6.1 (53)	56.6 (56)	265	18.1 (19)	52.8 (41)	46	* *	50	* *	255	4.3 (47)	255	7.8(48)
Woodson	24	* *	* *	23	* *	* *	4	* *	5	* *	24	* *	24	* *
Wyandotte	6,336	4.4 (29)	54.4 (43)	5,782	18.7 (27)	52.3 (37)	1,060	34.3 (18)	1,196	26.3 (22)	5,623	3.1 (20)	5,625	5.7 (20)
Unknown Clinics	3	* *	* *	3	* *	* *	0	* *	1	* *	3	* *	3	* *
Kansas	61,869	4.7	53.2	55,802	19.5	51.6	9,864	30.1	14,091	23.5	54,295	3.8	54,295	6.4

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6A
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Maternal Behavioral Indicators by Clinic
Combined 3 Years

Comparison of Maternal Behavioral Indicators (2, 3)										
Clinic	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	% (Rank)		Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Allen	265	91.7 (2)	1.9 (16)		261	42.5 (30)	160	13.1 (27)	162	54.9 (43)
Anderson	138	89.1 (11)	2.2 (21)		133	45.1 (21)	88	* *	86	* *
Atchison	367	89.6 (9)	2.5 (25)		366	35.8 (57)	236	7.6 (42)	221	62.9 (27)
Barber-Kio	4	* *	* *		5	* *	2	* *	2	* *
Barber-Med	94	* *	* *		95	* *	68	* *	65	* *
Barton	714	80.0 (31)	1.1 (9)		698	54.6 (5)	414	19.8 (5)	398	60.8 (32)
Bourbon	335	91.9 (1)	1.5 (12)		330	46.4 (18)	194	10.3 (37)	199	62.3 (30)
Brown	251	79.3 (35)	10.4 (61)		257	42.4 (31)	163	11.7 (32)	197	69.0 (8)
Butler	716	70.8 (58)	10.9 (62)		638	43.9 (23)	436	15.1 (18)	364	41.2 (50)
Chase	34	* *	* *		34	* *	25	* *	21	* *
Chautauqua	79	* *	* *		80	* *	48	* *	55	* *
Cherokee	463	77.3 (41)	6.7 (52)		515	39.6 (44)	327	13.1 (27)	296	58.4 (38)
Cheyenne	4	* *	* *		5	* *	3	* *	2	* *
Clark	6	* *	* *		8	* *	4	* *	5	* *
Clay	164	80.5 (29)	6.1 (49)		159	47.8 (15)	94	* *	92	* *
Cloud	214	83.2 (21)	2.8 (27)		205	49.8 (10)	133	16.5 (14)	113	59.3 (35)
Coffey	64	* *	* *		47	* *	39	* *	27	* *
Comanche	22	* *	* *		20	* *	14	* *	14	* *
Cowley-Ark	455	61.1 (67)	7.9 (56)		477	41.5 (36)	320	10.6 (36)	309	57.9 (40)
Cowley-Win	411	76.4 (45)	2.9 (29)		393	49.1 (12)	237	14.3 (25)	226	64.2 (22)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6A
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Maternal Behavioral Indicators by Clinic
Combined 3 Years

Comparison of Maternal Behavioral Indicators (2, 3)										
Clinic	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Crawford-G	1	* *	* *	* *	2	* *	1	* *	1	* *
Crawford-P	974	86.9 (12)	2.1 (19)		1,014	43.9 (23)	635	11.3 (33)	652	66.1 (15)
Decatur	22	* *	* *	* *	27	* *	15	* *	17	* *
DickinsonA	278	83.1 (23)	3.6 (36)		255	49.0 (13)	184	19.6 (6)	149	67.8 (10)
DickinsonH	40	* *	* *	* *	39	* *	18	* *	24	* *
Doniphan	106	78.3 (39)	5.7 (48)		103	39.8 (42)	64	* *	72	* *
Douglas	1,243	72.3 (54)	6.9 (55)		1,242	39.3 (45)	765	17.8 (9)	719	63.4 (25)
Edwards	56	* *	* *	* *	59	* *	44	* *	35	* *
Elk	42	* *	* *	* *	40	* *	25	* *	20	* *
Ellis	418	85.4 (17)	1.9 (16)		406	33.3 (61)	275	25.8 (1)	262	74.8 (2)
Ellsworth	92	* *	* *	* *	73	* *	49	* *	40	* *
Finney	1,721	64.4 (63)	18.3 (66)		1,741	37.7 (50)	1,131	10.3 (37)	1,052	39.5 (51)
Ford	1,668	72.1 (55)	14.0 (65)		1,712	45.7 (19)	1,033	6.8 (44)	1,074	57.3 (41)
Franklin	254	79.5 (33)	3.1 (30)		414	41.8 (35)	278	3.2 (49)	292	68.2 (9)
Geary	822	70.7 (59)	0.7 (3)		923	40.4 (41)	588	17.7 (10)	569	60.8 (32)
Gove	28	* *	* *	* *	22	* *	15	* *	12	* *
Graham	46	* *	* *	* *	44	* *	21	* *	23	* *
Grant	183	77.0 (42)	9.8 (60)		166	36.1 (56)	107	11.2 (34)	94	* *
Gray	133	79.7 (32)	2.3 (22)		133	43.6 (25)	77	* *	75	* *
Greeley	29	* *	* *	* *	27	* *	22	* *	17	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6A
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Maternal Behavioral Indicators by Clinic
Combined 3 Years

Comparison of Maternal Behavioral Indicators (2, 3)											
Clinic	Medical Care				WIC Enrollment			Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester			Prepregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	% (Rank)		Number	% (Rank)		Number	% (Rank)	Number	% (Rank)
Greenwood	118	61.9 (65)	29.7 (67)		124	54.8 (4)		69	* *	73	* *
Hamilton	61	* *	* *		62	* *		41	* *	40	* *
Harper	120	72.5 (52)	12.5 (63)		129	48.8 (14)		83	* *	85	* *
Harvey	659	72.1 (55)	13.8 (64)		688	47.2 (17)		422	14.7 (21)	417	70.5 (7)
Haskell	52	* *	* *		57	* *		36	* *	35	* *
Hodgeman	38	* *	* *		29	* *		23	* *	19	* *
Jackson	139	81.3 (28)	5.0 (46)		188	42.6 (29)		114	8.8 (39)	115	80.9 (1)
Jefferson	162	90.7 (4)	1.9 (16)		153	39.2 (46)		104	6.7 (45)	105	54.3 (44)
Jewell	38	* *	* *		31	* *		17	* *	21	* *
Johnson-M	2,113	66.7 (60)	1.8 (15)		2,098	32.7 (62)		1,304	14.4 (24)	1,216	72.0 (5)
Johnson-O	2,432	61.6 (66)	4.2 (41)		2,519	30.6 (65)		1,634	12.2 (31)	1,511	63.1 (26)
Kearny	54	* *	* *		53	* *		29	* *	27	* *
Kingman	142	81.7 (26)	4.9 (45)		147	49.7 (11)		98	* *	102	71.6 (6)
Kiowa	28	* *	* *		32	* *		22	* *	23	* *
Labette	434	85.5 (16)	3.2 (31)		471	42.0 (33)		286	15.4 (16)	300	63.7 (24)
Lane	31	* *	* *		34	* *		20	* *	21	* *
Leavenworth	915	85.4 (17)	0.7 (3)		823	36.5 (55)		561	18.9 (7)	485	72.4 (4)
Lincoln	55	* *	* *		46	* *		35	* *	31	* *
Linn	170	79.4 (34)	6.5 (50)		164	45.1 (21)		96	* *	109	66.1 (15)
Logan	14	* *	* *		32	* *		23	* *	23	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6A
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Maternal Behavioral Indicators by Clinic
Combined 3 Years

Comparison of Maternal Behavioral Indicators (2, 3)										
Clinic	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	% (Rank)		Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Lyon	927	81.6 (27)	1.1 (9)		841	41.5 (36)	516	14.9 (19)	476	64.7 (20)
Marion	120	72.5 (52)	3.3 (32)		109	42.2 (32)	74	* *	62	* *
Marshall	147	90.5 (5)	0.7 (3)		138	42.0 (33)	81	* *	72	* *
Mc Pherson	372	86.6 (14)	1.3 (11)		334	38.6 (48)	218	20.2 (4)	206	67.0 (12)
Meade	119	75.6 (49)	6.7 (52)		115	52.2 (7)	70	* *	72	* *
Miami	371	77.9 (40)	0.8 (6)		323	32.5 (63)	220	15.5 (15)	195	58.5 (37)
Mitchell	121	91.7 (2)	0.0 (1)		110	34.5 (59)	78	* *	72	* *
Montgome-C	338	80.5 (29)	5.6 (47)		384	37.2 (51)	241	16.6 (12)	233	67.8 (10)
Montgome-I	436	86.7 (13)	3.9 (37)		485	39.8 (42)	289	12.5 (30)	307	65.8 (19)
Morris	69	* *	* *		67	* *	37	* *	36	* *
Nemaha	84	* *	* *		72	* *	47	* *	46	* *
Neosho-Ch	346	89.3 (10)	4.3 (42)		348	42.8 (28)	222	7.2 (43)	205	66.8 (14)
Neosho-Er	31	* *	* *		34	* *	16	* *	18	* *
Ness	17	* *	* *		13	* *	10	* *	4	* *
Norton	79	* *	* *		68	* *	46	* *	41	* *
Osage	214	89.7 (8)	2.3 (22)		216	51.4 (8)	146	21.2 (3)	136	41.9 (49)
Osborne	48	* *	* *		43	* *	20	* *	27	* *
Ottawa	69	* *	* *		63	* *	39	* *	37	* *
Pawnee	144	81.9 (25)	2.8 (27)		148	57.4 (3)	90	* *	87	* *
Phillips	96	* *	* *		88	* *	56	* *	71	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6A
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Maternal Behavioral Indicators by Clinic
Combined 3 Years

Comparison of Maternal Behavioral Indicators (2, 3)										
Clinic	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Pottawat-S	54	* *	* *	* *	45	* *	22	* *	22	* *
Pottawat-W	55	* *	* *	* *	60	* *	35	* *	36	* *
Pratt	194	78.9 (37)	6.7 (52)		188	47.3 (16)	126	7.9 (41)	116	64.7 (20)
Rawlins	12	* *	* *	* *	27	* *	18	* *	19	* *
Reno	1,267	83.4 (20)	2.1 (19)		1,195	41.2 (39)	790	21.3 (2)	753	65.9 (18)
Republic	91	* *	* *	* *	92	* *	56	* *	51	* *
Rice	184	76.1 (46)	1.6 (14)		181	40.9 (40)	117	14.5 (23)	104	43.3 (48)
Riley-Ft	985	90.1 (7)	0.4 (2)		1,207	32.1 (64)	794	14.7 (21)	771	62.8 (28)
Riley-Manh	953	76.9 (43)	0.8 (6)		971	41.5 (36)	631	17.6 (11)	587	64.2 (22)
Rooks	119	83.2 (21)	3.4 (34)		113	43.4 (26)	67	* *	68	* *
Rush	32	* *	* *	* *	32	* *	17	* *	18	* *
Russell	116	90.5 (5)	0.9 (8)		105	51.4 (8)	65	* *	65	* *
Saline	1,209	74.0 (51)	4.8 (44)		1,205	33.9 (60)	802	5.2 (48)	775	57.0 (42)
Scott	127	78.7 (38)	8.7 (59)		122	36.9 (54)	69	* *	81	* *
Sedgwick-C	3,040	75.9 (47)	4.3 (42)		2,882	30.4 (66)	1,845	13.4 (26)	1,795	52.6 (46)
Sedgwick-M	5,376	71.5 (57)	3.9 (37)		4,857	26.5 (68)	3,141	12.9 (29)	2,760	52.9 (45)
Sedgwick-S	3,227	79.1 (36)	2.4 (24)		3,093	29.0 (67)	2,041	14.9 (19)	1,889	46.1 (47)
Seneca	27	* *	* *	* *	24	* *	12	* *	13	* *
Seward	1,260	66.4 (62)	8.4 (57)		1,279	37.1 (52)	799	8.8 (39)	740	66.9 (13)
ShawneeM&I	419	64.4 (63)	3.3 (32)		542	64.6 (1)	364	6.0 (47)	400	58.0 (39)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6A
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Maternal Behavioral Indicators by Clinic
Combined 3 Years

Comparison of Maternal Behavioral Indicators (2, 3)										
Clinic	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	% (Rank)		Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
ShawneeMa	3,183	76.8 (44)	3.9 (37)		3,164	45.7 (19)	1,931	6.7 (45)	1,890	58.6 (36)
Sheridan	23	* *	* *		20	* *	12	* *	12	* *
Sherman	106	84.9 (19)	8.5 (58)		168	52.4 (6)	99	* *	108	74.1 (3)
Smith	42	* *	* *		52	* *	29	* *	37	* *
Stafford	103	75.7 (48)	3.9 (37)		105	62.9 (2)	68	* *	72	* *
Stanton	54	* *	* *		47	* *	25	* *	22	* *
Stevens	224	74.6 (50)	2.7 (26)		210	35.2 (58)	127	18.9 (7)	115	60.0 (34)
Sumner	398	82.7 (24)	3.5 (35)		362	38.1 (49)	241	16.6 (12)	209	66.0 (17)
Thomas	98	* *	* *		129	38.8 (47)	83	* *	92	* *
Trego	45	* *	* *		41	* *	24	* *	28	* *
Wabaunsee	55	* *	* *		46	* *	35	* *	31	* *
Wallace	4	* *	* *		2	* *	1	* *	2	* *
Washington	74	* *	* *		74	* *	42	* *	47	* *
Wichita	42	* *	* *		40	* *	24	* *	24	* *
Wilson	260	85.8 (15)	1.5 (12)		254	42.9 (27)	151	15.2 (17)	156	62.8 (28)
Woodson	23	* *	* *		20	* *	8	* *	9	* *
Wyandotte	5,808	66.5 (61)	6.6 (51)		5,566	37.0 (53)	3,457	11.0 (35)	3,286	61.5 (31)
Unknown Clinics	3	* *	* *		3	* *	3	* *	1	* *
Kansas	54,171	75.0	4.9		53,570	38.0	34,056	13.7	32,418	60.0

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2007 - 2009 Pregnancy Nutrition Surveillance
Kansas
Comparison of Smoking Indicators by Clinic
Combined 3 Years

(1)

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Allen	291	44.3 (44)	249	28.5 (45)	82	* *	250	37.2 (55)	247	32.8 (53)
Anderson	148	56.8 (67)	126	32.5 (54)	57	* *	127	27.6 (43)	128	22.7 (42)
Atchison	298	47.0 (49)	195	28.7 (46)	65	* *	264	28.0 (44)	202	24.8 (44)
Barber-Kio	5	* *	4	* *	1	* *	5	* *	4	* *
Barber-Med	110	50.0 (57)	92	* *	40	* *	91	* *	93	* *
Barton	803	37.6 (23)	714	23.7 (27)	222	29.3 (18)	657	11.1 (12)	716	5.6 (6)
Bourbon	373	50.1 (58)	311	28.9 (47)	113	26.5 (25)	307	34.5 (54)	312	34.0 (55)
Brown	274	42.3 (38)	208	25.5 (31)	66	* *	234	44.4 (60)	210	44.8 (61)
Butler	771	50.5 (59)	639	31.0 (53)	235	30.6 (15)	558	14.3 (20)	641	8.6 (15)
Chase	38	* *	34	* *	16	* *	30	* *	34	* *
Chautauqua	78	* *	52	* *	15	* *	68	* *	52	* *
Cherokee	587	39.2 (27)	489	26.8 (35)	138	20.3 (30)	448	31.7 (51)	503	30.2 (52)
Cheyenne	6	* *	6	* *	0	* *	4	* *	6	* *
Clark	2	* *	1	* *	0	* *	5	* *	3	* *
Clay	181	54.7 (64)	157	32.5 (54)	74	* *	154	22.1 (34)	159	15.7 (25)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2007 - 2009 Pregnancy Nutrition Surveillance
Kansas
Comparison of Smoking Indicators by Clinic
Combined 3 Years

(1)

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Cloud	235	38.3 (25)	200	21.0 (23)	60	* *	193	6.7 (2)	202	2.0 (1)
Coffey	65	* *	62	* *	21	* *	38	* *	63	* *
Comanche	22	* *	21	* *	8	* *	18	* *	21	* *
Cowley-Ark	528	37.9 (24)	414	26.1 (33)	128	25.8 (27)	438	20.1 (30)	420	19.0 (34)
Cowley-Win	453	52.3 (61)	388	34.8 (60)	169	27.2 (23)	372	18.5 (27)	388	18.6 (31)
Crawford-G	2	* *	1	* *	0	* *	2	* *	1	* *
Crawford-P	1,085	43.4 (41)	865	25.5 (31)	304	34.9 (9)	935	40.1 (58)	867	39.3 (58)
Decatur	30	* *	27	* *	7	* *	24	* *	27	* *
DickinsonA	296	42.9 (39)	268	28.0 (43)	87	* *	223	9.9 (10)	270	7.0 (9)
DickinsonH	44	* *	37	* *	17	* *	38	* *	38	* *
Doniphan	98	* *	62	* *	23	* *	76	* *	65	* *
Douglas	1,407	43.7 (42)	1,195	20.6 (22)	392	42.9 (4)	1,121	10.1 (11)	1,200	6.5 (8)
Edwards	69	* *	59	* *	15	* *	54	* *	60	* *
Elk	47	* *	41	* *	12	* *	35	* *	40	* *
Ellis	463	37.1 (21)	396	14.1 (12)	117	48.7 (1)	366	8.5 (6)	394	7.4 (12)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2007 - 2009 Pregnancy Nutrition Surveillance
Kansas
Comparison of Smoking Indicators by Clinic
Combined 3 Years

(1)

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Ellsworth	91	* *	85	* *	29	* *	67	* *	87	* *
Finney	1,979	13.2 (5)	1,796	6.6 (4)	179	40.8 (6)	1,625	17.4 (25)	1,797	16.0 (26)
Ford	1,855	10.0 (4)	1,531	4.9 (3)	105	28.6 (20)	1,618	8.9 (7)	1,532	8.3 (14)
Franklin	504	46.0 (47)	444	27.3 (39)	136	31.6 (13)	378	14.8 (21)	446	10.5 (19)
Geary	1,032	39.2 (27)	842	20.4 (21)	265	36.6 (7)	850	39.8 (57)	844	37.0 (57)
Gove	30	* *	28	* *	3	* *	18	* *	28	* *
Graham	50	* *	43	* *	22	* *	41	* *	44	* *
Grant	207	7.7 (2)	199	4.0 (2)	9	* *	152	12.5 (15)	199	12.6 (23)
Gray	154	9.7 (3)	129	7.0 (5)	7	* *	127	11.8 (13)	132	9.8 (17)
Greeley	31	* *	29	* *	6	* *	22	* *	29	* *
Greenwood	124	54.8 (65)	72	* *	31	* *	110	31.8 (52)	104	26.0 (46)
Hamilton	72	* *	63	* *	3	* *	55	* *	63	* *
Harper	129	48.8 (55)	85	* *	32	* *	109	22.0 (33)	84	* *
Harvey	773	35.1 (19)	658	22.5 (24)	189	30.2 (16)	638	5.8 (1)	662	4.1 (3)
Haskell	61	* *	56	* *	7	* *	53	* *	56	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2007 - 2009 Pregnancy Nutrition Surveillance
Kansas
Comparison of Smoking Indicators by Clinic
Combined 3 Years

(1)

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Hodgeman	37	* *	34	* *	11	* *	25	* *	33	* *
Jackson	212	48.1 (54)	184	29.3 (50)	67	* *	176	19.9 (28)	187	18.2 (30)
Jefferson	70	* *	28	* *	9	* *	69	* *	32	* *
Jewell	39	* *	37	* *	11	* *	32	* *	37	* *
Johnson-M	2,471	26.5 (12)	2,156	10.8 (7)	394	44.4 (3)	1,894	20.6 (31)	2,168	20.4 (37)
Johnson-O	2,927	23.4 (8)	2,590	10.6 (6)	418	44.5 (2)	2,319	16.7 (24)	2,593	16.5 (28)
Kearny	60	* *	57	* *	8	* *	49	* *	57	* *
Kingman	146	40.4 (31)	118	19.5 (20)	29	* *	139	39.6 (56)	133	36.1 (56)
Kiowa	30	* *	21	* *	4	* *	25	* *	23	* *
Labette	521	38.6 (26)	455	25.1 (30)	133	15.8 (33)	443	28.0 (44)	467	27.0 (48)
Lane	18	* *	9	* *	1	* *	17	* *	10	* *
Leavenworth	1,010	47.6 (51)	853	27.1 (38)	281	29.9 (17)	742	42.7 (59)	862	40.5 (59)
Lincoln	56	* *	43	* *	16	* *	42	* *	43	* *
Linn	185	56.2 (66)	166	40.4 (62)	76	* *	155	14.2 (19)	168	10.7 (21)
Logan	36	* *	29	* *	9	* *	30	* *	29	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2007 - 2009 Pregnancy Nutrition Surveillance
Kansas
Comparison of Smoking Indicators by Clinic
Combined 3 Years

(1)

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Lyon	1,011	29.1 (14)	909	15.3 (14)	188	35.6 (8)	778	13.2 (17)	911	10.6 (20)
Marion	135	25.9 (11)	119	19.3 (19)	22	* *	93	* *	120	5.0 (5)
Marshall	166	50.6 (60)	132	32.6 (56)	46	* *	128	52.3 (63)	137	49.6 (62)
Mc Pherson	400	40.3 (30)	367	26.2 (34)	120	26.7 (24)	309	7.1 (3)	370	4.1 (3)
Meade	125	30.4 (16)	104	18.3 (17)	23	* *	106	22.6 (40)	106	19.8 (36)
Miami	359	47.9 (52)	287	30.3 (52)	99	* *	265	21.5 (32)	286	16.8 (29)
Mitchell	127	39.4 (29)	119	26.9 (36)	40	* *	101	8.9 (7)	121	9.1 (16)
Montgome-C	437	41.6 (37)	373	29.2 (48)	121	17.4 (31)	359	30.9 (50)	376	28.5 (50)
Montgome-I	538	40.5 (34)	442	29.2 (48)	137	16.1 (32)	445	29.0 (47)	451	27.7 (49)
Morris	78	* *	68	* *	24	* *	61	* *	68	* *
Nemaha	86	* *	76	* *	28	* *	65	* *	75	* *
Neosho-Ch	359	45.1 (45)	267	27.3 (39)	97	* *	304	24.3 (41)	272	22.4 (41)
Neosho-Er	34	* *	25	* *	5	* *	31	* *	25	* *
Ness	17	* *	15	* *	5	* *	11	* *	15	* *
Norton	84	* *	73	* *	30	* *	64	* *	74	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2007 - 2009 Pregnancy Nutrition Surveillance
Kansas
Comparison of Smoking Indicators by Clinic
Combined 3 Years

(1)

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Osage	252	46.0 (47)	210	29.5 (51)	73	* *	205	28.3 (46)	211	26.1 (47)
Osborne	52	* *	47	* *	18	* *	40	* *	47	* *
Ottawa	71	* *	58	* *	20	* *	58	* *	60	* *
Pawnee	161	47.2 (50)	138	27.5 (41)	49	* *	136	19.9 (28)	138	18.8 (32)
Phillips	81	* *	54	* *	18	* *	62	* *	54	* *
Pottawat-S	59	* *	54	* *	9	* *	40	* *	55	* *
Pottawat-W	70	* *	59	* *	20	* *	57	* *	61	* *
Pratt	223	41.3 (36)	167	28.1 (44)	60	* *	165	32.7 (53)	187	28.9 (51)
Rawlins	30	* *	27	* *	6	* *	26	* *	27	* *
Reno	1,401	45.9 (46)	1,214	24.9 (29)	424	34.4 (11)	1,107	17.5 (26)	1,222	16.4 (27)
Republic	100	43.0 (40)	85	* *	29	* *	82	* *	87	* *
Rice	206	49.0 (56)	181	33.7 (58)	68	* *	170	22.4 (36)	183	19.7 (35)
Riley-Ft	1,431	24.7 (9)	1,017	13.9 (11)	167	31.1 (14)	1,085	22.5 (39)	1,113	21.6 (39)
Riley-Manh	1,119	31.5 (18)	870	18.9 (18)	186	25.3 (29)	891	7.2 (4)	923	5.6 (6)
Rooks	127	59.1 (68)	107	38.3 (61)	53	* *	102	25.5 (42)	111	33.3 (54)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2007 - 2009 Pregnancy Nutrition Surveillance
Kansas
Comparison of Smoking Indicators by Clinic
Combined 3 Years

(1)

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Rush	34	* *	31	* *	14	* *	31	* *	31	* *
Russell	125	48.0 (53)	112	42.9 (63)	45	* *	98	* *	114	50.9 (64)
Saline	1,303	44.2 (43)	1,029	26.9 (36)	348	27.3 (22)	1,040	13.2 (17)	1,034	10.3 (18)
Scott	136	25.7 (10)	115	15.7 (16)	25	* *	119	29.4 (49)	119	22.7 (42)
Sedgwick-C	3,294	29.6 (15)	3,021	15.4 (15)	679	34.9 (9)	2,672	12.5 (15)	3,007	11.7 (22)
Sedgwick-M	5,751	28.6 (13)	5,197	13.1 (10)	1,027	41.6 (5)	4,440	8.2 (5)	5,205	7.1 (10)
Sedgwick-S	3,485	40.7 (35)	3,147	22.8 (25)	995	29.3 (18)	2,815	9.4 (9)	3,153	7.1 (10)
Seneca	27	* *	25	* *	13	* *	23	* *	25	* *
Seward	1,428	7.5 (1)	1,278	3.5 (1)	69	* *	1,188	15.7 (22)	1,292	14.4 (24)
ShawneeM&I	525	30.9 (17)	369	15.2 (13)	89	* *	495	16.2 (23)	372	3.5 (2)
ShawneeMa	3,465	37.3 (22)	2,795	23.2 (26)	767	25.4 (28)	2,836	12.4 (14)	2,805	8.1 (13)
Sheridan	22	* *	19	* *	2	* *	18	* *	19	* *
Sherman	183	40.4 (31)	155	27.7 (42)	54	* *	156	22.4 (36)	154	22.1 (40)
Smith	54	* *	36	* *	11	* *	49	* *	39	* *
Stafford	114	40.4 (31)	93	* *	33	* *	96	* *	95	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2007 - 2009 Pregnancy Nutrition Surveillance
Kansas
Comparison of Smoking Indicators by Clinic
Combined 3 Years

(1)

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Stanton	54	* *	50	* *	3	* *	41	* *	51	* *
Stevens	247	21.9 (7)	214	11.2 (8)	36	* *	192	22.4 (36)	217	18.9 (33)
Sumner	418	53.6 (63)	357	33.9 (59)	147	28.6 (20)	316	49.1 (62)	362	50.0 (63)
Thomas	156	36.5 (20)	134	24.6 (28)	33	* *	117	29.1 (48)	135	25.9 (45)
Trego	46	* *	33	* *	13	* *	39	* *	35	* *
Wabaunsee	61	* *	58	* *	17	* *	41	* *	58	* *
Wallace	2	* *	2	* *	1	* *	2	* *	2	* *
Washington	83	* *	72	* *	17	* *	67	* *	71	* *
Wichita	46	* *	37	* *	5	* *	37	* *	37	* *
Wilson	276	53.3 (62)	247	33.6 (57)	113	32.7 (12)	239	45.6 (61)	250	41.6 (60)
Woodson	24	* *	21	* *	6	* *	18	* *	22	* *
Wyandotte	6,305	21.6 (6)	5,597	12.7 (9)	906	26.3 (26)	5,183	22.2 (35)	5,641	21.0 (38)
Unknown Clinics	3	* *	3	* *	1	* *	1	* *	3	* *
Kansas	60,725	32.6	52,064	18.3	12,628	31.8	48,870	18.4	52,549	16.3

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Infant Health Indicators by Clinic
Combined 3 Years

Comparison of Infant Health Indicators (2, 3)									
Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)	High (5)							
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Allen	274	9.5 (64)	4.0 (7)	239	12.6 (53)	207	5.3 (50)	273	52.4 (63)
Anderson	132	9.1 (55)	5.3 (20)	122	9.0 (20)	101	7.9 (57)	137	53.3 (62)
Atchison	361	6.9 (27)	5.5 (22)	328	8.5 (13)	296	3.4 (23)	364	52.2 (64)
Barber-Kio	4	* * * *	*	4	* *	4	* *	4	* *
Barber-Med	89	* * * *	*	77	* *	61	* *	88	* *
Barton	693	6.5 (22)	4.6 (11)	607	12.2 (51)	529	3.0 (14)	675	73.5 (17)
Bourbon	343	7.6 (40)	5.2 (19)	293	11.6 (46)	257	3.9 (37)	339	58.7 (58)
Brown	223	7.2 (31)	7.2 (48)	231	7.8 (8)	195	3.6 (30)	235	50.6 (67)
Butler	727	8.1 (43)	5.6 (23)	575	10.4 (38)	509	3.7 (31)	700	72.1 (22)
Chase	33	* * * *	*	29	* *	27	* *	32	* *
Chautauqua	74	* * * *	*	72	* *	63	* *	73	* *
Cherokee	549	5.8 (12)	7.7 (57)	446	9.9 (29)	402	2.2 (6)	525	59.6 (56)
Cheyenne	5	* * * *	*	4	* *	4	* *	5	* *
Clark	7	* * * *	*	5	* *	4	* *	6	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Infant Health Indicators by Clinic
Combined 3 Years

Comparison of Infant Health Indicators (2, 3)										
Clinic	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)		Number	%	Number	%	Number	%
	Number	% (Rank)	% (Rank)	Number						
Clay	165	7.3 (33)	3.6 (4)	140	13.6 (55)	121	3.3 (21)	162	62.3 (48)	
Cloud	206	8.3 (46)	2.4 (1)	177	15.3 (61)	149	4.0 (40)	199	66.8 (38)	
Coffey	61	* * * *	* * * *	41	* *	38	* *	62	* *	
Comanche	22	* * * *	* * * *	18	* *	15	* *	22	* *	
Cowley-Ark	485	6.6 (25)	7.2 (48)	419	9.1 (21)	372	4.0 (40)	469	52.2 (64)	
Cowley-Win	409	9.3 (59)	7.3 (50)	352	13.1 (54)	303	5.9 (54)	406	65.3 (42)	
Crawford-G	1	* * * *	* * * *	1	* *	1	* *	1	* *	
Crawford-P	963	9.3 (59)	4.9 (13)	869	12.1 (50)	751	4.5 (46)	957	58.9 (57)	
Decatur	26	* * * *	* * * *	25	* *	24	* *	25	* *	
DickinsonA	263	7.2 (31)	9.9 (67)	225	8.9 (19)	203	3.4 (23)	262	73.3 (19)	
DickinsonH	38	* * * *	* * * *	35	* *	33	* *	39	* *	
Doniphan	104	5.8 (12)	5.8 (27)	87	* *	78	* *	102	52.0 (66)	
Douglas	1,234	6.1 (15)	6.6 (34)	1,060	10.6 (41)	933	3.0 (14)	1,188	77.2 (11)	
Edwards	55	* * * *	* * * *	50	* *	40	* *	56	* *	

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Infant Health Indicators by Clinic
Combined 3 Years

Comparison of Infant Health Indicators (2, 3)									
Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)	High (5)							
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Elk	45	* * * *		37	* *	31	* *	43	* *
Ellis	386	5.7 (10)	5.7 (24)	350	8.3 (11)	300	1.7 (3)	395	74.9 (14)
Ellsworth	89	* * * *		69	* *	61	* *	86	* *
Finney	1,799	6.4 (19)	6.6 (34)	1,587	10.8 (43)	1,388	2.8 (10)	1,347	74.8 (15)
Ford	1,658	5.1 (6)	5.7 (24)	1,519	10.4 (38)	1,335	1.9 (4)	1,467	72.1 (22)
Franklin	438	6.2 (17)	5.0 (15)	350	10.3 (35)	311	3.2 (19)	425	60.7 (53)
Geary	881	7.4 (37)	6.6 (34)	755	8.7 (14)	686	3.5 (27)	826	69.5 (31)
Gove	28	* * * *		20	* *	18	* *	26	* *
Graham	39	* * * *		36	* *	31	* *	43	* *
Grant	196	8.7 (51)	7.7 (57)	158	9.5 (25)	140	4.3 (44)	148	70.3 (29)
Gray	114	6.1 (15)	7.9 (61)	111	11.7 (48)	89	* *	124	87.1 (1)
Greeley	28	* * * *		24	* *	23	* *	27	* *
Greenwood	121	6.6 (25)	3.3 (2)	109	10.1 (31)	97	* *	116	60.3 (54)
Hamilton	59	* * * *		51	* *	46	* *	47	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Infant Health Indicators by Clinic
Combined 3 Years

Comparison of Infant Health Indicators (2, 3)									
Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)	High (5)		Number	%	Number	%	Number	%
	Number	% (Rank)	% (Rank)						
Harper	107	6.5 (22)	7.5 (52)	109	10.1 (31)	85	* *	121	76.9 (12)
Harvey	668	7.3 (33)	7.0 (44)	604	12.3 (52)	518	3.3 (21)	680	78.8 (5)
Haskell	56	* *	* *	51	* *	47	* *	39	* *
Hodgeman	38	* *	* *	28	* *	25	* *	36	* *
Jackson	196	9.2 (56)	9.7 (66)	167	14.4 (58)	143	5.6 (52)	190	78.4 (7)
Jefferson	172	8.1 (43)	7.6 (55)	138	16.7 (63)	114	5.3 (50)	170	61.8 (50)
Jewell	38	* *	* *	31	* *	26	* *	34	* *
Johnson-M	2,101	7.3 (33)	6.9 (41)	1,847	7.9 (9)	1,595	3.4 (23)	2,150	77.5 (8)
Johnson-O	2,527	5.8 (12)	8.1 (62)	2,219	8.0 (10)	1,956	2.7 (9)	2,462	80.6 (2)
Kearny	56	* *	* *	50	* *	43	* *	40	* *
Kingman	106	4.7 (1)	7.5 (52)	120	9.2 (22)	85	* *	134	80.6 (2)
Kiowa	28	* *	* *	27	* *	23	* *	27	* *
Labette	479	6.5 (22)	7.5 (52)	421	10.5 (40)	377	3.7 (31)	456	54.6 (61)
Lane	28	* *	* *	27	* *	22	* *	29	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Infant Health Indicators by Clinic
Combined 3 Years

Comparison of Infant Health Indicators (2, 3)										
Clinic	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)							
	Number	% (Rank)	% (Rank)		Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Leavenwort	862	9.4 (62)	5.0 (15)		713	9.4 (24)	631	4.8 (48)	892	63.2 (47)
Lincoln	54	* *	* *		41	* *	39	* *	51	* *
Linn	166	8.4 (49)	4.8 (12)		150	10.0 (30)	133	6.8 (56)	163	68.7 (35)
Logan	27	* *	* *		25	* *	23	* *	26	* *
Lyon	908	6.9 (27)	6.5 (31)		739	10.1 (31)	663	3.0 (14)	869	71.6 (27)
Marion	110	5.5 (9)	6.4 (30)		88	* *	80	* *	110	77.3 (10)
Marshall	138	9.4 (62)	5.1 (18)		113	10.6 (41)	99	* *	140	56.4 (59)
Mc Pherson	361	5.0 (4)	6.9 (41)		297	7.7 (7)	271	1.1 (2)	358	72.1 (22)
Meade	105	5.7 (10)	6.7 (38)		95	* *	77	* *	106	76.4 (13)
Miami	370	7.6 (40)	5.9 (28)		290	9.7 (27)	260	5.8 (53)	322	72.7 (20)
Mitchell	104	8.7 (51)	6.7 (38)		100	7.0 (5)	83	* *	111	65.8 (41)
Montgome-C	400	7.0 (29)	3.8 (6)		344	11.6 (46)	302	4.0 (40)	384	61.2 (51)
Montgome-I	486	8.6 (50)	3.3 (2)		416	11.3 (45)	368	3.0 (14)	469	59.9 (55)
Morris	69	* *	* *		58	* *	50	* *	69	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Infant Health Indicators by Clinic
Combined 3 Years

Comparison of Infant Health Indicators (2, 3)									
Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)	High (5)		Number	%	Number	%	Number	%
	Number	% (Rank)	% (Rank)						
Nemaha	68	* * * *	* *	68	* *	47	* *	82	* *
Neosho-Ch	275	8.7 (51)	6.5 (31)	322	13.7 (56)	206	3.9 (37)	372	55.1 (60)
Neosho-Er	27	* * * *	* *	33	* *	25	* *	35	* *
Ness	16	* * * *	* *	11	* *	9	* *	16	* *
Norton	75	* * * *	* *	61	* *	47	* *	72	* *
Osage	235	6.4 (19)	4.3 (9)	201	14.9 (60)	167	3.0 (14)	228	61.0 (52)
Osborne	46	* * * *	* *	38	* *	34	* *	45	* *
Ottawa	66	* * * *	* *	57	* *	52	* *	62	* *
Pawnee	137	8.8 (54)	4.4 (10)	127	8.7 (14)	116	2.6 (8)	134	72.4 (21)
Phillips	94	* * * *	* *	77	* *	62	* *	92	* *
Pottawat-S	53	* * * *	* *	41	* *	37	* *	53	* *
Pottawat-W	63	* * * *	* *	53	* *	48	* *	60	* *
Pratt	187	6.4 (19)	9.1 (63)	152	5.9 (4)	143	3.5 (27)	171	68.4 (37)
Rawlins	28	* * * *	* *	24	* *	22	* *	27	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Infant Health Indicators by Clinic
Combined 3 Years

Comparison of Infant Health Indicators (2, 3)										
Clinic	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)							
	Number	% (Rank)	% (Rank)		Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Reno	1,247	7.1 (30)	7.1 (46)		1,048	10.3 (35)	930	3.2 (19)	1,218	63.3 (46)
Republic	85	* *	* *		79	* *	65	* *	89	* *
Rice	181	8.3 (46)	7.7 (57)		157	14.6 (59)	133	3.8 (36)	174	71.8 (26)
Riley-Ft	1,164	5.0 (4)	9.1 (63)		914	5.5 (3)	859	2.3 (7)	1,140	74.2 (16)
Riley-Manh	1,002	6.3 (18)	6.3 (29)		820	8.8 (17)	745	2.8 (10)	969	80.1 (4)
Rooks	121	8.3 (46)	5.0 (15)		106	4.7 (2)	101	5.0 (49)	118	72.0 (25)
Rush	31	* *	* *		28	* *	24	* *	29	* *
Russell	109	9.2 (56)	3.7 (5)		93	* *	81	* *	103	64.1 (44)
Saline	1,283	9.3 (59)	6.5 (31)		1,095	11.0 (44)	957	4.4 (45)	1,285	62.3 (48)
Scott	118	5.1 (6)	4.2 (8)		108	3.7 (1)	102	2.9 (13)	121	69.4 (33)
Sedgwick-C	3,021	7.3 (33)	7.3 (50)		2,620	8.3 (11)	2,385	3.4 (23)	2,915	69.5 (31)
Sedgwick-M	5,281	9.6 (65)	6.6 (34)		4,393	9.3 (23)	3,947	4.7 (47)	4,651	66.4 (40)
Sedgwick-S	3,218	7.4 (37)	7.0 (44)		2,820	8.7 (14)	2,564	3.5 (27)	3,062	64.5 (43)
Seneca	21	* *	* *		23	* *	16	* *	27	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Infant Health Indicators by Clinic
Combined 3 Years

Comparison of Infant Health Indicators (2, 3)										
Clinic	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)		Number	%	Number	%	Number	%
	Number	% (Rank)	% (Rank)	Number						
Seward	1,270	5.2 (8)	7.1 (46)	1,148	9.7 (27)	998	2.8 (10)	1,298	69.4 (33)	
ShawneeM&I	402	9.2 (56)	7.7 (57)	393	15.8 (62)	322	3.7 (31)	396	78.5 (6)	
ShawneeMa	3,207	7.5 (39)	5.3 (20)	2,862	11.8 (49)	2,433	3.7 (31)	3,271	66.6 (39)	
Sheridan	24	* * *	* *	18	* *	15	* *	24	* *	
Sherman	163	11.0 (67)	4.9 (13)	145	7.6 (6)	134	6.7 (55)	155	69.7 (30)	
Smith	51	* * *	* *	43	* *	35	* *	52	* *	
Stafford	94	* * *	* *	90	* *	76	* *	89	* *	
Stanton	50	* * *	* *	46	* *	40	* *	35	* *	
Stevens	143	4.9 (3)	9.1 (63)	181	8.8 (17)	104	1.0 (1)	217	77.4 (9)	
Sumner	393	8.1 (43)	7.6 (55)	324	10.2 (34)	286	4.2 (43)	375	68.5 (36)	
Thomas	145	4.8 (2)	6.9 (41)	115	9.6 (26)	104	1.9 (4)	139	73.4 (18)	
Trego	44	* * *	* *	38	* *	35	* *	43	* *	
Wabaunsee	58	* * *	* *	44	* *	41	* *	55	* *	
Wallace	4	* * *	* *	2	* *	2	* *	4	* *	

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Infant Health Indicators by Clinic
Combined 3 Years

Comparison of Infant Health Indicators (2, 3)										
Clinic	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)							
	Number	% (Rank)	%	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Washington	62	* * * *	*	*	65	* *	50	* *	71	* *
Wichita	42	* * * *	*	*	36	* *	27	* *	42	* *
Wilson	261	9.6 (65)	5.7	(24)	240	14.2 (57)	205	3.9 (37)	235	63.8 (45)
Woodson	23	* * * *	*	*	19	* *	15	* *	22	* *
Wyandotte	5,635	8.0 (42)	6.8	(40)	4,952	10.3 (35)	4,358	3.7 (31)	5,555	70.7 (28)
Unknown Clinics	3	* * * *	*	*	3	* *	3	* *	3	* *
Kansas	54,513	7.4	6.6		47,348	10.0	41,621	3.5	52,595	69.2

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

