

Table 3A
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Clinic	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaska Native	Asian/Pacific Islander	Multiple Races	All Other/Unknown
		%	%	%	%	%	%	%
Allen	84	*	*	*	*	*	*	*
Anderson	56	*	*	*	*	*	*	*
Atchison	140	88.6	9.3	1.4	0.0	0.0	0.7	0.0
Barber-Kio	2	*	*	*	*	*	*	*
Barber-Med	54	*	*	*	*	*	*	*
Barton	255	68.6	2.4	28.2	0.0	0.8	0.0	0.0
Bourbon	127	93.7	1.6	2.4	0.8	0.8	0.8	0.0
Brown	91	*	*	*	*	*	*	*
Butler	285	89.8	1.4	7.0	0.7	0.0	1.1	0.0
Chase	9	*	*	*	*	*	*	*
Chautauqua	19	*	*	*	*	*	*	*
Cherokee	183	90.7	1.6	1.1	1.6	2.7	2.2	0.0
Cheyenne	10	*	*	*	*	*	*	*
Clark	2	*	*	*	*	*	*	*
Clay	51	*	*	*	*	*	*	*
Cloud	93	*	*	*	*	*	*	*
Coffey	33	*	*	*	*	*	*	*
Comanche	11	*	*	*	*	*	*	*
Cowley-Ark	212	67.5	1.9	28.8	0.9	0.9	0.0	0.0
Cowley-Win	151	86.8	2.6	5.3	0.7	2.6	2.0	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3A
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Clinic	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaska Native	Asian/Pacific Islander	Multiple Races	All Other/Unknown
		%	%	%	%	%	%	%
Crawford-G	2	*	*	*	*	*	*	*
Crawford-P	337	83.1	4.2	9.5	1.5	0.6	1.2	0.0
Decatur	14	*	*	*	*	*	*	*
DickinsonA	136	89.7	0.0	6.6	1.5	0.7	1.5	0.0
DickinsonH	19	*	*	*	*	*	*	*
Doniphan	53	*	*	*	*	*	*	*
Douglas	530	67.5	9.4	10.2	7.5	3.0	2.3	0.0
Edwards	25	*	*	*	*	*	*	*
Elk	24	*	*	*	*	*	*	*
Ellis	158	82.9	1.3	12.0	1.3	0.6	1.9	0.0
Ellsworth	37	*	*	*	*	*	*	*
Finney	703	27.9	1.6	65.6	0.0	4.6	0.4	0.0
Ford	697	21.8	2.3	75.0	0.3	0.6	0.0	0.0
Franklin	185	94.1	0.0	2.2	2.2	0.5	1.1	0.0
Geary	421	58.0	20.0	15.2	1.0	3.1	2.9	0.0
Gove	10	*	*	*	*	*	*	*
Graham	20	*	*	*	*	*	*	*
Grant	66	*	*	*	*	*	*	*
Gray	46	*	*	*	*	*	*	*
Greeley	12	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3A
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Clinic	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaska Native	Asian/Pacific Islander	Multiple Races	All Other/Unknown
		%	%	%	%	%	%	%
Greenwood	47	*	*	*	*	*	*	*
Hamilton	27	*	*	*	*	*	*	*
Harper	50	*	*	*	*	*	*	*
Harvey	283	70.0	3.2	21.9	1.4	1.4	2.1	0.0
Haskell	17	*	*	*	*	*	*	*
Hodgeman	11	*	*	*	*	*	*	*
Jackson	88	*	*	*	*	*	*	*
Jefferson	58	*	*	*	*	*	*	*
Jewell	7	*	*	*	*	*	*	*
Johnson-M	845	48.2	16.0	31.0	0.7	2.7	1.4	0.0
Johnson-O	1,145	47.6	10.7	36.2	0.3	3.7	1.6	0.0
Kearny	21	*	*	*	*	*	*	*
Kingman	49	*	*	*	*	*	*	*
Kiowa	8	*	*	*	*	*	*	*
Labette	176	89.2	3.4	4.0	1.7	0.0	1.7	0.0
Lane	11	*	*	*	*	*	*	*
Leavenworth	364	69.2	13.2	12.4	0.8	1.6	2.7	0.0
Lincoln	20	*	*	*	*	*	*	*
Linn	71	*	*	*	*	*	*	*
Logan	12	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3A
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Lyon	294	59.2	5.1	33.0	0.3	1.4	1.0	0.0
Marion	61	*	*	*	*	*	*	*
Marshall	52	*	*	*	*	*	*	*
Mc Pherson	125	82.4	1.6	13.6	0.8	1.6	0.0	0.0
Meade	38	*	*	*	*	*	*	*
Miami	151	90.1	2.6	4.0	0.7	0.7	2.0	0.0
Mitchell	36	*	*	*	*	*	*	*
Montgome-C	161	70.8	14.9	11.8	1.2	1.2	0.0	0.0
Montgome-I	184	79.9	7.6	7.1	2.2	1.1	2.2	0.0
Morris	39	*	*	*	*	*	*	*
Nemaha	29	*	*	*	*	*	*	*
Neosho-Ch	155	92.3	1.3	3.9	1.3	0.0	1.3	0.0
Neosho-Er	11	*	*	*	*	*	*	*
Ness	22	*	*	*	*	*	*	*
Norton	38	*	*	*	*	*	*	*
Osage	85	*	*	*	*	*	*	*
Osborne	26	*	*	*	*	*	*	*
Ottawa	29	*	*	*	*	*	*	*
Pawnee	58	*	*	*	*	*	*	*
Phillips	41	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3A
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Clinic	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaska Native	Asian/Pacific Islander	Multiple Races	All Other/Unknown
		%	%	%	%	%	%	%
Pottawat-S	18	*	*	*	*	*	*	*
Pottawat-W	18	*	*	*	*	*	*	*
Pratt	82	*	*	*	*	*	*	*
Rawlins	7	*	*	*	*	*	*	*
Reno	453	77.5	2.2	16.6	0.7	0.9	2.2	0.0
Republic	27	*	*	*	*	*	*	*
Rice	89	*	*	*	*	*	*	*
Riley-Ft	617	69.5	12.5	14.7	0.2	2.8	0.3	0.0
Riley-Manh	393	71.2	8.9	11.5	0.5	6.4	1.5	0.0
Rooks	50	*	*	*	*	*	*	*
Rush	10	*	*	*	*	*	*	*
Russell	44	*	*	*	*	*	*	*
Saline	546	72.5	3.7	20.3	0.0	1.6	1.8	0.0
Scott	36	*	*	*	*	*	*	*
Sedgwick-C	1,242	44.4	14.2	32.0	1.3	5.9	2.2	0.0
Sedgwick-M	1,913	32.7	27.9	33.8	1.2	2.6	1.9	0.0
Sedgwick-S	1,396	65.4	8.1	21.1	1.4	2.3	1.7	0.0
Seneca	9	*	*	*	*	*	*	*
Seward	428	15.0	2.1	82.0	0.5	0.5	0.0	0.0
ShawneeM&I	358	34.9	13.1	47.2	1.7	1.7	1.4	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3A
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Clinic	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaska Native	Asian/Pacific Islander	Multiple Races	All Other/Unknown
		%	%	%	%	%	%	%
ShawneeMa	1,146	60.3	15.9	17.7	3.1	0.8	2.2	0.0
Sheridan	8	*	*	*	*	*	*	*
Sherman	72	*	*	*	*	*	*	*
Smith	15	*	*	*	*	*	*	*
Stafford	33	*	*	*	*	*	*	*
Stanton	16	*	*	*	*	*	*	*
Stevens	70	*	*	*	*	*	*	*
Sumner	164	87.8	1.2	7.9	1.2	0.6	1.2	0.0
Thomas	50	*	*	*	*	*	*	*
Trego	14	*	*	*	*	*	*	*
Wabaunsee	17	*	*	*	*	*	*	*
Washington	22	*	*	*	*	*	*	*
Wichita	18	*	*	*	*	*	*	*
Wilson	75	*	*	*	*	*	*	*
Woodson	18	*	*	*	*	*	*	*
Wyandotte	2,100	26.8	29.4	38.3	0.5	4.0	1.1	0.0
Unknown Clinics	20	*	*	*	*	*	*	*
Kansas	21,902	57.3	11.2	26.7	1.2	2.3	1.4	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3A
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Nation (Prior Year)	1,304,320	41.6	24.1	28.3	1.1	3.2	1.2	0.5

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Age Distribution by Clinic

Comparison of Age Distribution ⁽²⁾								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Allen	84	*	*	*	*	*	*	*
Anderson	56	*	*	*	*	*	*	*
Atchison	140	0.0	7.9	12.9	61.4	17.1	0.7	0.0
Barber-Kio	2	*	*	*	*	*	*	*
Barber-Med	54	*	*	*	*	*	*	*
Barton	255	0.0	7.5	12.5	60.8	16.1	3.1	0.0
Bourbon	127	0.8	7.9	15.7	59.1	15.7	0.8	0.0
Brown	91	*	*	*	*	*	*	*
Butler	285	0.0	5.6	14.7	63.9	15.1	0.4	0.4
Chase	9	*	*	*	*	*	*	*
Chautauqua	19	*	*	*	*	*	*	*
Cherokee	183	0.5	10.9	22.4	54.6	11.5	0.0	0.0
Cheyenne	10	*	*	*	*	*	*	*
Clark	2	*	*	*	*	*	*	*
Clay	51	*	*	*	*	*	*	*
Cloud	93	*	*	*	*	*	*	*
Coffey	33	*	*	*	*	*	*	*
Comanche	11	*	*	*	*	*	*	*
Cowley-Ark	212	0.5	9.9	16.0	57.5	15.6	0.5	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Age Distribution by Clinic

Comparison of Age Distribution ⁽²⁾								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Cowley-Win	151	0.7	2.0	18.5	69.5	9.3	0.0	0.0
Crawford-G	2	*	*	*	*	*	*	*
Crawford-P	337	0.0	4.2	14.2	69.1	12.5	0.0	0.0
Decatur	14	*	*	*	*	*	*	*
DickinsonA	136	0.0	1.5	11.8	72.1	14.7	0.0	0.0
DickinsonH	19	*	*	*	*	*	*	*
Doniphan	53	*	*	*	*	*	*	*
Douglas	530	0.0	4.2	12.5	64.3	17.9	1.1	0.0
Edwards	25	*	*	*	*	*	*	*
Elk	24	*	*	*	*	*	*	*
Ellis	158	1.3	6.3	17.7	58.2	15.8	0.0	0.6
Ellsworth	37	*	*	*	*	*	*	*
Finney	703	0.4	6.7	14.1	57.6	18.9	2.3	0.0
Ford	697	0.3	6.0	12.1	53.8	24.8	3.0	0.0
Franklin	185	0.5	5.4	14.1	64.9	14.6	0.5	0.0
Geary	421	0.2	4.0	12.8	70.3	11.4	1.0	0.2
Gove	10	*	*	*	*	*	*	*
Graham	20	*	*	*	*	*	*	*
Grant	66	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Age Distribution by Clinic

Comparison of Age Distribution ⁽²⁾								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Gray	46	*	*	*	*	*	*	*
Greeley	12	*	*	*	*	*	*	*
Greenwood	47	*	*	*	*	*	*	*
Hamilton	27	*	*	*	*	*	*	*
Harper	50	*	*	*	*	*	*	*
Harvey	283	0.0	4.6	13.4	64.0	17.3	0.7	0.0
Haskell	17	*	*	*	*	*	*	*
Hodgeman	11	*	*	*	*	*	*	*
Jackson	88	*	*	*	*	*	*	*
Jefferson	58	*	*	*	*	*	*	*
Jewell	7	*	*	*	*	*	*	*
Johnson-M	845	0.1	3.4	10.1	63.3	21.9	1.2	0.0
Johnson-O	1,145	0.0	2.7	10.0	59.0	25.2	2.9	0.2
Kearny	21	*	*	*	*	*	*	*
Kingman	49	*	*	*	*	*	*	*
Kiowa	8	*	*	*	*	*	*	*
Labette	176	0.0	6.8	20.5	59.1	13.1	0.6	0.0
Lane	11	*	*	*	*	*	*	*
Leavenwort	364	0.3	5.5	10.2	69.8	13.7	0.5	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Age Distribution by Clinic

Comparison of Age Distribution ⁽²⁾								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Lincoln	20	*	*	*	*	*	*	*
Linn	71	*	*	*	*	*	*	*
Logan	12	*	*	*	*	*	*	*
Lyon	294	1.0	5.8	15.6	60.9	15.3	1.4	0.0
Marion	61	*	*	*	*	*	*	*
Marshall	52	*	*	*	*	*	*	*
Mc Pherson	125	0.8	3.2	12.8	62.4	18.4	2.4	0.0
Meade	38	*	*	*	*	*	*	*
Miami	151	0.7	4.0	18.5	61.6	13.9	1.3	0.0
Mitchell	36	*	*	*	*	*	*	*
Montgome-C	161	1.2	7.5	16.1	59.0	14.3	1.9	0.0
Montgome-I	184	0.0	6.5	19.0	61.4	10.3	2.7	0.0
Morris	39	*	*	*	*	*	*	*
Nemaha	29	*	*	*	*	*	*	*
Neosho-Ch	155	0.0	5.8	16.1	64.5	12.9	0.6	0.0
Neosho-Er	11	*	*	*	*	*	*	*
Ness	22	*	*	*	*	*	*	*
Norton	38	*	*	*	*	*	*	*
Osage	85	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Age Distribution by Clinic

Comparison of Age Distribution ⁽²⁾								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Osborne	26	*	*	*	*	*	*	*
Ottawa	29	*	*	*	*	*	*	*
Pawnee	58	*	*	*	*	*	*	*
Phillips	41	*	*	*	*	*	*	*
Pottawat-S	18	*	*	*	*	*	*	*
Pottawat-W	18	*	*	*	*	*	*	*
Pratt	82	*	*	*	*	*	*	*
Rawlins	7	*	*	*	*	*	*	*
Reno	453	0.2	7.7	14.3	63.1	14.1	0.4	0.0
Republic	27	*	*	*	*	*	*	*
Rice	89	*	*	*	*	*	*	*
Riley-Ft	617	0.0	1.1	10.4	75.9	12.5	0.2	0.0
Riley-Manh	393	0.3	3.1	12.5	70.5	13.5	0.3	0.0
Rooks	50	*	*	*	*	*	*	*
Rush	10	*	*	*	*	*	*	*
Russell	44	*	*	*	*	*	*	*
Saline	546	0.0	5.5	16.5	61.0	15.9	0.9	0.2
Scott	36	*	*	*	*	*	*	*
Sedgwick-C	1,242	0.2	6.4	12.8	58.9	20.3	1.4	0.1

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Age Distribution by Clinic

Comparison of Age Distribution ⁽²⁾								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Sedgwick-M	1,913	0.2	6.7	14.1	59.5	18.2	1.3	0.1
Sedgwick-S	1,396	0.4	6.8	13.8	62.2	15.6	1.1	0.0
Seneca	9	*	*	*	*	*	*	*
Seward	428	0.2	7.7	11.0	58.2	21.7	1.2	0.0
ShawneeM&I	358	0.8	7.8	17.0	53.1	19.8	1.4	0.0
ShawneeMa	1,146	0.1	5.8	13.0	62.6	17.5	1.0	0.1
Sheridan	8	*	*	*	*	*	*	*
Sherman	72	*	*	*	*	*	*	*
Smith	15	*	*	*	*	*	*	*
Stafford	33	*	*	*	*	*	*	*
Stanton	16	*	*	*	*	*	*	*
Stevens	70	*	*	*	*	*	*	*
Sumner	164	0.0	8.5	13.4	59.8	17.1	1.2	0.0
Thomas	50	*	*	*	*	*	*	*
Trego	14	*	*	*	*	*	*	*
Wabaunsee	17	*	*	*	*	*	*	*
Washington	22	*	*	*	*	*	*	*
Wichita	18	*	*	*	*	*	*	*
Wilson	75	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Age Distribution by Clinic

Comparison of Age Distribution ⁽²⁾								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Woodson	18	*	*	*	*	*	*	*
Wyandotte	2,100	0.5	9.0	13.3	55.5	20.1	1.7	0.0
Unknown Clinics	20	*	*	*	*	*	*	*
Kansas	21,902	0.3	5.9	13.5	61.3	17.7	1.3	0.0
Nation (Prior Year)	1,304,320	0.3	5.8	12.3	59.4	20.4	1.5	0.3

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators ^(2, 3)														
Clinic	Prepregnancy BMI ⁽⁴⁾			Weight Gain ⁽⁵⁾			Anemia (Low Hb/Hct) ⁽⁶⁾				Gestational Diabetes During Pregnancy	Hypertension During Pregnancy		
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)			Number	% (Rank)
Allen	82	* *	* *	72	* *	* *	9	* *	23	* *	75	* *	75	* *
Anderson	54	* *	* *	51	* *	* *	5	* *	11	* *	53	* *	53	* *
Atchison	138	4.3 (19)	48.6 (3)	119	12.6 (1)	63.0 (38)	18	* *	24	* *	105	4.8 (7)	106	11.3(19)
Barber-Kio	2	* *	* *	2	* *	* *	0	* *	1	* *	2	* *	2	* *
Barber-Med	54	* *	* *	45	* *	* *	6	* *	11	* *	45	* *	45	* *
Barton	244	5.7 (32)	57.8 (30)	213	23.9 (28)	51.2 (20)	23	* *	63	* *	222	8.6 (31)	222	14.0(30)
Bourbon	125	0.8 (1)	56.8 (26)	108	29.6 (38)	48.1 (10)	9	* *	20	* *	108	8.3 (29)	109	15.6(36)
Brown	90	* *	* *	74	* *	* *	11	* *	18	* *	75	* *	75	* *
Butler	280	6.1 (33)	53.9 (17)	253	21.3 (18)	57.3 (36)	41	* *	44	* *	260	4.6 (6)	260	11.2(18)
Chase	9	* *	* *	7	* *	* *	0	* *	0	* *	8	* *	8	* *
Chautauqua	18	* *	* *	15	* *	* *	2	* *	4	* *	16	* *	16	* *
Cherokee	170	7.6 (38)	48.8 (4)	152	21.7 (22)	53.3 (29)	24	* *	35	* *	163	8.0 (26)	162	11.7(23)
Cheyenne	9	* *	* *	6	* *	* *	3	* *	1	* *	7	* *	7	* *
Clark	2	* *	* *	2	* *	* *	1	* *	0	* *	2	* *	2	* *
Clay	48	* *	* *	41	* *	* *	8	* *	14	* *	45	* *	45	* *
Cloud	90	* *	* *	81	* *	* *	9	* *	23	* *	83	* *	84	* *
Coffey	31	* *	* *	23	* *	* *	5	* *	5	* *	25	* *	25	* *

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
- (4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
- (5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
- (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators ^(2, 3)														
Clinic	Prepregnancy BMI ⁽⁴⁾			Weight Gain ⁽⁵⁾			Anemia (Low Hb/Hct) ⁽⁶⁾				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Comanche	11	* * *	* *	9	* * *	* *	1	* * *	2	* * *	9	* * *	9	* * *
Cowley-Ark	207	4.3 (19)	50.7 (7)	185	22.7 (24)	47.0 (9)	27	* * *	34	* * *	190	9.5 (35)	191	9.4 (13)
Cowley-Win	148	2.0 (3)	60.8 (37)	122	21.3 (18)	56.6 (35)	13	* * *	23	* * *	125	8.0 (26)	125	10.4 (17)
Crawford-G	2	* * *	* *	2	* * *	* *	1	* * *	0	* * *	2	* * *	2	* * *
Crawford-P	335	6.6 (35)	51.9 (11)	281	23.8 (27)	46.6 (8)	44	* * *	43	* * *	288	10.1 (37)	288	12.5 (25)
Decatur	14	* * *	* *	14	* * *	* *	4	* * *	4	* * *	14	* * *	14	* * *
DickinsonA	131	0.8 (1)	66.4 (38)	108	27.8 (35)	45.4 (7)	18	* * *	20	* * *	113	6.2 (13)	113	15.0 (34)
DickinsonH	18	* * *	* *	16	* * *	* *	1	* * *	7	* * *	18	* * *	18	* * *
Doniphan	52	* * *	* *	44	* * *	* *	10	* * *	10	* * *	46	* * *	46	* * *
Douglas	513	2.9 (8)	48.3 (1)	423	19.4 (13)	51.3 (22)	62	* * *	130	32.3 (9)	439	7.5 (23)	439	9.6 (14)
Edwards	25	* * *	* *	21	* * *	* *	6	* * *	3	* * *	21	* * *	21	* * *
Elk	24	* * *	* *	20	* * *	* *	3	* * *	5	* * *	19	* * *	19	* * *
Ellis	155	6.5 (34)	51.0 (8)	135	15.6 (3)	49.6 (15)	28	* * *	27	* * *	139	2.9 (1)	139	6.5 (3)
Ellsworth	35	* * *	* *	31	* * *	* *	7	* * *	3	* * *	33	* * *	33	* * *
Finney	690	4.9 (25)	53.2 (14)	614	25.7 (32)	43.2 (2)	92	* * *	95	* * *	634	8.7 (32)	634	7.6 (5)
Ford	687	2.2 (4)	59.1 (33)	595	28.4 (36)	44.4 (5)	84	* * *	91	* * *	596	6.5 (14)	595	8.6 (7)
Franklin	172	5.2 (27)	48.3 (1)	148	18.2 (7)	50.0 (17)	20	* * *	48	* * *	139	5.0 (8)	139	10.1 (16)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators ^(2, 3)														
Clinic	Prepregnancy BMI ⁽⁴⁾			Weight Gain ⁽⁵⁾			Anemia (Low Hb/Hct) ⁽⁶⁾				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Geary	405	4.2 (18)	56.8 (26)	317	20.2 (14)	52.7 (27)	61	* *	64	* *	332	3.9 (2)	331	11.5(21)
Gove	8	* *	* *	6	* *	* *	1	* *	5	* *	8	* *	8	* *
Graham	20	* *	* *	16	* *	* *	1	* *	3	* *	16	* *	16	* *
Grant	64	* *	* *	59	* *	* *	7	* *	17	* *	62	* *	62	* *
Gray	42	* *	* *	33	* *	* *	6	* *	13	* *	37	* *	37	* *
Greeley	12	* *	* *	9	* *	* *	0	* *	1	* *	9	* *	9	* *
Greenwood	45	* *	* *	37	* *	* *	5	* *	9	* *	36	* *	36	* *
Hamilton	27	* *	* *	26	* *	* *	10	* *	7	* *	25	* *	25	* *
Harper	47	* *	* *	43	* *	* *	4	* *	4	* *	44	* *	44	* *
Harvey	276	4.7 (23)	59.4 (36)	244	18.9 (10)	51.2 (20)	38	* *	39	* *	251	4.4 (4)	250	11.6(22)
Haskell	17	* *	* *	16	* *	* *	0	* *	1	* *	16	* *	16	* *
Hodgeman	10	* *	* *	8	* *	* *	1	* *	1	* *	8	* *	8	* *
Jackson	84	* *	* *	77	* *	* *	9	* *	17	* *	78	* *	79	* *
Jefferson	56	* *	* *	48	* *	* *	10	* *	13	* *	50	* *	50	* *
Jewell	7	* *	* *	7	* *	* *	1	* *	2	* *	7	* *	7	* *
Johnson-M	808	3.1 (11)	58.4 (32)	695	19.3 (12)	49.4 (14)	126	27.0 (4)	201	16.9 (1)	716	6.7 (17)	714	7.4(4)
Johnson-O	1,108	3.2 (12)	52.2 (13)	956	18.3 (8)	52.2 (26)	155	18.7 (1)	261	18.0 (2)	985	8.5 (30)	981	6.3(1)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators ^(2, 3)														
Clinic	Prepregnancy BMI ⁽⁴⁾			Weight Gain ⁽⁵⁾			Anemia (Low Hb/Hct) ⁽⁶⁾				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Kearny	21	* *	* *	21	* *	* *	2	* *	1	* *	21	* *	21	* *
Kingman	49	* *	* *	42	* *	* *	2	* *	9	* *	43	* *	43	* *
Kiowa	7	* *	* *	7	* *	* *	2	* *	2	* *	8	* *	8	* *
Labette	172	4.7 (23)	59.3 (34)	154	26.0 (33)	50.0 (17)	35	* *	39	* *	154	6.5 (14)	154	14.3(32)
Lane	10	* *	* *	9	* *	* *	2	* *	3	* *	10	* *	10	* *
Leavenworth	348	4.0 (16)	56.0 (23)	292	20.5 (15)	56.2 (34)	51	* *	92	* *	304	5.6 (11)	303	13.2(27)
Lincoln	19	* *	* *	17	* *	* *	6	* *	3	* *	17	* *	17	* *
Linn	71	* *	* *	63	* *	* *	11	* *	9	* *	63	* *	64	* *
Logan	11	* *	* *	9	* *	* *	2	* *	1	* *	9	* *	9	* *
Lyon	288	2.4 (5)	56.6 (24)	249	25.3 (31)	44.2 (4)	34	* *	57	* *	252	6.7 (17)	252	13.5(28)
Marion	59	* *	* *	46	* *	* *	8	* *	11	* *	50	* *	51	* *
Marshall	51	* *	* *	49	* *	* *	5	* *	12	* *	49	* *	48	* *
Mc Pherson	122	3.3 (13)	58.2 (31)	105	29.5 (37)	43.8 (3)	14	* *	27	* *	109	9.2 (34)	110	9.1(10)
Meade	37	* *	* *	34	* *	* *	1	* *	6	* *	33	* *	33	* *
Miami	147	3.4 (14)	55.1 (20)	125	13.6 (2)	58.4 (37)	30	* *	34	* *	132	6.8 (19)	132	11.4(20)
Mitchell	35	* *	* *	30	* *	* *	2	* *	9	* *	32	* *	31	* *
Montgome-C	158	2.5 (7)	50.6 (6)	141	24.8 (30)	49.6 (15)	34	* *	38	* *	141	9.9 (36)	141	14.2(31)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators ^(2, 3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight	Over- weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Montgome-I	175	6.9 (36)	49.1 (5)	150	22.7 (24)	48.7 (11)	37	* *	32	* *	155	7.7 (25)	155	13.5(28)
Morris	38	* *	* *	34	* *	* *	5	* *	5	* *	35	* *	35	* *
Nemaha	26	* *	* *	23	* *	* *	7	* *	4	* *	26	* *	26	* *
Neosho-Ch	153	5.2 (27)	55.6 (22)	137	24.1 (29)	51.8 (25)	25	* *	25	* *	138	6.5 (14)	137	19.0(38)
Neosho-Er	11	* *	* *	10	* *	* *	1	* *	2	* *	10	* *	10	* *
Ness	21	* *	* *	17	* *	* *	3	* *	3	* *	17	* *	17	* *
Norton	38	* *	* *	35	* *	* *	8	* *	13	* *	35	* *	35	* *
Osage	82	* *	* *	73	* *	* *	11	* *	16	* *	74	* *	74	* *
Osborne	25	* *	* *	22	* *	* *	2	* *	5	* *	23	* *	23	* *
Ottawa	27	* *	* *	24	* *	* *	5	* *	2	* *	29	* *	29	* *
Pawnee	56	* *	* *	44	* *	* *	4	* *	8	* *	46	* *	45	* *
Phillips	39	* *	* *	28	* *	* *	5	* *	6	* *	34	* *	34	* *
Pottawat-S	18	* *	* *	16	* *	* *	1	* *	8	* *	16	* *	16	* *
Pottawat-W	18	* *	* *	17	* *	* *	3	* *	3	* *	17	* *	17	* *
Pratt	82	* *	* *	71	* *	* *	11	* *	14	* *	72	* *	72	* *
Rawlins	7	* *	* *	6	* *	* *	1	* *	2	* *	6	* *	6	* *
Reno	441	4.1 (17)	54.6 (19)	390	19.2 (11)	53.8 (30)	62	* *	76	* *	396	4.3 (3)	397	14.6(33)

- (1) Reporting period is January 1 through December 31.
 - (2) Excludes records with unknown data or errors.
 - (3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
 - (4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
 - (5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
 - (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators ^(2, 3)															
Clinic	Prepregnancy BMI ⁽⁴⁾				Weight Gain ⁽⁵⁾			Anemia (Low Hb/Hct) ⁽⁶⁾				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight		Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)		Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Republic	27	* *	* *		22	* *	* *	5	* *	4	* *	22	* *	22	* *
Rice	87	* *	* *		76	* *	* *	12	* *	16	* *	78	* *	78	* *
Riley-Ft	579	5.4 (30)	53.7 (16)		451	20.8 (16)	54.1 (31)	102	43.1 (7)	140	36.4 (10)	465	4.5 (5)	466	15.5 (35)
Riley-Manh	381	2.9 (8)	51.4 (9)		314	15.6 (3)	50.6 (19)	50	* *	109	22.0 (5)	326	5.8 (12)	326	8.3 (6)
Rooks	50	* *	* *		42	* *	* *	7	* *	8	* *	40	* *	40	* *
Rush	10	* *	* *		10	* *	* *	1	* *	3	* *	10	* *	10	* *
Russell	43	* *	* *		39	* *	* *	7	* *	4	* *	40	* *	40	* *
Saline	538	5.0 (26)	57.4 (29)		467	16.5 (5)	55.9 (33)	96	* *	77	* *	471	8.7 (32)	470	9.1 (10)
Scott	36	* *	* *		28	* *	* *	2	* *	5	* *	28	* *	28	* *
Sedgwick-C	1,207	4.4 (21)	53.3 (15)		1,077	21.7 (22)	51.3 (22)	214	26.6 (2)	244	18.9 (3)	1,114	7.3 (22)	1,115	9.1 (10)
Sedgwick-M	1,866	2.9 (8)	56.6 (24)		1,669	21.5 (21)	48.7 (11)	390	26.7 (3)	434	21.9 (4)	1,708	7.1 (20)	1,706	12.8 (26)
Sedgwick-S	1,353	5.5 (31)	51.9 (11)		1,201	18.3 (8)	55.4 (32)	230	36.5 (6)	308	29.5 (8)	1,248	5.0 (8)	1,242	12.2 (24)
Seneca	7	* *	* *		7	* *	* *	3	* *	2	* *	8	* *	8	* *
Seward	423	2.4 (5)	59.3 (34)		374	23.0 (26)	44.9 (6)	63	* *	69	* *	380	8.2 (28)	378	6.3 (1)
ShawneeM&I	356	3.9 (15)	55.3 (21)		264	17.0 (6)	49.2 (13)	2	* *	46	* *	266	11.7 (38)	265	8.7 (8)
ShawneeMa	1,108	5.2 (27)	54.3 (18)		982	21.3 (18)	52.7 (27)	133	50.4 (8)	193	28.0 (7)	1,016	7.1 (20)	1,017	8.8 (9)
Sheridan	8	* *	* *		6	* *	* *	0	* *	1	* *	7	* *	7	* *

- (1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators ^(2, 3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Sherman	69	* *	* *	57	* *	* *	7	* *	14	* *	60	* *	60	* *
Smith	15	* *	* *	12	* *	* *	2	* *	1	* *	12	* *	12	* *
Stafford	32	* *	* *	28	* *	* *	4	* *	5	* *	28	* *	29	* *
Stanton	15	* *	* *	14	* *	* *	1	* *	8	* *	15	* *	14	* *
Stevens	69	* *	* *	60	* *	* *	15	* *	17	* *	60	* *	60	* *
Sumner	161	7.5 (37)	51.6 (10)	144	26.4 (34)	40.3 (1)	23	* *	43	* *	146	7.5 (23)	147	17.7(37)
Thomas	50	* *	* *	46	* *	* *	5	* *	8	* *	47	* *	47	* *
Trego	14	* *	* *	10	* *	* *	2	* *	0	* *	10	* *	10	* *
Wabaunsee	16	* *	* *	15	* *	* *	4	* *	8	* *	17	* *	17	* *
Washington	21	* *	* *	19	* *	* *	2	* *	8	* *	20	* *	20	* *
Wichita	16	* *	* *	15	* *	* *	2	* *	0	* *	17	* *	17	* *
Wilson	74	* *	* *	69	* *	* *	13	* *	10	* *	69	* *	69	* *
Woodson	18	* *	* *	16	* *	* *	6	* *	8	* *	16	* *	16	* *
Wyandotte	2,037	4.5 (22)	57.1 (28)	1,785	20.8 (16)	51.5 (24)	310	31.0 (5)	390	27.4 (6)	1,811	5.4 (10)	1,809	9.8(15)
Unknown Clinics	18	* *	* *	17	* *	* *	1	* *	2	* *	19	* *	19	* *
Kansas	21,267	4.1	55.0	18,455	21.1	50.7	3,176	29.0	4,262	23.7	18,929	6.8	18,913	10.7

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators ^(2, 3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy	Hypertension During Pregnancy		
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)			Number	% (Rank)
Nation (Prior Year)	1,226,875	4.5	52.9	917,032	21.2	48.2	189,944	33.9	412,633	28.8	389,161	5.0	394,368	6.8

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6A
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators ^(2, 3)										
Clinic	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	% (Rank)		Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Allen	67	* *	* *		84	* *	84	* *	67	* *
Anderson	41	* *	* *		56	* *	56	* *	50	* *
Atchison	119	89.1 (3)	2.5 (14)		139	40.3 (7)	140	10.7 (33)	120	58.3 (24)
Barber-Kio	2	* *	* *		2	* *	2	* *	2	* *
Barber-Med	45	* *	* *		54	* *	54	* *	42	* *
Barton	225	87.6 (6)	0.9 (4)		254	48.8 (2)	255	21.2 (8)	201	59.2 (21)
Bourbon	99	* *	* *		127	40.2 (8)	127	11.8 (28)	94	* *
Brown	54	* *	* *		91	* *	91	* *	76	* *
Butler	255	74.9 (23)	3.1 (16)		285	33.3 (27)	285	14.0 (22)	224	37.9 (37)
Chase	8	* *	* *		7	* *	9	* *	9	* *
Chautauqua	16	* *	* *		19	* *	19	* *	16	* *
Cherokee	117	73.5 (25)	9.4 (31)		183	35.5 (23)	183	11.5 (29)	148	55.4 (28)
Cheyenne	6	* *	* *		10	* *	10	* *	10	* *
Clark	2	* *	* *		2	* *	2	* *	2	* *
Clay	46	* *	* *		51	* *	51	* *	46	* *
Cloud	85	* *	* *		92	* *	93	* *	82	* *
Coffey	25	* *	* *		33	* *	33	* *	31	* *
Comanche	9	* *	* *		11	* *	11	* *	9	* *
Cowley-Ark	175	69.1 (30)	4.6 (24)		210	36.2 (20)	212	14.6 (20)	174	57.5 (26)
Cowley-Win	122	77.0 (19)	9.0 (30)		150	48.7 (3)	151	11.3 (32)	129	54.3 (31)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6A
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators ^(2, 3)										
Clinic	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Crawford-G	2	* *	* *	* *	2	* *	2	* *	2	* *
Crawford-P	269	87.4 (7)	4.5 (23)		337	40.1 (9)	337	14.2 (21)	280	59.3 (20)
Decatur	13	* *	* *	* *	14	* *	14	* *	10	* *
DickinsonA	115	87.8 (5)	1.7 (11)		134	38.1 (16)	136	19.9 (10)	108	69.4 (5)
DickinsonH	18	* *	* *	* *	19	* *	19	* *	17	* *
Doniphan	47	* *	* *	* *	53	* *	53	* *	41	* *
Douglas	443	76.1 (21)	5.6 (28)		528	38.6 (14)	530	21.5 (7)	434	64.5 (13)
Edwards	18	* *	* *	* *	24	* *	25	* *	23	* *
Elk	19	* *	* *	* *	24	* *	24	* *	21	* *
Ellis	141	86.5 (9)	0.7 (3)		158	39.9 (10)	158	25.9 (3)	137	65.0 (11)
Ellsworth	33	* *	* *	* *	37	* *	37	* *	30	* *
Finney	603	52.4 (36)	29.2 (36)		694	35.2 (24)	703	12.4 (26)	589	48.6 (34)
Ford	533	72.2 (28)	12.4 (34)		695	44.7 (5)	697	13.9 (23)	591	52.1 (33)
Franklin	76	* *	* *	* *	185	31.4 (28)	185	8.6 (35)	127	63.8 (14)
Geary	317	76.0 (22)	1.6 (9)		421	30.2 (33)	421	22.1 (6)	342	58.5 (23)
Gove	8	* *	* *	* *	10	* *	10	* *	7	* *
Graham	16	* *	* *	* *	20	* *	20	* *	17	* *
Grant	58	* *	* *	* *	65	* *	66	* *	52	* *
Gray	37	* *	* *	* *	46	* *	46	* *	44	* *
Greeley	8	* *	* *	* *	12	* *	12	* *	10	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6A
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators ^(2, 3)										
Clinic	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Prepregnancy ⁽⁴⁾		During Pregnancy ⁽⁵⁾	
	Number	% (Rank)	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Greenwood	37	* *	* *	* *	47	* *	47	* *	43	* *
Hamilton	24	* *	* *	* *	27	* *	27	* *	25	* *
Harper	44	* *	* *	* *	48	* *	50	* *	43	* *
Harvey	220	73.6 (24)	11.4 (33)		281	48.4 (4)	283	19.4 (11)	235	66.0 (10)
Haskell	15	* *	* *	* *	16	* *	17	* *	15	* *
Hodgeman	8	* *	* *	* *	11	* *	11	* *	10	* *
Jackson	38	* *	* *	* *	88	* *	88	* *	74	* *
Jefferson	48	* *	* *	* *	58	* *	58	* *	49	* *
Jewell	7	* *	* *	* *	7	* *	7	* *	6	* *
Johnson-M	638	71.0 (29)	3.0 (15)		844	34.5 (25)	845	15.6 (19)	667	61.2 (17)
Johnson-O	880	61.6 (35)	4.7 (25)		1,139	33.9 (26)	1,145	12.0 (27)	931	60.5 (18)
Kearny	19	* *	* *	* *	21	* *	21	* *	16	* *
Kingman	43	* *	* *	* *	49	* *	49	* *	39	* *
Kiowa	8	* *	* *	* *	8	* *	8	* *	6	* *
Labette	128	86.7 (8)	1.6 (9)		175	30.9 (31)	176	11.4 (30)	146	67.8 (7)
Lane	10	* *	* *	* *	11	* *	11	* *	10	* *
Leavenwort	307	83.7 (13)	1.0 (5)		364	28.0 (36)	364	22.3 (5)	273	68.5 (6)
Lincoln	17	* *	* *	* *	20	* *	20	* *	20	* *
Linn	65	* *	* *	* *	71	* *	71	* *	60	* *
Logan	7	* *	* *	* *	12	* *	12	* *	11	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6A
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators ^(2, 3)										
Clinic	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	% (Rank)		Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Lyon	253	83.8 (12)	1.2 (7)		294	38.8 (13)	294	17.7 (16)	237	64.6 (12)
Marion	48	* *	* *		60	* *	61	* *	48	* *
Marshall	50	* *	* *		52	* *	52	* *	43	* *
Mc Pherson	109	84.4 (11)	0.0 (1)		125	38.4 (15)	125	28.0 (1)	106	73.6 (2)
Meade	33	* *	* *		38	* *	38	* *	36	* *
Miami	131	79.4 (16)	1.5 (8)		151	37.7 (18)	151	18.5 (14)	127	55.1 (29)
Mitchell	32	* *	* *		36	* *	36	* *	31	* *
Montgome-C	104	85.6 (10)	1.0 (5)		161	21.7 (38)	161	17.4 (17)	137	70.8 (4)
Montgome-I	113	89.4 (2)	1.8 (12)		184	28.3 (35)	184	11.4 (30)	150	67.3 (8)
Morris	32	* *	* *		38	* *	39	* *	32	* *
Nemaha	25	* *	* *		29	* *	29	* *	24	* *
Neosho-Ch	125	88.8 (4)	3.2 (17)		154	39.0 (12)	155	12.9 (24)	130	71.5 (3)
Neosho-Er	8	* *	* *		11	* *	11	* *	9	* *
Ness	13	* *	* *		22	* *	22	* *	20	* *
Norton	35	* *	* *		38	* *	38	* *	32	* *
Osage	59	* *	* *		85	* *	85	* *	70	* *
Osborne	21	* *	* *		26	* *	26	* *	18	* *
Ottawa	28	* *	* *		29	* *	29	* *	25	* *
Pawnee	45	* *	* *		57	* *	58	* *	49	* *
Phillips	34	* *	* *		41	* *	41	* *	36	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6A
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators ^(2, 3)										
Clinic	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Pottawat-S	13	* *	* *	* *	18	* *	18	* *	15	* *
Pottawat-W	17	* *	* *	* *	18	* *	18	* *	17	* *
Pratt	72	* *	* *	* *	82	* *	82	* *	66	* *
Rawlins	6	* *	* *	* *	7	* *	7	* *	5	* *
Reno	388	83.2 (14)	3.6 (20)		451	39.2 (11)	453	20.8 (9)	382	63.6 (15)
Republic	22	* *	* *	* *	27	* *	27	* *	23	* *
Rice	77	* *	* *	* *	88	* *	89	* *	69	* *
Riley-Ft	435	92.4 (1)	0.5 (2)		617	31.3 (29)	617	24.8 (4)	470	78.3 (1)
Riley-Manh	305	68.9 (32)	2.3 (13)		393	36.1 (21)	393	18.8 (13)	323	60.1 (19)
Rooks	41	* *	* *	* *	49	* *	50	* *	42	* *
Rush	10	* *	* *	* *	10	* *	10	* *	10	* *
Russell	40	* *	* *	* *	43	* *	44	* *	39	* *
Saline	402	76.4 (20)	3.2 (17)		544	37.9 (17)	546	4.6 (38)	478	54.6 (30)
Scott	27	* *	* *	* *	35	* *	36	* *	32	* *
Sedgwick-C	1,071	72.5 (27)	10.2 (32)		1,237	29.3 (34)	1,242	19.4 (11)	1,034	48.2 (35)
Sedgwick-M	1,700	73.0 (26)	5.1 (27)		1,910	26.3 (37)	1,913	18.2 (15)	1,581	59.1 (22)
Sedgwick-S	1,211	80.4 (15)	3.3 (19)		1,391	31.1 (30)	1,396	17.0 (18)	1,143	40.4 (36)
Seneca	8	* *	* *	* *	9	* *	9	* *	8	* *
Seward	336	66.4 (33)	13.7 (35)		423	30.7 (32)	428	8.4 (36)	345	67.0 (9)
ShawneeM&I	258	69.0 (31)	3.9 (22)		354	70.6 (1)	358	26.3 (2)	343	57.1 (27)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6A
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators ^(2, 3)										
Clinic	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	% (Rank)		Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
ShawneeMa	873	77.7 (18)	3.7 (21)		1,143	37.6 (19)	1,146	9.1 (34)	901	52.2 (32)
Sheridan	7	* *	* *		8	* *	8	* *	8	* *
Sherman	55	* *	* *		72	* *	72	* *	56	* *
Smith	11	* *	* *		15	* *	15	* *	14	* *
Stafford	29	* *	* *		33	* *	33	* *	26	* *
Stanton	15	* *	* *		16	* *	16	* *	12	* *
Stevens	60	* *	* *		69	* *	70	* *	60	* *
Sumner	147	78.2 (17)	4.8 (26)		164	42.1 (6)	164	12.8 (25)	136	62.5 (16)
Thomas	40	* *	* *		50	* *	50	* *	38	* *
Trego	11	* *	* *		14	* *	14	* *	13	* *
Wabaunsee	15	* *	* *		17	* *	17	* *	14	* *
Washington	19	* *	* *		21	* *	22	* *	18	* *
Wichita	17	* *	* *		18	* *	18	* *	13	* *
Wilson	68	* *	* *		74	* *	75	* *	63	* *
Woodson	18	* *	* *		18	* *	18	* *	15	* *
Wyandotte	1,726	64.1 (34)	8.6 (29)		2,082	36.0 (22)	2,100	6.6 (37)	1,778	57.9 (25)
Unknown Clinics	17	* *	* *		20	* *	20	* *	17	* *
Kansas	17,720	75.1	6.1		21,806	36.1	21,902	16.2	18,060	58.1

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6A
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators ^(2, 3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Pregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Nation (Prior Year)	527,184	82.2	3.1	1,213,982	34.1	360,369	18.7	372,743	83.5

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2010 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Allen	79	* *	60	* *	26	* *	66	* *	62	* *
Anderson	56	* *	52	* *	21	* *	50	* *	53	* *
Atchison	118	42.4 (30)	81	* *	26	* *	100	31.0 (29)	82	* *
Barber-Kio	2	* *	2	* *	0	* *	2	* *	2	* *
Barber-Med	53	* *	42	* *	14	* *	42	* *	43	* *
Barton	253	33.6 (14)	223	23.3 (22)	58	* *	201	10.0 (10)	222	5.9 (8)
Bourbon	118	43.2 (31)	93	* *	29	* *	88	* *	96	* *
Brown	85	* *	60	* *	20	* *	71	* *	62	* *
Butler	281	51.6 (37)	253	31.6 (34)	97	* *	223	13.0 (16)	258	9.7 (16)
Chase	9	* *	8	* *	4	* *	9	* *	8	* *
Chautauqua	18	* *	11	* *	5	* *	15	* *	11	* *
Cherokee	180	44.4 (34)	157	29.9 (31)	54	* *	146	38.4 (34)	159	40.9 (34)
Cheyenne	10	* *	7	* *	2	* *	10	* *	7	* *
Clark	2	* *	1	* *	0	* *	2	* *	1	* *
Clay	51	* *	44	* *	14	* *	46	* *	44	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2010 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Cloud	93	* *	84	* *	23	* *	82	* *	84	* *
Coffey	32	* *	25	* *	10	* *	30	* *	25	* *
Comanche	11	* *	9	* *	5	* *	9	* *	9	* *
Cowley-Ark	210	38.6 (20)	175	26.3 (27)	50	* *	171	24.0 (26)	175	22.9 (27)
Cowley-Win	145	44.1 (33)	103	31.1 (33)	38	* *	126	13.5 (17)	103	9.7 (16)
Crawford-G	2	* *	2	* *	1	* *	2	* *	2	* *
Crawford-P	318	49.1 (36)	245	30.6 (32)	98	* *	269	39.0 (36)	246	36.2 (32)
Decatur	14	* *	12	* *	3	* *	10	* *	13	* *
DickinsonA	133	42.1 (29)	112	22.3 (21)	36	* *	105	8.6 (8)	113	7.1 (12)
DickinsonH	19	* *	18	* *	10	* *	17	* *	18	* *
Doniphan	52	* *	37	* *	8	* *	40	* *	37	* *
Douglas	524	34.9 (17)	431	17.2 (12)	117	40.2 (5)	429	10.3 (12)	429	6.3 (11)
Edwards	25	* *	21	* *	9	* *	23	* *	21	* *
Elk	21	* *	16	* *	6	* *	19	* *	16	* *
Ellis	158	41.1 (26)	136	19.1 (15)	48	* *	136	7.4 (5)	138	3.6 (4)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2010 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Ellsworth	37	* *	33	* *	12	* *	30	* *	33	* *
Finney	696	9.3 (3)	616	5.2 (3)	40	* *	586	11.8 (15)	616	10.7 (19)
Ford	671	8.0 (2)	538	4.3 (2)	32	* *	581	8.4 (7)	540	8.0 (14)
Franklin	184	40.2 (25)	156	28.8 (29)	34	* *	126	15.1 (21)	157	10.8 (20)
Geary	415	35.2 (18)	328	21.6 (19)	77	* *	338	35.2 (31)	328	34.5 (30)
Gove	10	* *	8	* *	1	* *	7	* *	8	* *
Graham	20	* *	16	* *	6	* *	17	* *	16	* *
Grant	66	* *	62	* *	2	* *	52	* *	62	* *
Gray	46	* *	35	* *	2	* *	44	* *	35	* *
Greeley	12	* *	8	* *	1	* *	10	* *	8	* *
Greenwood	39	* *	13	* *	7	* *	39	* *	21	* *
Hamilton	27	* *	26	* *	3	* *	25	* *	26	* *
Harper	47	* *	41	* *	10	* *	40	* *	38	* *
Harvey	280	40.0 (23)	243	21.0 (17)	78	* *	233	5.6 (1)	242	1.7 (2)
Haskell	17	* *	16	* *	2	* *	15	* *	16	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2010 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Hodgeman	9	* *	7	* *	4	* *	9	* *	7	* *
Jackson	87	* *	76	* *	33	* *	74	* *	77	* *
Jefferson	15	* *	11	* *	3	* *	13	* *	12	* *
Jewell	7	* *	6	* *	2	* *	6	* *	6	* *
Johnson-M	826	25.9 (6)	712	11.9 (6)	121	44.6 (2)	658	19.9 (25)	712	18.8 (26)
Johnson-O	1,137	22.4 (5)	985	9.3 (4)	157	45.9 (1)	915	16.4 (23)	988	16.3 (24)
Kearny	21	* *	20	* *	4	* *	16	* *	20	* *
Kingman	43	* *	37	* *	13	* *	37	* *	38	* *
Kiowa	8	* *	8	* *	2	* *	6	* *	8	* *
Labette	170	37.1 (19)	150	27.3 (28)	44	* *	143	30.1 (28)	150	34.0 (29)
Lane	5	* *	1	* *	0	* *	5	* *	2	* *
Leavenwort	361	47.6 (35)	301	23.9 (23)	98	* *	271	38.4 (34)	299	38.8 (33)
Lincoln	20	* *	18	* *	10	* *	20	* *	18	* *
Linn	69	* *	63	* *	37	* *	59	* *	63	* *
Logan	11	* *	10	* *	4	* *	11	* *	10	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2010 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Lyon	292	31.2 (12)	257	18.7 (14)	61	* *	236	10.6 (13)	257	6.2 (10)
Marion	60	* *	50	* *	8	* *	48	* *	50	* *
Marshall	51	* *	45	* *	21	* *	41	* *	45	* *
Mc Pherson	125	33.6 (14)	110	17.3 (13)	31	* *	105	5.7 (2)	110	0.9 (1)
Meade	35	* *	32	* *	3	* *	36	* *	33	* *
Miami	143	39.9 (22)	121	19.8 (16)	36	* *	119	24.4 (27)	121	17.4 (25)
Mitchell	36	* *	31	* *	12	* *	31	* *	31	* *
Montgome-C	157	40.1 (24)	128	28.9 (30)	42	* *	135	37.8 (32)	128	35.9 (31)
Montgome-I	169	43.8 (32)	150	24.0 (24)	52	* *	147	38.1 (33)	149	30.2 (28)
Morris	39	* *	35	* *	14	* *	32	* *	34	* *
Nemaha	23	* *	14	* *	3	* *	20	* *	15	* *
Neosho-Ch	129	41.9 (28)	71	* *	25	* *	117	33.3 (30)	73	* *
Neosho-Er	10	* *	3	* *	2	* *	9	* *	3	* *
Ness	17	* *	14	* *	3	* *	17	* *	14	* *
Norton	37	* *	34	* *	13	* *	32	* *	34	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2010 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Osage	76	* *	57	* *	17	* *	62	* *	57	* *
Osborne	26	* *	23	* *	8	* *	18	* *	23	* *
Ottawa	28	* *	12	* *	6	* *	25	* *	23	* *
Pawnee	55	* *	37	* *	7	* *	48	* *	38	* *
Phillips	36	* *	23	* *	8	* *	31	* *	23	* *
Pottawat-S	18	* *	16	* *	2	* *	15	* *	16	* *
Pottawat-W	18	* *	16	* *	6	* *	17	* *	17	* *
Pratt	79	* *	67	* *	18	* *	66	* *	66	* *
Rawlins	7	* *	6	* *	1	* *	5	* *	6	* *
Reno	448	41.7 (27)	392	21.9 (20)	139	41.0 (4)	378	13.5 (17)	392	11.2 (21)
Republic	27	* *	21	* *	6	* *	22	* *	21	* *
Rice	85	* *	73	* *	20	* *	66	* *	74	* *
Riley-Ft	574	30.1 (11)	354	14.7 (9)	71	* *	455	19.6 (24)	354	13.8 (23)
Riley-Manh	387	27.6 (9)	310	12.6 (8)	69	* *	319	9.4 (9)	312	6.1 (9)
Rooks	48	* *	37	* *	19	* *	40	* *	39	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2010 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Rush	10	* *	10	* *	7	* *	10	* *	10	* *
Russell	44	* *	39	* *	16	* *	39	* *	39	* *
Saline	543	39.4 (21)	470	25.3 (26)	165	29.7 (8)	478	10.9 (14)	469	7.7 (13)
Scott	34	* *	22	* *	8	* *	30	* *	22	* *
Sedgwick-C	1,238	27.9 (10)	1,103	16.2 (11)	231	32.5 (6)	1,032	14.5 (19)	1,101	11.9 (22)
Sedgwick-M	1,890	27.5 (8)	1,711	12.2 (7)	363	42.7 (3)	1,570	7.1 (3)	1,708	4.4 (6)
Sedgwick-S	1,390	32.7 (13)	1,251	21.4 (18)	317	23.0 (9)	1,140	7.3 (4)	1,246	4.8 (7)
Seneca	8	* *	6	* *	2	* *	8	* *	6	* *
Seward	422	5.7 (1)	377	3.2 (1)	17	* *	342	10.2 (11)	382	8.9 (15)
ShawneeM&I	333	27.0 (7)	254	15.4 (10)	61	* *	288	15.3 (22)	255	3.1 (3)
ShawneeMa	1,053	34.0 (16)	825	24.0 (24)	205	21.5 (10)	831	8.2 (6)	837	4.3 (5)
Sheridan	8	* *	6	* *	2	* *	8	* *	6	* *
Sherman	70	* *	59	* *	20	* *	54	* *	59	* *
Smith	13	* *	10	* *	5	* *	12	* *	10	* *
Stafford	31	* *	27	* *	6	* *	25	* *	27	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2010 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Stanton	16	* *	15	* *	1	* *	12	* *	15	* *
Stevens	70	* *	60	* *	4	* *	60	* *	60	* *
Sumner	163	54.6 (38)	144	34.0 (35)	63	* *	135	44.4 (37)	144	44.4 (35)
Thomas	50	* *	46	* *	9	* *	38	* *	47	* *
Trego	14	* *	10	* *	4	* *	13	* *	10	* *
Wabaunsee	17	* *	16	* *	6	* *	14	* *	16	* *
Washington	22	* *	19	* *	1	* *	18	* *	19	* *
Wichita	17	* *	16	* *	4	* *	12	* *	16	* *
Wilson	74	* *	69	* *	35	* *	63	* *	69	* *
Woodson	15	* *	9	* *	0	* *	13	* *	10	* *
Wyandotte	2,056	20.6 (4)	1,754	10.9 (5)	271	32.5 (6)	1,761	14.7 (20)	1,807	10.6 (18)
Unknown Clinics	19	* *	19	* *	10	* *	17	* *	20	* *
Kansas	21,283	30.7	17,950	17.5	4,227	33.0	17,640	16.5	18,063	13.3
Nation (Prior Year)	1,040,568	22.9	838,781	13.0	156,709	36.7	650,648	17.1	559,995	16.8

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators ^(2, 3)										
Clinic	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)		Number	%	Number	%	Number	%
	Number	% (Rank)	% (Rank)	(Rank)						
Allen	69	* * * *	* * * *	73	* *	62	* *	72	* *	
Anderson	43	* * * *	* * * *	51	* *	39	* *	50	* *	
Atchison	115	9.6 (35)	3.5 (3)	122	6.6 (6)	106	7.5 (33)	121	48.8 (38)	
Barber-Kio	2	* * * *	* * * *	2	* *	2	* *	2	* *	
Barber-Med	41	* * * *	* * * *	41	* *	36	* *	41	* *	
Barton	215	7.4 (22)	4.2 (6)	216	5.6 (4)	202	5.9 (31)	217	77.9 (9)	
Bourbon	103	13.6 (37)	1.9 (1)	106	10.4 (33)	92	* *	105	62.9 (30)	
Brown	67	* * * *	* * * *	73	* *	54	* *	73	* *	
Butler	252	7.1 (18)	6.0 (17)	257	10.1 (32)	226	3.5 (12)	256	65.2 (28)	
Chase	8	* * * *	* * * *	6	* *	6	* *	8	* *	
Chautauqua	15	* * * *	* * * *	16	* *	15	* *	16	* *	
Cherokee	160	9.4 (33)	5.6 (13)	163	9.2 (26)	145	2.8 (5)	163	53.4 (36)	
Cheyenne	7	* * * *	* * * *	7	* *	5	* *	7	* *	
Clark	2	* * * *	* * * *	2	* *	2	* *	2	* *	

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators ^(2, 3)										
Clinic	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)		Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
	Number	% (Rank)	% (Rank)	Number						
Clay	45	* * * *	* * * *	45	* *	40	* *	45	* *	
Cloud	84	* * * *	* * * *	84	* *	76	* *	83	* *	
Coffey	25	* * * *	* * * *	25	* *	25	* *	25	* *	
Comanche	7	* * * *	* * * *	7	* *	7	* *	7	* *	
Cowley-Ark	188	5.9 (8)	6.9 (23)	191	9.9 (31)	168	3.6 (14)	191	49.7 (37)	
Cowley-Win	114	7.9 (23)	9.6 (35)	126	11.9 (36)	99	* *	126	67.5 (24)	
Crawford-G	2	* * * *	* * * *	2	* *	2	* *	2	* *	
Crawford-P	275	5.8 (7)	3.3 (2)	280	10.7 (34)	245	3.3 (8)	280	62.5 (31)	
Decatur	13	* * * *	* * * *	13	* *	12	* *	13	* *	
DickinsonA	113	6.2 (9)	8.8 (34)	111	11.7 (35)	98	* *	113	72.6 (16)	
DickinsonH	16	* * * *	* * * *	16	* *	14	* *	16	* *	
Doniphan	48	* * * *	* * * *	48	* *	39	* *	48	* *	
Douglas	435	8.3 (27)	8.5 (31)	441	7.7 (13)	400	5.3 (29)	438	79.5 (6)	
Edwards	19	* * * *	* * * *	18	* *	15	* *	19	* *	

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators ^(2, 3)										
Clinic	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)		Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
	Number	% (Rank)	% (Rank)	Number						
Elk	20	* * * *	* * * *	21	* *	19	* *	21	* *	
Ellis	130	6.2 (9)	6.2 (20)	136	8.8 (23)	118	3.4 (9)	136	80.1 (5)	
Ellsworth	32	* * * *	* * * *	32	* *	32	* *	32	* *	
Finney	598	7.2 (20)	6.0 (17)	611	9.2 (26)	539	4.5 (20)	613	77.8 (10)	
Ford	564	5.0 (3)	6.0 (17)	580	7.1 (8)	523	2.9 (6)	575	77.4 (11)	
Franklin	153	5.2 (5)	5.9 (16)	153	5.2 (3)	145	4.8 (25)	153	60.8 (33)	
Geary	326	6.4 (12)	8.3 (30)	327	8.9 (24)	297	2.0 (3)	324	71.0 (19)	
Gove	6	* * * *	* * * *	6	* *	5	* *	6	* *	
Graham	15	* * * *	* * * *	15	* *	12	* *	14	* *	
Grant	60	* * * *	* * * *	60	* *	56	* *	61	* *	
Gray	24	* * * *	* * * *	36	* *	23	* *	36	* *	
Greeley	9	* * * *	* * * *	9	* *	7	* *	9	* *	
Greenwood	39	* * * *	* * * *	39	* *	37	* *	39	* *	
Hamilton	25	* * * *	* * * *	26	* *	24	* *	26	* *	

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators ^(2, 3)										
Clinic	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)		Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
	Number	% (Rank)	% (Rank)	Number						
Harper	45	* * * *	* * * *	43	* *	41	* *	45	* *	
Harvey	237	8.4 (28)	4.2 (6)	240	6.7 (7)	219	4.6 (21)	241	83.4 (2)	
Haskell	15	* * * *	* * * *	15	* *	14	* *	15	* *	
Hodgeman	9	* * * *	* * * *	9	* *	9	* *	9	* *	
Jackson	76	* * * *	* * * *	80	* *	73	* *	79	* *	
Jefferson	49	* * * *	* * * *	49	* *	46	* *	48	* *	
Jewell	6	* * * *	* * * *	7	* *	5	* *	7	* *	
Johnson-M	660	7.9 (23)	5.5 (11)	716	7.3 (10)	607	4.1 (17)	712	78.7 (8)	
Johnson-O	941	6.6 (16)	7.5 (28)	974	7.3 (10)	869	3.1 (7)	971	84.0 (1)	
Kearny	19	* * * *	* * * *	20	* *	16	* *	20	* *	
Kingman	21	* * * *	* * * *	41	* *	20	* *	41	* *	
Kiowa	7	* * * *	* * * *	8	* *	7	* *	8	* *	
Labette	152	7.2 (20)	7.9 (29)	151	9.3 (29)	137	5.1 (26)	152	57.2 (35)	
Lane	10	* * * *	* * * *	10	* *	7	* *	10	* *	

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators ^(2, 3)										
Clinic	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)		Number	%	Number	%	Number	%
	Number	% (Rank)	% (Rank)	(Rank)						
Leavenwort	300	5.3 (6)	8.7 (32)		304	9.2 (26)	272	2.6 (4)	302	66.2 (26)
Lincoln	18	* *	* *		18	* *	14	* *	18	* *
Linn	58	* *	* *		62	* *	50	* *	62	* *
Logan	11	* *	* *		11	* *	11	* *	11	* *
Lyon	252	8.7 (30)	6.7 (22)		252	8.3 (17)	231	5.2 (27)	252	77.0 (13)
Marion	46	* *	* *		46	* *	42	* *	47	* *
Marshall	45	* *	* *		48	* *	40	* *	47	* *
Mc Pherson	105	4.8 (2)	4.8 (8)		107	7.5 (12)	97	* *	106	77.4 (11)
Meade	32	* *	* *		36	* *	28	* *	36	* *
Miami	132	3.0 (1)	10.6 (37)		132	4.5 (1)	126	0.8 (1)	132	75.0 (15)
Mitchell	28	* *	* *		32	* *	26	* *	31	* *
Montgome-C	140	7.1 (18)	3.6 (4)		142	8.5 (18)	128	3.9 (16)	142	63.4 (29)
Montgome-I	156	6.4 (12)	3.8 (5)		157	5.1 (2)	148	3.4 (9)	154	61.0 (32)
Morris	30	* *	* *		34	* *	27	* *	35	* *

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
- (4) <2500 g.
- (5) >4000 g.
- (6) <37 weeks gestation.
- (7) 37 or more weeks gestation and <2500 g.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators ^(2, 3)										
Clinic	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)		Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
	Number	% (Rank)	% (Rank)	Number						
Nemaha	12	* * * *	* * * *	25	* *	11	* *	22	* *	
Neosho-Ch	120	9.2 (32)	5.8 (15)	134	9.0 (25)	107	4.7 (24)	133	57.9 (34)	
Neosho-Er	10	* * * *	* * * *	10	* *	10	* *	10	* *	
Ness	16	* * * *	* * * *	18	* *	16	* *	18	* *	
Norton	34	* * * *	* * * *	34	* *	30	* *	33	* *	
Osage	70	* * * *	* * * *	76	* *	59	* *	76	* *	
Osborne	22	* * * *	* * * *	22	* *	20	* *	22	* *	
Ottawa	28	* * * *	* * * *	29	* *	26	* *	29	* *	
Pawnee	46	* * * *	* * * *	45	* *	40	* *	46	* *	
Phillips	31	* * * *	* * * *	32	* *	31	* *	30	* *	
Pottawat-S	15	* * * *	* * * *	15	* *	14	* *	15	* *	
Pottawat-W	17	* * * *	* * * *	17	* *	16	* *	17	* *	
Pratt	70	* * * *	* * * *	72	* *	65	* *	72	* *	
Rawlins	5	* * * *	* * * *	5	* *	5	* *	5	* *	

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators ^(2, 3)									
Clinic	Birthweight			Preterm ⁽⁶⁾		Full Term LBW ⁽⁷⁾		Ever Breastfed	
	Low ⁽⁴⁾		High ⁽⁵⁾						
	Number	% ^(Rank)	% ^(Rank)	Number	% ^(Rank)	Number	% ^(Rank)	Number	% ^(Rank)
Reno	388	9.0 (31)	7.0 (25)	395	8.6 (20)	354	5.4 (30)	394	70.3 (21)
Republic	21	* *	* *	22	* *	19	* *	22	* *
Rice	69	* *	* *	71	* *	63	* *	72	* *
Riley-Ft	470	6.2 (9)	8.7 (32)	482	7.1 (8)	436	3.4 (9)	478	75.9 (14)
Riley-Manh	322	5.0 (3)	10.2 (36)	327	6.4 (5)	301	1.0 (2)	324	82.7 (3)
Rooks	39	* *	* *	38	* *	35	* *	39	* *
Rush	10	* *	* *	10	* *	10	* *	10	* *
Russell	37	* *	* *	36	* *	30	* *	37	* *
Saline	471	6.4 (12)	5.5 (11)	476	8.0 (15)	432	4.2 (18)	475	65.9 (27)
Scott	26	* *	* *	26	* *	24	* *	26	* *
Sedgwick-C	1,079	9.4 (33)	7.2 (27)	1,091	8.6 (20)	983	5.2 (27)	1,087	69.0 (23)
Sedgwick-M	1,672	10.4 (36)	6.9 (23)	1,688	8.7 (22)	1,524	6.9 (32)	1,685	71.3 (18)
Sedgwick-S	1,225	8.6 (29)	7.1 (26)	1,226	8.2 (16)	1,120	4.6 (21)	1,221	70.4 (20)
Seneca	3	* *	* *	8	* *	2	* *	7	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators ^(2, 3)									
Clinic	Birthweight			Preterm ⁽⁶⁾		Full Term LBW ⁽⁷⁾		Ever Breastfed	
	Low ⁽⁴⁾	High ⁽⁵⁾		Number	%	Number	%	Number	%
	Number	% ^(Rank)	% ^(Rank)						
Seward	346	6.4 (12)	6.6 (21)	368	7.9 (14)	313	3.8 (15)	370	78.9 (7)
ShawneeM&I	262	8.0 (25)	11.8 (38)	260	12.3 (37)	227	3.5 (12)	260	81.9 (4)
ShawneeMa	987	7.0 (17)	4.8 (8)	1,002	8.5 (18)	901	4.3 (19)	994	67.5 (24)
Sheridan	7	*	*	7	*	6	*	7	*
Sherman	56	*	*	56	*	52	*	56	*
Smith	12	*	*	12	*	11	*	12	*
Stafford	27	*	*	27	*	25	*	27	*
Stanton	14	*	*	14	*	14	*	14	*
Stevens	39	*	*	59	*	38	*	58	*
Sumner	141	15.6 (38)	5.0 (10)	145	14.5 (38)	120	9.2 (34)	144	70.1 (22)
Thomas	44	*	*	44	*	41	*	44	*
Trego	11	*	*	11	*	9	*	11	*
Wabaunsee	16	*	*	16	*	16	*	16	*
Washington	16	*	*	18	*	15	*	18	*

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators ^(2, 3)										
Clinic	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)							
	Number	% (Rank)	% (Rank)		Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Wichita	15	* * * *	* *		15	* *	14	* *	15	* *
Wilson	71	* * * *	* *		71	* *	63	* *	71	* *
Woodson	15	* * * *	* *		18	* *	15	* *	18	* *
Wyandotte	1,715	8.2 (26)	5.7 (14)		1,796	9.6 (30)	1,541	4.6 (21)	1,794	71.9 (17)
Unknown Clinics	20	* * * *	* *		20	* *	20	* *	19	* *
Kansas	18,226	7.6	6.6		18,726	8.4	16,603	4.4	18,670	72.2
Nation (Prior Year)	1,018,145	8.2	6.7		1,017,333	11.1	848,957	3.5	997,136	66.7

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3A
2008 - 2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Racial and Ethnic Distribution by Clinic
Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Clinic	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaska Native	Asian/Pacific Islander	Multiple Races	All Other/Unknown
		%	%	%	%	%	%	%
Allen	287	92.0	3.8	1.4	1.4	0.0	1.4	0.0
Anderson	156	94.9	0.0	3.8	0.0	1.3	0.0	0.0
Atchison	413	88.9	7.5	1.5	0.7	0.0	1.5	0.0
Barber-Kio	4	*	*	*	*	*	*	*
Barber-Med	137	94.9	2.2	1.5	0.0	1.5	0.0	0.0
Barton	791	68.1	2.1	28.4	0.3	0.5	0.5	0.0
Bourbon	379	92.6	2.1	2.6	0.5	0.5	1.6	0.0
Brown	300	76.3	2.3	4.3	14.3	0.0	2.7	0.0
Butler	853	90.3	1.2	6.1	0.5	0.2	1.8	0.0
Chase	40	*	*	*	*	*	*	*
Chautauqua	83	*	*	*	*	*	*	*
Cherokee	614	90.1	1.3	3.3	1.6	2.3	1.5	0.0
Cheyenne	13	*	*	*	*	*	*	*
Clark	10	*	*	*	*	*	*	*
Clay	177	94.4	0.6	2.3	2.3	0.0	0.6	0.0
Cloud	254	92.5	1.6	2.8	1.6	0.8	0.8	0.0
Coffey	82	*	*	*	*	*	*	*
Comanche	28	*	*	*	*	*	*	*
Cowley-Ark	617	65.6	2.8	29.8	0.8	0.5	0.5	0.0
Cowley-Win	465	87.1	2.2	5.4	0.9	3.2	1.3	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3A
2008 - 2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Racial and Ethnic Distribution by Clinic
Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Clinic	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaska Native	Asian/Pacific Islander	Multiple Races	All Other/Unknown
		%	%	%	%	%	%	%
Crawford-G	4	*	*	*	*	*	*	*
Crawford-P	1,118	83.4	3.6	10.2	0.4	1.2	1.3	0.0
Decatur	33	*	*	*	*	*	*	*
DickinsonA	352	91.8	0.0	5.7	1.1	0.6	0.9	0.0
DickinsonH	42	*	*	*	*	*	*	*
Doniphan	133	88.0	2.3	6.8	0.8	0.0	2.3	0.0
Douglas	1,500	65.5	8.7	10.7	8.5	4.5	2.1	0.0
Edwards	77	*	*	*	*	*	*	*
Elk	57	*	*	*	*	*	*	*
Ellis	481	83.8	1.0	13.3	0.8	0.4	0.6	0.0
Ellsworth	98	*	*	*	*	*	*	*
Finney	2,109	29.7	1.0	65.5	0.0	3.3	0.4	0.0
Ford	2,029	21.8	1.5	76.0	0.2	0.4	0.0	0.0
Franklin	538	91.4	0.9	4.5	1.5	0.2	1.5	0.0
Geary	1,154	53.9	22.7	15.3	0.6	3.2	4.3	0.0
Gove	29	*	*	*	*	*	*	*
Graham	51	*	*	*	*	*	*	*
Grant	205	22.4	0.0	77.1	0.0	0.0	0.5	0.0
Gray	148	68.9	0.0	30.4	0.0	0.0	0.7	0.0
Greeley	37	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3A
2008 - 2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Racial and Ethnic Distribution by Clinic
Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Clinic	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaska Native	Asian/Pacific Islander	Multiple Races	All Other/Unknown
		%	%	%	%	%	%	%
Greenwood	140	93.6	0.7	4.3	0.0	0.0	1.4	0.0
Hamilton	76	*	*	*	*	*	*	*
Harper	156	80.1	0.0	17.9	0.6	0.0	1.3	0.0
Harvey	818	70.3	2.4	23.7	1.2	0.7	1.6	0.0
Haskell	61	*	*	*	*	*	*	*
Hodgeman	39	*	*	*	*	*	*	*
Jackson	233	76.8	0.0	5.6	15.5	0.0	2.1	0.0
Jefferson	191	94.2	0.5	2.1	2.6	0.0	0.5	0.0
Jewell	30	*	*	*	*	*	*	*
Johnson-M	2,557	47.1	15.1	32.5	0.7	2.8	1.8	0.0
Johnson-O	3,193	45.8	11.3	37.6	0.4	3.5	1.3	0.0
Kearny	61	*	*	*	*	*	*	*
Kingman	165	90.9	0.0	6.7	1.2	1.2	0.0	0.0
Kiowa	33	*	*	*	*	*	*	*
Labette	536	87.9	6.2	3.2	1.5	0.2	1.1	0.0
Lane	37	*	*	*	*	*	*	*
Leavenworth	1,079	69.0	14.6	10.6	0.6	1.7	3.7	0.0
Lincoln	63	*	*	*	*	*	*	*
Linn	191	95.8	0.0	2.1	1.0	0.0	1.0	0.0
Logan	40	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3A
2008 - 2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Racial and Ethnic Distribution by Clinic
Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Clinic	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaska Native	Asian/Pacific Islander	Multiple Races	All Other/Unknown
		%	%	%	%	%	%	%
Lyon	960	57.6	4.8	34.8	0.3	1.6	0.9	0.0
Marion	154	90.9	0.0	7.8	0.6	0.6	0.0	0.0
Marshall	163	92.0	0.0	2.5	3.1	0.0	2.5	0.0
Mc Pherson	414	87.4	1.7	9.2	0.7	0.5	0.5	0.0
Meade	131	67.2	0.8	31.3	0.0	0.0	0.8	0.0
Miami	433	91.0	2.8	3.5	0.2	0.9	1.6	0.0
Mitchell	133	92.5	0.0	7.5	0.0	0.0	0.0	0.0
Montgome-C	470	71.7	11.5	11.9	2.8	0.9	1.3	0.0
Montgome-I	549	82.3	6.7	7.1	2.2	0.9	0.7	0.0
Morris	94	*	*	*	*	*	*	*
Nemaha	87	*	*	*	*	*	*	*
Neosho-Ch	433	91.9	2.5	3.5	0.7	0.2	1.2	0.0
Neosho-Er	32	*	*	*	*	*	*	*
Ness	35	*	*	*	*	*	*	*
Norton	96	*	*	*	*	*	*	*
Osage	269	96.3	0.0	1.9	0.7	0.0	1.1	0.0
Osborne	59	*	*	*	*	*	*	*
Ottawa	79	*	*	*	*	*	*	*
Pawnee	163	78.5	2.5	17.2	0.6	0.0	1.2	0.0
Phillips	110	91.8	0.0	5.5	0.0	1.8	0.9	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3A
2008 - 2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Racial and Ethnic Distribution by Clinic
Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Clinic	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaska Native	Asian/Pacific Islander	Multiple Races	All Other/Unknown
		%	%	%	%	%	%	%
Pottawat-S	48	*	*	*	*	*	*	*
Pottawat-W	66	*	*	*	*	*	*	*
Pratt	235	86.0	0.0	13.6	0.0	0.4	0.0	0.0
Rawlins	29	*	*	*	*	*	*	*
Reno	1,446	79.7	2.2	15.4	0.6	0.7	1.5	0.0
Republic	103	91.3	1.0	5.8	0.0	0.0	1.9	0.0
Rice	234	73.9	2.1	22.2	0.9	0.9	0.0	0.0
Riley-Ft	1,586	70.6	12.1	13.9	0.4	2.9	0.1	0.0
Riley-Manh	1,203	73.3	10.4	9.7	0.7	4.9	1.0	0.0
Rooks	136	93.4	1.5	2.2	2.2	0.7	0.0	0.0
Rush	31	*	*	*	*	*	*	*
Russell	126	90.5	5.6	2.4	1.6	0.0	0.0	0.0
Saline	1,570	71.5	4.2	20.0	0.2	2.2	1.9	0.0
Scott	123	61.8	0.8	37.4	0.0	0.0	0.0	0.0
Sedgwick-C	3,550	42.4	13.6	33.8	1.2	6.9	2.1	0.0
Sedgwick-M	5,912	32.8	27.9	33.8	1.3	2.7	1.5	0.0
Sedgwick-S	3,959	66.2	8.0	19.5	2.1	2.4	1.9	0.0
Seneca	31	*	*	*	*	*	*	*
Seward	1,456	15.4	1.6	80.6	1.5	0.7	0.2	0.0
ShawneeM&I	788	35.8	13.8	46.6	1.6	1.3	0.9	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3A
2008 - 2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Racial and Ethnic Distribution by Clinic
Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Clinic	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaska Native	Asian/Pacific Islander	Multiple Races	All Other/Unknown
		%	%	%	%	%	%	%
ShawneeMa	3,577	59.1	15.8	20.1	2.4	0.8	1.9	0.0
Sheridan	25	*	*	*	*	*	*	*
Sherman	202	73.8	0.0	24.3	0.0	1.0	1.0	0.0
Smith	52	*	*	*	*	*	*	*
Stafford	116	68.1	0.9	30.2	0.0	0.9	0.0	0.0
Stanton	53	*	*	*	*	*	*	*
Stevens	225	45.3	0.0	52.9	1.3	0.4	0.0	0.0
Sumner	479	88.9	0.8	6.7	1.5	0.4	1.7	0.0
Thomas	160	87.5	1.9	8.1	0.6	0.6	1.3	0.0
Trego	46	*	*	*	*	*	*	*
Wabaunsee	60	*	*	*	*	*	*	*
Wallace	1	*	*	*	*	*	*	*
Washington	74	*	*	*	*	*	*	*
Wichita	47	*	*	*	*	*	*	*
Wilson	257	96.1	0.8	1.9	0.4	0.0	0.8	0.0
Woodson	28	*	*	*	*	*	*	*
Wyandotte	6,498	26.7	30.1	39.4	0.3	2.8	0.6	0.0
Unknown Clinics	23	*	*	*	*	*	*	*
Kansas	65,016	56.9	11.3	27.2	1.2	2.1	1.3	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2008 - 2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Age Distribution by Clinic
Combined 3 Years

Comparison of Age Distribution ⁽²⁾								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Allen	287	0.0	8.4	16.7	62.0	11.5	1.0	0.3
Anderson	156	0.6	5.1	15.4	62.8	16.0	0.0	0.0
Atchison	413	0.0	7.3	15.5	62.2	14.3	0.7	0.0
Barber-Kio	4	*	*	*	*	*	*	*
Barber-Med	137	0.0	5.8	20.4	62.8	10.2	0.7	0.0
Barton	791	0.0	7.5	15.4	59.9	15.7	1.5	0.0
Bourbon	379	0.3	6.6	18.2	61.5	12.9	0.5	0.0
Brown	300	0.3	7.3	13.7	63.7	14.0	1.0	0.0
Butler	853	0.1	6.8	16.9	61.9	13.6	0.5	0.2
Chase	40	*	*	*	*	*	*	*
Chautauqua	83	*	*	*	*	*	*	*
Cherokee	614	0.3	9.3	19.9	57.2	12.4	0.7	0.3
Cheyenne	13	*	*	*	*	*	*	*
Clark	10	*	*	*	*	*	*	*
Clay	177	1.1	8.5	16.4	59.9	13.6	0.6	0.0
Cloud	254	0.4	7.5	16.5	59.8	13.4	2.4	0.0
Coffey	82	*	*	*	*	*	*	*
Comanche	28	*	*	*	*	*	*	*
Cowley-Ark	617	0.6	9.2	16.7	58.8	13.8	0.8	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2008 - 2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Age Distribution by Clinic
Combined 3 Years

Comparison of Age Distribution ⁽²⁾								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Cowley-Win	465	0.6	6.0	15.9	65.4	12.0	0.0	0.0
Crawford-G	4	*	*	*	*	*	*	*
Crawford-P	1,118	0.2	6.8	14.8	65.0	12.7	0.4	0.1
Decatur	33	*	*	*	*	*	*	*
DickinsonA	352	0.3	4.8	13.1	65.3	16.2	0.3	0.0
DickinsonH	42	*	*	*	*	*	*	*
Doniphan	133	0.8	4.5	13.5	63.2	18.0	0.0	0.0
Douglas	1,500	0.0	4.1	12.5	63.7	18.4	1.4	0.0
Edwards	77	*	*	*	*	*	*	*
Elk	57	*	*	*	*	*	*	*
Ellis	481	0.4	4.6	13.1	67.6	12.7	1.5	0.2
Ellsworth	98	*	*	*	*	*	*	*
Finney	2,109	0.4	8.7	15.3	55.1	18.6	1.8	0.1
Ford	2,029	0.3	7.0	13.0	54.2	23.5	1.9	0.0
Franklin	538	0.4	5.9	18.0	60.0	14.7	0.7	0.2
Geary	1,154	0.3	4.8	14.9	67.9	11.4	0.6	0.1
Gove	29	*	*	*	*	*	*	*
Graham	51	*	*	*	*	*	*	*
Grant	205	0.0	6.3	13.2	63.4	14.1	2.9	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2008 - 2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Age Distribution by Clinic
Combined 3 Years

Comparison of Age Distribution ⁽²⁾								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Gray	148	0.0	5.4	10.1	60.1	23.0	1.4	0.0
Greeley	37	*	*	*	*	*	*	*
Greenwood	140	0.0	6.4	17.1	62.9	13.6	0.0	0.0
Hamilton	76	*	*	*	*	*	*	*
Harper	156	0.0	6.4	10.3	73.1	10.3	0.0	0.0
Harvey	818	0.2	5.9	13.4	63.4	15.8	1.2	0.0
Haskell	61	*	*	*	*	*	*	*
Hodgeman	39	*	*	*	*	*	*	*
Jackson	233	0.0	5.2	18.9	63.5	12.0	0.4	0.0
Jefferson	191	0.0	5.2	18.8	58.6	16.8	0.5	0.0
Jewell	30	*	*	*	*	*	*	*
Johnson-M	2,557	0.3	3.8	11.6	61.1	21.9	1.3	0.0
Johnson-O	3,193	0.1	3.8	10.2	59.8	24.1	1.9	0.1
Kearny	61	*	*	*	*	*	*	*
Kingman	165	0.0	7.3	15.8	63.6	12.1	1.2	0.0
Kiowa	33	*	*	*	*	*	*	*
Labette	536	0.0	7.5	20.1	61.2	10.4	0.7	0.0
Lane	37	*	*	*	*	*	*	*
Leavenworth	1,079	0.1	6.0	15.6	64.7	12.5	1.1	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2008 - 2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Age Distribution by Clinic
Combined 3 Years

Comparison of Age Distribution ⁽²⁾								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Lincoln	63	*	*	*	*	*	*	*
Linn	191	0.0	6.3	15.7	61.8	15.7	0.5	0.0
Logan	40	*	*	*	*	*	*	*
Lyon	960	0.6	6.3	14.2	62.3	14.5	2.2	0.0
Marion	154	0.0	3.2	13.6	64.9	14.3	3.9	0.0
Marshall	163	0.0	6.1	12.9	68.1	12.3	0.6	0.0
Mc Pherson	414	0.5	3.9	12.8	64.5	17.4	1.0	0.0
Meade	131	0.0	9.9	9.2	58.0	21.4	1.5	0.0
Miami	433	0.5	5.3	15.7	64.0	13.9	0.7	0.0
Mitchell	133	0.0	1.5	18.0	65.4	15.0	0.0	0.0
Montgome-C	470	0.6	5.3	17.9	61.9	13.2	0.9	0.2
Montgome-I	549	0.0	5.5	17.1	65.4	10.7	1.1	0.2
Morris	94	*	*	*	*	*	*	*
Nemaha	87	*	*	*	*	*	*	*
Neosho-Ch	433	0.2	6.5	18.5	62.4	12.0	0.5	0.0
Neosho-Er	32	*	*	*	*	*	*	*
Ness	35	*	*	*	*	*	*	*
Norton	96	*	*	*	*	*	*	*
Osage	269	0.0	5.2	20.4	62.1	10.8	1.1	0.4

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2008 - 2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Age Distribution by Clinic
Combined 3 Years

Comparison of Age Distribution ⁽²⁾								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Osborne	59	*	*	*	*	*	*	*
Ottawa	79	*	*	*	*	*	*	*
Pawnee	163	1.2	6.7	12.3	68.1	11.0	0.6	0.0
Phillips	110	0.0	3.6	17.3	62.7	15.5	0.9	0.0
Pottawat-S	48	*	*	*	*	*	*	*
Pottawat-W	66	*	*	*	*	*	*	*
Pratt	235	0.4	8.1	14.5	61.3	14.9	0.9	0.0
Rawlins	29	*	*	*	*	*	*	*
Reno	1,446	0.1	6.8	15.6	64.0	12.9	0.6	0.0
Republic	103	0.0	3.9	20.4	56.3	16.5	2.9	0.0
Rice	234	0.4	5.6	13.2	63.7	16.7	0.4	0.0
Riley-Ft	1,586	0.0	0.8	11.2	76.5	11.2	0.3	0.0
Riley-Manh	1,203	0.1	3.0	13.4	70.7	12.5	0.4	0.0
Rooks	136	0.0	2.2	15.4	62.5	17.6	1.5	0.7
Rush	31	*	*	*	*	*	*	*
Russell	126	0.0	10.3	16.7	62.7	10.3	0.0	0.0
Saline	1,570	0.1	6.7	15.9	62.4	14.0	1.0	0.1
Scott	123	0.8	3.3	13.0	68.3	13.8	0.8	0.0
Sedgwick-C	3,550	0.2	6.8	12.7	59.7	19.2	1.2	0.1

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2008 - 2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Age Distribution by Clinic
Combined 3 Years

Comparison of Age Distribution ⁽²⁾								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Sedgwick-M	5,912	0.3	6.9	14.0	60.4	17.0	1.4	0.0
Sedgwick-S	3,959	0.4	6.4	14.7	62.0	15.2	1.3	0.0
Seneca	31	*	*	*	*	*	*	*
Seward	1,456	0.2	7.3	13.6	56.6	20.9	1.4	0.0
ShawneeM&I	788	0.8	7.7	16.1	54.4	19.4	1.5	0.0
ShawneeMa	3,577	0.2	6.3	14.2	62.1	16.0	1.1	0.0
Sheridan	25	*	*	*	*	*	*	*
Sherman	202	0.0	5.4	15.8	59.4	17.3	2.0	0.0
Smith	52	*	*	*	*	*	*	*
Stafford	116	0.0	6.0	13.8	60.3	17.2	2.6	0.0
Stanton	53	*	*	*	*	*	*	*
Stevens	225	0.0	7.1	18.2	57.3	16.4	0.9	0.0
Sumner	479	0.0	6.7	16.7	61.2	14.6	0.8	0.0
Thomas	160	0.0	6.3	15.0	68.1	10.6	0.0	0.0
Trego	46	*	*	*	*	*	*	*
Wabaunsee	60	*	*	*	*	*	*	*
Wallace	1	*	*	*	*	*	*	*
Washington	74	*	*	*	*	*	*	*
Wichita	47	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2008 - 2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Age Distribution by Clinic
Combined 3 Years

Comparison of Age Distribution ⁽²⁾								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Wilson	257	0.4	8.2	15.6	62.3	13.6	0.0	0.0
Woodson	28	*	*	*	*	*	*	*
Wyandotte	6,498	0.5	8.4	13.5	57.2	19.0	1.4	0.0
Unknown Clinics	23	*	*	*	*	*	*	*
Kansas	65,016	0.3	6.2	14.2	61.3	16.7	1.2	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2008 - 2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Health Indicators by Clinic
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)														
Clinic	Prepregnancy BMI ⁽⁴⁾			Weight Gain ⁽⁵⁾			Anemia (Low Hb/Hct) ⁽⁶⁾				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Allen	283	4.9 (43)	56.9 (53)	260	18.1 (15)	55.4 (59)	48	* *	82	* *	254	7.5 (58)	254	15.7(66)
Anderson	153	6.5 (62)	51.0 (8)	141	23.4 (56)	46.8 (10)	22	* *	25	* *	136	2.9 (5)	136	14.7(63)
Atchison	401	5.2 (50)	53.9 (28)	357	16.2 (7)	57.4 (64)	61	* *	83	* *	329	5.5 (30)	329	12.8(57)
Barber-Kio	4	* *	* *	3	* *	* *	0	* *	1	* *	3	* *	3	* *
Barber-Med	135	3.0 (10)	51.9 (15)	110	30.9 (67)	48.2 (18)	15	* *	33	* *	109	0.9 (2)	109	16.5(67)
Barton	770	4.9 (43)	55.3 (38)	685	20.9 (38)	49.6 (25)	78	* *	161	26.1(23)	658	5.5 (30)	656	11.6(48)
Bourbon	373	2.1 (5)	56.6 (50)	329	23.7 (58)	48.9 (22)	34	* *	72	* *	317	6.3 (45)	318	11.9(51)
Brown	296	1.7 (4)	57.4 (54)	259	23.2 (55)	51.0 (38)	38	* *	66	* *	257	5.1 (22)	257	13.6(60)
Butler	826	5.9 (57)	51.0 (8)	755	19.7 (28)	53.4 (47)	114	43.9 (22)	188	29.3(28)	723	5.4 (28)	723	11.3(47)
Chase	38	* *	* *	31	* *	* *	4	* *	2	* *	34	* *	34	* *
Chautauqua	82	* *	* *	74	* *	* *	8	* *	13	* *	73	* *	73	* *
Cherokee	580	6.6 (64)	51.9 (15)	517	18.0 (13)	54.0 (50)	72	* *	119	25.2(20)	515	5.6 (33)	515	10.3(35)
Cheyenne	12	* *	* *	9	* *	* *	3	* *	2	* *	10	* *	10	* *
Clark	9	* *	* *	8	* *	* *	2	* *	2	* *	9	* *	9	* *
Clay	172	4.7 (42)	55.8 (47)	151	19.2 (23)	60.9 (67)	25	* *	43	* *	150	8.0 (63)	150	10.7(41)
Cloud	251	4.4 (37)	53.4 (23)	226	20.8 (36)	54.0 (50)	27	* *	66	* *	216	6.9 (53)	221	10.0(33)
Coffey	78	* *	* *	66	* *	* *	8	* *	21	* *	61	* *	61	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2008 - 2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Health Indicators by Clinic
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)																
Clinic	Prepregnancy BMI (4)				Weight Gain (5)				Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight		Over-weight		< Ideal		> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Comanche	28	* *	* *	* *	25	* *	* *	* *	6	* *	7	* *	23	* *	23	* *
Cowley-Ark	606	3.8 (24)	54.0 (31)		542	21.2 (42)	49.6 (25)		84	* *	87	* *	523	7.3 (56)	524	10.3(35)
Cowley-Win	460	5.4 (51)	55.7 (45)		410	20.0 (31)	56.1 (60)		48	* *	73	* *	394	4.1 (10)	395	9.4(29)
Crawford-G	4	* *	* *	* *	3	* *	* *	* *	2	* *	0	* *	3	* *	3	* *
Crawford-P	1,098	6.3 (61)	54.2 (32)		951	20.6 (33)	52.4 (43)		164	18.3 (2)	169	18.9 (7)	936	7.3 (56)	935	10.8(43)
Decatur	33	* *	* *	* *	32	* *	* *	* *	5	* *	8	* *	30	* *	30	* *
DickinsonA	340	3.2 (14)	57.4 (54)		298	22.1 (45)	49.7 (27)		45	* *	70	* *	294	6.8 (51)	290	12.1(52)
DickinsonH	41	* *	* *	* *	36	* *	* *	* *	4	* *	14	* *	38	* *	38	* *
Doniphan	130	0.8 (1)	62.3 (68)		110	13.6 (2)	65.5 (68)		23	* *	25	* *	107	5.6 (33)	107	12.1(52)
Douglas	1,463	3.8 (24)	48.4 (4)		1,253	17.6 (11)	54.7 (53)		199	26.6 (9)	384	26.8(24)	1,218	6.0 (37)	1,217	7.8(19)
Edwards	77	* *	* *	* *	67	* *	* *	* *	13	* *	14	* *	61	* *	61	* *
Elk	56	* *	* *	* *	50	* *	* *	* *	4	* *	11	* *	47	* *	47	* *
Ellis	462	3.9 (28)	51.1 (11)		400	18.3 (16)	49.0 (23)		103	23.3 (4)	111	17.1 (3)	397	2.5 (4)	397	6.3(8)
Ellsworth	96	* *	* *	* *	91	* *	* *	* *	14	* *	22	* *	84	* *	84	* *
Finney	2,072	6.5 (62)	50.9 (7)		1,882	24.5 (61)	45.2 (3)		290	29.7 (14)	297	21.9(13)	1,849	7.6 (61)	1,847	6.2(7)
Ford	1,999	2.2 (6)	57.5 (57)		1,777	23.7 (58)	46.7 (9)		254	21.3 (3)	230	13.5 (1)	1,729	5.3 (27)	1,727	7.6(15)
Franklin	516	3.7 (22)	51.6 (12)		453	16.8 (9)	53.0 (44)		52	* *	136	14.0 (2)	386	4.1 (10)	386	9.8(32)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2008 - 2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Health Indicators by Clinic
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)														
Clinic	Prepregnancy BMI ⁽⁴⁾			Weight Gain ⁽⁵⁾			Anemia (Low Hb/Hct) ⁽⁶⁾				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Geary	1,119	4.6 (40)	53.4 (23)	900	18.0 (13)	53.7 (48)	171	39.2 (20)	197	34.0 (30)	902	4.1 (10)	901	10.7 (41)
Gove	24	* *	* *	21	* *	* *	3	* *	9	* *	26	* *	26	* *
Graham	51	* *	* *	41	* *	* *	7	* *	6	* *	42	* *	42	* *
Grant	198	2.5 (8)	67.2 (71)	183	23.0 (53)	48.1 (17)	24	* *	59	* *	181	6.1 (39)	179	5.0 (2)
Gray	144	3.5 (17)	54.2 (32)	121	23.1 (54)	52.1 (42)	15	* *	39	* *	118	7.6 (61)	118	11.0 (45)
Greeley	35	* *	* *	31	* *	* *	3	* *	5	* *	33	* *	33	* *
Greenwood	135	2.2 (6)	60.0 (63)	123	22.0 (44)	55.3 (57)	13	* *	21	* *	120	5.0 (20)	121	18.2 (68)
Hamilton	73	* *	* *	65	* *	* *	15	* *	22	* *	66	* *	66	* *
Harper	150	6.0 (59)	52.7 (19)	130	18.5 (17)	48.5 (20)	16	* *	21	* *	125	5.6 (33)	124	8.1 (20)
Harvey	803	4.5 (39)	54.5 (36)	713	22.6 (49)	48.0 (16)	108	36.1 (19)	122	21.3 (10)	689	4.6 (16)	686	10.3 (35)
Haskell	61	* *	* *	55	* *	* *	8	* *	16	* *	52	* *	52	* *
Hodgeman	37	* *	* *	35	* *	* *	6	* *	7	* *	33	* *	33	* *
Jackson	224	3.6 (20)	55.4 (42)	206	19.9 (30)	56.8 (63)	32	* *	48	* *	203	8.4 (65)	204	14.7 (63)
Jefferson	188	3.7 (22)	58.0 (58)	169	16.0 (4)	58.0 (65)	31	* *	51	* *	165	6.7 (49)	165	9.1 (26)
Jewell	30	* *	* *	30	* *	* *	3	* *	5	* *	29	* *	29	* *
Johnson-M	2,463	4.2 (33)	55.3 (38)	2,184	18.7 (19)	50.2 (30)	398	25.1 (7)	624	17.1 (3)	2,119	6.5 (46)	2,116	6.4 (9)
Johnson-O	3,084	4.2 (33)	52.9 (20)	2,760	18.9 (22)	51.5 (39)	504	23.4 (5)	830	17.6 (5)	2,696	7.1 (54)	2,690	6.1 (6)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2008 - 2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Health Indicators by Clinic
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)														
Clinic	Prepregnancy BMI ⁽⁴⁾			Weight Gain ⁽⁵⁾			Anemia (Low Hb/Hct) ⁽⁶⁾				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Kearny	61	* *	* *	57	* *	* *	8	* *	12	* *	57	* *	57	* *
Kingman	165	3.6 (20)	46.1 (1)	144	22.2 (46)	50.0 (28)	16	* *	27	* *	135	6.7 (49)	135	14.1(61)
Kiowa	31	* *	* *	26	* *	* *	5	* *	7	* *	29	* *	29	* *
Labette	527	5.9 (57)	56.2 (49)	482	24.7 (63)	50.6 (34)	92	* *	118	24.6(19)	465	5.2 (26)	465	12.3(55)
Lane	34	* *	* *	30	* *	* *	6	* *	6	* *	30	* *	30	* *
Leavenwort	1,046	4.2 (33)	55.4 (42)	911	19.5 (25)	55.3 (57)	167	26.9 (10)	266	21.8(12)	860	5.6 (33)	859	9.7(31)
Lincoln	60	* *	* *	56	* *	* *	11	* *	15	* *	52	* *	52	* *
Linn	189	5.8 (55)	62.4 (69)	173	28.9 (66)	46.2 (7)	24	* *	29	* *	169	4.7 (18)	170	5.3(3)
Logan	39	* *	* *	32	* *	* *	6	* *	6	* *	31	* *	31	* *
Lyon	935	3.3 (16)	59.5 (61)	828	21.6 (43)	48.7 (21)	127	11.0 (1)	172	20.3 (9)	800	5.4 (28)	800	12.5(56)
Marion	149	7.4 (68)	48.3 (3)	124	18.5 (17)	48.4 (19)	19	* *	31	* *	123	2.4 (3)	124	8.9(25)
Marshall	159	5.7 (54)	53.5 (25)	144	20.8 (36)	50.0 (28)	21	* *	35	* *	137	5.1 (22)	136	9.6(30)
Mc Pherson	408	5.4 (51)	54.7 (37)	368	22.3 (48)	45.4 (5)	62	* *	92	* *	357	5.0 (20)	359	8.4(23)
Meade	130	3.1 (12)	61.5 (67)	115	22.6 (49)	45.2 (3)	12	* *	24	* *	109	8.3 (64)	109	9.2(27)
Miami	419	6.0 (59)	55.6 (44)	375	16.0 (4)	55.2 (55)	84	* *	120	25.8(21)	367	7.1 (54)	367	10.4(39)
Mitchell	130	3.1 (12)	58.5 (59)	122	32.0 (68)	45.9 (6)	14	* *	34	* *	120	9.2 (66)	119	14.3(62)
Montgome-C	463	3.5 (17)	55.3 (38)	418	23.4 (56)	47.4 (13)	95	* *	122	27.9(25)	398	6.8 (51)	398	9.3(28)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2008 - 2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Health Indicators by Clinic
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)														
Clinic	Prepregnancy BMI ⁽⁴⁾			Weight Gain ⁽⁵⁾			Anemia (Low Hb/Hct) ⁽⁶⁾				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Montgome-I	535	8.4 (71)	52.1 (17)	465	23.9 (60)	47.1 (12)	89	* *	129	22.5 (15)	454	4.6 (16)	454	10.8 (43)
Morris	92	* *	* *	82	* *	* *	13	* *	19	* *	81	* *	81	* *
Nemaha	80	* *	* *	74	* *	* *	21	* *	23	* *	77	* *	77	* *
Neosho-Ch	421	7.1 (66)	53.2 (22)	385	21.0 (40)	50.4 (32)	65	* *	65	* *	369	6.0 (37)	368	13.3 (59)
Neosho-Er	30	* *	* *	28	* *	* *	6	* *	6	* *	29	* *	29	* *
Ness	33	* *	* *	28	* *	* *	3	* *	11	* *	26	* *	26	* *
Norton	96	* *	* *	86	* *	* *	14	* *	27	* *	81	* *	81	* *
Osage	264	4.9 (43)	52.3 (18)	230	16.5 (8)	55.2 (55)	33	* *	56	* *	228	4.8 (19)	228	11.8 (50)
Osborne	58	* *	* *	50	* *	* *	5	* *	13	* *	50	* *	50	* *
Ottawa	75	* *	* *	67	* *	* *	11	* *	13	* *	70	* *	70	* *
Pawnee	159	5.0 (47)	56.6 (50)	132	28.0 (65)	46.2 (7)	13	* *	31	* *	134	7.5 (58)	133	7.5 (14)
Phillips	106	3.8 (24)	60.4 (65)	89	* *	* *	8	* *	20	* *	90	* *	90	* *
Pottawat-S	48	* *	* *	44	* *	* *	5	* *	25	* *	43	* *	43	* *
Pottawat-W	64	* *	* *	58	* *	* *	7	* *	21	* *	56	* *	56	* *
Pratt	230	3.5 (17)	53.5 (25)	196	20.9 (38)	49.0 (23)	29	* *	52	* *	189	0.5 (1)	189	7.4 (12)
Rawlins	28	* *	* *	25	* *	* *	3	* *	9	* *	25	* *	25	* *
Reno	1,423	4.6 (40)	53.9 (28)	1,278	20.6 (33)	50.7 (35)	182	27.5 (12)	242	23.1 (16)	1,210	3.4 (7)	1,211	13.1 (58)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2008 - 2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Health Indicators by Clinic
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Republic	101	6.9 (65)	57.4 (54)	87	* *	* *	15	* *	20	* *	86	* *	86	* *
Rice	228	3.9 (28)	60.1 (64)	198	19.2 (23)	50.5 (33)	28	* *	49	* *	199	6.5 (46)	198	7.6 (15)
Riley-Ft	1,507	5.0 (47)	51.0 (8)	1,184	18.8 (21)	56.3 (61)	254	30.3 (15)	332	32.8 (29)	1,153	3.7 (8)	1,153	12.1 (52)
Riley-Manh	1,167	3.9 (28)	50.6 (6)	993	16.0 (4)	51.7 (40)	164	25.6 (8)	300	21.3 (10)	969	5.1 (22)	969	7.6 (15)
Rooks	136	8.1 (70)	46.3 (2)	122	24.6 (62)	39.3 (1)	20	* *	21	* *	115	6.1 (39)	115	10.4 (39)
Rush	31	* *	* *	30	* *	* *	4	* *	8	* *	29	* *	29	* *
Russell	125	3.2 (14)	50.4 (5)	113	25.7 (64)	47.8 (15)	18	* *	18	* *	108	3.7 (8)	108	4.6 (1)
Saline	1,543	4.9 (43)	55.3 (38)	1,397	17.0 (10)	54.3 (52)	294	24.8 (6)	308	20.1 (8)	1,366	6.2 (43)	1,369	7.4 (12)
Scott	122	4.1 (31)	59.0 (60)	107	19.6 (26)	55.1 (54)	15	* *	19	* *	103	2.9 (5)	103	5.8 (5)
Sedgwick-C	3,466	4.4 (37)	53.5 (25)	3,170	20.1 (32)	50.9 (37)	635	32.1 (17)	754	24.4 (18)	3,125	6.2 (43)	3,125	8.2 (21)
Sedgwick-M	5,785	3.8 (24)	54.3 (34)	5,292	20.6 (33)	50.3 (31)	1,199	28.6 (13)	1,411	22.3 (14)	5,099	5.5 (30)	5,097	10.3 (35)
Sedgwick-S	3,869	5.8 (55)	51.7 (13)	3,536	19.7 (28)	53.9 (49)	736	35.7 (18)	965	28.4 (26)	3,471	5.1 (22)	3,468	11.0 (45)
Seneca	26	* *	* *	25	* *	* *	6	* *	9	* *	28	* *	28	* *
Seward	1,432	2.5 (8)	56.7 (52)	1,290	22.2 (46)	47.7 (14)	238	27.3 (11)	214	24.3 (17)	1,251	7.5 (58)	1,251	6.5 (10)
ShawneeM&I	783	4.1 (31)	53.9 (28)	587	22.8 (51)	47.0 (11)	13	* *	82	* *	574	9.2 (66)	573	6.5 (10)
ShawneeMa	3,477	5.1 (49)	54.4 (35)	3,145	18.7 (19)	53.3 (46)	413	43.3 (21)	626	29.1 (27)	3,074	6.5 (46)	3,080	7.6 (15)
Sheridan	25	* *	* *	21	* *	* *	5	* *	4	* *	22	* *	22	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2008 - 2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Health Indicators by Clinic
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Sherman	198	3.0 (10)	59.6 (62)	171	21.1 (41)	56.7 (62)	18	* *	36	* *	165	4.2 (13)	165	8.5(24)
Smith	52	* *	* *	43	* *	* *	9	* *	13	* *	43	* *	43	* *
Stafford	114	0.9 (2)	65.8 (70)	99	* *	* *	11	* *	16	* *	93	* *	94	* *
Stanton	52	* *	* *	49	* *	* *	5	* *	21	* *	50	* *	49	* *
Stevens	221	0.9 (2)	61.1 (66)	191	15.7 (3)	50.8 (36)	37	* *	60	* *	183	14.2 (68)	183	5.5(4)
Sumner	466	5.6 (53)	51.7 (13)	426	22.8 (51)	43.9 (2)	68	* *	123	17.9 (6)	407	6.1 (39)	408	15.0(65)
Thomas	155	7.1 (66)	52.9 (20)	139	11.5 (1)	59.7 (66)	21	* *	29	* *	138	4.3 (14)	138	11.6(48)
Trego	44	* *	* *	37	* *	* *	5	* *	5	* *	38	* *	38	* *
Wabaunsee	58	* *	* *	57	* *	* *	7	* *	24	* *	56	* *	56	* *
Wallace	1	* *	* *	1	* *	* *	0	* *	1	* *	0	* *	0	* *
Washington	71	* *	* *	64	* *	* *	5	* *	21	* *	63	* *	62	* *
Wichita	45	* *	* *	41	* *	* *	6	* *	6	* *	42	* *	42	* *
Wilson	254	7.9 (69)	55.9 (48)	239	17.6 (11)	53.1 (45)	43	* *	38	* *	228	6.1 (39)	228	10.1(34)
Woodson	28	* *	* *	26	* *	* *	7	* *	12	* *	26	* *	26	* *
Wyandotte	6,341	4.3 (36)	55.7 (45)	5,681	19.6 (26)	51.9 (41)	981	30.9 (16)	1,141	26.0(22)	5,503	4.5 (15)	5,503	8.3(22)
Unknown Clinics	21	* *	* *	20	* *	* *	1	* *	3	* *	22	* *	22	* *
Kansas	63,394	4.5	54.2	56,431	20.2	51.1	9,783	29.3	13,286	23.5	54,852	5.7	54,836	9.2

- (1) Reporting period is January 1 through December 31.
 - (2) Excludes records with unknown data or errors.
 - (3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
 - (4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
 - (5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
 - (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6A
2008 - 2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Behavioral Indicators by Clinic
Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2, 3)										
Clinic	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	% (Rank)		Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Allen	246	92.7 (1)	1.6 (16)		271	38.7 (42)	244	14.3 (46)	214	52.3 (52)
Anderson	125	87.2 (12)	4.8 (40)		146	48.6 (12)	144	20.1 (24)	124	63.7 (25)
Atchison	359	88.6 (9)	2.5 (21)		389	37.3 (48)	376	8.8 (63)	318	58.5 (41)
Barber-Kio	3	* *	* *		4	* *	4	* *	4	* *
Barber-Med	114	80.7 (34)	8.8 (59)		129	43.4 (24)	122	33.6 (5)	101	68.3 (7)
Barton	693	84.4 (23)	1.2 (11)		728	51.9 (7)	669	20.3 (22)	548	56.6 (45)
Bourbon	311	90.0 (6)	1.6 (16)		357	45.1 (21)	321	10.9 (56)	276	59.1 (40)
Brown	236	75.8 (45)	11.0 (61)		284	39.4 (39)	254	12.6 (49)	244	64.8 (22)
Butler	750	70.1 (59)	9.9 (60)		760	38.9 (41)	721	14.7 (43)	548	35.6 (63)
Chase	33	* *	* *		35	* *	34	* *	29	* *
Chautauqua	75	* *	* *		78	* *	67	* *	63	* *
Cherokee	437	76.4 (43)	8.2 (58)		560	37.3 (48)	510	12.5 (50)	407	53.6 (49)
Cheyenne	7	* *	* *		13	* *	13	* *	12	* *
Clark	8	* *	* *		8	* *	6	* *	6	* *
Clay	158	84.8 (20)	5.7 (45)		165	43.0 (26)	145	20.0 (25)	134	67.9 (11)
Cloud	230	83.5 (27)	4.8 (40)		238	48.3 (13)	226	20.8 (21)	195	68.2 (9)
Coffey	69	* *	* *		72	* *	72	* *	56	* *
Comanche	25	* *	* *		25	* *	25	* *	20	* *
Cowley-Ark	495	62.8 (65)	7.1 (53)		563	39.4 (39)	532	12.2 (52)	448	54.5 (48)
Cowley-Win	399	77.2 (42)	5.3 (43)		424	50.5 (10)	388	13.1 (48)	334	58.1 (42)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6A
2008 - 2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Behavioral Indicators by Clinic
Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2, 3)										
Clinic	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Crawford-G	3	* *	* *	* *	4	* *	3	* *	3	* *
Crawford-P	927	86.0 (15)	3.0 (23)		1,048	42.3 (28)	972	12.3 (51)	841	60.2 (36)
Decatur	24	* *	* *	* *	30	* *	29	* *	21	* *
DickinsonA	311	87.5 (11)	1.3 (12)		329	44.4 (22)	320	19.7 (26)	248	67.3 (12)
DickinsonH	38	* *	* *	* *	41	* *	37	* *	34	* *
Doniphan	116	77.6 (41)	7.8 (55)		123	39.8 (35)	117	35.0 (4)	92	* *
Douglas	1,274	74.0 (49)	5.7 (45)		1,397	37.6 (47)	1,295	19.3 (28)	1,054	60.4 (35)
Edwards	59	* *	* *	* *	68	* *	69	* *	57	* *
Elk	48	* *	* *	* *	53	* *	49	* *	41	* *
Ellis	423	85.6 (17)	1.7 (18)		449	34.5 (56)	433	25.9 (12)	360	68.3 (7)
Ellsworth	93	* *	* *	* *	86	* *	86	* *	67	* *
Finney	1,762	59.0 (67)	24.7 (67)		1,957	35.4 (54)	1,834	11.1 (55)	1,539	39.0 (61)
Ford	1,666	71.6 (57)	14.4 (64)		1,915	45.3 (20)	1,730	9.7 (59)	1,538	51.8 (55)
Franklin	203	76.4 (43)	3.0 (23)		498	38.2 (45)	463	5.4 (68)	370	62.4 (27)
Geary	876	72.1 (53)	1.3 (12)		1,097	35.3 (55)	1,009	19.5 (27)	846	56.9 (44)
Gove	25	* *	* *	* *	27	* *	25	* *	18	* *
Graham	42	* *	* *	* *	49	* *	41	* *	36	* *
Grant	172	68.6 (60)	16.3 (66)		186	33.3 (62)	173	9.2 (62)	135	45.9 (58)
Gray	122	84.4 (23)	3.3 (30)		133	46.6 (17)	123	52.8 (2)	107	68.2 (9)
Greeley	32	* *	* *	* *	34	* *	34	* *	27	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6A
2008 - 2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Behavioral Indicators by Clinic
Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2, 3)											
Clinic	Medical Care				WIC Enrollment			Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester			Prepregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	% (Rank)		Number	% (Rank)		Number	% (Rank)	Number	% (Rank)
Greenwood	119	56.3 (68)	36.1 (68)		132	52.3 (5)		116	14.7 (43)	104	70.2 (4)
Hamilton	66	* *	* *		72	* *		68	* *	60	* *
Harper	129	73.6 (52)	14.7 (65)		144	52.1 (6)		133	26.3 (11)	117	70.1 (5)
Harvey	657	74.0 (49)	12.9 (63)		764	47.8 (14)		705	16.6 (34)	601	66.2 (17)
Haskell	51	* *	* *		57	* *		53	* *	47	* *
Hodgeman	36	* *	* *		35	* *		34	* *	27	* *
Jackson	118	82.2 (30)	4.2 (35)		225	44.0 (23)		202	10.4 (57)	178	77.0 (1)
Jefferson	160	85.6 (17)	6.9 (52)		174	39.7 (37)		162	6.2 (67)	136	50.7 (56)
Jewell	30	* *	* *		28	* *		24	* *	25	* *
Johnson-M	2,027	68.1 (61)	2.3 (20)		2,330	33.5 (61)		2,149	14.9 (41)	1,729	65.3 (19)
Johnson-O	2,534	62.4 (66)	4.8 (40)		2,964	32.3 (64)		2,779	12.1 (53)	2,287	59.6 (38)
Kearny	53	* *	* *		57	* *		50	* *	40	* *
Kingman	141	78.0 (39)	4.3 (36)		154	51.9 (7)		147	22.4 (17)	128	67.2 (15)
Kiowa	29	* *	* *		33	* *		30	* *	26	* *
Labette	407	86.5 (14)	3.2 (29)		500	36.6 (52)		462	13.9 (47)	401	61.1 (34)
Lane	31	* *	* *		34	* *		31	* *	29	* *
Leavenwort	931	85.5 (19)	0.6 (5)		965	32.5 (63)		925	20.2 (23)	699	68.5 (6)
Lincoln	58	* *	* *		56	* *		55	* *	45	* *
Linn	175	80.0 (37)	7.4 (54)		180	47.8 (14)		167	21.0 (20)	148	61.5 (30)
Logan	15	* *	* *		37	* *		35	* *	30	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6A
2008 - 2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Behavioral Indicators by Clinic
Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2, 3)										
Clinic	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	% (Rank)		Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Lyon	844	83.9 (25)	0.8 (6)		881	40.4 (34)	810	15.9 (37)	671	62.4 (27)
Marion	128	71.9 (55)	3.9 (32)		141	41.8 (30)	135	32.6 (6)	100	65.0 (20)
Marshall	148	92.6 (3)	0.0 (1)		153	41.8 (30)	133	30.8 (7)	108	65.7 (18)
Mc Pherson	373	87.1 (13)	0.5 (4)		383	38.1 (46)	343	23.0 (16)	294	67.3 (12)
Meade	114	71.9 (55)	7.9 (57)		116	57.8 (4)	108	37.0 (3)	97	* *
Miami	389	81.2 (32)	1.0 (8)		391	34.3 (57)	371	16.7 (33)	296	53.4 (50)
Mitchell	123	92.7 (1)	0.0 (1)		123	34.1 (59)	114	24.6 (14)	97	* *
Montgome-C	331	82.2 (30)	5.4 (44)		435	30.3 (67)	402	16.9 (32)	341	66.3 (16)
Montgome-I	381	88.2 (10)	3.4 (31)		509	35.8 (53)	473	12.1 (53)	398	61.3 (32)
Morris	76	* *	* *		87	* *	76	* *	64	* *
Nemaha	80	* *	* *		82	* *	76	* *	66	* *
Neosho-Ch	361	88.9 (8)	4.4 (37)		402	38.3 (44)	377	9.5 (60)	321	67.3 (12)
Neosho-Er	25	* *	* *		31	* *	27	* *	22	* *
Ness	26	* *	* *		30	* *	32	* *	24	* *
Norton	87	* *	* *		85	* *	84	* *	69	* *
Osage	208	91.3 (5)	1.0 (8)		240	50.0 (11)	231	19.0 (30)	196	38.8 (62)
Osborne	49	* *	* *		54	* *	46	* *	40	* *
Ottawa	72	* *	* *		73	* *	68	* *	59	* *
Pawnee	136	84.6 (22)	2.2 (19)		154	50.6 (9)	148	27.7 (9)	124	61.3 (32)
Phillips	97	* *	* *		102	58.8 (3)	97	* *	89	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6A
2008 - 2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Behavioral Indicators by Clinic
Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2, 3)										
Clinic	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Pottawat-S	40	* *	* *	* *	46	* *	40	* *	33	* *
Pottawat-W	52	* *	* *	* *	61	* *	53	* *	50	* *
Pratt	202	80.7 (34)	6.4 (49)		212	43.4 (24)	208	7.2 (66)	158	61.4 (31)
Rawlins	11	* *	* *	* *	28	* *	25	* *	23	* *
Reno	1,262	83.6 (26)	2.5 (21)		1,320	39.8 (35)	1,243	21.1 (19)	1,043	62.0 (29)
Republic	89	* *	* *	* *	96	* *	83	* *	72	* *
Rice	203	83.3 (28)	1.0 (8)		220	42.3 (28)	206	16.0 (36)	171	44.4 (59)
Riley-Ft	1,126	91.9 (4)	0.4 (3)		1,458	31.3 (66)	1,411	19.1 (29)	1,087	64.2 (24)
Riley-Manh	966	73.9 (51)	1.3 (12)		1,103	38.4 (43)	1,024	18.1 (31)	837	59.5 (39)
Rooks	120	82.5 (29)	5.8 (47)		129	39.5 (38)	117	30.8 (7)	100	65.0 (20)
Rush	30	* *	* *	* *	29	* *	27	* *	24	* *
Russell	114	89.5 (7)	0.9 (7)		115	47.0 (16)	109	10.1 (58)	94	* *
Saline	1,234	74.9 (48)	4.1 (34)		1,451	34.1 (59)	1,348	5.0 (69)	1,151	52.2 (54)
Scott	104	80.8 (33)	6.7 (50)		112	45.5 (19)	105	79.0 (1)	95	* *
Sedgwick-C	3,166	75.1 (47)	6.3 (48)		3,317	29.4 (69)	3,087	15.8 (38)	2,629	47.2 (57)
Sedgwick-M	5,367	71.6 (57)	4.6 (38)		5,419	26.4 (70)	5,054	14.9 (41)	4,102	52.5 (51)
Sedgwick-S	3,547	79.6 (38)	3.0 (23)		3,718	30.2 (68)	3,437	15.8 (38)	2,852	40.4 (60)
Seneca	29	* *	* *	* *	28	* *	21	* *	20	* *
Seward	1,183	65.4 (64)	11.7 (62)		1,349	34.3 (57)	1,227	8.6 (64)	1,021	64.8 (22)
ShawneeM&I	563	67.7 (62)	3.0 (23)		761	67.1 (1)	722	16.1 (35)	702	55.1 (47)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6A
2008 - 2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Maternal Behavioral Indicators by Clinic
Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2, 3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
ShawneeMa	2,897	77.8 (40)	4.0 (33)	3,307	42.6 (27)	3,077	7.6 (65)	2,545	52.3 (52)
Sheridan	22	* *	* *	22	* *	20	* *	17	* *
Sherman	118	84.7 (21)	6.8 (51)	190	46.3 (18)	171	25.7 (13)	146	70.5 (3)
Smith	38	* *	* *	47	* *	44	* *	42	* *
Stafford	100	72.0 (54)	3.0 (23)	109	60.6 (2)	101	23.8 (15)	87	* *
Stanton	50	* *	* *	51	* *	41	* *	31	* *
Stevens	194	75.3 (46)	3.1 (28)	206	32.0 (65)	197	22.3 (18)	161	60.2 (36)
Sumner	435	80.2 (36)	4.6 (38)	440	40.5 (33)	405	15.1 (40)	325	62.5 (26)
Thomas	96	* *	* *	146	37.0 (51)	133	27.1 (10)	113	76.1 (2)
Trego	39	* *	* *	41	* *	38	* *	34	* *
Wabaunsee	52	* *	* *	52	* *	52	* *	41	* *
Wallace	1	* *	* *	0	* *	1	* *	0	* *
Washington	66	* *	* *	68	* *	64	* *	52	* *
Wichita	43	* *	* *	45	* *	42	* *	33	* *
Wilson	236	86.0 (15)	1.3 (12)	246	41.5 (32)	226	14.6 (45)	198	56.1 (46)
Woodson	28	* *	* *	27	* *	26	* *	21	* *
Wyandotte	5,629	65.5 (63)	7.8 (55)	6,006	37.2 (50)	5,557	9.3 (61)	4,735	57.5 (43)
Unknown Clinics	20	* *	* *	23	* *	23	* *	18	* *
Kansas	54,276	75.1	5.7	60,184	37.0	55,958	14.7	46,711	56.0

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2008 - 2010 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by Clinic
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Allen	276	48.2 (56)	228	30.3 (52)	86	* *	242	39.7 (56)	228	38.2 (54)
Anderson	152	50.7 (64)	132	29.5 (51)	56	* *	133	23.3 (39)	134	20.1 (38)
Atchison	301	48.8 (59)	200	31.5 (55)	73	* *	256	27.3 (43)	204	22.5 (43)
Barber-Kio	4	* *	3	* *	0	* *	4	* *	3	* *
Barber-Med	136	50.0 (62)	111	32.4 (57)	46	* *	113	41.6 (60)	112	45.5 (60)
Barton	780	36.8 (25)	687	24.0 (27)	206	26.7 (25)	634	11.4 (16)	688	5.8 (8)
Bourbon	356	46.3 (52)	277	27.8 (47)	97	* *	296	38.9 (55)	281	34.9 (52)
Brown	285	46.0 (50)	215	26.5 (41)	71	* *	244	42.2 (62)	217	39.6 (57)
Butler	818	49.4 (61)	685	30.7 (53)	252	32.1 (18)	618	13.4 (20)	690	8.4 (16)
Chase	39	* *	34	* *	17	* *	33	* *	34	* *
Chautauqua	71	* *	46	* *	16	* *	62	* *	45	* *
Cherokee	585	40.3 (36)	502	27.7 (46)	157	23.6 (31)	454	33.7 (52)	511	34.2 (51)
Cheyenne	13	* *	10	* *	2	* *	12	* *	10	* *
Clark	3	* *	2	* *	0	* *	5	* *	4	* *
Clay	175	47.4 (55)	154	25.3 (32)	64	* *	155	18.1 (30)	156	10.3 (21)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2008 - 2010 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by Clinic
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Cloud	252	35.7 (21)	224	19.2 (19)	63	* *	216	7.4 (4)	226	2.2 (1)
Coffey	77	* *	68	* *	26	* *	60	* *	68	* *
Comanche	27	* *	24	* *	10	* *	21	* *	24	* *
Cowley-Ark	590	37.1 (29)	459	25.3 (32)	128	26.6 (26)	486	21.6 (37)	462	21.0 (41)
Cowley-Win	451	49.2 (60)	360	32.5 (58)	146	26.0 (29)	378	20.1 (32)	360	19.4 (35)
Crawford-G	4	* *	3	* *	1	* *	4	* *	3	* *
Crawford-P	1,041	43.8 (45)	811	25.6 (35)	288	36.5 (7)	905	39.9 (57)	814	39.3 (56)
Decatur	33	* *	28	* *	6	* *	23	* *	29	* *
DickinsonA	343	43.1 (43)	299	26.8 (44)	99	* *	269	10.4 (13)	302	7.6 (14)
DickinsonH	42	* *	38	* *	18	* *	38	* *	38	* *
Doniphan	117	41.9 (39)	85	* *	24	* *	87	* *	88	* *
Douglas	1,468	40.9 (37)	1,234	19.8 (20)	385	42.1 (4)	1,179	10.2 (12)	1,236	6.4 (10)
Edwards	77	* *	65	* *	19	* *	63	* *	66	* *
Elk	53	* *	43	* *	13	* *	43	* *	42	* *
Ellis	477	36.1 (23)	408	14.0 (11)	120	46.7 (1)	386	7.0 (2)	410	5.6 (6)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2008 - 2010 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by Clinic
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Ellsworth	97	* *	90	* *	29	* *	73	* *	91	* *
Finney	2,076	11.5 (5)	1,861	6.0 (4)	162	38.9 (6)	1,730	15.4 (25)	1,861	14.1 (27)
Ford	1,956	9.0 (4)	1,639	4.5 (3)	105	31.4 (19)	1,704	8.9 (9)	1,637	8.4 (16)
Franklin	533	44.1 (46)	468	28.4 (49)	142	27.5 (23)	403	13.4 (20)	470	9.6 (19)
Geary	1,134	36.9 (26)	905	20.7 (21)	257	33.9 (12)	942	37.4 (54)	908	34.1 (50)
Gove	28	* *	26	* *	2	* *	19	* *	26	* *
Graham	51	* *	41	* *	19	* *	43	* *	42	* *
Grant	202	6.9 (2)	191	2.6 (1)	8	* *	158	9.5 (11)	191	11.0 (22)
Gray	146	8.2 (3)	119	6.7 (5)	6	* *	125	10.4 (13)	120	7.5 (13)
Greeley	37	* *	32	* *	7	* *	27	* *	32	* *
Greenwood	118	56.8 (68)	60	* *	27	* *	110	30.9 (50)	87	* *
Hamilton	75	* *	68	* *	6	* *	63	* *	68	* *
Harper	139	44.6 (48)	99	* *	33	* *	116	21.6 (37)	92	* *
Harvey	802	37.2 (30)	687	21.7 (22)	209	34.9 (10)	667	7.0 (2)	689	4.2 (4)
Haskell	61	* *	56	* *	8	* *	53	* *	56	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2008 - 2010 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by Clinic
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Hodgeman	37	* *	32	* *	14	* *	29	* *	31	* *
Jackson	231	48.5 (57)	203	31.0 (54)	80	* *	197	20.8 (36)	205	19.5 (36)
Jefferson	48	* *	21	* *	4	* *	48	* *	25	* *
Jewell	30	* *	28	* *	11	* *	27	* *	28	* *
Johnson-M	2,501	26.1 (11)	2,180	11.2 (8)	382	42.7 (3)	1,947	20.6 (35)	2,186	19.9 (37)
Johnson-O	3,151	22.1 (8)	2,779	10.0 (7)	439	43.3 (2)	2,543	16.6 (26)	2,786	16.8 (30)
Kearny	61	* *	57	* *	9	* *	48	* *	57	* *
Kingman	149	47.0 (54)	122	27.0 (45)	39	* *	136	41.9 (61)	131	38.9 (55)
Kiowa	31	* *	22	* *	5	* *	25	* *	24	* *
Labette	520	36.0 (22)	458	25.1 (31)	129	17.1 (33)	439	30.5 (49)	463	30.7 (47)
Lane	11	* *	2	* *	0	* *	12	* *	3	* *
Leavenwort	1,059	46.8 (53)	892	25.8 (37)	288	34.7 (11)	788	41.5 (59)	895	40.1 (58)
Lincoln	61	* *	50	* *	21	* *	51	* *	50	* *
Linn	188	56.4 (67)	170	37.6 (61)	84	* *	159	15.1 (24)	172	12.8 (25)
Logan	39	* *	33	* *	11	* *	34	* *	33	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2008 - 2010 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by Clinic
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Lyon	949	29.7 (17)	840	16.7 (17)	183	32.8 (14)	761	11.8 (17)	841	8.6 (18)
Marion	153	22.2 (9)	131	15.3 (14)	22	* *	113	5.3 (1)	131	4.6 (5)
Marshall	159	50.3 (63)	134	32.1 (56)	53	* *	132	53.8 (66)	136	51.5 (62)
Mc Pherson	411	37.7 (32)	371	24.0 (27)	121	29.8 (20)	339	7.4 (4)	372	3.8 (3)
Meade	117	24.8 (10)	101	15.8 (15)	20	* *	104	25.0 (40)	103	22.3 (42)
Miami	395	44.8 (49)	335	25.7 (36)	113	33.6 (13)	306	20.3 (33)	334	16.8 (30)
Mitchell	131	38.2 (33)	120	26.7 (43)	40	* *	109	9.2 (10)	122	8.2 (15)
Montgome-C	460	40.2 (35)	387	28.9 (50)	123	15.4 (34)	383	33.7 (52)	390	31.0 (48)
Montgome-I	521	41.3 (38)	439	28.0 (48)	142	19.0 (32)	433	32.8 (51)	440	30.0 (46)
Morris	93	* *	80	* *	29	* *	75	* *	79	* *
Nemaha	79	* *	62	* *	25	* *	65	* *	63	* *
Neosho-Ch	366	42.3 (41)	224	24.1 (30)	74	* *	320	28.1 (45)	231	18.6 (34)
Neosho-Er	29	* *	13	* *	3	* *	26	* *	13	* *
Ness	29	* *	25	* *	6	* *	23	* *	25	* *
Norton	94	* *	83	* *	31	* *	75	* *	85	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2008 - 2010 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by Clinic
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Osage	252	46.0 (50)	207	26.6 (42)	70	* *	209	29.7 (47)	208	26.4 (44)
Osborne	58	* *	51	* *	20	* *	43	* *	51	* *
Ottawa	73	* *	46	* *	19	* *	64	* *	59	* *
Pawnee	157	36.9 (26)	124	25.8 (37)	35	* *	133	17.3 (28)	125	14.4 (28)
Phillips	83	* *	50	* *	14	* *	67	* *	50	* *
Pottawat-S	48	* *	42	* *	7	* *	38	* *	43	* *
Pottawat-W	66	* *	55	* *	16	* *	57	* *	58	* *
Pratt	228	38.2 (33)	194	25.3 (32)	60	* *	174	40.2 (58)	194	37.6 (53)
Rawlins	29	* *	25	* *	6	* *	25	* *	25	* *
Reno	1,416	43.6 (44)	1,224	23.4 (26)	433	36.5 (7)	1,157	16.7 (27)	1,230	15.0 (29)
Republic	101	48.5 (57)	83	* *	29	* *	81	* *	85	* *
Rice	230	42.6 (42)	198	25.8 (37)	61	* *	186	17.7 (29)	200	13.5 (26)
Riley-Ft	1,518	27.5 (13)	1,037	14.1 (12)	183	32.2 (17)	1,174	20.5 (34)	1,077	17.5 (33)
Riley-Manh	1,163	29.8 (18)	910	16.7 (17)	198	29.3 (21)	951	8.4 (8)	959	6.5 (11)
Rooks	132	62.1 (69)	105	39.0 (63)	55	* *	110	29.1 (46)	111	33.3 (49)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2008 - 2010 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by Clinic
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Rush	31	* *	29	* *	16	* *	28	* *	29	* *
Russell	126	44.4 (47)	111	37.8 (62)	43	* *	103	53.4 (65)	112	52.7 (63)
Saline	1,479	42.0 (40)	1,233	26.4 (40)	430	26.5 (27)	1,235	12.6 (18)	1,237	9.9 (20)
Scott	119	26.1 (11)	95	* *	24	* *	103	30.1 (48)	97	* *
Sedgwick-C	3,512	28.5 (16)	3,193	14.9 (13)	693	35.9 (9)	2,902	12.6 (18)	3,179	11.2 (23)
Sedgwick-M	5,830	28.4 (15)	5,320	13.1 (10)	1,101	41.0 (5)	4,673	7.6 (6)	5,321	6.1 (9)
Sedgwick-S	3,925	37.5 (31)	3,584	22.1 (24)	1,050	27.2 (24)	3,191	7.7 (7)	3,583	5.6 (6)
Seneca	29	* *	25	* *	12	* *	26	* *	25	* *
Seward	1,418	6.8 (1)	1,264	3.2 (2)	66	* *	1,177	13.6 (22)	1,284	11.7 (24)
ShawneeM&I	723	27.9 (14)	529	15.9 (16)	125	32.8 (14)	648	15.0 (23)	534	3.4 (2)
ShawneeMa	3,305	36.6 (24)	2,642	24.0 (27)	715	24.6 (30)	2,689	11.1 (15)	2,660	6.6 (12)
Sheridan	24	* *	21	* *	4	* *	20	* *	21	* *
Sherman	198	34.8 (20)	169	21.9 (23)	50	* *	161	25.5 (41)	168	20.8 (40)
Smith	46	* *	35	* *	12	* *	41	* *	37	* *
Stafford	111	36.9 (26)	93	* *	30	* *	92	* *	94	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2008 - 2010 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by Clinic
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Stanton	51	* *	47	* *	4	* *	40	* *	47	* *
Stevens	223	19.3 (6)	189	8.5 (6)	27	* *	179	25.7 (42)	191	20.4 (39)
Sumner	466	53.9 (65)	414	33.8 (59)	175	29.1 (22)	367	47.7 (63)	416	49.0 (61)
Thomas	156	34.6 (19)	135	22.2 (25)	34	* *	122	27.9 (44)	136	26.5 (45)
Trego	42	* *	32	* *	13	* *	38	* *	33	* *
Wabaunsee	60	* *	57	* *	18	* *	44	* *	57	* *
Wallace	0	* *	0	* *	0	* *	0	* *	0	* *
Washington	74	* *	65	* *	11	* *	58	* *	64	* *
Wichita	46	* *	37	* *	8	* *	36	* *	37	* *
Wilson	252	56.0 (66)	230	34.3 (60)	110	32.7 (16)	220	47.7 (63)	230	43.0 (59)
Woodson	25	* *	19	* *	2	* *	20	* *	20	* *
Wyandotte	6,335	21.1 (7)	5,508	12.4 (9)	870	26.3 (28)	5,304	19.6 (31)	5,600	17.3 (32)
Unknown Clinics	22	* *	22	* *	11	* *	18	* *	23	* *
Kansas	62,786	31.6	53,620	17.9	12,870	31.8	51,403	17.6	54,019	15.1

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2008 - 2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Infant Health Indicators by Clinic
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)									
Clinic	Birthweight			Preterm ⁽⁶⁾		Full Term LBW ⁽⁷⁾		Ever Breastfed	
	Low ⁽⁴⁾	High ⁽⁵⁾							
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Allen	257	8.6 (52)	3.5 (3)	246	9.3 (32)	218	6.0 (54)	260	57.7 (61)
Anderson	122	8.2 (47)	8.2 (54)	131	8.4 (19)	105	6.7 (57)	136	57.4 (62)
Atchison	352	7.4 (30)	5.1 (20)	345	6.1 (3)	312	4.8 (45)	365	52.3 (65)
Barber-Kio	3	* * *	* *	3	* *	3	* *	3	* *
Barber-Med	106	11.3 (64)	3.8 (5)	100	15.0 (64)	84	* *	107	76.6 (14)
Barton	662	6.3 (17)	4.4 (13)	624	9.5 (35)	559	4.1 (33)	658	75.4 (17)
Bourbon	318	9.7 (60)	3.8 (5)	306	10.8 (51)	268	6.3 (55)	320	60.9 (55)
Brown	233	9.0 (58)	7.7 (50)	249	10.0 (44)	204	4.9 (49)	251	49.8 (67)
Butler	762	7.6 (34)	6.3 (32)	687	9.5 (35)	612	3.4 (17)	744	69.4 (37)
Chase	33	* * *	* *	29	* *	28	* *	32	* *
Chautauqua	71	* * *	* *	70	* *	62	* *	71	* *
Cherokee	539	7.4 (30)	6.5 (39)	493	8.7 (21)	447	2.9 (10)	528	58.0 (60)
Cheyenne	9	* * *	* *	9	* *	7	* *	9	* *
Clark	9	* * *	* *	7	* *	6	* *	8	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2008 - 2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Infant Health Indicators by Clinic
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)										
Clinic	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)							
	Number	% (Rank)	%	(Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Clay	157	7.6 (34)	6.4	(37)	146	13.0 (62)	127	3.9 (29)	156	61.5 (50)
Cloud	225	8.0 (44)	3.1	(2)	213	11.7 (56)	187	4.8 (45)	220	71.4 (31)
Coffey	67	* * *	* *		58	* *	55	* *	68	* *
Comanche	23	* * *	* *		20	* *	18	* *	23	* *
Cowley-Ark	539	5.9 (12)	6.9	(43)	499	10.4 (47)	438	3.7 (22)	535	52.1 (66)
Cowley-Win	389	9.8 (61)	8.5	(55)	372	12.6 (58)	310	6.5 (56)	403	67.0 (43)
Crawford-G	3	* * *	* *		3	* *	3	* *	3	* *
Crawford-P	928	8.1 (45)	4.8	(16)	893	10.4 (47)	782	4.1 (33)	937	59.0 (59)
Decatur	28	* * *	* *		27	* *	25	* *	27	* *
DickinsonA	297	7.1 (26)	9.8	(61)	282	9.6 (38)	254	2.8 (7)	299	72.9 (29)
DickinsonH	35	* * *	* *		35	* *	32	* *	35	* *
Doniphan	116	12.1 (65)	10.3	(62)	107	13.1 (63)	92	* *	114	61.4 (52)
Douglas	1,264	6.7 (20)	7.0	(44)	1,184	8.9 (25)	1,063	4.0 (31)	1,254	77.7 (11)
Edwards	62	* * *	* *		55	* *	43	* *	61	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2008 - 2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Infant Health Indicators by Clinic
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)									
Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Elk	51	* * * *	*	48	* *	44	* *	51	* *
Ellis	384	6.3 (17)	6.0 (27)	382	8.6 (20)	322	2.8 (7)	401	77.6 (12)
Ellsworth	89	* * * *	*	77	* *	72	* *	88	* *
Finney	1,827	6.7 (20)	6.4 (37)	1,750	9.8 (41)	1,540	3.6 (21)	1,575	78.0 (10)
Ford	1,694	4.8 (4)	6.0 (27)	1,651	8.9 (25)	1,465	2.0 (5)	1,602	74.6 (21)
Franklin	462	5.6 (8)	5.4 (22)	426	8.2 (16)	389	3.9 (29)	457	61.5 (50)
Geary	927	6.9 (23)	6.3 (32)	874	8.9 (25)	794	3.3 (15)	896	70.5 (34)
Gove	24	* * * *	*	22	* *	19	* *	22	* *
Graham	38	* * * *	*	38	* *	32	* *	39	* *
Grant	187	7.0 (24)	4.3 (10)	173	8.1 (15)	156	4.5 (43)	163	74.2 (22)
Gray	97	* * * *	*	108	11.1 (52)	81	* *	117	82.1 (2)
Greeley	32	* * * *	*	29	* *	26	* *	32	* *
Greenwood	119	5.0 (5)	5.0 (18)	114	8.8 (23)	104	4.8 (45)	117	60.7 (57)
Hamilton	64	* * * *	*	63	* *	57	* *	57	* *

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
- (4) <2500 g.
- (5) >4000 g.
- (6) <37 weeks gestation.
- (7) 37 or more weeks gestation and <2500 g.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2008 - 2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Infant Health Indicators by Clinic
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)										
Clinic	Birthweight				Preterm ⁽⁶⁾		Full Term LBW ⁽⁷⁾		Ever Breastfed	
	Low ⁽⁴⁾		High ⁽⁵⁾		Number	% ^(Rank)	Number	% ^(Rank)	Number	% ^(Rank)
	Number	% ^(Rank)	% ^(Rank)							
Harper	125	6.4 (19)	9.6 (60)		123	8.9 (25)	106	2.8 (7)	132	75.8 (15)
Harvey	685	7.9 (41)	6.1 (30)		654	10.2 (46)	579	3.8 (25)	695	80.7 (6)
Haskell	53	* * *	* *		51	* *	47	* *	43	* *
Hodgeman	36	* * *	* *		32	* *	29	* *	34	* *
Jackson	207	8.7 (55)	12.6 (65)		205	9.8 (41)	181	5.0 (50)	206	75.7 (16)
Jefferson	169	8.3 (49)	6.5 (39)		155	12.9 (60)	134	5.2 (52)	166	65.1 (46)
Jewell	28	* * *	* *		28	* *	21	* *	28	* *
Johnson-M	2,053	7.7 (37)	6.3 (32)		2,024	7.6 (12)	1,724	3.7 (22)	2,170	78.2 (9)
Johnson-O	2,677	6.1 (15)	7.8 (52)		2,598	7.7 (13)	2,296	3.1 (11)	2,707	81.8 (4)
Kearny	55	* * *	* *		53	* *	44	* *	46	* *
Kingman	98	* * *	* *		129	7.0 (8)	85	* *	137	83.2 (1)
Kiowa	28	* * *	* *		29	* *	24	* *	29	* *
Labette	468	5.6 (8)	7.9 (53)		436	9.9 (43)	393	3.3 (15)	451	54.5 (64)
Lane	30	* * *	* *		28	* *	23	* *	31	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2008 - 2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Infant Health Indicators by Clinic
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)									
Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Leavenwort	897	7.9 (41)	7.0 (44)	823	8.9 (25)	739	4.3 (40)	910	63.5 (48)
Lincoln	57	* *	* *	50	* *	45	* *	54	* *
Linn	163	9.2 (59)	4.9 (17)	161	11.2 (53)	137	7.3 (59)	167	74.9 (19)
Logan	33	* *	* *	31	* *	29	* *	32	* *
Lyon	826	7.6 (34)	6.2 (31)	760	8.9 (25)	691	4.1 (33)	810	74.2 (22)
Marion	117	5.1 (6)	7.7 (50)	108	7.4 (11)	97	* *	119	79.8 (8)
Marshall	139	8.6 (52)	4.3 (10)	135	9.6 (38)	118	5.1 (51)	140	60.0 (58)
Mc Pherson	360	4.4 (3)	5.6 (23)	336	6.3 (4)	311	1.3 (3)	361	73.4 (26)
Meade	100	6.0 (14)	5.0 (18)	99	* *	80	* *	105	80.0 (7)
Miami	386	6.2 (16)	8.5 (55)	349	7.7 (13)	320	3.8 (25)	358	74.0 (25)
Mitchell	104	5.8 (10)	5.8 (24)	112	7.1 (9)	90	* *	115	73.0 (28)
Montgome-C	414	7.2 (28)	4.1 (8)	386	10.6 (50)	341	3.8 (25)	405	63.0 (49)
Montgome-I	463	8.6 (52)	3.5 (3)	430	8.8 (23)	390	4.1 (33)	454	60.8 (56)
Morris	74	* *	* *	75	* *	61	* *	80	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2008 - 2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Infant Health Indicators by Clinic
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)										
Clinic	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)		Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
	Number	% (Rank)	% (Rank)	Number						
Nemaha	54	* * * *	* *	75	* *	43	* *	76	* *	
Neosho-Ch	295	7.1 (26)	5.8 (24)	360	11.7 (56)	242	3.7 (22)	387	57.4 (62)	
Neosho-Er	24	* * * *	* *	29	* *	23	* *	30	* *	
Ness	28	* * * *	* *	25	* *	21	* *	30	* *	
Norton	82	* * * *	* *	74	* *	62	* *	79	* *	
Osage	238	6.7 (20)	3.8 (5)	222	17.6 (65)	175	3.4 (17)	237	61.2 (53)	
Osborne	48	* * * *	* *	45	* *	42	* *	47	* *	
Ottawa	72	* * * *	* *	68	* *	63	* *	69	* *	
Pawnee	133	7.5 (33)	4.5 (14)	127	8.7 (21)	116	1.7 (4)	132	72.7 (30)	
Phillips	92	* * * *	* *	87	* *	77	* *	90	* *	
Pottawat-S	41	* * * *	* *	39	* *	34	* *	41	* *	
Pottawat-W	59	* * * *	* *	54	* *	51	* *	58	* *	
Pratt	193	2.1 (1)	10.4 (63)	176	5.7 (2)	164	1.2 (2)	190	68.4 (39)	
Rawlins	25	* * * *	* *	24	* *	23	* *	25	* *	

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2008 - 2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Infant Health Indicators by Clinic
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)									
Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Reno	1,254	7.7 (37)	6.0 (27)	1,160	9.5 (35)	1,037	4.4 (42)	1,243	65.2 (45)
Republic	85	* * *	* *	81	* *	68	* *	88	* *
Rice	193	7.8 (40)	10.4 (63)	183	11.5 (55)	158	3.2 (14)	192	73.4 (26)
Riley-Ft	1,234	5.5 (7)	8.6 (58)	1,124	6.4 (6)	1,038	3.1 (11)	1,234	74.7 (20)
Riley-Manh	1,013	5.9 (12)	7.3 (48)	928	7.2 (10)	853	2.3 (6)	1,006	81.6 (5)
Rooks	117	8.5 (51)	4.3 (10)	112	6.3 (4)	105	4.8 (45)	116	69.0 (38)
Rush	30	* * *	* *	28	* *	25	* *	30	* *
Russell	108	7.4 (30)	2.8 (1)	98	* *	84	* *	105	61.0 (54)
Saline	1,384	7.9 (41)	6.3 (32)	1,307	9.6 (38)	1,159	4.2 (38)	1,395	63.9 (47)
Scott	99	* * *	* *	95	* *	89	* *	98	* *
Sedgwick-C	3,170	8.1 (45)	7.6 (49)	2,983	8.3 (18)	2,709	4.3 (40)	3,119	70.1 (35)
Sedgwick-M	5,265	10.1 (62)	6.6 (41)	4,867	9.1 (31)	4,380	5.6 (53)	4,983	69.6 (36)
Sedgwick-S	3,556	7.7 (37)	7.0 (44)	3,346	8.2 (16)	3,057	4.1 (33)	3,464	67.5 (42)
Seneca	20	* * *	* *	26	* *	15	* *	28	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2008 - 2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Infant Health Indicators by Clinic
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)									
Clinic	Birthweight			Preterm ⁽⁶⁾		Full Term LBW ⁽⁷⁾		Ever Breastfed	
	Low ⁽⁴⁾	High ⁽⁵⁾		Number	%	Number	%	Number	%
	Number	% (Rank)	% (Rank)						
Seward	1,212	5.8 (10)	6.7 (42)	1,190	9.4 (33)	1,022	3.1 (11)	1,266	70.8 (33)
ShawneeM&I	562	8.4 (50)	9.1 (59)	555	12.8 (59)	477	3.8 (25)	557	82.0 (3)
ShawneeMa	3,131	7.3 (29)	5.1 (20)	2,957	10.4 (47)	2,609	4.0 (31)	3,140	68.1 (41)
Sheridan	23	* * *	* *	20	* *	16	* *	23	* *
Sherman	170	8.8 (56)	4.7 (15)	159	6.9 (7)	148	3.4 (17)	166	74.1 (24)
Smith	42	* * *	* *	38	* *	32	* *	43	* *
Stafford	92	* * *	* *	91	* *	79	* *	89	* *
Stanton	47	* * *	* *	47	* *	42	* *	37	* *
Stevens	117	3.4 (2)	8.5 (55)	172	5.2 (1)	101	1.0 (1)	189	76.7 (13)
Sumner	422	10.9 (63)	7.1 (47)	394	11.4 (54)	340	6.8 (58)	427	68.4 (39)
Thomas	142	7.0 (24)	4.2 (9)	128	9.4 (33)	116	3.4 (17)	138	75.4 (17)
Trego	38	* * *	* *	36	* *	31	* *	38	* *
Wabaunsee	57	* * *	* *	50	* *	48	* *	56	* *
Wallace	1	* * *	* *	0	* *	0	* *	1	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2008 - 2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Kansas
Comparison of Infant Health Indicators by Clinic
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)										
Clinic	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)							
	Number	% (Rank)	% (Rank)		Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Washington	60	* * * *	* *		60	* *	48	* *	63	* *
Wichita	41	* * * *	* *		39	* *	33	* *	41	* *
Wilson	240	8.8 (56)	5.8 (24)		232	12.9 (60)	200	4.5 (43)	226	66.4 (44)
Woodson	25	* * * *	* *		27	* *	23	* *	28	* *
Wyandotte	5,495	8.2 (47)	6.3 (32)		5,247	10.0 (44)	4,570	4.2 (38)	5,549	71.2 (32)
Unknown Clinics	23	* * * *	* *		23	* *	23	* *	22	* *
Kansas	55,301	7.5	6.5		52,556	9.2	46,451	4.0	54,920	70.7

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

