

**Table 3A**  
**2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Racial and Ethnic Distribution by Clinic**

<b>Comparison of Racial and Ethnic Distribution</b> <sup>(2)</sup>								
<b>Clinic</b>	<b>Number Included in Analysis</b>	<b>White, Not Hispanic</b>	<b>Black, Not Hispanic</b>	<b>Hispanic</b>	<b>American Indian/Alaska Native</b>	<b>Asian/Pacific Islander</b>	<b>Multiple Races</b>	<b>All Other/Unknown</b>
		%	%	%	%	%	%	%
Allen	71	*	*	*	*	*	*	*
Anderson	47	*	*	*	*	*	*	*
Atchison	148	83.8	12.2	0.7	0.7	0.0	2.7	0.0
Barber-Kio	1	*	*	*	*	*	*	*
Barber-Med	36	*	*	*	*	*	*	*
Barton	259	68.7	0.0	30.1	0.0	0.0	1.2	0.0
Bourbon	111	90.1	4.5	1.8	1.8	0.9	0.9	0.0
Brown	89	*	*	*	*	*	*	*
Butler	253	86.6	0.8	7.9	1.6	0.0	3.2	0.0
Chase	16	*	*	*	*	*	*	*
Chautauqua	32	*	*	*	*	*	*	*
CherokeeBS	75	*	*	*	*	*	*	*
CherokeeCo	93	*	*	*	*	*	*	*
Cheyenne	20	*	*	*	*	*	*	*
Clark	4	*	*	*	*	*	*	*
Clay	50	*	*	*	*	*	*	*
Cloud	60	*	*	*	*	*	*	*
Coffey	26	*	*	*	*	*	*	*
Comanche	9	*	*	*	*	*	*	*
Cowley-Ark	187	67.9	1.1	28.9	1.1	0.5	0.5	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 3A**  
**2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Racial and Ethnic Distribution by Clinic**

<b>Comparison of Racial and Ethnic Distribution</b> <sup>(2)</sup>								
<b>Clinic</b>	<b>Number Included in Analysis</b>	<b>White, Not Hispanic</b>	<b>Black, Not Hispanic</b>	<b>Hispanic</b>	<b>American Indian/Alaska Native</b>	<b>Asian/Pacific Islander</b>	<b>Multiple Races</b>	<b>All Other/Unknown</b>
		%	%	%	%	%	%	%
Cowley-Win	174	86.2	1.7	6.3	0.6	3.4	1.7	0.0
Crawford-G	1	*	*	*	*	*	*	*
Crawford-P	367	85.0	3.0	8.7	1.1	1.4	0.8	0.0
Decatur	16	*	*	*	*	*	*	*
DickinsonA	103	91.3	1.0	5.8	0.0	1.9	0.0	0.0
DickinsonH	19	*	*	*	*	*	*	*
Doniphan	40	*	*	*	*	*	*	*
Douglas	525	64.8	8.0	10.7	8.6	5.9	2.1	0.0
Edwards	21	*	*	*	*	*	*	*
Elk	12	*	*	*	*	*	*	*
Ellis	154	76.6	1.9	18.8	1.3	0.6	0.6	0.0
Ellsworth	34	*	*	*	*	*	*	*
Finney	684	26.8	1.3	66.8	0.0	5.0	0.1	0.0
Ford	674	19.6	3.0	76.0	0.3	0.9	0.3	0.0
Franklin	189	92.1	1.1	3.7	2.1	0.5	0.5	0.0
Geary	406	55.9	22.2	15.3	0.7	2.7	3.2	0.0
Gove	13	*	*	*	*	*	*	*
Graham	21	*	*	*	*	*	*	*
Grant	72	*	*	*	*	*	*	*
Gray	54	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 3A**  
**2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Racial and Ethnic Distribution by Clinic**

<b>Comparison of Racial and Ethnic Distribution</b> <sup>(2)</sup>								
<b>Clinic</b>	<b>Number Included in Analysis</b>	<b>White, Not Hispanic</b>	<b>Black, Not Hispanic</b>	<b>Hispanic</b>	<b>American Indian/Alaska Native</b>	<b>Asian/Pacific Islander</b>	<b>Multiple Races</b>	<b>All Other/Unknown</b>
		%	%	%	%	%	%	%
Greeley	14	*	*	*	*	*	*	*
Greenwood	35	*	*	*	*	*	*	*
Hamilton	19	*	*	*	*	*	*	*
Harper	60	*	*	*	*	*	*	*
Harvey	251	73.3	3.2	21.5	0.8	0.4	0.8	0.0
Haskell	22	*	*	*	*	*	*	*
Hodgeman	12	*	*	*	*	*	*	*
JacksonHol	70	*	*	*	*	*	*	*
JacksonPBP	7	*	*	*	*	*	*	*
Jefferson	57	*	*	*	*	*	*	*
Jewell	6	*	*	*	*	*	*	*
Johnson-M	808	48.5	17.8	29.8	0.1	2.6	1.1	0.0
Johnson-O	1,090	46.8	12.2	36.4	0.4	2.9	1.3	0.0
Kearny	16	*	*	*	*	*	*	*
Kingman	47	*	*	*	*	*	*	*
Kiowa	15	*	*	*	*	*	*	*
Labette	160	80.6	10.6	4.4	2.5	0.0	1.9	0.0
Lane	16	*	*	*	*	*	*	*
Leavenwort	367	66.8	16.3	11.2	1.4	1.1	3.3	0.0
Lincoln	23	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 3A**  
**2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Racial and Ethnic Distribution by Clinic**

<b>Comparison of Racial and Ethnic Distribution</b> <sup>(2)</sup>								
<b>Clinic</b>	<b>Number Included in Analysis</b>	<b>White, Not Hispanic</b>	<b>Black, Not Hispanic</b>	<b>Hispanic</b>	<b>American Indian/Alaska Native</b>	<b>Asian/Pacific Islander</b>	<b>Multiple Races</b>	<b>All Other/Unknown</b>
		<b>%</b>	<b>%</b>	<b>%</b>	<b>%</b>	<b>%</b>	<b>%</b>	<b>%</b>
Linn	69	*	*	*	*	*	*	*
Logan	11	*	*	*	*	*	*	*
Lyon	267	57.3	6.0	34.8	0.4	1.1	0.4	0.0
Marion	53	*	*	*	*	*	*	*
Marshall	60	*	*	*	*	*	*	*
Mc Pherson	140	85.0	2.1	10.7	0.0	0.7	1.4	0.0
Meade	36	*	*	*	*	*	*	*
Miami	158	91.1	1.3	5.7	0.0	0.6	1.3	0.0
Mitchell	43	*	*	*	*	*	*	*
Montgome-C	158	70.9	11.4	14.6	0.6	1.3	1.3	0.0
Montgome-I	162	80.9	5.6	11.1	1.9	0.6	0.0	0.0
Morris	23	*	*	*	*	*	*	*
Nemaha	30	*	*	*	*	*	*	*
Neosho-Ch	146	92.5	1.4	4.8	0.0	0.0	1.4	0.0
Neosho-Er	8	*	*	*	*	*	*	*
Ness	8	*	*	*	*	*	*	*
Norton	50	*	*	*	*	*	*	*
Osage	102	96.1	1.0	2.0	0.0	0.0	1.0	0.0
Osborne	17	*	*	*	*	*	*	*
Ottawa	14	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 3A**  
**2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Racial and Ethnic Distribution by Clinic**

<b>Comparison of Racial and Ethnic Distribution</b> <sup>(2)</sup>								
<b>Clinic</b>	<b>Number Included in Analysis</b>	<b>White, Not Hispanic</b>	<b>Black, Not Hispanic</b>	<b>Hispanic</b>	<b>American Indian/Alaska Native</b>	<b>Asian/Pacific Islander</b>	<b>Multiple Races</b>	<b>All Other/Unknown</b>
		<b>%</b>	<b>%</b>	<b>%</b>	<b>%</b>	<b>%</b>	<b>%</b>	<b>%</b>
Pawnee	49	*	*	*	*	*	*	*
Phillips	32	*	*	*	*	*	*	*
Pottawat-S	12	*	*	*	*	*	*	*
Pottawat-W	9	*	*	*	*	*	*	*
Pratt	98	*	*	*	*	*	*	*
Rawlins	8	*	*	*	*	*	*	*
Reno	460	79.6	3.0	14.3	0.2	0.2	2.6	0.0
Republic	28	*	*	*	*	*	*	*
Rice	73	*	*	*	*	*	*	*
Riley-Ft	611	67.6	11.5	13.3	2.8	3.8	1.1	0.0
Riley-Manh	407	69.0	11.1	11.3	0.7	5.7	2.2	0.0
Rooks	50	*	*	*	*	*	*	*
Rush	11	*	*	*	*	*	*	*
Russell	47	*	*	*	*	*	*	*
Saline	486	71.8	5.6	19.3	0.2	0.6	2.5	0.0
Scott	48	*	*	*	*	*	*	*
Sedgwick-C	1,225	39.1	15.0	35.2	1.0	7.1	2.6	0.0
Sedgwick-M	1,891	33.8	24.7	35.0	1.4	3.1	2.1	0.0
Sedgwick-S	1,420	65.5	8.5	19.6	1.8	1.9	2.7	0.0
Seneca	12	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 3A**  
**2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Racial and Ethnic Distribution by Clinic**

<b>Comparison of Racial and Ethnic Distribution</b> <sup>(2)</sup>								
<b>Clinic</b>	<b>Number Included in Analysis</b>	<b>White, Not Hispanic</b>	<b>Black, Not Hispanic</b>	<b>Hispanic</b>	<b>American Indian/Alaska Native</b>	<b>Asian/Pacific Islander</b>	<b>Multiple Races</b>	<b>All Other/Unknown</b>
		%	%	%	%	%	%	%
Seward	447	15.7	2.2	81.0	0.4	0.4	0.2	0.0
ShawneeM&I	293	35.8	16.0	42.7	0.7	1.7	3.1	0.0
ShawneeMa	1,166	58.1	17.2	19.0	1.9	0.8	2.9	0.0
Sheridan	8	*	*	*	*	*	*	*
Sherman	57	*	*	*	*	*	*	*
Smith	18	*	*	*	*	*	*	*
Stafford	24	*	*	*	*	*	*	*
Stanton	11	*	*	*	*	*	*	*
Stevens	72	*	*	*	*	*	*	*
Sumner	136	83.8	1.5	8.8	2.9	0.0	2.9	0.0
Thomas	56	*	*	*	*	*	*	*
Trego	12	*	*	*	*	*	*	*
Wabaunsee	19	*	*	*	*	*	*	*
Washington	26	*	*	*	*	*	*	*
Wichita	8	*	*	*	*	*	*	*
Wilson	84	*	*	*	*	*	*	*
Woodson	14	*	*	*	*	*	*	*
Wyandotte	2,101	26.2	27.0	39.3	0.6	6.1	0.9	0.0
Unknown Clinics	25	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 3A**  
**2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Racial and Ethnic Distribution by Clinic**

<b>Comparison of Racial and Ethnic Distribution</b> <sup>(2)</sup>								
<b>Clinic</b>	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
<b>Kansas</b>	<b>21,370</b>	<b>56.0</b>	<b>11.3</b>	<b>27.2</b>	<b>1.2</b>	<b>2.6</b>	<b>1.7</b>	<b>0.0</b>
<b>Nation (Prior Year)</b>	<b>1,162,961</b>	<b>41.6</b>	<b>22.7</b>	<b>29.2</b>	<b>1.1</b>	<b>3.4</b>	<b>1.3</b>	<b>0.7</b>

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 4A**  
**2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Age Distribution by Clinic**

<b>Comparison of Age Distribution</b> <sup>(2)</sup>								
<b>Clinic</b>	<b>Number Included in Analysis</b>	<b>&lt; 15 Years %</b>	<b>15 - 17 Years %</b>	<b>18 - 19 Years %</b>	<b>20 - 29 Years %</b>	<b>30 - 39 Years %</b>	<b>&gt;= 40 Years %</b>	<b>Unknown %</b>
Allen	71	*	*	*	*	*	*	*
Anderson	47	*	*	*	*	*	*	*
Atchison	148	0.0	8.1	16.9	61.5	13.5	0.0	0.0
Barber-Kio	1	*	*	*	*	*	*	*
Barber-Med	36	*	*	*	*	*	*	*
Barton	259	0.0	7.3	16.2	64.1	11.2	1.2	0.0
Bourbon	111	0.0	4.5	18.0	59.5	17.1	0.0	0.9
Brown	89	*	*	*	*	*	*	*
Butler	253	0.0	3.6	13.0	68.0	14.2	1.2	0.0
Chase	16	*	*	*	*	*	*	*
Chautauqua	32	*	*	*	*	*	*	*
CherokeeBS	75	*	*	*	*	*	*	*
CherokeeCo	93	*	*	*	*	*	*	*
Cheyenne	20	*	*	*	*	*	*	*
Clark	4	*	*	*	*	*	*	*
Clay	50	*	*	*	*	*	*	*
Cloud	60	*	*	*	*	*	*	*
Coffey	26	*	*	*	*	*	*	*
Comanche	9	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.





**Table 4A**  
**2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Age Distribution by Clinic**

<b>Comparison of Age Distribution</b> <sup>(2)</sup>								
<b>Clinic</b>	<b>Number Included in Analysis</b>	<b>&lt; 15 Years %</b>	<b>15 - 17 Years %</b>	<b>18 - 19 Years %</b>	<b>20 - 29 Years %</b>	<b>30 - 39 Years %</b>	<b>&gt;= 40 Years %</b>	<b>Unknown %</b>
Cowley-Ark	187	0.0	6.4	12.3	64.2	16.0	1.1	0.0
Cowley-Win	174	0.0	6.9	19.0	61.5	12.1	0.6	0.0
Crawford-G	1	*	*	*	*	*	*	*
Crawford-P	367	0.5	2.7	16.3	62.9	16.1	1.4	0.0
Decatur	16	*	*	*	*	*	*	*
DickinsonA	103	0.0	4.9	16.5	66.0	12.6	0.0	0.0
DickinsonH	19	*	*	*	*	*	*	*
Doniphan	40	*	*	*	*	*	*	*
Douglas	525	0.2	3.2	9.3	63.8	21.3	2.1	0.0
Edwards	21	*	*	*	*	*	*	*
Elk	12	*	*	*	*	*	*	*
Ellis	154	0.6	2.6	16.9	64.3	14.9	0.6	0.0
Ellsworth	34	*	*	*	*	*	*	*
Finney	684	0.4	5.6	16.7	55.6	20.3	1.5	0.0
Ford	674	0.0	5.2	12.8	58.8	20.9	2.4	0.0
Franklin	189	0.0	4.8	13.8	61.4	19.0	1.1	0.0
Geary	406	0.2	3.7	16.0	66.3	12.8	1.0	0.0
Gove	13	*	*	*	*	*	*	*
Graham	21	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 4A**  
**2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Age Distribution by Clinic**

<b>Comparison of Age Distribution</b> <sup>(2)</sup>								
<b>Clinic</b>	<b>Number Included in Analysis</b>	<b>&lt; 15 Years %</b>	<b>15 - 17 Years %</b>	<b>18 - 19 Years %</b>	<b>20 - 29 Years %</b>	<b>30 - 39 Years %</b>	<b>&gt;= 40 Years %</b>	<b>Unknown %</b>
Grant	72	*	*	*	*	*	*	*
Gray	54	*	*	*	*	*	*	*
Greeley	14	*	*	*	*	*	*	*
Greenwood	35	*	*	*	*	*	*	*
Hamilton	19	*	*	*	*	*	*	*
Harper	60	*	*	*	*	*	*	*
Harvey	251	0.4	6.8	13.9	54.2	22.7	2.0	0.0
Haskell	22	*	*	*	*	*	*	*
Hodgeman	12	*	*	*	*	*	*	*
JacksonHol	70	*	*	*	*	*	*	*
JacksonPBP	7	*	*	*	*	*	*	*
Jefferson	57	*	*	*	*	*	*	*
Jewell	6	*	*	*	*	*	*	*
Johnson-M	808	0.0	2.6	9.2	65.0	21.9	1.2	0.1
Johnson-O	1,090	0.2	3.2	9.7	60.1	24.6	2.1	0.1
Kearny	16	*	*	*	*	*	*	*
Kingman	47	*	*	*	*	*	*	*
Kiowa	15	*	*	*	*	*	*	*
Labette	160	1.3	6.9	11.9	66.3	13.1	0.6	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 4A**  
**2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Age Distribution by Clinic**

<b>Comparison of Age Distribution</b> <sup>(2)</sup>								
<b>Clinic</b>	<b>Number Included in Analysis</b>	<b>&lt; 15 Years %</b>	<b>15 - 17 Years %</b>	<b>18 - 19 Years %</b>	<b>20 - 29 Years %</b>	<b>30 - 39 Years %</b>	<b>&gt;= 40 Years %</b>	<b>Unknown %</b>
Lane	16	*	*	*	*	*	*	*
Leavenworth	367	0.3	3.5	15.8	66.8	12.8	0.8	0.0
Lincoln	23	*	*	*	*	*	*	*
Linn	69	*	*	*	*	*	*	*
Logan	11	*	*	*	*	*	*	*
Lyon	267	0.4	7.5	11.6	66.3	13.1	1.1	0.0
Marion	53	*	*	*	*	*	*	*
Marshall	60	*	*	*	*	*	*	*
Mc Pherson	140	0.7	4.3	15.0	62.1	16.4	1.4	0.0
Meade	36	*	*	*	*	*	*	*
Miami	158	0.0	3.8	17.1	70.9	8.2	0.0	0.0
Mitchell	43	*	*	*	*	*	*	*
Montgome-C	158	0.0	5.1	13.9	65.2	14.6	0.6	0.6
Montgome-I	162	1.2	8.6	15.4	55.6	17.9	1.2	0.0
Morris	23	*	*	*	*	*	*	*
Nemaha	30	*	*	*	*	*	*	*
Neosho-Ch	146	0.0	4.8	15.8	66.4	11.0	0.7	1.4
Neosho-Er	8	*	*	*	*	*	*	*
Ness	8	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 4A**  
**2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Age Distribution by Clinic**

Comparison of Age Distribution <sup>(2)</sup>								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Norton	50	*	*	*	*	*	*	*
Osage	102	0.0	1.0	13.7	62.7	21.6	1.0	0.0
Osborne	17	*	*	*	*	*	*	*
Ottawa	14	*	*	*	*	*	*	*
Pawnee	49	*	*	*	*	*	*	*
Phillips	32	*	*	*	*	*	*	*
Pottawat-S	12	*	*	*	*	*	*	*
Pottawat-W	9	*	*	*	*	*	*	*
Pratt	98	*	*	*	*	*	*	*
Rawlins	8	*	*	*	*	*	*	*
Reno	460	0.4	4.6	15.9	65.7	13.0	0.4	0.0
Republic	28	*	*	*	*	*	*	*
Rice	73	*	*	*	*	*	*	*
Riley-Ft	611	0.0	0.5	9.5	78.4	11.6	0.0	0.0
Riley-Manh	407	0.0	2.5	12.8	69.3	15.2	0.2	0.0
Rooks	50	*	*	*	*	*	*	*
Rush	11	*	*	*	*	*	*	*
Russell	47	*	*	*	*	*	*	*
Saline	486	0.6	7.2	13.8	61.3	15.6	1.4	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 4A**  
**2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Age Distribution by Clinic**

Comparison of Age Distribution <sup>(2)</sup>								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Scott	48	*	*	*	*	*	*	*
Sedgwick-C	1,225	0.7	4.8	11.6	60.7	20.5	1.7	0.0
Sedgwick-M	1,891	0.3	5.0	12.7	61.0	19.4	1.5	0.1
Sedgwick-S	1,420	0.4	5.9	14.2	60.1	18.1	1.3	0.1
Seneca	12	*	*	*	*	*	*	*
Seward	447	0.2	9.4	12.5	53.5	22.6	1.6	0.2
ShawneeM&I	293	0.3	7.8	16.4	59.7	15.0	0.7	0.0
ShawneeMa	1,166	0.6	5.5	13.8	63.6	15.4	1.1	0.0
Sheridan	8	*	*	*	*	*	*	*
Sherman	57	*	*	*	*	*	*	*
Smith	18	*	*	*	*	*	*	*
Stafford	24	*	*	*	*	*	*	*
Stanton	11	*	*	*	*	*	*	*
Stevens	72	*	*	*	*	*	*	*
Sumner	136	0.0	3.7	13.2	64.7	16.9	1.5	0.0
Thomas	56	*	*	*	*	*	*	*
Trego	12	*	*	*	*	*	*	*
Wabaunsee	19	*	*	*	*	*	*	*
Washington	26	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 4A**  
**2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Age Distribution by Clinic**

<b>Comparison of Age Distribution</b> <sup>(2)</sup>								
<b>Clinic</b>	<b>Number Included in Analysis</b>	<b>&lt; 15 Years %</b>	<b>15 - 17 Years %</b>	<b>18 - 19 Years %</b>	<b>20 - 29 Years %</b>	<b>30 - 39 Years %</b>	<b>&gt;= 40 Years %</b>	<b>Unknown %</b>
Wichita	8	*	*	*	*	*	*	*
Wilson	84	*	*	*	*	*	*	*
Woodson	14	*	*	*	*	*	*	*
Wyandotte	2,101	0.4	6.9	12.9	57.8	20.4	1.6	0.0
Unknown Clinics	25	*	*	*	*	*	*	*
<b>Kansas</b>	<b>21,370</b>	<b>0.3</b>	<b>5.1</b>	<b>13.2</b>	<b>62.1</b>	<b>17.9</b>	<b>1.3</b>	<b>0.1</b>
<b>Nation (Prior Year)</b>	<b>1,162,961</b>	<b>0.3</b>	<b>5.3</b>	<b>11.5</b>	<b>59.4</b>	<b>21.7</b>	<b>1.7</b>	<b>0.2</b>

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 5A**  
**2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Maternal Health Indicators by Clinic**

Comparison of Maternal Health Indicators <sup>(2, 3)</sup>														
Clinic	Prepregnancy BMI <sup>(4)</sup>			Weight Gain <sup>(5)</sup>			Anemia (Low Hb/Hct) <sup>(6)</sup>				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Allen	70	* *	* *	62	* *	* *	9	* *	20	* *	64	* *	66	* *
Anderson	46	* *	* *	38	* *	* *	4	* *	11	* *	39	* *	39	* *
Atchison	142	8.5 (38)	47.9 (3)	120	28.3 (36)	46.7 (12)	16	* *	26	* *	120	5.0 (6)	120	7.5(6)
Barber-Kio	1	* *	* *	0	* *	* *	0	* *	0	* *	0	* *	0	* *
Barber-Med	36	* *	* *	29	* *	* *	3	* *	4	* *	30	* *	30	* *
Barton	256	4.3 (24)	57.8 (32)	226	26.5 (33)	42.9 (4)	24	* *	30	* *	231	6.9 (14)	231	12.6(26)
Bourbon	110	2.7 (3)	60.9 (36)	101	21.8 (14)	52.5 (33)	16	* *	15	* *	101	14.9 (36)	101	14.9(32)
Brown	87	* *	* *	72	* *	* *	12	* *	12	* *	70	* *	70	* *
Butler	248	4.8 (29)	55.2 (18)	224	24.6 (29)	48.7 (19)	36	* *	38	* *	231	6.1 (11)	231	13.4(29)
Chase	16	* *	* *	15	* *	* *	3	* *	2	* *	15	* *	15	* *
Chautauqua	32	* *	* *	28	* *	* *	7	* *	4	* *	28	* *	28	* *
CherokeeBS	70	* *	* *	65	* *	* *	12	* *	12	* *	67	* *	67	* *
CherokeeCo	89	* *	* *	77	* *	* *	9	* *	13	* *	78	* *	78	* *
Cheyenne	19	* *	* *	16	* *	* *	3	* *	8	* *	17	* *	17	* *
Clark	4	* *	* *	4	* *	* *	1	* *	1	* *	4	* *	4	* *
Clay	49	* *	* *	42	* *	* *	7	* *	9	* *	44	* *	44	* *
Cloud	58	* *	* *	53	* *	* *	3	* *	12	* *	55	* *	56	* *

(1) Reporting period is January 1 through December 31.  
(2) Excludes records with unknown data or errors.  
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.  
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).  
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).  
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.  
\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 5A**  
**2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Maternal Health Indicators by Clinic**

Comparison of Maternal Health Indicators <sup>(2, 3)</sup>															
Clinic	Prepregnancy BMI <sup>(4)</sup>				Weight Gain <sup>(5)</sup>			Anemia (Low Hb/Hct) <sup>(6)</sup>				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight		Over-weight		< Ideal		> Ideal		3rd Trimester		Postpartum				
	Number	% (Rank)	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Coffey	26	* *	* *	* *	22	* *	* *	5	* *	4	* *	22	* *	22	* *
Comanche	9	* *	* *	* *	9	* *	* *	0	* *	3	* *	9	* *	9	* *
Cowley-Ark	184	3.3 (8)	61.4 (38)		158	21.5 (12)	50.0 (24)	23	* *	30	* *	163	9.8 (28)	163	12.9(27)
Cowley-Win	170	2.9 (4)	56.5 (26)		148	22.3 (21)	44.6 (8)	18	* *	17	* *	153	2.0 (2)	153	11.1(21)
Crawford-G	1	* *	* *	* *	1	* *	* *	0	* *	0	* *	1	* *	1	* *
Crawford-P	361	6.6 (33)	56.8 (27)		315	22.2 (20)	47.3 (14)	46	* *	57	* *	320	10.3 (30)	323	13.3(28)
Decatur	16	* *	* *	* *	15	* *	* *	3	* *	4	* *	15	* *	15	* *
DickinsonA	102	3.9 (19)	55.9 (21)		87	* *	* *	24	* *	16	* *	88	* *	88	* *
DickinsonH	19	* *	* *	* *	17	* *	* *	3	* *	3	* *	17	* *	17	* *
Doniphan	38	* *	* *	* *	26	* *	* *	5	* *	10	* *	29	* *	30	* *
Douglas	516	3.3 (8)	52.9 (7)		432	20.1 (7)	51.9 (31)	67	* *	124	29.0 (7)	426	9.9 (29)	426	7.7(7)
Edwards	21	* *	* *	* *	19	* *	* *	1	* *	4	* *	18	* *	18	* *
Elk	11	* *	* *	* *	10	* *	* *	2	* *	1	* *	11	* *	11	* *
Ellis	150	3.3 (8)	50.0 (5)		125	20.0 (6)	41.6 (2)	24	* *	35	* *	132	1.5 (1)	131	5.3(2)
Ellsworth	34	* *	* *	* *	31	* *	* *	4	* *	1	* *	30	* *	30	* *
Finney	671	7.7 (37)	53.4 (12)		598	27.1 (35)	38.5 (1)	82	* *	94	* *	606	8.9 (26)	606	5.9(4)
Ford	669	2.4 (2)	61.1 (37)		589	25.0 (30)	47.5 (15)	77	* *	70	* *	593	7.4 (18)	591	8.6(10)

(1) Reporting period is January 1 through December 31.  
(2) Excludes records with unknown data or errors.  
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.  
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).  
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).  
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.  
\* Percentages are not calculated if <100 records are available for analysis after exclusions.





**Table 5A**  
**2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Maternal Health Indicators by Clinic**

Comparison of Maternal Health Indicators <sup>(2, 3)</sup>														
Clinic	Prepregnancy BMI <sup>(4)</sup>			Weight Gain <sup>(5)</sup>			Anemia (Low Hb/Hct) <sup>(6)</sup>				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Franklin	183	3.3 (8)	58.5 (33)	157	22.9 (23)	51.6 (29)	21	* *	44	* *	137	7.3 (16)	137	13.9(30)
Geary	387	3.6 (15)	51.4 (6)	310	21.9 (16)	51.0 (28)	52	* *	64	* *	323	6.2 (12)	322	8.7(11)
Gove	12	* *	* *	10	* *	* *	3	* *	3	* *	11	* *	11	* *
Graham	20	* *	* *	18	* *	* *	0	* *	1	* *	19	* *	19	* *
Grant	72	* *	* *	67	* *	* *	13	* *	17	* *	67	* *	67	* *
Gray	54	* *	* *	44	* *	* *	3	* *	10	* *	44	* *	44	* *
Greeley	14	* *	* *	12	* *	* *	2	* *	3	* *	12	* *	12	* *
Greenwood	35	* *	* *	33	* *	* *	4	* *	4	* *	33	* *	33	* *
Hamilton	19	* *	* *	17	* *	* *	3	* *	4	* *	17	* *	17	* *
Harper	59	* *	* *	50	* *	* *	5	* *	11	* *	52	* *	52	* *
Harvey	246	2.0 (1)	57.3 (30)	218	22.9 (23)	47.2 (13)	31	* *	43	* *	226	3.1 (3)	226	15.9(33)
Haskell	21	* *	* *	18	* *	* *	3	* *	2	* *	19	* *	19	* *
Hodgeman	12	* *	* *	12	* *	* *	2	* *	2	* *	12	* *	12	* *
JacksonHol	70	* *	* *	60	* *	* *	8	* *	14	* *	61	* *	61	* *
JacksonPBP	7	* *	* *	4	* *	* *	0	* *	3	* *	5	* *	5	* *
Jefferson	56	* *	* *	50	* *	* *	9	* *	18	* *	51	* *	50	* *
Jewell	6	* *	* *	5	* *	* *	1	* *	4	* *	5	* *	5	* *

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
- (4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
- (5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
- (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- \* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 5A**  
**2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Maternal Health Indicators by Clinic**

Comparison of Maternal Health Indicators <sup>(2, 3)</sup>														
Clinic	Prepregnancy BMI <sup>(4)</sup>			Weight Gain <sup>(5)</sup>			Anemia (Low Hb/Hct) <sup>(6)</sup>				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Johnson-M	784	3.2 (6)	57.0 (28)	688	21.8 (14)	48.1 (18)	109	22.9 (2)	194	14.9 (1)	702	11.1 (34)	702	8.3(9)
Johnson-O	1,052	2.9 (4)	52.9 (7)	928	22.3 (21)	46.1 (11)	154	20.1 (1)	252	17.9 (2)	963	10.8 (32)	962	7.4(5)
Kearny	16	* *	* *	14	* *	* *	4	* *	3	* *	14	* *	14	* *
Kingman	46	* *	* *	34	* *	* *	4	* *	11	* *	36	* *	36	* *
Kiowa	15	* *	* *	14	* *	* *	3	* *	6	* *	14	* *	14	* *
Labette	157	5.1 (30)	56.1 (23)	140	25.7 (32)	44.3 (7)	22	* *	32	* *	139	9.4 (27)	140	19.3(35)
Lane	15	* *	* *	11	* *	* *	1	* *	1	* *	12	* *	12	* *
Leavenwort	350	4.3 (24)	52.9 (7)	290	17.9 (3)	49.3 (22)	54	* *	80	* *	302	5.6 (7)	301	11.0(20)
Lincoln	23	* *	* *	19	* *	* *	1	* *	2	* *	19	* *	19	* *
Linn	69	* *	* *	60	* *	* *	4	* *	6	* *	61	* *	61	* *
Logan	11	* *	* *	10	* *	* *	2	* *	2	* *	10	* *	10	* *
Lyon	264	3.8 (17)	59.8 (35)	230	23.9 (27)	48.7 (19)	26	* *	49	* *	232	10.3 (30)	231	14.7(31)
Marion	53	* *	* *	42	* *	* *	8	* *	5	* *	45	* *	45	* *
Marshall	57	* *	* *	51	* *	* *	9	* *	18	* *	55	* *	55	* *
Mc Pherson	136	5.9 (31)	52.9 (7)	118	21.2 (11)	44.9 (10)	14	* *	37	* *	124	3.2 (4)	125	10.4(19)
Meade	36	* *	* *	36	* *	* *	5	* *	8	* *	36	* *	35	* *
Miami	155	3.2 (6)	54.8 (15)	141	12.1 (1)	60.3 (36)	28	* *	35	* *	148	3.4 (5)	148	11.5(22)

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
- (4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
- (5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
- (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- \* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 5A**  
**2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Maternal Health Indicators by Clinic**

Comparison of Maternal Health Indicators <sup>(2, 3)</sup>														
Clinic	Prepregnancy BMI <sup>(4)</sup>			Weight Gain <sup>(5)</sup>			Anemia (Low Hb/Hct) <sup>(6)</sup>				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Mitchell	42	* * *	* *	37	* *	* *	7	* *	3	* *	37	* *	38	* *
Montgome-C	153	7.2 (35)	53.6 (13)	139	23.0 (25)	44.6 (8)	40	* *	37	* *	139	8.6 (24)	139	18.0(34)
Montgome-I	158	7.6 (36)	44.9 (1)	145	26.9 (34)	42.8 (3)	21	* *	34	* *	148	10.8 (32)	147	11.6(23)
Morris	21	* *	* *	21	* *	* *	0	* *	9	* *	21	* *	22	* *
Nemaha	30	* *	* *	26	* *	* *	2	* *	5	* *	26	* *	26	* *
Neosho-Ch	140	3.6 (15)	55.0 (16)	126	20.6 (9)	50.0 (24)	22	* *	22	* *	134	8.2 (19)	134	9.7(16)
Neosho-Er	7	* *	* *	6	* *	* *	1	* *	2	* *	7	* *	7	* *
Ness	8	* *	* *	7	* *	* *	0	* *	3	* *	7	* *	7	* *
Norton	48	* *	* *	42	* *	* *	3	* *	12	* *	44	* *	44	* *
Osage	100	7.0 (34)	55.0 (16)	88	* *	* *	13	* *	25	* *	93	* *	93	* *
Osborne	17	* *	* *	15	* *	* *	3	* *	2	* *	15	* *	15	* *
Ottawa	14	* *	* *	12	* *	* *	1	* *	3	* *	12	* *	12	* *
Pawnee	48	* *	* *	43	* *	* *	9	* *	9	* *	45	* *	45	* *
Phillips	32	* *	* *	26	* *	* *	8	* *	6	* *	26	* *	26	* *
Pottawat-S	12	* *	* *	8	* *	* *	2	* *	2	* *	9	* *	9	* *
Pottawat-W	9	* *	* *	8	* *	* *	0	* *	4	* *	8	* *	8	* *
Pratt	95	* *	* *	86	* *	* *	14	* *	22	* *	89	* *	89	* *

(1) Reporting period is January 1 through December 31.  
(2) Excludes records with unknown data or errors.  
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.  
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).  
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).  
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.  
\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 5A**  
**2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Maternal Health Indicators by Clinic**

<b>Comparison of Maternal Health Indicators</b> <sup>(2, 3)</sup>														
<b>Clinic</b>	<b>Prepregnancy BMI</b> (4)			<b>Weight Gain</b> (5)			<b>Anemia</b> (Low Hb/Hct) (6)				<b>Gestational Diabetes</b> During Pregnancy		<b>Hypertension</b> During Pregnancy	
	<b>Under-weight</b>		<b>Over-weight</b>	<b>&lt; Ideal</b>		<b>&gt; Ideal</b>	<b>3rd Trimester</b>		<b>Postpartum</b>					
	<b>Number</b>	<b>% (Rank)</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>
Rawlins	7	* *	* *	5	* *	* *	0	* *	2	* *	5	* *	5	* *
Reno	451	3.8 (17)	56.1 (23)	393	19.1 (4)	52.4 (32)	54	* *	69	* *	402	6.2 (12)	402	10.2(17)
Republic	27	* *	* *	23	* *	* *	0	* *	8	* *	25	* *	25	* *
Rice	73	* *	* *	64	* *	* *	11	* *	15	* *	65	* *	65	* *
Riley-Ft	576	3.5 (14)	53.3 (11)	473	19.7 (5)	52.6 (34)	96	* *	138	31.2 (9)	499	7.0 (15)	499	10.2(17)
Riley-Manh	396	4.0 (21)	46.7 (2)	336	22.0 (17)	50.6 (27)	53	* *	87	* *	345	11.6 (35)	350	8.9(13)
Rooks	48	* *	* *	44	* *	* *	4	* *	11	* *	43	* *	43	* *
Rush	10	* *	* *	7	* *	* *	1	* *	2	* *	8	* *	8	* *
Russell	45	* *	* *	42	* *	* *	11	* *	9	* *	43	* *	43	* *
Saline	478	4.4 (27)	55.6 (20)	434	15.9 (2)	53.7 (35)	67	* *	65	* *	439	7.3 (16)	438	9.6(15)
Scott	48	* *	* *	47	* *	* *	11	* *	14	* *	45	* *	45	* *
Sedgwick-C	1,200	4.7 (28)	53.7 (14)	1,085	21.0 (10)	48.0 (17)	245	27.8 (3)	279	18.3 (3)	1,112	8.5 (23)	1,113	9.5(14)
Sedgwick-M	1,827	3.4 (12)	57.1 (29)	1,660	23.5 (26)	48.9 (21)	357	28.3 (4)	413	21.3 (4)	1,711	8.7 (25)	1,711	12.1(24)
Sedgwick-S	1,375	3.9 (19)	56.3 (25)	1,241	20.5 (8)	51.6 (29)	234	32.5 (5)	324	27.2 (6)	1,289	8.4 (21)	1,289	12.4(25)
Seneca	12	* *	* *	10	* *	* *	2	* *	1	* *	10	* *	10	* *
Seward	440	3.4 (12)	59.5 (34)	402	25.1 (31)	43.8 (6)	74	* *	55	* *	408	8.3 (20)	407	4.7(1)
ShawneeM&I	288	4.2 (23)	49.3 (4)	245	22.0 (17)	47.8 (16)	11	* *	28	* *	249	8.4 (21)	249	5.6(3)

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
- (4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
- (5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
- (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- \* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 5A**  
**2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Maternal Health Indicators by Clinic**

Comparison of Maternal Health Indicators <sup>(2, 3)</sup>														
Clinic	Prepregnancy BMI <sup>(4)</sup>			Weight Gain <sup>(5)</sup>			Anemia (Low Hb/Hct) <sup>(6)</sup>				Gestational Diabetes During Pregnancy	Hypertension During Pregnancy		
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
							Number	% (Rank)	% (Rank)	Number			% (Rank)	Number
ShawneeMa	1,143	4.3 (24)	56.0 (22)	1,015	21.7 (13)	50.3 (26)	135	48.9 (7)	177	29.4 (8)	1,038	5.9 (10)	1,038	8.7(11)
Sheridan	7	* *	* *	7	* *	* *	2	* *	0	* *	8	* *	8	* *
Sherman	55	* *	* *	48	* *	* *	5	* *	15	* *	50	* *	50	* *
Smith	18	* *	* *	17	* *	* *	3	* *	2	* *	17	* *	17	* *
Stafford	24	* *	* *	18	* *	* *	2	* *	3	* *	18	* *	18	* *
Stanton	11	* *	* *	10	* *	* *	1	* *	4	* *	10	* *	10	* *
Stevens	70	* *	* *	62	* *	* *	15	* *	18	* *	62	* *	63	* *
Sumner	132	6.1 (32)	57.6 (31)	119	24.4 (28)	42.9 (4)	20	* *	32	* *	123	5.7 (8)	123	19.5(36)
Thomas	55	* *	* *	44	* *	* *	9	* *	7	* *	46	* *	46	* *
Trego	12	* *	* *	10	* *	* *	3	* *	2	* *	8	* *	10	* *
Wabaunsee	18	* *	* *	17	* *	* *	1	* *	10	* *	18	* *	18	* *
Washington	26	* *	* *	19	* *	* *	6	* *	4	* *	19	* *	19	* *
Wichita	8	* *	* *	8	* *	* *	1	* *	0	* *	8	* *	8	* *
Wilson	81	* *	* *	80	* *	* *	12	* *	13	* *	82	* *	82	* *
Woodson	14	* *	* *	14	* *	* *	2	* *	3	* *	14	* *	14	* *
Wyandotte	2,034	4.0 (21)	55.5 (19)	1,821	22.0 (17)	49.4 (23)	310	36.5 (6)	371	22.4 (5)	1,851	5.7 (8)	1,849	8.0(8)
Unknown Clinics	25	* *	* *	24	* *	* *	2	* *	3	* *	24	* *	24	* *

(1) Reporting period is January 1 through December 31.  
(2) Excludes records with unknown data or errors.  
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.  
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).  
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).  
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.  
\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 5A**  
**2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Maternal Health Indicators by Clinic**

<b>Comparison of Maternal Health Indicators</b> <sup>(2, 3)</sup>														
<b>Clinic</b>	<b>Prepregnancy BMI</b> (4)			<b>Weight Gain</b> (5)			<b>Anemia</b> (Low Hb/Hct) (6)				<b>Gestational Diabetes</b> <b>During</b> <b>Pregnancy</b>	<b>Hypertension</b> <b>During</b> <b>Pregnancy</b>		
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
		Number	% (Rank)		% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)			Number	% (Rank)
<b>Kansas</b>	<b>20,811</b>	<b>4.1</b>	<b>55.5</b>	<b>18,364</b>	<b>22.5</b>	<b>48.6</b>	<b>3,063</b>	<b>30.9</b>	<b>4,062</b>	<b>22.1</b>	<b>18,787</b>	<b>7.6</b>	<b>18,793</b>	<b>10.1</b>
<b>Nation (Prior Year)</b>	<b>1,106,370</b>	<b>4.5</b>	<b>53.4</b>	<b>835,927</b>	<b>21.5</b>	<b>48.0</b>	<b>164,201</b>	<b>33.9</b>	<b>346,981</b>	<b>28.1</b>	<b>419,956</b>	<b>5.2</b>	<b>467,675</b>	<b>6.3</b>

(1) Reporting period is January 1 through December 31.  
(2) Excludes records with unknown data or errors.  
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.  
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).  
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).  
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.  
\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 6A**  
**2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Maternal Behavioral Indicators by Clinic**

<b>Comparison of Maternal Behavioral Indicators</b> <sup>(2, 3)</sup>										
<b>Clinic</b>	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	%	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Allen	60	* *	* *	* *	71	* *	71	* *	58	* *
Anderson	37	* *	* *	* *	47	* *	47	* *	43	* *
Atchison	121	90.9 (3)	3.3 (22)		147	46.3 (4)	148	12.8 (26)	122	77.9 (2)
Barber-Kio	0	* *	* *	* *	1	* *	1	* *	1	* *
Barber-Med	30	* *	* *	* *	36	* *	36	* *	32	* *
Barton	230	89.6 (8)	0.4 (4)		257	45.1 (8)	259	17.8 (17)	199	69.8 (10)
Bourbon	97	* *	* *	* *	111	38.7 (17)	111	16.2 (20)	96	* *
Brown	54	* *	* *	* *	88	* *	89	* *	79	* *
Butler	231	86.1 (14)	0.9 (8)		252	36.9 (21)	253	19.0 (14)	198	67.2 (13)
Chase	15	* *	* *	* *	16	* *	16	* *	15	* *
Chautauqua	28	* *	* *	* *	31	* *	32	* *	29	* *
CherokeeBS	49	* *	* *	* *	75	* *	75	* *	60	* *
CherokeeCo	57	* *	* *	* *	93	* *	93	* *	76	* *
Cheyenne	17	* *	* *	* *	20	* *	20	* *	16	* *
Clark	4	* *	* *	* *	4	* *	4	* *	4	* *
Clay	44	* *	* *	* *	50	* *	50	* *	44	* *
Cloud	56	* *	* *	* *	60	* *	60	* *	48	* *
Coffey	21	* *	* *	* *	26	* *	26	* *	23	* *
Comanche	8	* *	* *	* *	9	* *	9	* *	8	* *
Cowley-Ark	134	65.7 (31)	1.5 (11)		186	38.2 (19)	187	20.3 (12)	147	61.2 (19)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 6A**  
**2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Maternal Behavioral Indicators by Clinic**

Comparison of Maternal Behavioral Indicators <sup>(2, 3)</sup>									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy <sup>(4)</sup>		During Pregnancy <sup>(5)</sup>	
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Cowley-Win	136	78.7 (22)	10.3 (31)	171	46.8 (3)	174	11.5 (33)	145	57.9 (28)
Crawford-G	1	*	*	1	*	1	*	1	*
Crawford-P	274	86.5 (13)	4.0 (23)	364	45.9 (7)	367	13.6 (24)	303	55.8 (32)
Decatur	15	*	*	16	*	16	*	14	*
DickinsonA	88	*	*	101	43.6 (10)	103	22.3 (7)	87	*
DickinsonH	17	*	*	19	*	19	*	17	*
Doniphan	31	*	*	40	*	40	*	30	*
Douglas	391	70.6 (30)	11.0 (32)	522	42.7 (12)	525	18.7 (15)	429	64.1 (17)
Edwards	17	*	*	21	*	21	*	17	*
Elk	11	*	*	12	*	12	*	9	*
Ellis	125	88.0 (11)	3.2 (21)	153	28.8 (37)	154	27.9 (2)	129	61.2 (19)
Ellsworth	31	*	*	34	*	34	*	33	*
Finney	557	49.9 (35)	29.4 (35)	682	38.6 (18)	684	7.9 (37)	555	41.3 (34)
Ford	562	76.3 (24)	9.6 (30)	667	46.0 (6)	674	12.6 (27)	588	60.4 (23)
Franklin	127	83.5 (15)	7.1 (28)	188	44.1 (9)	189	13.8 (23)	150	65.3 (15)
Geary	309	79.0 (21)	1.0 (9)	405	42.0 (13)	406	20.4 (11)	334	59.9 (24)
Gove	12	*	*	13	*	13	*	8	*
Graham	19	*	*	20	*	21	*	18	*
Grant	56	*	*	70	*	72	*	61	*
Gray	43	*	*	52	*	54	*	44	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.





**Table 6A**  
**2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Maternal Behavioral Indicators by Clinic**

Comparison of Maternal Behavioral Indicators <sup>(2, 3)</sup>										
Clinic	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Prepregnancy <sup>(4)</sup>		During Pregnancy <sup>(5)</sup>	
	Number	% (Rank)	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Greeley	7	* *	* *	* *	14	* *	14	* *	11	* *
Greenwood	31	* *	* *	* *	35	* *	35	* *	28	* *
Hamilton	15	* *	* *	* *	19	* *	19	* *	15	* *
Harper	52	* *	* *	* *	59	* *	60	* *	48	* *
Harvey	224	81.7 (19)	4.0 (23)		251	48.2 (2)	251	23.5 (5)	212	67.5 (11)
Haskell	17	* *	* *	* *	22	* *	22	* *	18	* *
Hodgeman	12	* *	* *	* *	12	* *	12	* *	11	* *
JacksonHol	37	* *	* *	* *	70	* *	70	* *	54	* *
JacksonPBP	2	* *	* *	* *	7	* *	7	* *	5	* *
Jefferson	50	* *	* *	* *	57	* *	57	* *	49	* *
Jewell	6	* *	* *	* *	6	* *	6	* *	5	* *
Johnson-M	670	73.4 (27)	2.7 (18)		804	34.2 (27)	808	19.7 (13)	646	66.1 (14)
Johnson-O	906	62.5 (33)	2.2 (14)		1,088	35.8 (25)	1,090	12.6 (27)	906	56.6 (31)
Kearny	13	* *	* *	* *	16	* *	16	* *	15	* *
Kingman	36	* *	* *	* *	47	* *	47	* *	39	* *
Kiowa	14	* *	* *	* *	15	* *	15	* *	11	* *
Labette	101	90.1 (5)	2.0 (13)		160	36.3 (23)	160	11.9 (31)	131	71.0 (6)
Lane	12	* *	* *	* *	16	* *	16	* *	13	* *
Leavenwort	302	88.1 (10)	1.3 (10)		365	31.0 (34)	367	21.5 (8)	292	74.3 (4)
Lincoln	19	* *	* *	* *	23	* *	23	* *	22	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 6A**  
**2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Maternal Behavioral Indicators by Clinic**

Comparison of Maternal Behavioral Indicators <sup>(2, 3)</sup>										
Clinic	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Prepregnancy <sup>(4)</sup>		During Pregnancy <sup>(5)</sup>	
	Number	% (Rank)	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Linn	61	* *	* *	* *	69	* *	69	* *	64	* *
Logan	9	* *	* *	* *	11	* *	11	* *	11	* *
Lyon	219	90.0 (6)	0.5 (5)		267	46.1 (5)	267	21.3 (10)	226	58.8 (27)
Marion	45	* *	* *	* *	53	* *	53	* *	48	* *
Marshall	54	* *	* *	* *	60	* *	60	* *	50	* *
Mc Pherson	123	87.0 (12)	2.4 (16)		140	35.7 (26)	140	24.3 (4)	107	70.1 (8)
Meade	34	* *	* *	* *	35	* *	36	* *	33	* *
Miami	146	82.2 (18)	0.0 (1)		158	34.2 (27)	158	12.0 (30)	120	64.2 (16)
Mitchell	38	* *	* *	* *	43	* *	43	* *	35	* *
Montgome-C	109	89.0 (9)	0.0 (1)		157	33.1 (31)	158	13.3 (25)	139	78.4 (1)
Montgome-I	108	89.8 (7)	2.8 (19)		159	31.4 (33)	162	14.2 (22)	131	64.1 (17)
Morris	16	* *	* *	* *	22	* *	23	* *	16	* *
Nemaha	26	* *	* *	* *	30	* *	30	* *	27	* *
Neosho-Ch	123	91.1 (2)	2.4 (16)		145	40.7 (14)	146	17.1 (18)	120	70.0 (9)
Neosho-Er	7	* *	* *	* *	8	* *	8	* *	5	* *
Ness	7	* *	* *	* *	7	* *	8	* *	7	* *
Norton	44	* *	* *	* *	49	* *	50	* *	41	* *
Osage	69	* *	* *	* *	102	36.3 (23)	102	8.8 (35)	83	* *
Osborne	14	* *	* *	* *	17	* *	17	* *	15	* *
Ottawa	12	* *	* *	* *	14	* *	14	* *	11	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 6A**  
**2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Maternal Behavioral Indicators by Clinic**

Comparison of Maternal Behavioral Indicators <sup>(2, 3)</sup>										
Clinic	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Prepregnancy <sup>(4)</sup>		During Pregnancy <sup>(5)</sup>	
	Number	% (Rank)	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Pawnee	44	* *	* *	* *	49	* *	49	* *	41	* *
Phillips	26	* *	* *	* *	30	* *	32	* *	28	* *
Pottawat-S	8	* *	* *	* *	12	* *	12	* *	8	* *
Pottawat-W	8	* *	* *	* *	9	* *	9	* *	9	* *
Pratt	89	* *	* *	* *	98	* *	98	* *	81	* *
Rawlins	5	* *	* *	* *	8	* *	8	* *	7	* *
Reno	397	82.9 (16)	2.8 (19)		457	42.9 (11)	460	21.5 (8)	365	59.7 (25)
Republic	25	* *	* *	* *	27	* *	28	* *	26	* *
Rice	65	* *	* *	* *	72	* *	73	* *	62	* *
Riley-Ft	473	93.0 (1)	0.8 (7)		611	33.7 (30)	611	28.8 (1)	482	75.9 (3)
Riley-Manh	312	73.1 (28)	0.6 (6)		406	39.2 (16)	407	22.4 (6)	340	56.8 (30)
Rooks	45	* *	* *	* *	50	* *	50	* *	39	* *
Rush	8	* *	* *	* *	11	* *	11	* *	8	* *
Russell	43	* *	* *	* *	46	* *	47	* *	40	* *
Saline	305	79.7 (20)	5.2 (26)		486	34.2 (27)	486	8.6 (36)	406	67.5 (11)
Scott	47	* *	* *	* *	48	* *	48	* *	44	* *
Sedgwick-C	1,055	71.9 (29)	7.1 (28)		1,223	29.3 (36)	1,225	14.6 (21)	1,029	60.8 (22)
Sedgwick-M	1,696	73.7 (26)	4.2 (25)		1,889	30.2 (35)	1,891	18.0 (16)	1,531	60.9 (21)
Sedgwick-S	1,159	82.9 (16)	1.9 (12)		1,416	28.0 (38)	1,420	16.8 (19)	1,139	32.2 (35)
Seneca	10	* *	* *	* *	12	* *	12	* *	11	* *

(1) Reporting period is January 1 through December 31.  
(2) Excludes records with unknown data or errors.  
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.  
(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.  
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.  
\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 6A**  
**2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Maternal Behavioral Indicators by Clinic**

<b>Comparison of Maternal Behavioral Indicators</b> <sup>(2, 3)</sup>										
<b>Clinic</b>	<b>Medical Care</b>				<b>WIC Enrollment</b>		<b>Multivitamin Consumption</b>			
	<b>1st Trimester</b>		<b>No Care</b>		<b>1st Trimester</b>		<b>Prepregnancy</b> (4)		<b>During Pregnancy</b> (5)	
	<b>Number</b>	<b>% (Rank)</b>	<b>% (Rank)</b>		<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>
Seward	355	63.4 (32)	18.0 (33)		446	37.4 (20)	447	12.1 (29)	380	70.3 (7)
ShawneeM&I	217	76.0 (25)	2.3 (15)		293	57.7 (1)	293	25.3 (3)	264	59.5 (26)
ShawneeMa	931	77.0 (23)	5.2 (26)		1,162	39.5 (15)	1,166	10.5 (34)	983	52.1 (33)
Sheridan	8	* *	* *		8	* *	8	* *	4	* *
Sherman	49	* *	* *		57	* *	57	* *	44	* *
Smith	15	* *	* *		18	* *	18	* *	16	* *
Stafford	16	* *	* *		24	* *	24	* *	21	* *
Stanton	7	* *	* *		11	* *	11	* *	9	* *
Stevens	62	* *	* *		71	* *	72	* *	56	* *
Sumner	123	90.2 (4)	0.0 (1)		136	33.1 (31)	136	11.8 (32)	114	71.9 (5)
Thomas	45	* *	* *		56	* *	56	* *	49	* *
Trego	10	* *	* *		12	* *	12	* *	11	* *
Wabaunsee	18	* *	* *		19	* *	19	* *	13	* *
Washington	19	* *	* *		26	* *	26	* *	24	* *
Wichita	8	* *	* *		8	* *	8	* *	7	* *
Wilson	75	* *	* *		83	* *	84	* *	73	* *
Woodson	14	* *	* *		14	* *	14	* *	13	* *
Wyandotte	1,452	57.9 (34)	19.5 (34)		2,093	36.8 (22)	2,101	5.9 (38)	1,782	57.5 (29)
Unknown Clinics	17	* *	* *		25	* *	25	* *	22	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 6A**  
**2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Maternal Behavioral Indicators by Clinic**

<b>Comparison of Maternal Behavioral Indicators</b> <sup>(2, 3)</sup>									
<b>Clinic</b>	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Pregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
<b>Kansas</b>	<b>17,166</b>	<b>76.1</b>	<b>6.3</b>	<b>21,285</b>	<b>37.4</b>	<b>21,370</b>	<b>16.4</b>	<b>17,671</b>	<b>59.5</b>
<b>Nation (Prior Year)</b>	<b>429,770</b>	<b>83.1</b>	<b>4.1</b>	<b>1,074,863</b>	<b>36.0</b>	<b>380,154</b>	<b>19.2</b>	<b>418,448</b>	<b>85.0</b>

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 7A**  
**2011 Pregnancy Nutrition Surveillance** (1)  
**Kansas**  
**Comparison of Smoking Indicators by Clinic**

<b>Comparison of Smoking Indicators</b> (2, 3)										
<b>Clinic</b>	<b>Smoking</b>				<b>Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)</b>	<b>Smoking in Household</b>				
	<b>3 Months Prior to Pregnancy</b>		<b>Last 3 Months of Pregnancy</b>			<b>Prenatal</b>		<b>Postpartum</b>		
	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>
Allen	69	* *	56	* *	13	* *	58	* *	57	* *
Anderson	47	* *	36	* *	11	* *	42	* *	36	* *
Atchison	102	41.2 (25)	54	* *	18	* *	90	* *	54	* *
Barber-Kio	1	* *	0	* *	0	* *	1	* *	0	* *
Barber-Med	36	* *	30	* *	13	* *	32	* *	30	* *
Barton	258	38.8 (22)	230	23.0 (20)	62	* *	195	13.8 (21)	230	7.4 (15)
Bourbon	109	44.0 (30)	96	* *	38	* *	95	* *	96	* *
Brown	85	* *	56	* *	20	* *	77	* *	59	* *
Butler	251	53.4 (37)	229	33.6 (33)	102	28.4 (8)	196	7.1 (6)	229	2.6 (4)
Chase	16	* *	14	* *	9	* *	15	* *	14	* *
Chautauqua	30	* *	26	* *	8	* *	28	* *	26	* *
CherokeeBS	72	* *	64	* *	20	* *	57	* *	64	* *
CherokeeCo	89	* *	77	* *	27	* *	75	* *	77	* *
Cheyenne	20	* *	17	* *	4	* *	16	* *	17	* *
Clark	4	* *	4	* *	0	* *	4	* *	4	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 7A**  
**2011 Pregnancy Nutrition Surveillance** (1)  
**Kansas**  
**Comparison of Smoking Indicators by Clinic**

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Clay	50	* *	44	* *	21	* *	44	* *	44	* *
Cloud	60	* *	56	* *	15	* *	48	* *	56	* *
Coffey	26	* *	21	* *	9	* *	23	* *	21	* *
Comanche	9	* *	9	* *	2	* *	8	* *	9	* *
Cowley-Ark	183	38.3 (19)	126	19.0 (14)	40	* *	143	21.0 (25)	123	21.1 (25)
Cowley-Win	168	38.7 (20)	131	32.8 (32)	45	* *	141	17.7 (23)	116	16.4 (24)
Crawford-G	1	* *	1	* *	0	* *	1	* *	1	* *
Crawford-P	350	44.6 (32)	268	25.7 (26)	91	* *	296	41.2 (34)	268	40.3 (33)
Decatur	14	* *	11	* *	1	* *	13	* *	11	* *
DickinsonA	102	51.0 (36)	88	* *	34	* *	86	* *	88	* *
DickinsonH	19	* *	15	* *	6	* *	17	* *	15	* *
Doniphan	35	* *	26	* *	4	* *	26	* *	26	* *
Douglas	511	33.5 (15)	396	17.4 (12)	106	46.2 (4)	422	11.1 (17)	414	8.7 (17)
Edwards	21	* *	17	* *	4	* *	17	* *	18	* *
Elk	12	* *	11	* *	6	* *	9	* *	11	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 7A**  
**2011 Pregnancy Nutrition Surveillance** (1)  
**Kansas**  
**Comparison of Smoking Indicators by Clinic**

<b>Comparison of Smoking Indicators</b> (2, 3)										
<b>Clinic</b>	<b>Smoking</b>				<b>Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)</b>		<b>Smoking in Household</b>			
	<b>3 Months Prior to Pregnancy</b>		<b>Last 3 Months of Pregnancy</b>				<b>Prenatal</b>		<b>Postpartum</b>	
	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>
Ellis	152	42.1 (28)	127	22.8 (18)	44	* *	129	8.5 (10)	127	7.1 (14)
Ellsworth	33	* *	28	* *	13	* *	33	* *	28	* *
Finney	680	10.4 (3)	581	4.5 (2)	44	* *	553	10.1 (14)	580	10.5 (20)
Ford	646	7.3 (2)	529	5.1 (3)	27	* *	565	7.4 (7)	529	5.5 (11)
Franklin	188	41.5 (26)	159	28.9 (29)	50	* *	150	10.0 (13)	161	6.8 (13)
Geary	399	40.6 (24)	307	19.5 (15)	95	* *	332	30.4 (29)	314	22.3 (26)
Gove	12	* *	10	* *	1	* *	8	* *	10	* *
Graham	20	* *	18	* *	10	* *	18	* *	18	* *
Grant	72	* *	65	* *	10	* *	61	* *	65	* *
Gray	54	* *	44	* *	4	* *	44	* *	44	* *
Greeley	13	* *	11	* *	1	* *	11	* *	11	* *
Greenwood	33	* *	30	* *	10	* *	28	* *	31	* *
Hamilton	18	* *	15	* *	1	* *	15	* *	15	* *
Harper	59	* *	50	* *	18	* *	46	* *	50	* *
Harvey	249	37.8 (18)	215	22.8 (18)	64	* *	210	2.9 (1)	214	1.9 (2)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.





**Table 7A**  
**2011 Pregnancy Nutrition Surveillance** (1)  
**Kansas**  
**Comparison of Smoking Indicators by Clinic**

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Haskell	22	* *	19	* *	2	* *	18	* *	19	* *
Hodgeman	12	* *	12	* *	4	* *	11	* *	12	* *
JacksonHol	69	* *	59	* *	22	* *	53	* *	59	* *
JacksonPBP	7	* *	3	* *	0	* *	5	* *	4	* *
Jefferson	49	* *	28	* *	11	* *	43	* *	29	* *
Jewell	5	* *	2	* *	0	* *	4	* *	3	* *
Johnson-M	798	25.4 (6)	694	9.1 (4)	131	54.2 (1)	638	22.1 (26)	694	23.5 (27)
Johnson-O	1,085	21.7 (5)	960	9.2 (5)	154	48.1 (3)	899	14.5 (22)	961	14.4 (22)
Kearny	16	* *	13	* *	3	* *	15	* *	13	* *
Kingman	46	* *	32	* *	8	* *	38	* *	33	* *
Kiowa	15	* *	12	* *	3	* *	11	* *	14	* *
Labette	154	41.6 (27)	135	31.1 (30)	49	* *	126	34.9 (32)	136	36.8 (32)
Lane	10	* *	1	* *	0	* *	9	* *	2	* *
Leavenwort	356	44.4 (31)	284	24.3 (22)	94	* *	287	34.8 (31)	286	35.3 (30)
Lincoln	22	* *	18	* *	12	* *	22	* *	18	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 7A**  
**2011 Pregnancy Nutrition Surveillance** (1)  
**Kansas**  
**Comparison of Smoking Indicators by Clinic**

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Linn	68	* *	59	* *	23	* *	63	* *	60	* *
Logan	11	* *	10	* *	2	* *	11	* *	10	* *
Lyon	260	30.0 (12)	224	12.9 (9)	56	* *	224	8.9 (11)	208	2.4 (3)
Marion	53	* *	43	* *	10	* *	48	* *	43	* *
Marshall	60	* *	53	* *	20	* *	50	* *	53	* *
Mc Pherson	140	32.9 (13)	125	21.6 (17)	31	* *	106	5.7 (3)	125	1.6 (1)
Meade	34	* *	32	* *	11	* *	32	* *	34	* *
Miami	157	46.5 (33)	143	32.2 (31)	48	* *	115	20.9 (24)	143	16.1 (23)
Mitchell	42	* *	37	* *	8	* *	35	* *	37	* *
Montgome-C	155	38.7 (20)	137	25.5 (25)	43	* *	138	29.0 (28)	136	32.4 (28)
Montgome-I	157	36.3 (17)	143	28.0 (28)	43	* *	130	33.1 (30)	142	34.5 (29)
Morris	22	* *	21	* *	4	* *	16	* *	22	* *
Nemaha	11	* *	10	* *	4	* *	7	* *	10	* *
Neosho-Ch	138	43.5 (29)	98	* *	34	* *	115	27.0 (27)	98	* *
Neosho-Er	7	* *	5	* *	0	* *	5	* *	5	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 7A**  
**2011 Pregnancy Nutrition Surveillance** (1)  
**Kansas**  
**Comparison of Smoking Indicators by Clinic**

<b>Comparison of Smoking Indicators</b> (2, 3)										
<b>Clinic</b>	<b>Smoking</b>				<b>Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)</b>		<b>Smoking in Household</b>			
	<b>3 Months Prior to Pregnancy</b>		<b>Last 3 Months of Pregnancy</b>				<b>Prenatal</b>		<b>Postpartum</b>	
	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>
Ness	8	* *	7	* *	3	* *	6	* *	7	* *
Norton	48	* *	34	* *	10	* *	40	* *	36	* *
Osage	97	* *	86	* *	32	* *	80	* *	87	* *
Osborne	16	* *	9	* *	2	* *	15	* *	9	* *
Ottawa	13	* *	10	* *	2	* *	10	* *	10	* *
Pawnee	49	* *	40	* *	11	* *	41	* *	40	* *
Phillips	30	* *	21	* *	7	* *	25	* *	22	* *
Pottawat-S	12	* *	9	* *	0	* *	8	* *	9	* *
Pottawat-W	9	* *	8	* *	4	* *	9	* *	8	* *
Pratt	96	* *	85	* *	31	* *	76	* *	80	* *
Rawlins	7	* *	5	* *	1	* *	7	* *	5	* *
Reno	453	48.1 (34)	391	26.9 (27)	140	37.9 (6)	359	12.5 (19)	393	8.9 (19)
Republic	28	* *	24	* *	10	* *	26	* *	25	* *
Rice	72	* *	64	* *	28	* *	62	* *	63	* *
Riley-Ft	589	29.2 (11)	479	11.7 (6)	119	52.9 (2)	475	8.2 (8)	481	5.2 (9)

(1) Reporting period is January 1 through December 31.  
(2) Excludes records with unknown data or errors.  
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.  
(4) Among women who smoked 3 months prior to pregnancy.  
\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 7A**  
**2011 Pregnancy Nutrition Surveillance** (1)  
**Kansas**  
**Comparison of Smoking Indicators by Clinic**

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Riley-Manh	402	26.4 (7)	335	16.4 (11)	72	* *	338	5.3 (2)	339	2.9 (6)
Rooks	46	* *	36	* *	13	* *	34	* *	37	* *
Rush	11	* *	8	* *	4	* *	8	* *	8	* *
Russell	46	* *	43	* *	14	* *	40	* *	42	* *
Saline	476	40.1 (23)	429	24.5 (23)	135	25.9 (11)	401	6.7 (5)	431	4.6 (8)
Scott	40	* *	32	* *	9	* *	36	* *	32	* *
Sedgwick-C	1,215	27.5 (9)	1,101	16.3 (10)	238	33.6 (7)	1,023	11.3 (18)	1,103	8.3 (16)
Sedgwick-M	1,869	27.1 (8)	1,677	12.1 (8)	344	43.9 (5)	1,510	6.6 (4)	1,683	3.9 (7)
Sedgwick-S	1,416	33.5 (15)	1,277	23.1 (21)	323	23.2 (12)	1,137	9.2 (12)	1,277	6.1 (12)
Seneca	6	* *	1	* *	0	* *	4	* *	3	* *
Seward	442	4.3 (1)	398	1.8 (1)	12	* *	376	10.4 (16)	402	8.7 (17)
ShawneeM&I	281	28.5 (10)	225	17.8 (13)	46	* *	239	8.4 (9)	226	2.7 (5)
ShawneeMa	1,089	33.3 (14)	791	19.5 (15)	203	26.6 (10)	924	10.3 (15)	775	5.4 (10)
Sheridan	7	* *	5	* *	1	* *	4	* *	7	* *
Sherman	56	* *	50	* *	10	* *	43	* *	50	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 7A**  
**2011 Pregnancy Nutrition Surveillance** (1)  
**Kansas**  
**Comparison of Smoking Indicators by Clinic**

<b>Comparison of Smoking Indicators</b> (2, 3)										
<b>Clinic</b>	<b>Smoking</b>				<b>Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)</b>		<b>Smoking in Household</b>			
	<b>3 Months Prior to Pregnancy</b>		<b>Last 3 Months of Pregnancy</b>				<b>Prenatal</b>		<b>Postpartum</b>	
	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>
Smith	18	* *	13	* *	4	* *	16	* *	13	* *
Stafford	23	* *	17	* *	3	* *	20	* *	16	* *
Stanton	11	* *	10	* *	1	* *	9	* *	9	* *
Stevens	71	* *	62	* *	15	* *	56	* *	63	* *
Sumner	135	49.6 (35)	121	24.8 (24)	49	* *	111	37.8 (33)	120	35.8 (31)
Thomas	55	* *	45	* *	12	* *	46	* *	46	* *
Trego	12	* *	10	* *	4	* *	10	* *	10	* *
Wabaunsee	19	* *	18	* *	3	* *	13	* *	18	* *
Washington	26	* *	18	* *	6	* *	24	* *	19	* *
Wichita	8	* *	6	* *	1	* *	7	* *	6	* *
Wilson	83	* *	81	* *	30	* *	72	* *	80	* *
Woodson	14	* *	14	* *	5	* *	13	* *	14	* *
Wyandotte	2,062	19.4 (4)	1,800	11.9 (7)	271	27.7 (9)	1,760	12.5 (19)	1,845	11.2 (21)
Unknown Clinics	25	* *	23	* *	8	* *	22	* *	22	* *
<b>Kansas</b>	<b>20,850</b>	<b>30.4</b>	<b>17,728</b>	<b>17.5</b>	<b>4,187</b>	<b>32.7</b>	<b>17,287</b>	<b>15.1</b>	<b>17,791</b>	<b>12.6</b>

(1) Reporting period is January 1 through December 31.  
(2) Excludes records with unknown data or errors.  
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.  
(4) Among women who smoked 3 months prior to pregnancy.  
\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 7A**  
**2011 Pregnancy Nutrition Surveillance** (1)  
**Kansas**  
**Comparison of Smoking Indicators by Clinic**

<b>Comparison of Smoking Indicators</b> (2, 3)										
<b>Clinic</b>	<b>Smoking</b>				<b>Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)</b>	<b>Smoking in Household</b>				
	<b>3 Months Prior to Pregnancy</b>		<b>Last 3 Months of Pregnancy</b>			<b>Prenatal</b>	<b>Postpartum</b>			
	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>		
<b>Nation (Prior Year)</b>	<b>814,631</b>	<b>23.0</b>	<b>715,034</b>	<b>12.1</b>	<b>137,883</b>	<b>38.1</b>	<b>660,570</b>	<b>16.3</b>	<b>562,126</b>	<b>15.6</b>

(1) Reporting period is January 1 through December 31.  
(2) Excludes records with unknown data or errors.  
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.  
(4) Among women who smoked 3 months prior to pregnancy.  
\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 8A**  
**2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Infant Health Indicators by Clinic**

<b>Comparison of Infant Health Indicators</b> <sup>(2, 3)</sup>										
<b>Clinic</b>	<b>Birthweight</b>				<b>Preterm</b> (6)		<b>Full Term LBW</b> (7)		<b>Ever Breastfed</b>	
	<b>Low</b> (4)		<b>High</b> (5)		<b>Number</b>	<b>%</b> (Rank)	<b>Number</b>	<b>%</b> (Rank)	<b>Number</b>	<b>%</b> (Rank)
	<b>Number</b>	<b>%</b> (Rank)	<b>%</b> (Rank)	<b>Number</b>						
Allen	59	* * * *	* * * *	65	* *	58	* *	65	* *	
Anderson	36	* * * *	* * * *	38	* *	33	* *	38	* *	
Atchison	114	13.2 (35)	1.8 (1)	125	10.4 (30)	102	8.8 (35)	125	52.0 (35)	
Barber-Kio	0	* * * *	* * * *	0	* *	0	* *	0	* *	
Barber-Med	27	* * * *	* * * *	28	* *	25	* *	27	* *	
Barton	220	5.0 (4)	5.0 (9)	220	6.4 (7)	204	3.4 (9)	221	74.2 (19)	
Bourbon	99	* * * *	* * * *	99	* *	87	* *	99	* *	
Brown	69	* * * *	* * * *	75	* *	60	* *	75	* *	
Butler	220	7.7 (22)	5.0 (9)	223	7.6 (15)	202	5.4 (29)	223	76.7 (12)	
Chase	15	* * * *	* * * *	15	* *	15	* *	15	* *	
Chautauqua	25	* * * *	* * * *	25	* *	23	* *	26	* *	
CherokeeBS	68	* * * *	* * * *	68	* *	62	* *	68	* *	
CherokeeCo	78	* * * *	* * * *	79	* *	68	* *	79	* *	
Cheyenne	15	* * * *	* * * *	15	* *	12	* *	15	* *	

(1) Reporting period is January 1 through December 31.  
(2) Excludes records with unknown data or errors.  
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.  
(4) <2500 g.  
(5) >4000 g.  
(6) <37 weeks gestation.  
(7) 37 or more weeks gestation and <2500 g.  
\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 8A**  
**2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Infant Health Indicators by Clinic**

Comparison of Infant Health Indicators <sup>(2, 3)</sup>										
Clinic	Birthweight				Preterm <sup>(6)</sup>		Full Term LBW <sup>(7)</sup>		Ever Breastfed	
	Low <sup>(4)</sup>		High <sup>(5)</sup>		Number	% <sup>(Rank)</sup>	Number	% <sup>(Rank)</sup>	Number	% <sup>(Rank)</sup>
	Number	% <sup>(Rank)</sup>	% <sup>(Rank)</sup>							
Clark	4	* * * *			4	* *	4	* *	4	* *
Clay	41	* * * *			42	* *	36	* *	42	* *
Cloud	52	* * * *			53	* *	45	* *	53	* *
Coffey	21	* * * *			21	* *	21	* *	21	* *
Comanche	7	* * * *			8	* *	6	* *	8	* *
Cowley-Ark	151	6.0 (6)	5.3 (14)		157	10.8 (33)	134	3.7 (12)	155	58.1 (33)
Cowley-Win	132	7.6 (18)	6.1 (19)		146	6.8 (10)	119	5.9 (32)	148	70.3 (24)
Crawford-G	1	* * * *			1	* *	1	* *	1	* *
Crawford-P	301	7.6 (18)	5.0 (9)		312	9.6 (24)	268	3.0 (5)	315	66.3 (28)
Decatur	14	* * * *			14	* *	13	* *	14	* *
DickinsonA	83	* * * *			81	* *	74	* *	83	* *
DickinsonH	17	* * * *			17	* *	16	* *	17	* *
Doniphan	28	* * * *			28	* *	23	* *	27	* *
Douglas	411	5.6 (5)	6.3 (21)		422	7.1 (11)	379	3.2 (7)	423	80.4 (6)

(1) Reporting period is January 1 through December 31.  
(2) Excludes records with unknown data or errors.  
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.  
(4) <2500 g.  
(5) >4000 g.  
(6) <37 weeks gestation.  
(7) 37 or more weeks gestation and <2500 g.  
\* Percentages are not calculated if <100 records are available for analysis after exclusions.





**Table 8A**  
**2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Infant Health Indicators by Clinic**

<b>Comparison of Infant Health Indicators</b> <sup>(2, 3)</sup>										
<b>Clinic</b>	<b>Birthweight</b>				<b>Preterm</b> (6)		<b>Full Term LBW</b> (7)		<b>Ever Breastfed</b>	
	<b>Low</b> (4)		<b>High</b> (5)		<b>Number</b>	<b>%</b> (Rank)	<b>Number</b>	<b>%</b> (Rank)	<b>Number</b>	<b>%</b> (Rank)
	<b>Number</b>	<b>%</b> (Rank)	<b>%</b> (Rank)	<b>Number</b>						
Edwards	18	* * * *	* * * *	19	* *	13	* *	17	* *	
Elk	10	* * * *	* * * *	11	* *	9	* *	11	* *	
Ellis	121	7.4 (16)	7.4 (27)	127	3.1 (1)	116	5.2 (28)	128	81.3 (5)	
Ellsworth	31	* * * *	* * * *	31	* *	30	* *	31	* *	
Finney	593	6.6 (8)	5.1 (13)	600	10.2 (28)	533	3.6 (11)	596	75.3 (17)	
Ford	549	7.3 (15)	4.9 (8)	558	8.4 (18)	496	4.2 (18)	557	77.6 (10)	
Franklin	156	7.1 (11)	4.5 (5)	158	12.7 (34)	135	3.0 (5)	159	64.2 (30)	
Geary	314	8.0 (27)	9.6 (35)	316	6.0 (4)	294	5.4 (29)	315	75.6 (14)	
Gove	12	* * * *	* * * *	12	* *	12	* *	12	* *	
Graham	16	* * * *	* * * *	17	* *	13	* *	17	* *	
Grant	65	* * * *	* * * *	63	* *	60	* *	64	* *	
Gray	41	* * * *	* * * *	40	* *	37	* *	41	* *	
Greeley	12	* * * *	* * * *	12	* *	11	* *	12	* *	
Greenwood	31	* * * *	* * * *	32	* *	30	* *	32	* *	

(1) Reporting period is January 1 through December 31.  
(2) Excludes records with unknown data or errors.  
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.  
(4) <2500 g.  
(5) >4000 g.  
(6) <37 weeks gestation.  
(7) 37 or more weeks gestation and <2500 g.  
\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 8A**  
**2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Infant Health Indicators by Clinic**

<b>Comparison of Infant Health Indicators</b> <sup>(2, 3)</sup>										
<b>Clinic</b>	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)							
	Number	% (Rank)	%	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Hamilton	17	* * * *	* *	* *	17	* *	16	* *	17	* *
Harper	50	* * * *	* *	* *	50	* *	45	* *	49	* *
Harvey	217	7.8 (25)	7.8 (31)		219	10.0 (27)	195	5.1 (27)	219	88.1 (1)
Haskell	17	* * * *	* *	* *	17	* *	15	* *	17	* *
Hodgeman	11	* * * *	* *	* *	11	* *	10	* *	11	* *
JacksonHol	56	* * * *	* *	* *	59	* *	52	* *	59	* *
JacksonPBP	4	* * * *	* *	* *	4	* *	4	* *	4	* *
Jefferson	49	* * * *	* *	* *	49	* *	40	* *	49	* *
Jewell	6	* * * *	* *	* *	6	* *	5	* *	6	* *
Johnson-M	649	7.6 (18)	8.5 (34)		696	6.3 (5)	601	3.8 (13)	695	79.4 (8)
Johnson-O	917	6.1 (7)	6.2 (20)		934	6.3 (5)	858	4.0 (17)	932	85.2 (2)
Kearny	14	* * * *	* *	* *	14	* *	10	* *	14	* *
Kingman	24	* * * *	* *	* *	32	* *	22	* *	32	* *
Kiowa	14	* * * *	* *	* *	14	* *	12	* *	13	* *

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
- (4) <2500 g.
- (5) >4000 g.
- (6) <37 weeks gestation.
- (7) 37 or more weeks gestation and <2500 g.
- \* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 8A**  
**2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Infant Health Indicators by Clinic**

Comparison of Infant Health Indicators <sup>(2, 3)</sup>										
Clinic	Birthweight				Preterm <sup>(6)</sup>		Full Term LBW <sup>(7)</sup>		Ever Breastfed	
	Low <sup>(4)</sup>		High <sup>(5)</sup>		Number	%	Number	%	Number	%
	Number	% (Rank)	% (Rank)	(Rank)						
Labette	139	7.2 (13)	5.8 (16)		139	12.9 (35)	121	5.0 (24)	138	55.1 (34)
Lane	10	*	*	*	10	*	9	*	10	*
Leavenworth	281	10.0 (34)	5.3 (14)		304	7.9 (16)	257	6.6 (33)	304	66.1 (29)
Lincoln	17	*	*	*	18	*	14	*	18	*
Linn	61	*	*	*	61	*	51	*	61	*
Logan	10	*	*	*	10	*	9	*	10	*
Lyon	229	7.9 (26)	7.4 (27)		229	9.6 (24)	207	2.9 (4)	228	79.4 (8)
Marion	44	*	*	*	44	*	39	*	44	*
Marshall	52	*	*	*	52	*	47	*	52	*
Mc Pherson	118	3.4 (1)	2.5 (3)		120	4.2 (2)	113	1.8 (2)	120	77.5 (11)
Meade	32	*	*	*	34	*	27	*	34	*
Miami	147	6.8 (9)	7.5 (30)		148	9.5 (23)	133	3.8 (13)	147	66.7 (27)
Mitchell	34	*	*	*	37	*	34	*	37	*
Montgome-C	140	9.3 (33)	3.6 (4)		141	7.1 (11)	130	5.4 (29)	140	60.7 (31)

(1) Reporting period is January 1 through December 31.  
(2) Excludes records with unknown data or errors.  
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.  
(4) <2500 g.  
(5) >4000 g.  
(6) <37 weeks gestation.  
(7) 37 or more weeks gestation and <2500 g.  
\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 8A**  
**2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Infant Health Indicators by Clinic**

Comparison of Infant Health Indicators <sup>(2, 3)</sup>									
Clinic	Birthweight			Preterm <sup>(6)</sup>		Full Term LBW <sup>(7)</sup>		Ever Breastfed	
	Low <sup>(4)</sup>	High <sup>(5)</sup>		Number	%	Number	%	Number	%
	Number	% <sup>(Rank)</sup>	% <sup>(Rank)</sup>						
Montgome-I	139	8.6 (29)	2.2 (2)	139	7.2 (13)	127	7.1 (34)	141	60.3 (32)
Morris	21	* *	* *	21	* *	15	* *	22	* *
Nemaha	23	* *	* *	26	* *	22	* *	26	* *
Neosho-Ch	124	4.0 (2)	6.5 (22)	133	5.3 (3)	116	1.7 (1)	133	72.9 (20)
Neosho-Er	6	* *	* *	6	* *	5	* *	6	* *
Ness	6	* *	* *	6	* *	4	* *	7	* *
Norton	39	* *	* *	39	* *	35	* *	39	* *
Osage	86	* *	* *	91	* *	79	* *	91	* *
Osborne	14	* *	* *	14	* *	14	* *	14	* *
Ottawa	10	* *	* *	12	* *	9	* *	12	* *
Pawnee	41	* *	* *	43	* *	35	* *	42	* *
Phillips	24	* *	* *	23	* *	22	* *	24	* *
Pottawat-S	7	* *	* *	7	* *	7	* *	7	* *
Pottawat-W	8	* *	* *	8	* *	7	* *	8	* *

(1) Reporting period is January 1 through December 31.  
(2) Excludes records with unknown data or errors.  
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.  
(4) <2500 g.  
(5) >4000 g.  
(6) <37 weeks gestation.  
(7) 37 or more weeks gestation and <2500 g.  
\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 8A**  
**2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Infant Health Indicators by Clinic**

<b>Comparison of Infant Health Indicators</b> <sup>(2, 3)</sup>										
<b>Clinic</b>	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)							
	Number	% (Rank)	%	(Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Pratt	87	* * * *	* *	* *	87	* *	81	* *	87	* *
Rawlins	6	* * * *	* *	* *	6	* *	6	* *	6	* *
Reno	385	6.8 (9)	6.8 (25)		386	7.3 (14)	354	4.2 (18)	387	72.6 (21)
Republic	23	* * * *	* *	* *	23	* *	19	* *	24	* *
Rice	63	* * * *	* *	* *	62	* *	58	* *	63	* *
Riley-Ft	485	7.2 (13)	8.2 (32)		489	6.5 (8)	454	4.4 (21)	484	81.4 (4)
Riley-Manh	340	7.1 (11)	4.7 (7)		341	6.5 (8)	317	5.0 (24)	341	83.9 (3)
Rooks	45	* * * *	* *	* *	46	* *	40	* *	45	* *
Rush	7	* * * *	* *	* *	7	* *	7	* *	7	* *
Russell	42	* * * *	* *	* *	41	* *	33	* *	42	* *
Saline	400	8.0 (27)	4.5 (5)		430	10.7 (31)	355	3.4 (9)	429	70.4 (23)
Scott	45	* * * *	* *	* *	46	* *	39	* *	46	* *
Sedgwick-C	1,078	8.8 (30)	6.5 (22)		1,080	9.6 (24)	972	3.8 (13)	1,080	75.0 (18)
Sedgwick-M	1,666	9.1 (32)	5.9 (18)		1,675	8.2 (17)	1,528	5.0 (24)	1,670	75.5 (15)

(1) Reporting period is January 1 through December 31.  
(2) Excludes records with unknown data or errors.  
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.  
(4) <2500 g.  
(5) >4000 g.  
(6) <37 weeks gestation.  
(7) 37 or more weeks gestation and <2500 g.  
\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 8A**  
**2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Infant Health Indicators by Clinic**

Comparison of Infant Health Indicators <sup>(2, 3)</sup>									
Clinic	Birthweight			Preterm <sup>(6)</sup>		Full Term LBW <sup>(7)</sup>		Ever Breastfed	
	Low <sup>(4)</sup>	High <sup>(5)</sup>		Number	%	Number	%	Number	%
	Number	% (Rank)	% (Rank)						
Sedgwick-S	1,254	7.7 (22)	7.2 (26)	1,255	8.5 (19)	1,145	4.5 (22)	1,254	68.7 (25)
Seneca	9	* * *	* *	10	* *	9	* *	10	* *
Seward	368	4.1 (3)	8.4 (33)	381	9.2 (22)	333	2.1 (3)	382	80.1 (7)
ShawneeM&I	243	7.4 (16)	7.4 (27)	243	10.3 (29)	218	3.2 (7)	243	76.1 (13)
ShawneeMa	1,000	8.8 (30)	5.0 (9)	1,009	9.1 (20)	905	4.9 (23)	1,006	68.4 (26)
Sheridan	8	* * *	* *	8	* *	8	* *	8	* *
Sherman	47	* * *	* *	47	* *	43	* *	47	* *
Smith	17	* * *	* *	17	* *	16	* *	17	* *
Stafford	16	* * *	* *	16	* *	13	* *	16	* *
Stanton	10	* * *	* *	10	* *	10	* *	10	* *
Stevens	36	* * *	* *	62	* *	32	* *	63	* *
Sumner	119	7.6 (18)	6.7 (24)	122	10.7 (31)	106	3.8 (13)	122	75.4 (16)
Thomas	43	* * *	* *	43	* *	40	* *	43	* *
Trego	8	* * *	* *	10	* *	7	* *	10	* *

(1) Reporting period is January 1 through December 31.  
(2) Excludes records with unknown data or errors.  
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.  
(4) <2500 g.  
(5) >4000 g.  
(6) <37 weeks gestation.  
(7) 37 or more weeks gestation and <2500 g.  
\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 8A**  
**2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Infant Health Indicators by Clinic**

<b>Comparison of Infant Health Indicators</b> <sup>(2, 3)</sup>										
<b>Clinic</b>	<b>Birthweight</b>				<b>Preterm</b> (6)		<b>Full Term LBW</b> (7)		<b>Ever Breastfed</b>	
	<b>Low</b> (4)		<b>High</b> (5)		<b>Number</b>	<b>%</b> (Rank)	<b>Number</b>	<b>%</b> (Rank)	<b>Number</b>	<b>%</b> (Rank)
	<b>Number</b>	<b>%</b> (Rank)	<b>%</b> (Rank)	<b>Number</b>						
Wabaunsee	18	* * * *	* * * *	18	* *	17	* *	18	* *	
Washington	17	* * * *	* * * *	17	* *	17	* *	17	* *	
Wichita	8	* * * *	* * * *	8	* *	8	* *	8	* *	
Wilson	75	* * * *	* * * *	77	* *	68	* *	77	* *	
Woodson	14	* * * *	* * * *	14	* *	13	* *	14	* *	
Wyandotte	1,786	7.7 (22)	5.8 (16)	1,832	9.1 (20)	1,619	4.3 (20)	1,817	72.2 (22)	
Unknown Clinics	24	* * * *	* * * *	24	* *	24	* *	24	* *	
<b>Kansas</b>	<b>17,976</b>	<b>7.6</b>	<b>6.2</b>	<b>18,356</b>	<b>8.5</b>	<b>16,366</b>	<b>4.3</b>	<b>18,328</b>	<b>74.2</b>	
<b>Nation (Prior Year)</b>	<b>880,933</b>	<b>8.0</b>	<b>6.7</b>	<b>948,286</b>	<b>10.7</b>	<b>749,107</b>	<b>3.4</b>	<b>871,811</b>	<b>68.2</b>	

(1) Reporting period is January 1 through December 31.  
(2) Excludes records with unknown data or errors.  
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.  
(4) <2500 g.  
(5) >4000 g.  
(6) <37 weeks gestation.  
(7) 37 or more weeks gestation and <2500 g.  
\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 3A**  
**2009 - 2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Racial and Ethnic Distribution by Clinic**  
**Combined 3 Years**

<b>Comparison of Racial and Ethnic Distribution</b> <sup>(2)</sup>								
<b>Clinic</b>	<b>Number Included in Analysis</b>	<b>White, Not Hispanic</b>	<b>Black, Not Hispanic</b>	<b>Hispanic</b>	<b>American Indian/Alaska Native</b>	<b>Asian/Pacific Islander</b>	<b>Multiple Races</b>	<b>All Other/Unknown</b>
		<b>%</b>	<b>%</b>	<b>%</b>	<b>%</b>	<b>%</b>	<b>%</b>	<b>%</b>
Allen	246	91.9	2.8	1.2	1.6	0.0	2.4	0.0
Anderson	153	96.7	0.0	3.3	0.0	0.0	0.0	0.0
Atchison	443	86.9	9.5	1.1	0.9	0.0	1.6	0.0
Barber-Kio	3	*	*	*	*	*	*	*
Barber-Med	136	94.9	1.5	1.5	0.0	1.5	0.7	0.0
Barton	771	67.4	1.3	29.8	0.1	0.5	0.8	0.0
Bourbon	355	92.1	2.5	2.5	1.1	0.6	1.1	0.0
Brown	279	74.2	1.8	5.7	16.5	0.0	1.8	0.0
Butler	820	88.3	1.1	7.3	1.0	0.2	2.1	0.0
Chase	41	*	*	*	*	*	*	*
Chautauqua	83	*	*	*	*	*	*	*
CherokeeBS	75	*	*	*	*	*	*	*
CherokeeCo	488	89.3	1.6	2.7	1.8	3.1	1.4	0.0
Cheyenne	33	*	*	*	*	*	*	*
Clark	10	*	*	*	*	*	*	*
Clay	160	93.8	0.6	1.9	1.9	0.0	1.9	0.0
Cloud	242	92.6	1.7	2.1	2.1	0.8	0.8	0.0
Coffey	84	*	*	*	*	*	*	*
Comanche	31	*	*	*	*	*	*	*
Cowley-Ark	603	66.0	2.2	29.9	1.0	0.5	0.5	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.





**Table 3A**  
**2009 - 2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Racial and Ethnic Distribution by Clinic**  
**Combined 3 Years**

<b>Comparison of Racial and Ethnic Distribution</b> <sup>(2)</sup>								
<b>Clinic</b>	<b>Number Included in Analysis</b>	<b>White, Not Hispanic</b>	<b>Black, Not Hispanic</b>	<b>Hispanic</b>	<b>American Indian/Alaska Native</b>	<b>Asian/Pacific Islander</b>	<b>Multiple Races</b>	<b>All Other/Unknown</b>
		<b>%</b>	<b>%</b>	<b>%</b>	<b>%</b>	<b>%</b>	<b>%</b>	<b>%</b>
Cowley-Win	484	86.4	2.1	6.2	0.6	3.1	1.7	0.0
Crawford-G	4	*	*	*	*	*	*	*
Crawford-P	1,104	84.1	3.3	9.4	0.8	1.2	1.2	0.0
Decatur	40	*	*	*	*	*	*	*
DickinsonA	361	90.9	0.3	6.1	0.8	1.1	0.8	0.0
DickinsonH	50	*	*	*	*	*	*	*
Doniphan	130	90.8	2.3	4.6	0.8	0.0	1.5	0.0
Douglas	1,562	65.2	9.3	10.8	8.1	4.7	1.9	0.0
Edwards	71	*	*	*	*	*	*	*
Elk	56	*	*	*	*	*	*	*
Ellis	492	81.5	1.0	14.8	1.2	0.6	0.8	0.0
Ellsworth	104	90.4	0.0	6.7	1.0	0.0	1.9	0.0
Finney	2,105	28.5	1.1	65.9	0.0	4.1	0.3	0.0
Ford	2,057	21.0	2.1	75.9	0.2	0.7	0.1	0.0
Franklin	561	92.2	0.5	4.3	1.6	0.4	1.1	0.0
Geary	1,220	54.7	22.3	15.3	0.8	3.1	3.8	0.0
Gove	35	*	*	*	*	*	*	*
Graham	58	*	*	*	*	*	*	*
Grant	209	23.9	0.0	75.6	0.0	0.0	0.5	0.0
Gray	145	73.1	0.0	26.2	0.7	0.0	0.0	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 3A**  
**2009 - 2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Racial and Ethnic Distribution by Clinic**  
**Combined 3 Years**

<b>Comparison of Racial and Ethnic Distribution</b> <sup>(2)</sup>								
<b>Clinic</b>	<b>Number Included in Analysis</b>	<b>White, Not Hispanic</b>	<b>Black, Not Hispanic</b>	<b>Hispanic</b>	<b>American Indian/Alaska Native</b>	<b>Asian/Pacific Islander</b>	<b>Multiple Races</b>	<b>All Other/Unknown</b>
		%	%	%	%	%	%	%
Greeley	41	*	*	*	*	*	*	*
Greenwood	131	93.1	0.8	4.6	0.8	0.0	0.8	0.0
Hamilton	77	*	*	*	*	*	*	*
Harper	165	80.6	0.0	17.0	0.6	0.0	1.8	0.0
Harvey	818	71.5	2.9	21.9	1.3	0.9	1.5	0.0
Haskell	59	*	*	*	*	*	*	*
Hodgeman	39	*	*	*	*	*	*	*
JacksonHol	233	76.4	0.4	6.0	14.2	0.0	3.0	0.0
JacksonPBP	7	*	*	*	*	*	*	*
Jefferson	183	94.0	0.5	2.2	2.7	0.0	0.5	0.0
Jewell	22	*	*	*	*	*	*	*
Johnson-M	2,503	48.2	16.2	30.7	0.5	2.8	1.6	0.0
Johnson-O	3,301	46.0	11.6	37.3	0.4	3.4	1.2	0.0
Kearny	56	*	*	*	*	*	*	*
Kingman	155	92.3	0.0	5.8	0.6	1.3	0.0	0.0
Kiowa	39	*	*	*	*	*	*	*
Labette	530	85.7	7.2	3.2	2.1	0.2	1.7	0.0
Lane	40	*	*	*	*	*	*	*
Leavenwort	1,090	69.2	14.2	11.4	0.8	1.2	3.2	0.0
Lincoln	64	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 3A**  
**2009 - 2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Racial and Ethnic Distribution by Clinic**  
**Combined 3 Years**

<b>Comparison of Racial and Ethnic Distribution</b> <sup>(2)</sup>								
<b>Clinic</b>	<b>Number Included in Analysis</b>	<b>White, Not Hispanic</b>	<b>Black, Not Hispanic</b>	<b>Hispanic</b>	<b>American Indian/Alaska Native</b>	<b>Asian/Pacific Islander</b>	<b>Multiple Races</b>	<b>All Other/Unknown</b>
		%	%	%	%	%	%	%
Linn	206	95.6	0.0	2.4	1.5	0.0	0.5	0.0
Logan	38	*	*	*	*	*	*	*
Lyon	884	57.0	5.1	34.8	0.5	1.7	0.9	0.0
Marion	161	91.9	0.0	7.5	0.0	0.6	0.0	0.0
Marshall	160	89.4	0.6	3.1	1.9	0.0	5.0	0.0
Mc Pherson	409	85.8	1.7	10.3	0.5	0.7	1.0	0.0
Meade	118	65.3	0.8	33.1	0.0	0.0	0.8	0.0
Miami	455	91.2	2.4	4.0	0.2	0.7	1.5	0.0
Mitchell	133	90.2	0.0	7.5	0.0	1.5	0.8	0.0
Montgome-C	476	70.6	12.2	12.8	2.5	0.8	1.1	0.0
Montgome-I	533	81.1	6.4	9.0	2.1	0.8	0.8	0.0
Morris	85	*	*	*	*	*	*	*
Nemaha	93	*	*	*	*	*	*	*
Neosho-Ch	436	92.0	2.1	4.1	0.5	0.0	1.4	0.0
Neosho-Er	30	*	*	*	*	*	*	*
Ness	35	*	*	*	*	*	*	*
Norton	120	90.0	1.7	6.7	0.0	0.8	0.8	0.0
Osage	282	96.5	0.4	1.8	0.4	0.0	1.1	0.0
Osborne	57	*	*	*	*	*	*	*
Ottawa	71	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 3A**  
**2009 - 2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Racial and Ethnic Distribution by Clinic**  
**Combined 3 Years**

<b>Comparison of Racial and Ethnic Distribution</b> <sup>(2)</sup>								
<b>Clinic</b>	<b>Number Included in Analysis</b>	<b>White, Not Hispanic</b>	<b>Black, Not Hispanic</b>	<b>Hispanic</b>	<b>American Indian/Alaska Native</b>	<b>Asian/Pacific Islander</b>	<b>Multiple Races</b>	<b>All Other/Unknown</b>
		<b>%</b>	<b>%</b>	<b>%</b>	<b>%</b>	<b>%</b>	<b>%</b>	<b>%</b>
Pawnee	169	78.7	4.1	14.2	1.8	0.6	0.6	0.0
Phillips	109	93.6	0.0	3.7	0.0	1.8	0.9	0.0
Pottawat-S	48	*	*	*	*	*	*	*
Pottawat-W	47	*	*	*	*	*	*	*
Pratt	265	84.9	0.8	14.0	0.0	0.0	0.4	0.0
Rawlins	27	*	*	*	*	*	*	*
Reno	1,412	79.2	2.6	15.4	0.4	0.5	1.9	0.0
Republic	98	*	*	*	*	*	*	*
Rice	237	75.9	2.1	20.3	0.8	0.8	0.0	0.0
Riley-Ft	1,769	69.9	11.6	13.7	1.1	3.1	0.5	0.0
Riley-Manh	1,212	71.9	10.1	10.6	0.5	5.1	1.7	0.0
Rooks	150	94.7	1.3	2.0	1.3	0.7	0.0	0.0
Rush	30	*	*	*	*	*	*	*
Russell	133	89.5	4.5	5.3	0.8	0.0	0.0	0.0
Saline	1,563	70.9	4.7	20.3	0.2	1.7	2.2	0.0
Scott	135	61.5	0.0	38.5	0.0	0.0	0.0	0.0
Sedgwick-C	3,697	41.4	14.1	33.9	1.2	7.0	2.3	0.0
Sedgwick-M	5,863	32.9	26.7	34.3	1.4	2.8	1.8	0.0
Sedgwick-S	4,248	65.5	8.3	20.0	1.8	2.1	2.2	0.0
Seneca	30	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 3A**  
**2009 - 2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Racial and Ethnic Distribution by Clinic**  
**Combined 3 Years**

<b>Comparison of Racial and Ethnic Distribution</b> <sup>(2)</sup>								
<b>Clinic</b>	<b>Number Included in Analysis</b>	<b>White, Not Hispanic</b>	<b>Black, Not Hispanic</b>	<b>Hispanic</b>	<b>American Indian/Alaska Native</b>	<b>Asian/Pacific Islander</b>	<b>Multiple Races</b>	<b>All Other/Unknown</b>
		<b>%</b>	<b>%</b>	<b>%</b>	<b>%</b>	<b>%</b>	<b>%</b>	<b>%</b>
Seward	1,407	15.2	2.0	81.7	0.5	0.4	0.1	0.0
ShawneeM&I	913	35.9	14.5	45.0	1.3	1.5	1.8	0.0
ShawneeMa	3,560	59.4	16.3	18.9	2.4	0.7	2.2	0.0
Sheridan	26	*	*	*	*	*	*	*
Sherman	189	75.1	0.5	23.3	0.0	0.5	0.5	0.0
Smith	53	*	*	*	*	*	*	*
Stafford	100	65.0	1.0	33.0	0.0	1.0	0.0	0.0
Stanton	41	*	*	*	*	*	*	*
Stevens	231	46.8	0.4	51.5	1.3	0.0	0.0	0.0
Sumner	457	86.9	1.3	7.0	2.0	0.2	2.6	0.0
Thomas	166	83.7	1.2	13.3	0.6	1.2	0.0	0.0
Trego	44	*	*	*	*	*	*	*
Wabaunsee	61	*	*	*	*	*	*	*
Washington	75	*	*	*	*	*	*	*
Wichita	44	*	*	*	*	*	*	*
Wilson	260	94.2	1.5	1.5	0.4	0.0	2.3	0.0
Woodson	40	*	*	*	*	*	*	*
Wyandotte	6,487	26.4	28.8	39.3	0.5	4.2	0.8	0.0
Unknown Clinics	48	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 3A**  
**2009 - 2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Racial and Ethnic Distribution by Clinic**  
**Combined 3 Years**

<b>Comparison of Racial and Ethnic Distribution</b> <sup>(2)</sup>								
<b>Clinic</b>	Number Included in Analysis	White, Not Hispanic  %	Black, Not Hispanic  %	Hispanic  %	American Indian/Alaska Native  %	Asian/Pacific Islander  %	Multiple Races  %	All Other/Unknown  %
<b>Kansas</b>	<b>65,683</b>	<b>56.6</b>	<b>11.2</b>	<b>27.2</b>	<b>1.2</b>	<b>2.3</b>	<b>1.5</b>	<b>0.0</b>

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 4A**  
**2009 - 2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Age Distribution by Clinic**  
**Combined 3 Years**

<b>Comparison of Age Distribution</b> <sup>(2)</sup>								
<b>Clinic</b>	<b>Number Included in Analysis</b>	<b>&lt; 15 Years %</b>	<b>15 - 17 Years %</b>	<b>18 - 19 Years %</b>	<b>20 - 29 Years %</b>	<b>30 - 39 Years %</b>	<b>&gt;= 40 Years %</b>	<b>Unknown %</b>
Allen	246	0.4	8.5	16.3	63.4	10.6	0.8	0.0
Anderson	153	0.7	4.6	17.6	61.4	15.7	0.0	0.0
Atchison	443	0.0	7.7	15.6	61.9	14.7	0.2	0.0
Barber-Kio	3	*	*	*	*	*	*	*
Barber-Med	136	0.0	5.9	20.6	59.6	13.2	0.7	0.0
Barton	771	0.0	7.8	14.4	62.3	13.7	1.8	0.0
Bourbon	355	0.3	6.8	17.5	59.2	15.5	0.6	0.3
Brown	279	0.4	7.2	11.5	65.6	14.3	1.1	0.0
Butler	820	0.0	5.7	16.0	63.8	13.7	0.6	0.2
Chase	41	*	*	*	*	*	*	*
Chautauqua	83	*	*	*	*	*	*	*
CherokeeBS	75	*	*	*	*	*	*	*
CherokeeCo	488	0.4	8.4	19.7	59.8	10.9	0.6	0.2
Cheyenne	33	*	*	*	*	*	*	*
Clark	10	*	*	*	*	*	*	*
Clay	160	1.3	9.4	18.1	56.9	14.4	0.0	0.0
Cloud	242	0.4	6.6	15.3	59.5	15.7	2.5	0.0
Coffey	84	*	*	*	*	*	*	*
Comanche	31	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 4A**  
**2009 - 2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Age Distribution by Clinic**  
**Combined 3 Years**

<b>Comparison of Age Distribution</b> <sup>(2)</sup>								
<b>Clinic</b>	<b>Number Included in Analysis</b>	<b>&lt; 15 Years %</b>	<b>15 - 17 Years %</b>	<b>18 - 19 Years %</b>	<b>20 - 29 Years %</b>	<b>30 - 39 Years %</b>	<b>&gt;= 40 Years %</b>	<b>Unknown %</b>
Cowley-Ark	603	0.5	8.0	15.9	59.9	14.8	1.0	0.0
Cowley-Win	484	0.4	6.0	16.9	65.1	11.4	0.2	0.0
Crawford-G	4	*	*	*	*	*	*	*
Crawford-P	1,104	0.3	4.7	15.8	64.9	13.7	0.6	0.1
Decatur	40	*	*	*	*	*	*	*
DickinsonA	361	0.0	5.3	13.6	66.8	14.1	0.3	0.0
DickinsonH	50	*	*	*	*	*	*	*
Doniphan	130	0.8	2.3	11.5	66.2	18.5	0.0	0.8
Douglas	1,562	0.1	3.8	11.1	63.8	19.8	1.5	0.0
Edwards	71	*	*	*	*	*	*	*
Elk	56	*	*	*	*	*	*	*
Ellis	492	0.6	4.1	14.8	65.9	13.6	0.8	0.2
Ellsworth	104	0.0	2.9	11.5	57.7	24.0	3.8	0.0
Finney	2,105	0.5	7.1	15.5	55.7	19.3	1.9	0.0
Ford	2,057	0.2	6.4	12.9	55.5	22.5	2.4	0.0
Franklin	561	0.2	5.2	16.6	61.0	16.0	1.1	0.0
Geary	1,220	0.2	4.2	14.4	68.2	12.1	0.7	0.1
Gove	35	*	*	*	*	*	*	*
Graham	58	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.





**Table 4A**  
**2009 - 2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Age Distribution by Clinic**  
**Combined 3 Years**

<b>Comparison of Age Distribution</b> <sup>(2)</sup>								
<b>Clinic</b>	<b>Number Included in Analysis</b>	<b>&lt; 15 Years %</b>	<b>15 - 17 Years %</b>	<b>18 - 19 Years %</b>	<b>20 - 29 Years %</b>	<b>30 - 39 Years %</b>	<b>&gt;= 40 Years %</b>	<b>Unknown %</b>
Grant	209	0.0	5.3	12.9	62.7	17.7	1.4	0.0
Gray	145	0.0	4.1	12.4	60.7	21.4	1.4	0.0
Greeley	41	*	*	*	*	*	*	*
Greenwood	131	0.0	4.6	16.0	62.6	16.0	0.0	0.8
Hamilton	77	*	*	*	*	*	*	*
Harper	165	0.0	5.5	12.1	67.9	13.9	0.6	0.0
Harvey	818	0.4	6.2	15.3	59.5	17.6	1.0	0.0
Haskell	59	*	*	*	*	*	*	*
Hodgeman	39	*	*	*	*	*	*	*
JacksonHol	233	0.0	5.6	18.5	63.5	11.6	0.9	0.0
JacksonPBP	7	*	*	*	*	*	*	*
Jefferson	183	0.0	3.8	16.4	56.3	23.0	0.5	0.0
Jewell	22	*	*	*	*	*	*	*
Johnson-M	2,503	0.2	3.3	10.1	62.4	22.5	1.4	0.0
Johnson-O	3,301	0.1	3.2	10.1	59.7	24.7	2.1	0.1
Kearny	56	*	*	*	*	*	*	*
Kingman	155	0.0	6.5	16.8	64.5	10.3	1.9	0.0
Kiowa	39	*	*	*	*	*	*	*
Labette	530	0.4	8.1	17.4	61.3	12.1	0.8	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 4A**  
**2009 - 2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Age Distribution by Clinic**  
**Combined 3 Years**

<b>Comparison of Age Distribution</b> <sup>(2)</sup>								
<b>Clinic</b>	<b>Number Included in Analysis</b>	<b>&lt; 15 Years %</b>	<b>15 - 17 Years %</b>	<b>18 - 19 Years %</b>	<b>20 - 29 Years %</b>	<b>30 - 39 Years %</b>	<b>&gt;= 40 Years %</b>	<b>Unknown %</b>
Lane	40	*	*	*	*	*	*	*
Leavenwort	1,090	0.2	5.2	14.4	66.4	13.1	0.6	0.0
Lincoln	64	*	*	*	*	*	*	*
Linn	206	0.0	6.3	13.1	62.1	18.0	0.5	0.0
Logan	38	*	*	*	*	*	*	*
Lyon	884	0.7	5.5	13.6	64.5	14.4	1.4	0.0
Marion	161	0.0	1.2	16.1	59.6	19.9	3.1	0.0
Marshall	160	0.0	5.0	17.5	63.8	13.1	0.6	0.0
Mc Pherson	409	0.5	4.2	12.5	64.5	17.1	1.2	0.0
Meade	118	0.0	10.2	11.0	55.9	22.0	0.8	0.0
Miami	455	0.4	4.2	17.6	65.3	11.9	0.7	0.0
Mitchell	133	0.0	1.5	14.3	67.7	16.5	0.0	0.0
Montgome-C	476	0.6	4.8	16.0	63.4	14.1	0.8	0.2
Montgome-I	533	0.4	5.6	16.7	62.5	13.3	1.3	0.2
Morris	85	*	*	*	*	*	*	*
Nemaha	93	*	*	*	*	*	*	*
Neosho-Ch	436	0.0	6.7	16.7	64.0	11.5	0.7	0.5
Neosho-Er	30	*	*	*	*	*	*	*
Ness	35	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 4A**  
**2009 - 2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Age Distribution by Clinic**  
**Combined 3 Years**

<b>Comparison of Age Distribution</b> <sup>(2)</sup>								
<b>Clinic</b>	<b>Number Included in Analysis</b>	<b>&lt; 15 Years %</b>	<b>15 - 17 Years %</b>	<b>18 - 19 Years %</b>	<b>20 - 29 Years %</b>	<b>30 - 39 Years %</b>	<b>&gt;= 40 Years %</b>	<b>Unknown %</b>
Norton	120	0.0	1.7	15.8	57.5	24.2	0.8	0.0
Osage	282	0.0	3.9	18.4	61.0	14.9	1.4	0.4
Osborne	57	*	*	*	*	*	*	*
Ottawa	71	*	*	*	*	*	*	*
Pawnee	169	0.6	5.3	13.0	69.2	11.2	0.6	0.0
Phillips	109	0.9	1.8	14.7	66.1	15.6	0.9	0.0
Pottawat-S	48	*	*	*	*	*	*	*
Pottawat-W	47	*	*	*	*	*	*	*
Pratt	265	0.4	7.2	14.0	64.2	13.6	0.8	0.0
Rawlins	27	*	*	*	*	*	*	*
Reno	1,412	0.2	6.4	15.8	63.7	13.4	0.4	0.0
Republic	98	*	*	*	*	*	*	*
Rice	237	0.0	7.6	12.2	62.0	17.3	0.8	0.0
Riley-Ft	1,769	0.0	0.8	10.2	77.4	11.5	0.2	0.0
Riley-Manh	1,212	0.1	3.0	13.3	69.8	13.6	0.2	0.0
Rooks	150	0.0	4.7	14.0	64.0	15.3	1.3	0.7
Rush	30	*	*	*	*	*	*	*
Russell	133	0.0	7.5	15.0	60.2	15.8	1.5	0.0
Saline	1,563	0.3	6.6	15.0	62.3	14.7	1.2	0.1

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 4A**  
**2009 - 2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Age Distribution by Clinic**  
**Combined 3 Years**

Comparison of Age Distribution <sup>(2)</sup>								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Scott	135	0.0	3.7	13.3	64.4	18.5	0.0	0.0
Sedgwick-C	3,697	0.4	6.3	12.3	59.4	20.2	1.4	0.1
Sedgwick-M	5,863	0.3	6.3	13.3	60.6	18.1	1.3	0.1
Sedgwick-S	4,248	0.4	6.2	14.5	61.3	16.4	1.2	0.0
Seneca	30	*	*	*	*	*	*	*
Seward	1,407	0.1	8.2	12.7	55.4	22.0	1.4	0.1
ShawneeM&I	913	0.7	7.4	17.0	55.8	18.1	1.1	0.0
ShawneeMa	3,560	0.3	6.0	14.2	62.5	15.8	1.2	0.0
Sheridan	26	*	*	*	*	*	*	*
Sherman	189	0.0	3.2	18.5	61.4	15.3	1.6	0.0
Smith	53	*	*	*	*	*	*	*
Stafford	100	0.0	5.0	11.0	68.0	14.0	2.0	0.0
Stanton	41	*	*	*	*	*	*	*
Stevens	231	0.0	6.9	15.2	56.7	19.5	1.7	0.0
Sumner	457	0.0	6.1	15.5	61.5	15.5	1.3	0.0
Thomas	166	0.0	10.2	15.7	60.8	12.0	1.2	0.0
Trego	44	*	*	*	*	*	*	*
Wabaunsee	61	*	*	*	*	*	*	*
Washington	75	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 4A**  
**2009 - 2011 Pregnancy Nutrition Surveillance** (1)  
**Kansas**  
**Comparison of Age Distribution by Clinic**  
**Combined 3 Years**

<b>Comparison of Age Distribution</b> (2)								
<b>Clinic</b>	<b>Number Included in Analysis</b>	<b>&lt; 15 Years %</b>	<b>15 - 17 Years %</b>	<b>18 - 19 Years %</b>	<b>20 - 29 Years %</b>	<b>30 - 39 Years %</b>	<b>&gt;= 40 Years %</b>	<b>Unknown %</b>
Wichita	44	*	*	*	*	*	*	*
Wilson	260	0.4	7.7	14.2	65.8	11.5	0.4	0.0
Woodson	40	*	*	*	*	*	*	*
Wyandotte	6,487	0.4	7.8	13.3	57.1	19.8	1.6	0.0
Unknown Clinics	48	*	*	*	*	*	*	*
<b>Kansas</b>	<b>65,683</b>	<b>0.3</b>	<b>5.8</b>	<b>13.8</b>	<b>61.5</b>	<b>17.4</b>	<b>1.2</b>	<b>0.1</b>

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 5A**  
**2009 - 2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Maternal Health Indicators by Clinic**  
**Combined 3 Years**

<b>Comparison of Maternal Health Indicators</b> <sup>(2, 3)</sup>														
<b>Clinic</b>	<b>Prepregnancy BMI</b> (4)			<b>Weight Gain</b> (5)			<b>Anemia</b> (Low Hb/Hct) (6)				<b>Gestational Diabetes</b> During Pregnancy		<b>Hypertension</b> During Pregnancy	
	Under- weight	Over- weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
							Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)
Allen	241	6.2 (63)	58.1 (53)	214	19.2 (15)	58.4 (67)	37	* *	67	* *	221	5.4 (17)	223	15.2(63)
Anderson	149	6.0 (60)	49.7 (5)	132	25.0 (60)	48.5 (23)	17	* *	28	* *	137	5.1 (15)	137	11.7(38)
Atchison	427	6.6 (65)	52.2 (12)	372	18.5 (11)	55.4 (62)	57	* *	79	* *	359	5.8 (22)	359	11.7(38)
Barber-Kio	3	* *	* *	2	* *	* *	0	* *	1	* *	2	* *	2	* *
Barber-Med	134	3.7 (23)	52.2 (12)	107	31.8 (68)	48.6 (24)	16	* *	25	* *	112	0.0 (1)	112	14.3(59)
Barton	752	4.9 (46)	57.0 (50)	652	22.1 (38)	48.8 (27)	75	* *	133	21.1(14)	670	6.4 (26)	668	13.3(52)
Bourbon	349	2.3 (5)	58.7 (56)	308	24.0 (51)	51.3 (42)	42	* *	56	* *	310	9.7 (65)	311	14.1(56)
Brown	274	2.2 (3)	58.4 (54)	233	24.0 (51)	45.5 (12)	31	* *	56	* *	232	6.9 (38)	232	17.2(66)
Butler	791	5.9 (57)	53.6 (25)	711	21.2 (30)	51.9 (46)	118	44.1 (21)	137	30.7(29)	746	6.4 (26)	746	12.5(46)
Chase	40	* *	* *	35	* *	* *	4	* *	2	* *	37	* *	37	* *
Chautauqua	82	* *	* *	75	* *	* *	13	* *	12	* *	76	* *	76	* *
CherokeeBS	70	* *	* *	65	* *	* *	12	* *	12	* *	67	* *	67	* *
CherokeeCo	451	7.5 (69)	52.8 (18)	398	18.3 (9)	55.0 (60)	57	* *	93	* *	425	5.4 (17)	425	12.9(49)
Cheyenne	31	* *	* *	25	* *	* *	6	* *	10	* *	27	* *	27	* *
Clark	10	* *	* *	9	* *	* *	3	* *	1	* *	9	* *	9	* *
Clay	156	5.1 (50)	53.2 (20)	134	21.6 (32)	53.0 (54)	19	* *	39	* *	142	7.0 (39)	142	10.6(30)
Cloud	237	5.5 (54)	52.3 (14)	214	23.8 (50)	50.5 (36)	25	* *	62	* *	216	7.4 (46)	221	10.4(28)

(1) Reporting period is January 1 through December 31.  
(2) Excludes records with unknown data or errors.  
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.  
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).  
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).  
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.  
\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 5A**  
**2009 - 2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Maternal Health Indicators by Clinic**  
**Combined 3 Years**

Comparison of Maternal Health Indicators <sup>(2, 3)</sup>														
Clinic	Prepregnancy BMI <sup>(4)</sup>			Weight Gain <sup>(5)</sup>			Anemia (Low Hb/Hct) <sup>(6)</sup>				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Coffey	80	* *	* *	64	* *	* *	11	* *	18	* *	67	* *	67	* *
Comanche	31	* *	* *	29	* *	* *	5	* *	7	* *	29	* *	29	* *
Cowley-Ark	589	3.9 (28)	55.2 (34)	515	21.9 (36)	50.1 (33)	81	* *	83	* *	528	9.1 (59)	529	11.5(34)
Cowley-Win	476	3.6 (21)	56.1 (45)	411	21.7 (33)	51.1 (39)	52	* *	66	* *	419	4.1 (7)	420	11.0(31)
Crawford-G	4	* *	* *	3	* *	* *	1	* *	0	* *	3	* *	3	* *
Crawford-P	1,084	6.8 (67)	54.0 (29)	932	21.7 (33)	50.3 (35)	144	22.2 (4)	154	15.6 (3)	956	9.1 (59)	959	13.2(51)
Decatur	40	* *	* *	38	* *	* *	8	* *	9	* *	38	* *	38	* *
DickinsonA	349	3.2 (14)	60.5 (63)	299	23.4 (47)	47.8 (19)	54	* *	66	* *	312	6.4 (26)	308	11.7(38)
DickinsonH	49	* *	* *	43	* *	* *	5	* *	13	* *	45	* *	45	* *
Doniphan	125	0.0 (1)	59.2 (57)	101	16.8 (6)	58.4 (67)	20	* *	29	* *	108	6.5 (29)	109	8.3(14)
Douglas	1,520	3.3 (15)	49.8 (6)	1,276	18.8 (13)	52.7 (52)	199	34.2 (16)	391	27.4(23)	1,297	8.0 (50)	1,297	8.8(18)
Edwards	71	* *	* *	65	* *	* *	10	* *	11	* *	64	* *	64	* *
Elk	54	* *	* *	48	* *	* *	6	* *	10	* *	49	* *	49	* *
Ellis	471	4.2 (32)	51.0 (9)	400	18.3 (9)	46.5 (15)	94	* *	105	16.2 (5)	428	2.1 (3)	427	5.9(2)
Ellsworth	102	3.9 (28)	63.7 (68)	94	* *	* *	15	* *	9	* *	94	* *	94	* *
Finney	2,062	6.7 (66)	52.4 (15)	1,842	25.4 (61)	42.8 (3)	273	26.4 (8)	293	18.1 (7)	1,885	8.3 (53)	1,883	6.9(8)
Ford	2,033	2.4 (7)	58.6 (55)	1,788	25.4 (61)	45.8 (14)	241	19.1 (2)	222	15.3 (1)	1,790	6.8 (37)	1,786	9.0(19)

(1) Reporting period is January 1 through December 31.  
(2) Excludes records with unknown data or errors.  
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.  
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).  
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).  
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.  
\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 5A**  
**2009 - 2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Maternal Health Indicators by Clinic**  
**Combined 3 Years**

<b>Comparison of Maternal Health Indicators</b> <sup>(2, 3)</sup>														
<b>Clinic</b>	<b>Prepregnancy BMI</b> (4)			<b>Weight Gain</b> (5)			<b>Anemia</b> (Low Hb/Hct) (6)				<b>Gestational Diabetes</b> During Pregnancy		<b>Hypertension</b> During Pregnancy	
	<b>Under-weight</b>		<b>Over-weight</b>	<b>&lt; Ideal</b>		<b>&gt; Ideal</b>	<b>3rd Trimester</b>		<b>Postpartum</b>					
	<b>Number</b>	<b>% (Rank)</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>
Franklin	535	3.6 (21)	54.2 (30)	465	19.8 (22)	52.9 (53)	60	* *	148	15.5 (2)	404	5.7 (21)	404	11.6(36)
Geary	1,170	4.4 (38)	53.2 (20)	928	19.4 (17)	52.0 (47)	168	38.7 (20)	211	35.1 (30)	969	4.5 (12)	967	10.2(26)
Gove	29	* *	* *	24	* *	* *	4	* *	12	* *	30	* *	30	* *
Graham	57	* *	* *	48	* *	* *	3	* *	6	* *	49	* *	49	* *
Grant	205	3.4 (17)	65.4 (71)	189	27.5 (66)	43.4 (5)	27	* *	56	* *	195	6.7 (33)	193	6.7(4)
Gray	141	5.0 (47)	52.5 (16)	117	23.1 (44)	56.4 (65)	11	* *	30	* *	121	9.1 (59)	121	11.6(36)
Greeley	39	* *	* *	33	* *	* *	3	* *	6	* *	35	* *	35	* *
Greenwood	127	2.4 (7)	64.6 (70)	113	19.5 (20)	54.9 (59)	14	* *	18	* *	113	6.2 (24)	114	17.5(67)
Hamilton	74	* *	* *	68	* *	* *	16	* *	21	* *	69	* *	69	* *
Harper	159	3.1 (12)	56.0 (43)	137	15.3 (4)	49.6 (30)	16	* *	24	* *	139	6.5 (29)	139	9.4(23)
Harvey	799	3.9 (28)	56.6 (47)	703	22.8 (43)	47.9 (21)	109	37.6 (19)	119	20.2(12)	723	4.6 (13)	722	13.0(50)
Haskell	58	* *	* *	51	* *	* *	6	* *	10	* *	52	* *	52	* *
Hodgeman	37	* *	* *	35	* *	* *	6	* *	7	* *	36	* *	36	* *
JacksonHol	224	4.5 (39)	57.6 (51)	199	19.1 (14)	56.3 (64)	32	* *	50	* *	203	8.9 (58)	204	15.2(63)
JacksonPBP	7	* *	* *	4	* *	* *	0	* *	3	* *	5	* *	5	* *
Jefferson	179	3.4 (17)	55.3 (36)	159	18.2 (8)	54.7 (58)	30	* *	51	* *	163	6.7 (33)	162	12.3(44)
Jewell	22	* *	* *	21	* *	* *	3	* *	7	* *	21	* *	21	* *

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
- (4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
- (5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
- (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- \* Percentages are not calculated if <100 records are available for analysis after exclusions.





**Table 5A**  
**2009 - 2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Maternal Health Indicators by Clinic**  
**Combined 3 Years**

<b>Comparison of Maternal Health Indicators</b> <sup>(2, 3)</sup>														
<b>Clinic</b>	<b>Prepregnancy BMI</b> (4)			<b>Weight Gain</b> (5)			<b>Anemia</b> (Low Hb/Hct) (6)				<b>Gestational Diabetes</b> During Pregnancy		<b>Hypertension</b> During Pregnancy	
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
							Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)
Johnson-M	2,396	3.8 (24)	56.9 (49)	2,083	20.1 (25)	49.3 (28)	369	25.5 (7)	623	16.1 (4)	2,151	8.7 (56)	2,147	7.9(11)
Johnson-O	3,168	3.4 (17)	53.2 (20)	2,793	20.0 (24)	50.0 (32)	482	20.5 (3)	877	18.4 (8)	2,901	9.2 (62)	2,895	6.8(7)
Kearny	56	* *	* *	52	* *	* *	8	* *	8	* *	52	* *	52	* *
Kingman	154	5.2 (51)	49.4 (4)	127	26.0 (64)	47.2 (18)	13	* *	26	* *	130	5.4 (17)	130	14.6(60)
Kiowa	37	* *	* *	33	* *	* *	6	* *	12	* *	35	* *	35	* *
Labette	519	6.0 (60)	55.9 (42)	472	25.8 (63)	45.6 (13)	88	* *	114	23.7(18)	472	7.0 (39)	473	14.6(60)
Lane	38	* *	* *	32	* *	* *	3	* *	6	* *	34	* *	34	* *
Leavenwort	1,045	4.3 (34)	55.4 (37)	879	19.2 (15)	52.3 (50)	166	27.1 (9)	247	19.0 (9)	909	6.7 (33)	906	11.4(33)
Lincoln	61	* *	* *	53	* *	* *	11	* *	12	* *	55	* *	55	* *
Linn	206	6.3 (64)	60.7 (64)	185	24.3 (55)	48.1 (22)	25	* *	25	* *	185	7.6 (49)	186	7.0(9)
Logan	37	* *	* *	32	* *	* *	6	* *	5	* *	32	* *	32	* *
Lyon	858	2.7 (10)	59.9 (61)	746	24.3 (55)	45.4 (11)	105	15.2 (1)	157	19.7(11)	766	8.0 (50)	765	14.1(56)
Marion	156	5.8 (55)	50.6 (7)	126	19.8 (22)	46.8 (17)	21	* *	30	* *	137	1.5 (2)	138	6.5(3)
Marshall	155	5.2 (51)	55.5 (38)	141	27.7 (67)	43.3 (4)	19	* *	40	* *	146	9.6 (64)	145	13.8(53)
Mc Pherson	400	5.0 (47)	53.5 (24)	349	23.2 (45)	45.0 (8)	46	* *	99	* *	361	5.8 (22)	364	9.9(24)
Meade	117	1.7 (2)	63.2 (67)	107	18.7 (12)	48.6 (24)	10	* *	20	* *	106	11.3 (68)	105	10.5(29)
Miami	439	4.3 (34)	55.8 (41)	386	14.5 (1)	56.2 (63)	80	* *	113	26.5(22)	411	6.3 (25)	411	11.9(41)

(1) Reporting period is January 1 through December 31.  
(2) Excludes records with unknown data or errors.  
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.  
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).  
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).  
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.  
\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 5A**  
**2009 - 2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Maternal Health Indicators by Clinic**  
**Combined 3 Years**

Comparison of Maternal Health Indicators <sup>(2, 3)</sup>														
Clinic	Prepregnancy BMI <sup>(4)</sup>			Weight Gain <sup>(5)</sup>			Anemia (Low Hb/Hct) <sup>(6)</sup>				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Mitchell	129	3.1 (12)	62.8 (66)	116	35.3 (69)	44.0 (7)	13	* *	24	* *	119	10.1 (66)	119	17.6(68)
Montgome-C	464	4.5 (39)	55.2 (34)	419	22.0 (37)	48.7 (26)	111	33.3 (15)	114	26.3(20)	420	8.8 (57)	420	13.8(53)
Montgome-I	517	7.5 (69)	50.7 (8)	450	24.4 (57)	45.3 (10)	82	* *	114	22.8(17)	461	7.4 (46)	460	12.0(42)
Morris	81	* *	* *	75	* *	* *	8	* *	22	* *	77	* *	78	* *
Nemaha	86	* *	* *	77	* *	* *	16	* *	18	* *	84	* *	84	* *
Neosho-Ch	424	5.9 (57)	53.8 (27)	383	22.5 (40)	51.7 (45)	67	* *	64	* *	395	8.1 (52)	394	14.2(58)
Neosho-Er	27	* *	* *	24	* *	* *	4	* *	6	* *	27	* *	27	* *
Ness	33	* *	* *	28	* *	* *	3	* *	9	* *	29	* *	29	* *
Norton	118	3.4 (17)	48.3 (2)	106	22.6 (42)	43.4 (5)	15	* *	32	* *	108	4.6 (13)	108	14.8(62)
Osage	276	5.8 (55)	51.8 (10)	243	15.2 (3)	56.4 (65)	33	* *	59	* *	253	4.3 (9)	253	11.5(34)
Osborne	56	* *	* *	47	* *	* *	7	* *	12	* *	48	* *	48	* *
Ottawa	67	* *	* *	59	* *	* *	9	* *	9	* *	66	* *	66	* *
Pawnee	164	3.0 (11)	59.8 (59)	133	24.8 (59)	51.1 (39)	18	* *	29	* *	143	11.2 (67)	142	11.3(32)
Phillips	105	3.8 (24)	61.0 (65)	84	* *	* *	15	* *	21	* *	92	* *	92	* *
Pottawat-S	48	* *	* *	41	* *	* *	4	* *	21	* *	42	* *	42	* *
Pottawat-W	46	* *	* *	42	* *	* *	3	* *	14	* *	43	* *	43	* *
Pratt	258	5.4 (53)	56.6 (47)	224	21.4 (31)	51.3 (42)	35	* *	53	* *	232	3.0 (5)	232	8.2(12)

(1) Reporting period is January 1 through December 31.  
(2) Excludes records with unknown data or errors.  
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.  
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).  
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).  
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.  
\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 5A**  
**2009 - 2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Maternal Health Indicators by Clinic**  
**Combined 3 Years**

Comparison of Maternal Health Indicators <sup>(2, 3)</sup>														
Clinic	Prepregnancy BMI <sup>(4)</sup>			Weight Gain <sup>(5)</sup>			Anemia (Low Hb/Hct) <sup>(6)</sup>				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Rawlins	26	* *	* *	22	* *	* *	2	* *	7	* *	22	* *	22	* *
Reno	1,381	4.3 (34)	55.5 (38)	1,209	19.4 (17)	52.4 (51)	170	30.6 (13)	209	26.3 (20)	1,236	4.3 (9)	1,236	13.8 (53)
Republic	95	* *	* *	77	* *	* *	10	* *	23	* *	81	* *	81	* *
Rice	232	4.7 (43)	63.8 (69)	203	23.2 (45)	50.2 (34)	30	* *	49	* *	211	6.6 (31)	210	9.0 (19)
Riley-Ft	1,659	4.7 (43)	51.8 (10)	1,307	19.4 (17)	55.1 (61)	294	37.4 (18)	392	29.3 (26)	1,379	5.3 (16)	1,379	12.5 (46)
Riley-Manh	1,171	3.8 (24)	48.8 (3)	983	17.6 (7)	51.1 (39)	167	25.1 (6)	290	20.3 (13)	1,021	7.1 (43)	1,026	8.7 (17)
Rooks	148	8.1 (71)	45.9 (1)	131	24.4 (57)	39.7 (1)	16	* *	26	* *	128	7.0 (39)	128	10.2 (26)
Rush	29	* *	* *	26	* *	* *	3	* *	8	* *	27	* *	27	* *
Russell	130	2.3 (5)	53.8 (27)	118	23.7 (48)	53.4 (55)	24	* *	22	* *	120	3.3 (6)	120	6.7 (4)
Saline	1,532	4.5 (39)	55.7 (40)	1,374	16.4 (5)	54.2 (57)	252	22.6 (5)	247	19.0 (9)	1,392	7.2 (44)	1,394	8.2 (12)
Scott	134	6.0 (60)	53.0 (19)	120	22.5 (40)	46.7 (16)	21	* *	25	* *	118	2.5 (4)	118	7.6 (10)
Sedgwick-C	3,597	4.5 (39)	53.6 (25)	3,236	20.9 (29)	49.7 (31)	684	28.7 (12)	750	21.2 (15)	3,338	7.4 (46)	3,340	9.3 (22)
Sedgwick-M	5,683	3.3 (15)	56.0 (43)	5,131	21.7 (33)	49.5 (29)	1,167	27.6 (11)	1,287	21.8 (16)	5,276	7.2 (44)	5,275	12.2 (43)
Sedgwick-S	4,122	5.0 (47)	53.4 (23)	3,716	19.7 (21)	53.6 (56)	733	35.3 (17)	953	28.2 (24)	3,846	6.6 (31)	3,842	12.3 (44)
Seneca	26	* *	* *	23	* *	* *	6	* *	4	* *	26	* *	26	* *
Seward	1,381	2.5 (9)	59.2 (57)	1,244	24.1 (54)	45.0 (8)	230	27.4 (10)	188	28.7 (25)	1,266	8.5 (54)	1,267	5.8 (1)
ShawneeM&I	903	4.0 (31)	52.5 (16)	696	22.1 (38)	47.8 (19)	17	* *	102	29.4 (27)	704	9.5 (63)	703	6.7 (4)

(1) Reporting period is January 1 through December 31.  
(2) Excludes records with unknown data or errors.  
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.  
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).  
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).  
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.  
\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 5A**  
**2009 - 2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Maternal Health Indicators by Clinic**  
**Combined 3 Years**

Comparison of Maternal Health Indicators <sup>(2, 3)</sup>														
Clinic	Prepregnancy BMI <sup>(4)</sup>			Weight Gain <sup>(5)</sup>			Anemia (Low Hb/Hct) <sup>(6)</sup>				Gestational Diabetes During Pregnancy	Hypertension During Pregnancy		
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
		Number	% (Rank)		% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)			Number	% (Rank)
ShawneeMa	3,453	4.7 (43)	54.9 (33)	3,083	20.2 (26)	52.1 (48)	416	47.6 (22)	578	29.4(27)	3,168	6.7 (33)	3,171	8.5(16)
Sheridan	25	* *	* *	23	* *	* *	4	* *	3	* *	25	* *	25	* *
Sherman	183	3.8 (24)	54.6 (32)	156	26.3 (65)	51.3 (42)	17	* *	34	* *	162	4.3 (9)	162	9.9(24)
Smith	53	* *	* *	46	* *	* *	9	* *	7	* *	45	* *	45	* *
Stafford	99	* *	* *	84	* *	* *	10	* *	13	* *	84	* *	85	* *
Stanton	40	* *	* *	38	* *	* *	3	* *	19	* *	39	* *	38	* *
Stevens	225	2.2 (3)	60.4 (62)	198	20.2 (26)	50.5 (36)	44	* *	61	* *	201	15.9 (69)	202	8.4(15)
Sumner	440	5.9 (57)	54.3 (31)	397	23.7 (48)	41.3 (2)	63	* *	112	17.9 (6)	413	7.0 (39)	414	17.9(69)
Thomas	162	4.3 (34)	58.0 (52)	142	14.8 (2)	59.9 (69)	25	* *	26	* *	147	4.1 (7)	147	15.6(65)
Trego	42	* *	* *	34	* *	* *	8	* *	5	* *	34	* *	36	* *
Wabaunsee	58	* *	* *	56	* *	* *	8	* *	28	* *	59	* *	59	* *
Washington	72	* *	* *	59	* *	* *	9	* *	20	* *	62	* *	62	* *
Wichita	42	* *	* *	38	* *	* *	6	* *	2	* *	40	* *	40	* *
Wilson	256	7.0 (68)	59.8 (59)	242	24.0 (51)	52.1 (48)	42	* *	37	* *	244	8.6 (55)	244	12.7(48)
Woodson	40	* *	* *	38	* *	* *	9	* *	15	* *	38	* *	38	* *
Wyandotte	6,299	4.2 (32)	56.1 (45)	5,579	20.7 (28)	50.7 (38)	971	33.1 (14)	1,154	24.1(19)	5,666	5.4 (17)	5,663	9.1(21)
Unknown Clinics	46	* *	* *	44	* *	* *	3	* *	6	* *	46	* *	46	* *

(1) Reporting period is January 1 through December 31.  
(2) Excludes records with unknown data or errors.  
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.  
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).  
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).  
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.  
\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 5A**  
**2009 - 2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Maternal Health Indicators by Clinic**  
**Combined 3 Years**

<b>Comparison of Maternal Health Indicators</b> <sup>(2, 3)</sup>														
<b>Clinic</b>	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
<b>Kansas</b>	<b>63,724</b>	<b>4.3</b>	<b>54.9</b>	<b>55,948</b>	<b>21.2</b>	<b>50.1</b>	<b>9,633</b>	<b>29.8</b>	<b>12,815</b>	<b>22.8</b>	<b>57,413</b>	<b>6.9</b>	<b>57,405</b>	<b>10.4</b>

(1) Reporting period is January 1 through December 31.  
(2) Excludes records with unknown data or errors.  
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.  
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).  
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).  
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.  
\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 6A**  
**2009 - 2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Maternal Behavioral Indicators by Clinic**  
**Combined 3 Years**

<b>Comparison of Maternal Behavioral Indicators</b> <sup>(2, 3)</sup>										
<b>Clinic</b>	<b>Medical Care</b>				<b>WIC Enrollment</b>		<b>Multivitamin Consumption</b>			
	<b>1st Trimester</b>		<b>No Care</b>		<b>1st Trimester</b>		<b>Prepregnancy</b> (4)		<b>During Pregnancy</b> (5)	
	<b>Number</b>	<b>% (Rank)</b>	<b>% (Rank)</b>		<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>
Allen	204	92.6 (3)	2.0 (18)		246	35.4 (54)	246	17.5 (36)	208	48.1 (60)
Anderson	114	85.1 (23)	7.0 (54)		153	48.4 (6)	153	21.6 (24)	138	67.4 (24)
Atchison	376	89.6 (6)	2.7 (26)		441	39.0 (40)	443	11.1 (62)	369	67.8 (22)
Barber-Kio	2	* *	* *		3	* *	3	* *	3	* *
Barber-Med	112	81.3 (37)	8.0 (58)		135	45.2 (17)	136	35.3 (6)	114	75.4 (2)
Barton	669	87.4 (14)	0.9 (7)		768	48.3 (7)	771	19.6 (32)	615	59.3 (44)
Bourbon	293	88.4 (10)	1.7 (15)		355	42.0 (31)	355	13.5 (53)	294	59.9 (42)
Brown	194	79.9 (42)	10.8 (63)		277	39.0 (40)	279	15.4 (47)	240	65.8 (28)
Butler	727	75.1 (53)	5.8 (49)		819	36.1 (51)	820	16.5 (43)	629	45.2 (62)
Chase	37	* *	* *		39	* *	41	* *	36	* *
Chautauqua	76	* *	* *		81	* *	83	* *	75	* *
CherokeeBS	49	* *	* *		75	* *	75	* *	60	* *
CherokeeCo	317	78.5 (45)	6.9 (53)		486	32.9 (63)	488	12.7 (57)	375	56.8 (54)
Cheyenne	24	* *	* *		33	* *	33	* *	28	* *
Clark	9	* *	* *		10	* *	10	* *	9	* *
Clay	143	86.0 (20)	3.5 (33)		160	43.8 (23)	160	21.3 (25)	146	68.5 (20)
Cloud	222	84.7 (25)	4.5 (41)		240	47.1 (12)	242	25.6 (14)	207	68.6 (19)
Coffey	66	* *	* *		84	* *	84	* *	71	* *
Comanche	28	* *	* *		31	* *	31	* *	26	* *
Cowley-Ark	461	64.4 (64)	4.8 (46)		598	38.3 (43)	603	15.8 (46)	492	57.1 (53)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 6A**  
**2009 - 2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Maternal Behavioral Indicators by Clinic**  
**Combined 3 Years**

<b>Comparison of Maternal Behavioral Indicators</b> <sup>(2, 3)</sup>									
<b>Clinic</b>	<b>Medical Care</b>			<b>WIC Enrollment</b>		<b>Multivitamin Consumption</b>			
	<b>1st Trimester</b>		<b>No Care</b>	<b>1st Trimester</b>		<b>Prepregnancy</b> (4)		<b>During Pregnancy</b> (5)	
	<b>Number</b>	<b>% (Rank)</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>
Cowley-Win	389	77.1 (48)	7.7 (57)	477	46.5 (13)	484	12.0 (60)	410	58.8 (47)
Crawford-G	3	* *	* *	4	* *	4	* *	4	* *
Crawford-P	872	85.6 (21)	4.0 (37)	1,101	42.8 (27)	1,104	13.7 (51)	932	58.4 (48)
Decatur	32	* *	* *	40	* *	40	* *	31	* *
DickinsonA	315	85.1 (23)	1.6 (14)	356	42.4 (29)	361	20.5 (28)	286	69.6 (16)
DickinsonH	45	* *	* *	50	* *	50	* *	45	* *
Doniphan	111	82.0 (34)	4.5 (41)	130	33.8 (60)	130	36.2 (5)	96	* *
Douglas	1,262	74.2 (54)	7.0 (54)	1,557	38.2 (44)	1,562	20.2 (30)	1,260	63.9 (34)
Edwards	58	* *	* *	69	* *	71	* *	61	* *
Elk	48	* *	* *	56	* *	56	* *	46	* *
Ellis	420	86.2 (19)	2.1 (20)	490	32.9 (63)	492	26.8 (11)	405	67.2 (26)
Ellsworth	96	* *	* *	104	48.1 (8)	104	23.1 (20)	90	* *
Finney	1,741	53.8 (69)	28.1 (68)	2,088	35.9 (52)	2,105	11.1 (62)	1,737	41.6 (65)
Ford	1,679	72.8 (59)	12.9 (65)	2,044	44.6 (20)	2,057	11.3 (61)	1,790	55.0 (56)
Franklin	235	79.1 (43)	6.0 (50)	557	39.5 (38)	561	8.9 (67)	429	60.6 (40)
Geary	911	74.2 (54)	1.3 (13)	1,218	35.1 (55)	1,220	20.4 (29)	997	59.7 (43)
Gove	30	* *	* *	35	* *	35	* *	23	* *
Graham	49	* *	* *	57	* *	58	* *	49	* *
Grant	174	63.8 (65)	19.5 (67)	205	34.1 (59)	209	10.0 (65)	169	44.4 (63)
Gray	116	87.9 (11)	1.7 (15)	142	47.2 (11)	145	61.4 (2)	127	74.8 (3)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 6A**  
**2009 - 2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Maternal Behavioral Indicators by Clinic**  
**Combined 3 Years**

<b>Comparison of Maternal Behavioral Indicators</b> <sup>(2, 3)</sup>										
<b>Clinic</b>	<b>Medical Care</b>				<b>WIC Enrollment</b>		<b>Multivitamin Consumption</b>			
	<b>1st Trimester</b>		<b>No Care</b>		<b>1st Trimester</b>		<b>Prepregnancy</b> (4)		<b>During Pregnancy</b> (5)	
	<b>Number</b>	<b>% (Rank)</b>	<b>% (Rank)</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>
Greeley	29	* *	* *	* *	41	* *	41	* *	31	* *
Greenwood	110	60.9 (68)	30.9 (69)		131	43.5 (25)	131	16.8 (41)	111	74.8 (3)
Hamilton	66	* *	* *	* *	77	* *	77	* *	64	* *
Harper	140	77.1 (48)	10.0 (61)		160	45.6 (15)	165	30.3 (9)	136	73.5 (5)
Harvey	669	76.7 (51)	10.0 (61)		813	47.8 (9)	818	19.6 (32)	685	67.4 (24)
Haskell	47	* *	* *	* *	58	* *	59	* *	50	* *
Hodgeman	36	* *	* *	* *	39	* *	39	* *	34	* *
JacksonHol	113	81.4 (36)	3.5 (33)		232	42.7 (28)	233	13.3 (54)	190	76.3 (1)
JacksonPBP	2	* *	* *	* *	7	* *	7	* *	5	* *
Jefferson	158	83.5 (27)	9.5 (60)		183	44.3 (22)	183	9.8 (66)	152	41.4 (66)
Jewell	22	* *	* *	* *	22	* *	22	* *	20	* *
Johnson-M	1,968	71.4 (61)	2.8 (27)		2,494	33.4 (61)	2,503	17.0 (38)	1,973	65.8 (28)
Johnson-O	2,644	62.7 (67)	3.8 (35)		3,286	33.2 (62)	3,301	12.4 (58)	2,704	58.9 (46)
Kearny	47	* *	* *	* *	56	* *	56	* *	46	* *
Kingman	128	80.5 (40)	4.7 (44)		154	54.5 (4)	155	21.9 (23)	131	70.2 (13)
Kiowa	35	* *	* *	* *	39	* *	39	* *	29	* *
Labette	380	87.6 (12)	3.2 (31)		528	35.6 (53)	530	13.2 (56)	443	64.1 (33)
Lane	34	* *	* *	* *	40	* *	40	* *	33	* *
Leavenwort	913	86.7 (16)	0.8 (4)		1,086	31.2 (67)	1,090	22.1 (21)	840	71.8 (10)
Lincoln	55	* *	* *	* *	64	* *	64	* *	58	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.





**Table 6A**  
**2009 - 2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Maternal Behavioral Indicators by Clinic**  
**Combined 3 Years**

Comparison of Maternal Behavioral Indicators <sup>(2, 3)</sup>									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Pregpregnancy <sup>(4)</sup>		During Pregpregnancy <sup>(5)</sup>	
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Linn	188	80.3 (41)	8.0 (58)	206	47.6 (10)	206	23.3 (18)	181	65.2 (30)
Logan	18	* *	* *	38	* *	38	* *	34	* *
Lyon	751	86.7 (16)	0.8 (4)	883	41.0 (34)	884	18.0 (35)	728	62.4 (36)
Marion	133	78.9 (44)	2.3 (23)	160	39.4 (39)	161	39.1 (4)	127	67.7 (23)
Marshall	146	93.2 (1)	0.0 (1)	160	44.4 (21)	160	45.6 (3)	133	69.9 (15)
Mc Pherson	358	86.6 (18)	1.1 (11)	409	37.4 (46)	409	24.9 (15)	334	69.5 (17)
Meade	104	77.9 (47)	6.7 (51)	116	56.0 (3)	118	33.9 (7)	105	73.3 (7)
Miami	406	82.3 (31)	1.0 (10)	455	34.3 (58)	455	14.9 (49)	354	57.9 (49)
Mitchell	118	91.5 (4)	0.8 (4)	132	37.1 (48)	133	23.3 (18)	110	61.8 (37)
Montgome-C	322	85.4 (22)	2.2 (21)	474	28.1 (70)	476	16.0 (45)	408	72.5 (9)
Montgome-I	353	87.3 (15)	3.1 (30)	529	32.1 (65)	533	12.4 (58)	441	64.2 (32)
Morris	66	* *	* *	83	* *	85	* *	67	* *
Nemaha	83	* *	* *	93	* *	93	* *	81	* *
Neosho-Ch	365	89.3 (7)	4.1 (40)	434	38.5 (42)	436	13.3 (54)	368	70.1 (14)
Neosho-Er	22	* *	* *	30	* *	30	* *	23	* *
Ness	25	* *	* *	34	* *	35	* *	30	* *
Norton	108	90.7 (5)	0.9 (7)	119	46.2 (14)	120	17.5 (36)	102	13.7 (68)
Osage	201	89.1 (8)	2.0 (18)	277	43.7 (24)	282	14.2 (50)	242	42.1 (64)
Osborne	45	* *	* *	57	* *	57	* *	45	* *
Ottawa	65	* *	* *	71	* *	71	* *	60	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 6A**  
**2009 - 2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Maternal Behavioral Indicators by Clinic**  
**Combined 3 Years**

Comparison of Maternal Behavioral Indicators <sup>(2, 3)</sup>										
Clinic	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Pregnancy <sup>(4)</sup>		During Pregnancy <sup>(5)</sup>	
	Number	% (Rank)	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Pawnee	139	82.0 (34)	2.2 (21)		167	45.5 (16)	169	24.9 (15)	140	62.9 (35)
Phillips	92	* *	* *		107	52.3 (5)	109	22.0 (22)	96	* *
Pottawat-S	38	* *	* *		48	* *	48	* *	36	* *
Pottawat-W	42	* *	* *		47	* *	47	* *	43	* *
Pratt	231	82.7 (30)	3.9 (36)		264	41.3 (33)	265	7.2 (71)	210	68.1 (21)
Rawlins	11	* *	* *		27	* *	27	* *	23	* *
Reno	1,209	83.4 (28)	2.8 (27)		1,403	40.7 (35)	1,412	20.7 (27)	1,161	60.9 (39)
Republic	81	* *	* *		97	* *	98	* *	84	* *
Rice	209	82.3 (31)	0.5 (2)		233	42.1 (30)	237	16.9 (40)	192	45.3 (61)
Riley-Ft	1,291	92.9 (2)	0.6 (3)		1,768	31.3 (66)	1,769	23.6 (17)	1,376	71.6 (11)
Riley-Manh	946	72.9 (58)	1.2 (12)		1,211	37.9 (45)	1,212	19.5 (34)	1,002	59.3 (44)
Rooks	130	84.6 (26)	4.6 (43)		149	45.0 (18)	150	32.7 (8)	121	60.3 (41)
Rush	27	* *	* *		29	* *	30	* *	25	* *
Russell	120	83.3 (29)	2.5 (24)		131	45.0 (18)	133	8.3 (69)	113	70.8 (12)
Saline	1,126	76.4 (52)	4.7 (44)		1,559	35.0 (56)	1,563	6.0 (72)	1,315	57.6 (50)
Scott	118	77.1 (48)	6.8 (52)		132	43.2 (26)	135	84.4 (1)	121	50.4 (59)
Sedgwick-C	3,223	73.9 (56)	7.2 (56)		3,685	29.2 (69)	3,697	15.4 (47)	3,086	51.9 (57)
Sedgwick-M	5,243	72.5 (60)	4.8 (46)		5,851	27.4 (72)	5,863	16.6 (42)	4,770	57.4 (51)
Sedgwick-S	3,651	81.2 (38)	2.5 (24)		4,235	29.8 (68)	4,248	17.0 (38)	3,448	37.4 (67)
Seneca	26	* *	* *		30	* *	30	* *	28	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 6A**  
**2009 - 2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Maternal Behavioral Indicators by Clinic**  
**Combined 3 Years**

Comparison of Maternal Behavioral Indicators <sup>(2, 3)</sup>										
Clinic	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Pregpregnancy (4)		During Pregpregnancy (5)	
	Number	% (Rank)	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Seward	1,111	64.5 (63)	13.8 (66)		1,397	34.4 (57)	1,407	10.2 (64)	1,163	66.7 (27)
ShawneeM&I	648	71.3 (62)	2.9 (29)		906	65.2 (1)	913	20.8 (26)	853	55.6 (55)
ShawneeMa	2,798	78.3 (46)	4.0 (37)		3,548	40.2 (36)	3,560	8.9 (67)	2,902	51.2 (58)
Sheridan	25	* *	* *		26	* *	26	* *	20	* *
Sherman	120	80.8 (39)	5.0 (48)		189	40.2 (36)	189	25.9 (13)	152	73.0 (8)
Smith	41	* *	* *		52	* *	53	* *	48	* *
Stafford	83	* *	* *		100	58.0 (2)	100	20.0 (31)	85	* *
Stanton	36	* *	* *		41	* *	41	* *	28	* *
Stevens	201	73.6 (57)	4.0 (37)		228	28.1 (70)	231	26.0 (12)	188	69.1 (18)
Sumner	413	82.1 (33)	3.4 (32)		457	37.2 (47)	457	13.6 (52)	371	64.7 (31)
Thomas	105	87.6 (12)	1.9 (17)		166	36.7 (49)	166	28.3 (10)	136	73.5 (5)
Trego	36	* *	* *		44	* *	44	* *	39	* *
Wabaunsee	54	* *	* *		61	* *	61	* *	47	* *
Washington	61	* *	* *		74	* *	75	* *	62	* *
Wichita	40	* *	* *		44	* *	44	* *	35	* *
Wilson	235	88.9 (9)	0.9 (7)		258	41.5 (32)	260	16.5 (43)	224	61.6 (38)
Woodson	40	* *	* *		40	* *	40	* *	34	* *
Wyandotte	5,159	63.0 (66)	11.4 (64)		6,442	36.2 (50)	6,487	7.7 (70)	5,470	57.4 (51)
Unknown Clinics	37	* *	* *		48	* *	48	* *	40	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 6A**  
**2009 - 2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Maternal Behavioral Indicators by Clinic**  
**Combined 3 Years**

<b>Comparison of Maternal Behavioral Indicators</b> <sup>(2, 3)</sup>									
<b>Clinic</b>	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Pregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
<b>Kansas</b>	<b>53,378</b>	<b>75.5</b>	<b>6.0</b>	<b>65,378</b>	<b>36.4</b>	<b>65,683</b>	<b>15.8</b>	<b>54,117</b>	<b>57.7</b>

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 7A**  
**2009 - 2011 Pregnancy Nutrition Surveillance** (1)  
**Kansas**  
**Comparison of Smoking Indicators by Clinic**  
**Combined 3 Years**

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Allen	234	45.3 (53)	184	26.6 (36)	61	* *	204	41.2 (60)	187	35.8 (55)
Anderson	151	47.0 (60)	125	27.2 (39)	50	* *	136	23.5 (41)	127	20.5 (39)
Atchison	329	44.1 (45)	216	29.6 (50)	71	* *	279	29.0 (47)	220	24.1 (44)
Barber-Kio	3	* *	2	* *	0	* *	3	* *	2	* *
Barber-Med	135	47.4 (61)	109	32.1 (57)	45	* *	114	48.2 (66)	110	50.0 (64)
Barton	766	36.4 (26)	668	23.2 (28)	188	28.2 (24)	611	12.6 (20)	666	6.0 (13)
Bourbon	337	46.6 (56)	271	28.0 (44)	101	25.7 (29)	280	40.7 (59)	274	37.6 (56)
Brown	265	43.8 (44)	186	31.2 (56)	65	* *	230	45.2 (63)	191	42.9 (62)
Butler	807	49.6 (62)	716	30.6 (54)	276	31.5 (20)	623	11.1 (16)	723	6.8 (16)
Chase	41	* *	36	* *	19	* *	36	* *	36	* *
Chautauqua	71	* *	50	* *	18	* *	65	* *	50	* *
CherokeeBS	72	* *	64	* *	20	* *	57	* *	64	* *
CherokeeCo	473	41.4 (40)	416	27.6 (42)	134	22.4 (31)	369	35.5 (54)	419	35.6 (54)
Cheyenne	33	* *	27	* *	6	* *	28	* *	27	* *
Clark	6	* *	6	* *	0	* *	8	* *	8	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 7A**  
**2009 - 2011 Pregnancy Nutrition Surveillance** (1)  
**Kansas**  
**Comparison of Smoking Indicators by Clinic**  
**Combined 3 Years**

<b>Comparison of Smoking Indicators</b> (2, 3)										
<b>Clinic</b>	<b>Smoking</b>				<b>Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)</b>		<b>Smoking in Household</b>			
	<b>3 Months Prior to Pregnancy</b>		<b>Last 3 Months of Pregnancy</b>				<b>Prenatal</b>		<b>Postpartum</b>	
	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>
Clay	160	50.0 (64)	141	29.1 (49)	60	* *	146	14.4 (25)	141	5.7 (12)
Cloud	242	36.4 (26)	219	20.1 (22)	67	* *	207	8.7 (10)	221	2.7 (1)
Coffey	80	* *	66	* *	27	* *	69	* *	66	* *
Comanche	31	* *	29	* *	12	* *	26	* *	29	* *
Cowley-Ark	587	36.8 (28)	434	23.3 (29)	125	32.0 (18)	479	21.5 (38)	432	21.1 (41)
Cowley-Win	466	44.8 (49)	350	32.9 (61)	134	21.6 (32)	399	19.0 (34)	334	17.7 (37)
Crawford-G	4	* *	3	* *	1	* *	4	* *	3	* *
Crawford-P	1,031	44.8 (49)	795	26.5 (35)	282	31.9 (19)	892	40.2 (58)	797	39.8 (59)
Decatur	38	* *	32	* *	6	* *	30	* *	33	* *
DickinsonA	356	45.2 (52)	310	26.1 (34)	106	35.8 (11)	281	7.8 (7)	311	6.1 (14)
DickinsonH	50	* *	43	* *	19	* *	45	* *	43	* *
Doniphan	121	38.0 (32)	89	* *	19	* *	90	* *	90	* *
Douglas	1,532	37.5 (30)	1,249	18.7 (19)	373	42.4 (5)	1,243	10.5 (13)	1,268	6.9 (17)
Edwards	71	* *	62	* *	18	* *	60	* *	63	* *
Elk	52	* *	43	* *	17	* *	43	* *	42	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 7A**  
**2009 - 2011 Pregnancy Nutrition Surveillance** (1)  
**Kansas**  
**Comparison of Smoking Indicators by Clinic**  
**Combined 3 Years**

<b>Comparison of Smoking Indicators</b> (2, 3)										
<b>Clinic</b>	<b>Smoking</b>				<b>Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)</b>		<b>Smoking in Household</b>			
	<b>3 Months Prior to Pregnancy</b>		<b>Last 3 Months of Pregnancy</b>				<b>Prenatal</b>		<b>Postpartum</b>	
	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>
Ellis	489	39.9 (36)	414	18.1 (18)	130	40.8 (6)	404	7.7 (4)	416	5.3 (9)
Ellsworth	103	46.6 (56)	91	* *	32	* *	90	* *	91	* *
Finney	2,082	10.6 (5)	1,815	5.2 (4)	138	39.9 (7)	1,727	13.1 (22)	1,815	12.5 (29)
Ford	1,990	8.8 (2)	1,635	5.0 (3)	103	29.1 (23)	1,747	8.8 (11)	1,639	8.1 (18)
Franklin	558	43.2 (43)	480	29.6 (50)	145	21.4 (33)	427	12.9 (21)	484	8.7 (19)
Geary	1,200	36.9 (29)	942	20.2 (23)	254	35.0 (14)	990	34.3 (53)	949	28.9 (48)
Gove	33	* *	29	* *	3	* *	22	* *	29	* *
Graham	57	* *	47	* *	22	* *	49	* *	48	* *
Grant	209	10.0 (4)	194	4.1 (2)	13	* *	169	11.8 (17)	194	12.4 (28)
Gray	145	9.7 (3)	116	7.8 (5)	7	* *	126	7.9 (8)	117	5.1 (6)
Greeley	40	* *	33	* *	5	* *	31	* *	33	* *
Greenwood	110	53.6 (68)	57	* *	23	* *	102	31.4 (51)	83	* *
Hamilton	75	* *	68	* *	5	* *	64	* *	68	* *
Harper	153	44.4 (46)	124	27.4 (41)	41	* *	121	26.4 (43)	116	25.0 (46)
Harvey	804	37.9 (31)	692	21.1 (25)	212	37.7 (9)	674	6.1 (2)	692	2.9 (3)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 7A**  
**2009 - 2011 Pregnancy Nutrition Surveillance** (1)  
**Kansas**  
**Comparison of Smoking Indicators by Clinic**  
**Combined 3 Years**

<b>Comparison of Smoking Indicators</b> (2, 3)										
<b>Clinic</b>	<b>Smoking</b>				<b>Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)</b>	<b>Smoking in Household</b>				
	<b>3 Months Prior to Pregnancy</b>		<b>Last 3 Months of Pregnancy</b>			<b>Prenatal</b>		<b>Postpartum</b>		
	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>
Haskell	59	* *	52	* *	7	* *	50	* *	52	* *
Hodgeman	37	* *	35	* *	15	* *	32	* *	34	* *
JacksonHol	231	49.8 (63)	198	32.8 (60)	75	* *	188	20.2 (35)	199	19.6 (38)
JacksonPBP	7	* *	3	* *	0	* *	5	* *	4	* *
Jefferson	81	* *	40	* *	14	* *	75	* *	43	* *
Jewell	21	* *	17	* *	8	* *	19	* *	18	* *
Johnson-M	2,460	26.1 (11)	2,132	10.4 (7)	383	48.3 (1)	1,951	21.3 (37)	2,132	20.8 (40)
Johnson-O	3,274	21.4 (7)	2,884	9.4 (6)	446	44.8 (2)	2,676	15.5 (29)	2,889	15.4 (33)
Kearny	56	* *	50	* *	10	* *	46	* *	50	* *
Kingman	145	46.9 (59)	112	32.1 (57)	38	* *	127	39.4 (57)	116	33.6 (52)
Kiowa	38	* *	30	* *	6	* *	28	* *	33	* *
Labette	513	38.4 (33)	454	28.2 (46)	145	18.6 (36)	434	31.8 (52)	456	32.9 (51)
Lane	18	* *	3	* *	0	* *	17	* *	5	* *
Leavenwort	1,068	46.0 (55)	879	24.5 (32)	287	36.2 (10)	832	37.7 (56)	880	38.4 (58)
Lincoln	63	* *	51	* *	27	* *	58	* *	51	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.





**Table 7A**  
**2009 - 2011 Pregnancy Nutrition Surveillance** (1)  
**Kansas**  
**Comparison of Smoking Indicators by Clinic**  
**Combined 3 Years**

<b>Comparison of Smoking Indicators</b> (2, 3)										
<b>Clinic</b>	<b>Smoking</b>				<b>Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)</b>		<b>Smoking in Household</b>			
	<b>3 Months Prior to Pregnancy</b>		<b>Last 3 Months of Pregnancy</b>				<b>Prenatal</b>		<b>Postpartum</b>	
	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>
Linn	203	52.7 (67)	182	32.4 (59)	86	* *	179	14.5 (26)	183	10.4 (23)
Logan	37	* *	33	* *	9	* *	34	* *	33	* *
Lyon	871	29.6 (18)	760	15.8 (14)	176	35.8 (11)	725	10.6 (14)	743	6.1 (14)
Marion	160	23.8 (9)	134	17.2 (16)	26	* *	127	3.9 (1)	134	4.5 (5)
Marshall	156	44.9 (51)	135	26.7 (38)	49	* *	129	46.5 (65)	136	41.2 (60)
Mc Pherson	408	35.0 (23)	364	20.9 (24)	105	33.3 (15)	331	6.9 (3)	364	2.7 (1)
Meade	107	24.3 (10)	96	* *	23	* *	101	24.8 (42)	100	23.0 (43)
Miami	438	42.9 (41)	380	26.6 (36)	122	27.0 (26)	336	20.8 (36)	380	16.1 (35)
Mitchell	131	33.6 (21)	116	23.3 (29)	34	* *	110	10.9 (15)	117	9.4 (21)
Montgome-C	468	40.0 (37)	399	28.3 (47)	128	20.3 (35)	403	31.3 (50)	398	31.7 (49)
Montgome-I	507	40.2 (38)	434	27.6 (42)	139	20.9 (34)	432	36.1 (55)	433	33.9 (53)
Morris	84	* *	74	* *	22	* *	65	* *	74	* *
Nemaha	66	* *	51	* *	20	* *	53	* *	52	* *
Neosho-Ch	370	44.6 (48)	221	28.5 (48)	80	* *	324	30.9 (49)	227	22.9 (42)
Neosho-Er	26	* *	10	* *	2	* *	22	* *	10	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 7A**  
**2009 - 2011 Pregnancy Nutrition Surveillance** (1)  
**Kansas**  
**Comparison of Smoking Indicators by Clinic**  
**Combined 3 Years**

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Ness	30	* *	25	* *	7	* *	26	* *	25	* *
Norton	116	39.7 (35)	94	* *	35	* *	100	27.0 (45)	98	* *
Osage	264	46.6 (56)	217	27.2 (39)	78	* *	225	28.0 (46)	219	26.5 (47)
Osborne	56	* *	42	* *	15	* *	45	* *	42	* *
Ottawa	68	* *	41	* *	17	* *	59	* *	54	* *
Pawnee	164	34.8 (22)	123	19.5 (21)	31	* *	138	18.1 (33)	124	14.5 (32)
Phillips	96	* *	68	* *	18	* *	75	* *	69	* *
Pottawat-S	48	* *	40	* *	5	* *	36	* *	41	* *
Pottawat-W	47	* *	41	* *	13	* *	43	* *	43	* *
Pratt	258	39.5 (34)	221	28.1 (45)	74	* *	204	42.6 (61)	214	37.9 (57)
Rawlins	26	* *	22	* *	6	* *	23	* *	22	* *
Reno	1,391	44.4 (46)	1,198	24.1 (31)	422	38.9 (8)	1,144	14.9 (28)	1,204	12.1 (27)
Republic	97	* *	76	* *	26	* *	81	* *	78	* *
Rice	232	43.1 (42)	204	29.9 (52)	71	* *	189	14.8 (27)	205	10.7 (24)
Riley-Ft	1,694	27.9 (14)	1,220	12.5 (11)	252	44.0 (3)	1,352	15.7 (30)	1,229	11.6 (26)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 7A**  
**2009 - 2011 Pregnancy Nutrition Surveillance** (1)  
**Kansas**  
**Comparison of Smoking Indicators by Clinic**  
**Combined 3 Years**

<b>Comparison of Smoking Indicators</b> (2, 3)										
<b>Clinic</b>	<b>Smoking</b>				<b>Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)</b>		<b>Smoking in Household</b>			
	<b>3 Months Prior to Pregnancy</b>		<b>Last 3 Months of Pregnancy</b>				<b>Prenatal</b>		<b>Postpartum</b>	
	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>
Riley-Manh	1,195	28.8 (17)	963	15.9 (15)	211	29.9 (22)	994	7.7 (4)	979	5.2 (7)
Rooks	143	55.2 (69)	115	36.5 (62)	51	* *	114	29.8 (48)	119	31.9 (50)
Rush	30	* *	27	* *	16	* *	25	* *	27	* *
Russell	132	45.5 (54)	118	39.8 (63)	46	* *	113	54.9 (67)	117	53.0 (65)
Saline	1,522	40.9 (39)	1,305	25.4 (33)	435	26.9 (27)	1,289	11.8 (17)	1,308	9.1 (20)
Scott	125	29.6 (18)	98	* *	26	* *	110	22.7 (40)	100	16.0 (34)
Sedgwick-C	3,673	27.8 (13)	3,308	15.5 (13)	715	35.2 (13)	3,072	13.7 (23)	3,304	11.2 (25)
Sedgwick-M	5,797	27.3 (12)	5,237	12.4 (10)	1,067	42.5 (4)	4,732	7.7 (4)	5,243	5.4 (11)
Sedgwick-S	4,228	35.0 (23)	3,825	22.5 (26)	1,034	24.8 (30)	3,435	7.9 (8)	3,824	5.2 (7)
Seneca	22	* *	13	* *	4	* *	20	* *	15	* *
Seward	1,378	5.3 (1)	1,232	2.7 (1)	46	* *	1,153	12.0 (19)	1,247	9.9 (22)
ShawneeM&I	848	28.4 (16)	637	17.4 (17)	149	31.5 (20)	741	14.0 (24)	643	3.4 (4)
ShawneeMa	3,319	35.4 (25)	2,561	22.8 (27)	681	25.8 (28)	2,732	9.8 (12)	2,563	5.3 (9)
Sheridan	24	* *	21	* *	4	* *	20	* *	23	* *
Sherman	185	28.1 (15)	161	14.9 (12)	40	* *	147	22.4 (39)	161	13.7 (30)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 7A**  
**2009 - 2011 Pregnancy Nutrition Surveillance** (1)  
**Kansas**  
**Comparison of Smoking Indicators by Clinic**  
**Combined 3 Years**

<b>Comparison of Smoking Indicators</b> (2, 3)										
<b>Clinic</b>	<b>Smoking</b>				<b>Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)</b>	<b>Smoking in Household</b>				
	<b>3 Months Prior to Pregnancy</b>		<b>Last 3 Months of Pregnancy</b>			<b>Prenatal</b>		<b>Postpartum</b>		
	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>
Smith	50	* *	37	* *	13	* *	45	* *	38	* *
Stafford	95	* *	80	* *	24	* *	80	* *	79	* *
Stanton	40	* *	37	* *	3	* *	28	* *	36	* *
Stevens	228	21.5 (8)	199	11.1 (8)	32	* *	186	26.9 (44)	202	24.3 (45)
Sumner	451	52.5 (66)	400	30.8 (55)	167	32.3 (17)	363	43.0 (62)	402	43.5 (63)
Thomas	164	32.3 (20)	144	18.8 (20)	36	* *	132	16.7 (31)	146	16.4 (36)
Trego	42	* *	32	* *	13	* *	37	* *	32	* *
Wabaunsee	61	* *	58	* *	16	* *	47	* *	58	* *
Washington	75	* *	59	* *	12	* *	62	* *	60	* *
Wichita	43	* *	34	* *	7	* *	34	* *	34	* *
Wilson	256	51.2 (65)	238	30.3 (53)	105	33.3 (15)	222	45.9 (64)	237	42.2 (61)
Woodson	37	* *	31	* *	7	* *	32	* *	32	* *
Wyandotte	6,373	19.9 (6)	5,554	11.6 (9)	841	27.8 (25)	5,418	16.7 (31)	5,656	13.9 (31)
Unknown Clinics	47	* *	45	* *	19	* *	40	* *	45	* *
<b>Kansas</b>	<b>63,865</b>	<b>30.9</b>	<b>54,208</b>	<b>17.6</b>	<b>12,825</b>	<b>32.5</b>	<b>52,872</b>	<b>16.8</b>	<b>54,494</b>	<b>14.0</b>

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 8A**  
**2009 - 2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Infant Health Indicators by Clinic**  
**Combined 3 Years**

<b>Comparison of Infant Health Indicators</b> <sup>(2, 3)</sup>									
<b>Clinic</b>	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Allen	209	8.6 (50)	2.4 (3)	220	8.2 (26)	191	5.8 (54)	219	62.1 (59)
Anderson	112	10.7 (63)	6.3 (33)	132	9.8 (47)	102	5.9 (56)	131	61.1 (62)
Atchison	365	9.6 (59)	3.8 (8)	386	7.8 (21)	335	6.3 (58)	385	51.9 (68)
Barber-Kio	2	* * *	* *	2	* *	2	* *	2	* *
Barber-Med	103	7.8 (37)	4.9 (21)	103	11.7 (62)	90	* *	103	79.6 (8)
Barton	639	5.6 (7)	4.1 (12)	642	7.0 (13)	591	4.2 (31)	642	74.8 (24)
Bourbon	300	8.7 (52)	3.0 (4)	304	10.9 (58)	267	5.6 (53)	303	61.7 (60)
Brown	216	8.8 (56)	6.5 (36)	236	11.0 (60)	190	5.3 (52)	236	55.1 (66)
Butler	718	6.8 (23)	6.1 (31)	730	7.7 (20)	661	4.1 (27)	727	70.4 (41)
Chase	36	* * *	* *	34	* *	33	* *	36	* *
Chautauqua	70	* * *	* *	72	* *	65	* *	74	* *
CherokeeBS	68	* * *	* *	68	* *	62	* *	68	* *
CherokeeCo	423	7.3 (28)	7.3 (47)	425	8.0 (24)	387	2.6 (4)	425	57.6 (64)
Cheyenne	24	* * *	* *	24	* *	19	* *	24	* *

(1) Reporting period is January 1 through December 31.  
(2) Excludes records with unknown data or errors.  
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.  
(4) <2500 g.  
(5) >4000 g.  
(6) <37 weeks gestation.  
(7) 37 or more weeks gestation and <2500 g.  
\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 8A**  
**2009 - 2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Infant Health Indicators by Clinic**  
**Combined 3 Years**

<b>Comparison of Infant Health Indicators</b> <sup>(2, 3)</sup>									
<b>Clinic</b>	<b>Birthweight</b>			<b>Preterm</b> (6)		<b>Full Term LBW</b> (7)		<b>Ever Breastfed</b>	
	Low (4)		High (5)	Number	%	Number	%	Number	%
	Number	% (Rank)	% (Rank)						
Clark	9	* * * *	*	9	* *	8	* *	9	* *
Clay	138	8.7 (52)	9.4 (61)	139	11.5 (61)	122	1.6 (3)	139	67.6 (48)
Cloud	214	8.4 (47)	4.2 (13)	214	10.3 (51)	190	4.7 (42)	214	78.0 (15)
Coffey	65	* * * *	*	65	* *	64	* *	65	* *
Comanche	25	* * * *	*	26	* *	22	* *	26	* *
Cowley-Ark	510	5.9 (10)	6.3 (33)	521	10.4 (53)	453	3.5 (14)	518	55.2 (65)
Cowley-Win	381	9.7 (60)	8.1 (56)	408	10.5 (55)	333	6.3 (58)	412	69.2 (45)
Crawford-G	3	* * * *	*	3	* *	3	* *	3	* *
Crawford-P	906	7.7 (35)	4.5 (17)	934	10.5 (55)	805	3.9 (22)	935	61.3 (61)
Decatur	35	* * * *	*	35	* *	32	* *	35	* *
DickinsonA	302	7.0 (26)	10.3 (63)	297	10.1 (50)	267	3.7 (16)	302	74.5 (26)
DickinsonH	43	* * * *	*	43	* *	39	* *	43	* *
Doniphan	107	12.1 (66)	10.3 (63)	108	13.9 (66)	92	* *	107	64.5 (55)
Douglas	1,268	6.8 (23)	6.7 (40)	1,293	7.5 (18)	1,169	4.1 (27)	1,288	79.2 (10)

(1) Reporting period is January 1 through December 31.  
(2) Excludes records with unknown data or errors.  
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.  
(4) <2500 g.  
(5) >4000 g.  
(6) <37 weeks gestation.  
(7) 37 or more weeks gestation and <2500 g.  
\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 8A**  
**2009 - 2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Infant Health Indicators by Clinic**  
**Combined 3 Years**

<b>Comparison of Infant Health Indicators</b> <sup>(2, 3)</sup>										
<b>Clinic</b>	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)							
	Number	% (Rank)	%	(Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Edwards	60	* * * *	* *	* *	60	* *	47	* *	59	* *
Elk	49	* * * *	* *	* *	51	* *	45	* *	51	* *
Ellis	387	5.9 (10)	8.0 (55)		413	6.5 (9)	358	3.4 (11)	414	77.5 (16)
Ellsworth	92	* * * *	* *	* *	92	* *	88	* *	92	* *
Finney	1,802	7.0 (26)	5.9 (28)		1,836	9.8 (47)	1,616	4.0 (24)	1,830	77.0 (17)
Ford	1,681	5.6 (7)	5.7 (26)		1,726	8.5 (31)	1,529	2.9 (6)	1,719	75.6 (23)
Franklin	473	5.9 (10)	5.1 (22)		474	9.1 (37)	426	3.8 (20)	477	64.6 (54)
Geary	954	7.8 (37)	7.7 (53)		956	8.3 (27)	873	4.2 (31)	950	72.6 (31)
Gove	29	* * * *	* *	* *	29	* *	27	* *	27	* *
Graham	43	* * * *	* *	* *	45	* *	37	* *	44	* *
Grant	190	6.8 (23)	4.2 (13)		189	5.8 (5)	175	4.0 (24)	192	73.4 (29)
Gray	101	5.0 (4)	10.9 (65)		114	8.8 (36)	92	* *	114	78.1 (14)
Greeley	34	* * * *	* *	* *	34	* *	30	* *	34	* *
Greenwood	110	3.6 (2)	4.5 (17)		111	4.5 (2)	105	3.8 (20)	111	69.4 (44)

(1) Reporting period is January 1 through December 31.  
(2) Excludes records with unknown data or errors.  
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.  
(4) <2500 g.  
(5) >4000 g.  
(6) <37 weeks gestation.  
(7) 37 or more weeks gestation and <2500 g.  
\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 8A**  
**2009 - 2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Infant Health Indicators by Clinic**  
**Combined 3 Years**

Comparison of Infant Health Indicators <sup>(2, 3)</sup>									
Clinic	Birthweight			Preterm <sup>(6)</sup>		Full Term LBW <sup>(7)</sup>		Ever Breastfed	
	Low <sup>(4)</sup>	High <sup>(5)</sup>		Number	%	Number	%	Number	%
	Number	% (Rank)	% (Rank)						
Hamilton	68	* * * *	* *	69	* *	63	* *	69	* *
Harper	138	8.0 (40)	10.1 (62)	136	7.4 (16)	123	4.1 (27)	139	74.8 (24)
Harvey	688	7.4 (29)	6.8 (42)	693	8.4 (29)	625	4.5 (38)	695	83.7 (2)
Haskell	47	* * * *	* *	48	* *	43	* *	48	* *
Hodgeman	35	* * * *	* *	35	* *	32	* *	35	* *
JacksonHol	197	7.6 (33)	12.2 (66)	204	7.4 (16)	182	4.9 (47)	203	72.4 (32)
JacksonPBP	4	* * * *	* *	4	* *	4	* *	4	* *
Jefferson	159	6.3 (14)	4.4 (16)	160	15.0 (68)	135	3.7 (16)	158	67.1 (49)
Jewell	21	* * * *	* *	22	* *	18	* *	22	* *
Johnson-M	1,986	7.5 (32)	6.7 (40)	2,141	6.7 (11)	1,839	4.0 (24)	2,135	78.7 (11)
Johnson-O	2,742	6.3 (14)	7.5 (49)	2,841	7.0 (13)	2,540	3.6 (15)	2,838	83.5 (3)
Kearny	50	* * * *	* *	51	* *	41	* *	51	* *
Kingman	79	* * * *	* *	120	5.0 (3)	74	* *	121	79.3 (9)
Kiowa	34	* * * *	* *	35	* *	29	* *	34	* *

(1) Reporting period is January 1 through December 31.  
(2) Excludes records with unknown data or errors.  
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.  
(4) <2500 g.  
(5) >4000 g.  
(6) <37 weeks gestation.  
(7) 37 or more weeks gestation and <2500 g.  
\* Percentages are not calculated if <100 records are available for analysis after exclusions.





**Table 8A**  
**2009 - 2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Infant Health Indicators by Clinic**  
**Combined 3 Years**

<b>Comparison of Infant Health Indicators</b> <sup>(2, 3)</sup>									
<b>Clinic</b>	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Labette	458	6.3 (14)	6.6 (37)	458	10.3 (51)	411	4.4 (35)	456	54.4 (67)
Lane	30	* *	* *	31	* *	25	* *	31	* *
Leavenwort	879	8.1 (44)	7.5 (49)	911	8.6 (33)	801	5.1 (51)	910	64.8 (53)
Lincoln	52	* *	* *	53	* *	45	* *	52	* *
Linn	178	6.7 (22)	3.9 (9)	183	10.9 (58)	158	4.4 (35)	183	76.5 (18)
Logan	33	* *	* *	33	* *	31	* *	33	* *
Lyon	754	8.4 (47)	6.6 (37)	753	9.2 (39)	684	4.7 (42)	749	76.2 (20)
Marion	122	7.4 (29)	6.6 (37)	124	7.3 (15)	112	4.5 (38)	124	85.5 (1)
Marshall	137	8.8 (56)	2.2 (2)	141	9.9 (49)	123	6.5 (60)	140	64.3 (57)
Mc Pherson	345	4.3 (3)	4.6 (19)	350	6.0 (8)	324	1.2 (1)	349	74.5 (26)
Meade	94	* *	* *	103	9.7 (46)	82	* *	103	78.6 (13)
Miami	407	6.6 (20)	8.8 (58)	410	7.6 (19)	377	3.7 (16)	407	70.0 (42)
Mitchell	98	* *	* *	116	4.3 (1)	94	* *	115	75.7 (22)
Montgome-C	420	7.6 (33)	3.6 (6)	422	8.1 (25)	385	4.4 (35)	422	62.3 (58)

(1) Reporting period is January 1 through December 31.  
(2) Excludes records with unknown data or errors.  
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.  
(4) <2500 g.  
(5) >4000 g.  
(6) <37 weeks gestation.  
(7) 37 or more weeks gestation and <2500 g.  
\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 8A**  
**2009 - 2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Infant Health Indicators by Clinic**  
**Combined 3 Years**

Comparison of Infant Health Indicators <sup>(2, 3)</sup>										
Clinic	Birthweight				Preterm <sup>(6)</sup>		Full Term LBW <sup>(7)</sup>		Ever Breastfed	
	Low <sup>(4)</sup>		High <sup>(5)</sup>		Number	%	Number	%	Number	%
	Number	% (Rank)	% (Rank)	(Rank)						
Montgome-I	448	8.0 (40)	3.1 (5)		449	7.8 (21)	410	4.9 (47)	448	60.3 (63)
Morris	66	* *	* *		73	* *	57	* *	74	* *
Nemaha	56	* *	* *		83	* *	52	* *	79	* *
Neosho-Ch	310	6.5 (19)	5.5 (25)		390	9.2 (39)	280	3.2 (8)	388	64.4 (56)
Neosho-Er	20	* *	* *		26	* *	19	* *	26	* *
Ness	26	* *	* *		28	* *	23	* *	29	* *
Norton	101	10.9 (64)	4.0 (11)		101	13.9 (66)	87	* *	99	* *
Osage	241	8.7 (52)	3.7 (7)		249	12.9 (65)	205	5.9 (56)	254	65.4 (52)
Osborne	45	* *	* *		46	* *	43	* *	45	* *
Ottawa	63	* *	* *		66	* *	59	* *	66	* *
Pawnee	135	7.4 (29)	5.9 (28)		135	10.4 (53)	119	3.4 (11)	136	70.6 (39)
Phillips	87	* *	* *		87	* *	78	* *	86	* *
Pottawat-S	38	* *	* *		38	* *	34	* *	38	* *
Pottawat-W	43	* *	* *		43	* *	40	* *	43	* *

(1) Reporting period is January 1 through December 31.  
(2) Excludes records with unknown data or errors.  
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.  
(4) <2500 g.  
(5) >4000 g.  
(6) <37 weeks gestation.  
(7) 37 or more weeks gestation and <2500 g.  
\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 8A**  
**2009 - 2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Infant Health Indicators by Clinic**  
**Combined 3 Years**

<b>Comparison of Infant Health Indicators</b> <sup>(2, 3)</sup>										
<b>Clinic</b>	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)							
	Number	% (Rank)	%	(Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Pratt	223	3.1 (1)	9.0 (60)		224	6.7 (11)	207	1.4 (2)	225	71.6 (35)
Rawlins	22	* *	* *	*	22	* *	21	* *	22	* *
Reno	1,201	8.0 (40)	5.7 (26)		1,208	8.5 (31)	1,092	4.9 (47)	1,212	69.7 (43)
Republic	76	* *	* *	*	78	* *	70	* *	79	* *
Rice	197	6.6 (20)	7.6 (52)		196	9.2 (39)	175	2.9 (6)	200	76.0 (21)
Riley-Ft	1,366	6.1 (13)	8.8 (58)		1,383	5.8 (5)	1,286	3.7 (16)	1,374	76.3 (19)
Riley-Manh	1,007	6.3 (14)	7.0 (44)		1,015	6.5 (9)	940	3.3 (9)	1,012	83.4 (4)
Rooks	129	7.8 (37)	4.7 (20)		129	7.8 (21)	119	4.2 (31)	129	72.9 (30)
Rush	26	* *	* *	*	25	* *	23	* *	26	* *
Russell	114	5.3 (5)	5.3 (24)		112	15.2 (69)	95	* *	114	65.8 (51)
Saline	1,337	7.7 (35)	6.2 (32)		1,384	9.5 (44)	1,205	4.3 (34)	1,380	66.4 (50)
Scott	116	8.6 (50)	1.7 (1)		115	8.7 (35)	103	4.9 (47)	117	70.9 (38)
Sedgwick-C	3,246	8.7 (52)	7.1 (46)		3,260	8.4 (29)	2,964	4.5 (38)	3,257	70.6 (39)
Sedgwick-M	5,153	9.9 (61)	6.3 (33)		5,192	8.6 (33)	4,702	5.8 (54)	5,177	72.2 (33)

(1) Reporting period is January 1 through December 31.  
(2) Excludes records with unknown data or errors.  
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.  
(4) <2500 g.  
(5) >4000 g.  
(6) <37 weeks gestation.  
(7) 37 or more weeks gestation and <2500 g.  
\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 8A**  
**2009 - 2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Infant Health Indicators by Clinic**  
**Combined 3 Years**

Comparison of Infant Health Indicators <sup>(2, 3)</sup>									
Clinic	Birthweight			Preterm <sup>(6)</sup>		Full Term LBW <sup>(7)</sup>		Ever Breastfed	
	Low <sup>(4)</sup>	High <sup>(5)</sup>		Number	%	Number	%	Number	%
	Number	% (Rank)	% (Rank)						
Sedgwick-S	3,757	8.2 (46)	7.0 (44)	3,761	8.3 (27)	3,434	4.7 (42)	3,753	69.1 (46)
Seneca	17	* * *	* *	26	* *	16	* *	25	* *
Seward	1,156	5.4 (6)	7.9 (54)	1,215	9.1 (37)	1,040	3.4 (11)	1,216	73.6 (28)
ShawneeM&I	681	8.5 (49)	8.5 (57)	679	12.5 (63)	590	3.9 (22)	681	79.7 (7)
ShawneeMa	3,084	8.0 (40)	5.1 (22)	3,122	9.5 (44)	2,780	4.7 (42)	3,107	69.0 (47)
Sheridan	25	* * *	* *	25	* *	22	* *	25	* *
Sherman	155	9.0 (58)	3.9 (9)	155	5.8 (5)	146	4.1 (27)	155	80.0 (6)
Smith	44	* * *	* *	44	* *	39	* *	45	* *
Stafford	78	* * *	* *	80	* *	69	* *	77	* *
Stanton	36	* * *	* *	38	* *	36	* *	38	* *
Stevens	121	5.8 (9)	7.4 (48)	196	5.1 (4)	112	2.7 (5)	197	78.7 (11)
Sumner	395	11.6 (65)	6.8 (42)	407	12.5 (63)	344	6.7 (61)	406	71.7 (34)
Thomas	141	9.9 (61)	4.3 (15)	141	9.2 (39)	128	4.7 (42)	141	81.6 (5)
Trego	35	* * *	* *	37	* *	32	* *	37	* *

(1) Reporting period is January 1 through December 31.  
(2) Excludes records with unknown data or errors.  
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.  
(4) <2500 g.  
(5) >4000 g.  
(6) <37 weeks gestation.  
(7) 37 or more weeks gestation and <2500 g.  
\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 8A**  
**2009 - 2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Infant Health Indicators by Clinic**  
**Combined 3 Years**

<b>Comparison of Infant Health Indicators</b> <sup>(2, 3)</sup>										
<b>Clinic</b>	<b>Birthweight</b>				<b>Preterm</b> (6)		<b>Full Term LBW</b> (7)		<b>Ever Breastfed</b>	
	<b>Low</b> (4)		<b>High</b> (5)		<b>Number</b>	<b>%</b> (Rank)	<b>Number</b>	<b>%</b> (Rank)	<b>Number</b>	<b>%</b> (Rank)
	<b>Number</b>	<b>%</b> (Rank)	<b>%</b> (Rank)	<b>Number</b>						
Wabaunsee	58	* * * *	* * * *	58	* *	57	* *	58	* *	
Washington	54	* * * *	* * * *	57	* *	48	* *	57	* *	
Wichita	38	* * * *	* * * *	38	* *	34	* *	38	* *	
Wilson	239	6.3 (14)	7.5 (49)	242	10.7 (57)	212	3.3 (9)	241	71.0 (37)	
Woodson	37	* * * *	* * * *	40	* *	35	* *	40	* *	
Wyandotte	5,426	8.1 (44)	6.0 (30)	5,594	9.2 (39)	4,902	4.6 (41)	5,581	71.4 (36)	
Unknown Clinics	47	* * * *	* * * *	47	* *	47	* *	46	* *	
<b>Kansas</b>	<b>55,067</b>	<b>7.6</b>	<b>6.4</b>	<b>56,409</b>	<b>8.6</b>	<b>50,073</b>	<b>4.4</b>	<b>56,310</b>	<b>72.2</b>	

(1) Reporting period is January 1 through December 31.  
(2) Excludes records with unknown data or errors.  
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.  
(4) <2500 g.  
(5) >4000 g.  
(6) <37 weeks gestation.  
(7) 37 or more weeks gestation and <2500 g.  
\* Percentages are not calculated if <100 records are available for analysis after exclusions.

