

**Table 3B**  
**2008 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Racial and Ethnic Distribution by County**

<b>Comparison of Racial and Ethnic Distribution</b> <sup>(2)</sup>								
<b>County</b>	<b>Number Included in Analysis</b>	<b>White, Not Hispanic %</b>	<b>Black, Not Hispanic %</b>	<b>Hispanic %</b>	<b>American Indian/Alaska Native %</b>	<b>Asian/Pacific Islander %</b>	<b>Multiple Races %</b>	<b>All Other/Unknown %</b>
Allen	112	92.0	5.4	1.8	0.9	0.0	0.0	0.0
Anderson	50	*	*	*	*	*	*	*
Atchison	118	89.8	5.9	1.7	0.0	0.0	2.5	0.0
Barber	39	*	*	*	*	*	*	*
Barton	279	70.6	2.5	26.2	0.4	0.0	0.4	0.0
Bourbon	135	91.9	3.0	2.2	0.0	0.7	2.2	0.0
Brown	110	76.4	2.7	3.6	12.7	0.0	4.5	0.0
Butler	286	92.7	1.0	4.2	0.0	0.0	2.1	0.0
Chase	15	*	*	*	*	*	*	*
Chautauqua	32	*	*	*	*	*	*	*
Cherokee	219	90.9	1.4	4.6	1.4	0.5	1.4	0.0
Clark	4	*	*	*	*	*	*	*
Clay	67	*	*	*	*	*	*	*
Cloud	72	*	*	*	*	*	*	*
Coffey	24	*	*	*	*	*	*	*
Comanche	6	*	*	*	*	*	*	*
Cowley	356	76.1	2.5	18.0	0.8	2.0	0.6	0.0
Crawford	382	82.7	3.9	11.0	0.0	1.3	1.0	0.0
Decatur	9	*	*	*	*	*	*	*
Dickinson	105	94.3	0.0	4.8	1.0	0.0	0.0	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 3B**  
**2008 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Racial and Ethnic Distribution by County**

<b>Comparison of Racial and Ethnic Distribution</b> <sup>(2)</sup>								
County	Number Included in Analysis	White, Not Hispanic  %	Black, Not Hispanic  %	Hispanic  %	American Indian/Alaska Native  %	Asian/Pacific Islander  %	Multiple Races  %	All Other/Unknown  %
Doniphan	43	*	*	*	*	*	*	*
Douglas	463	65.7	6.0	10.2	10.2	5.2	2.8	0.0
Edwards	27	*	*	*	*	*	*	*
Elk	13	*	*	*	*	*	*	*
Ellis	143	83.9	2.1	14.0	0.0	0.0	0.0	0.0
Ellsworth	28	*	*	*	*	*	*	*
Finney	688	30.5	0.9	65.7	0.0	2.5	0.4	0.0
Ford	646	22.1	1.2	76.3	0.2	0.2	0.0	0.0
Franklin	166	89.8	2.4	4.2	1.8	0.0	1.8	0.0
Geary	340	53.5	23.5	15.0	0.0	2.9	5.0	0.0
Gove	7	*	*	*	*	*	*	*
Graham	14	*	*	*	*	*	*	*
Grant	68	*	*	*	*	*	*	*
Gray	57	*	*	*	*	*	*	*
Greeley	10	*	*	*	*	*	*	*
Greenwood	44	*	*	*	*	*	*	*
Hamilton	18	*	*	*	*	*	*	*
Harper	51	*	*	*	*	*	*	*
Harvey	251	69.3	1.6	27.5	0.4	0.0	1.2	0.0
Haskell	24	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 3B**  
**2008 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Racial and Ethnic Distribution by County**

<b>Comparison of Racial and Ethnic Distribution</b> <sup>(2)</sup>								
<b>County</b>	<b>Number Included in Analysis</b>	<b>White, Not Hispanic</b>	<b>Black, Not Hispanic</b>	<b>Hispanic</b>	<b>American Indian/Alaska Native</b>	<b>Asian/Pacific Islander</b>	<b>Multiple Races</b>	<b>All Other/Unknown</b>
		%	%	%	%	%	%	%
Hodgeman	12	*	*	*	*	*	*	*
Jackson	70	*	*	*	*	*	*	*
Jefferson	65	*	*	*	*	*	*	*
Jewell	14	*	*	*	*	*	*	*
Johnson	1,844	45.7	12.7	36.3	0.6	2.9	1.7	0.0
Kearny	21	*	*	*	*	*	*	*
Kingman	57	*	*	*	*	*	*	*
Kiowa	9	*	*	*	*	*	*	*
Labette	166	88.0	7.2	4.2	0.6	0.0	0.0	0.0
Lane	13	*	*	*	*	*	*	*
Leavenworth	356	66.0	17.4	8.7	0.6	2.5	4.8	0.0
Lincoln	22	*	*	*	*	*	*	*
Linn	54	*	*	*	*	*	*	*
Logan	13	*	*	*	*	*	*	*
Lyon	343	58.9	5.0	34.7	0.0	0.9	0.6	0.0
Marion	145	89.7	2.1	7.6	0.7	0.0	0.0	0.0
Marshall	46	*	*	*	*	*	*	*
Mcperson	63	*	*	*	*	*	*	*
Meade	49	*	*	*	*	*	*	*
Miami	136	90.4	2.2	4.4	0.0	1.5	1.5	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 3B**  
**2008 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Racial and Ethnic Distribution by County**

<b>Comparison of Racial and Ethnic Distribution</b> <sup>(2)</sup>								
County	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Mitchell	43	*	*	*	*	*	*	*
Montgomery	330	80.0	7.9	8.2	1.8	1.2	0.9	0.0
Morris	32	*	*	*	*	*	*	*
Nemaha	37	*	*	*	*	*	*	*
Neosho	153	91.5	2.6	3.9	0.7	0.7	0.7	0.0
Ness	8	*	*	*	*	*	*	*
Norton	26	*	*	*	*	*	*	*
Osage	89	*	*	*	*	*	*	*
Osborne	19	*	*	*	*	*	*	*
Ottawa	22	*	*	*	*	*	*	*
Pawnee	43	*	*	*	*	*	*	*
Phillips	33	*	*	*	*	*	*	*
Pottawatom	40	*	*	*	*	*	*	*
Pratt	68	*	*	*	*	*	*	*
Rawlins	10	*	*	*	*	*	*	*
Reno	494	81.0	1.8	14.4	0.8	0.8	1.2	0.0
Republic	33	*	*	*	*	*	*	*
Rice	70	*	*	*	*	*	*	*
Riley	826	71.2	12.5	11.1	1.0	4.1	0.1	0.0
Rooks	36	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 3B**  
**2008 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Racial and Ethnic Distribution by County**

<b>Comparison of Racial and Ethnic Distribution</b> <sup>(2)</sup>								
County	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Rush	12	*	*	*	*	*	*	*
Russell	40	*	*	*	*	*	*	*
Saline	493	73.8	4.1	18.3	0.2	2.2	1.4	0.0
Scott	36	*	*	*	*	*	*	*
Sedgwick	4,149	45.0	18.7	29.5	1.5	3.8	1.5	0.0
Seward	496	16.1	1.0	77.8	3.4	1.2	0.4	0.0
Shawnee	1,351	54.4	15.4	25.8	1.8	0.8	1.9	0.0
Sheridan	7	*	*	*	*	*	*	*
Sherman	70	*	*	*	*	*	*	*
Smith	17	*	*	*	*	*	*	*
Stafford	40	*	*	*	*	*	*	*
Stanton	23	*	*	*	*	*	*	*
Stevens	66	*	*	*	*	*	*	*
Sumner	158	90.5	0.0	7.6	1.3	0.6	0.0	0.0
Thomas	50	*	*	*	*	*	*	*
Trego	14	*	*	*	*	*	*	*
Wabaunsee	18	*	*	*	*	*	*	*
Wallace	1	*	*	*	*	*	*	*
Washington	25	*	*	*	*	*	*	*
Wichita	11	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 3B**  
**2008 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Racial and Ethnic Distribution by County**

<b>Comparison of Racial and Ethnic Distribution</b> <sup>(2)</sup>								
<b>County</b>	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Wilson	81	*	*	*	*	*	*	*
Woodson	2	*	*	*	*	*	*	*
Wyandotte	2,112	27.3	31.0	39.3	0.2	1.8	0.4	0.0
<b>Kansas</b>	<b>20,703</b>	<b>56.9</b>	<b>11.3</b>	<b>27.3</b>	<b>1.2</b>	<b>2.0</b>	<b>1.2</b>	<b>0.0</b>
<b>Nation (Prior Year)</b>	<b>1,252,728</b>	<b>41.7</b>	<b>23.3</b>	<b>27.9</b>	<b>1.1</b>	<b>3.2</b>	<b>1.0</b>	<b>1.8</b>

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 4B**  
**2008 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Age Distribution by County**

<b>Comparison of Age Distribution</b> <sup>(2)</sup>								
<b>County</b>	<b>Number Included in Analysis</b>	<b>&lt; 15 Years %</b>	<b>15 - 17 Years %</b>	<b>18 - 19 Years %</b>	<b>20 - 29 Years %</b>	<b>30 - 39 Years %</b>	<b>&gt;= 40 Years %</b>	<b>Unknown %</b>
Allen	112	0.0	6.3	15.2	63.4	13.4	0.9	0.9
Anderson	50	*	*	*	*	*	*	*
Atchison	118	0.0	6.8	16.9	62.7	11.9	1.7	0.0
Barber	39	*	*	*	*	*	*	*
Barton	279	0.0	6.5	19.0	57.3	16.8	0.4	0.0
Bourbon	135	0.0	4.4	20.0	65.9	9.6	0.0	0.0
Brown	110	0.0	7.3	16.4	65.5	10.0	0.9	0.0
Butler	286	0.3	7.0	16.1	61.9	14.0	0.7	0.0
Chase	15	*	*	*	*	*	*	*
Chautauqua	32	*	*	*	*	*	*	*
Cherokee	219	0.5	9.1	17.8	56.6	14.6	0.9	0.5
Clark	4	*	*	*	*	*	*	*
Clay	67	*	*	*	*	*	*	*
Cloud	72	*	*	*	*	*	*	*
Coffey	24	*	*	*	*	*	*	*
Comanche	6	*	*	*	*	*	*	*
Cowley	356	0.6	9.0	15.4	61.2	13.5	0.3	0.0
Crawford	382	0.3	8.9	13.6	63.6	13.1	0.5	0.0
Decatur	9	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 4B**  
**2008 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Age Distribution by County**

<b>Comparison of Age Distribution</b> <sup>(2)</sup>								
<b>County</b>	<b>Number Included in Analysis</b>	<b>&lt; 15 Years %</b>	<b>15 - 17 Years %</b>	<b>18 - 19 Years %</b>	<b>20 - 29 Years %</b>	<b>30 - 39 Years %</b>	<b>&gt;= 40 Years %</b>	<b>Unknown %</b>
Dickinson	105	1.0	2.9	14.3	63.8	18.1	0.0	0.0
Doniphan	43	*	*	*	*	*	*	*
Douglas	463	0.0	4.1	13.6	63.5	17.1	1.7	0.0
Edwards	27	*	*	*	*	*	*	*
Elk	13	*	*	*	*	*	*	*
Ellis	143	0.0	4.2	11.2	69.9	11.9	2.8	0.0
Ellsworth	28	*	*	*	*	*	*	*
Finney	688	0.1	10.5	16.0	53.8	18.2	1.2	0.3
Ford	646	0.3	7.3	12.8	54.8	24.0	0.8	0.0
Franklin	166	0.6	7.2	18.1	58.4	15.1	0.0	0.6
Geary	340	0.6	5.6	17.9	65.0	10.3	0.6	0.0
Gove	7	*	*	*	*	*	*	*
Graham	14	*	*	*	*	*	*	*
Grant	68	*	*	*	*	*	*	*
Gray	57	*	*	*	*	*	*	*
Greeley	10	*	*	*	*	*	*	*
Greenwood	44	*	*	*	*	*	*	*
Hamilton	18	*	*	*	*	*	*	*
Harper	51	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.





**Table 4B**  
**2008 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Age Distribution by County**

<b>Comparison of Age Distribution</b> <sup>(2)</sup>								
<b>County</b>	<b>Number Included in Analysis</b>	<b>&lt; 15 Years %</b>	<b>15 - 17 Years %</b>	<b>18 - 19 Years %</b>	<b>20 - 29 Years %</b>	<b>30 - 39 Years %</b>	<b>&gt;= 40 Years %</b>	<b>Unknown %</b>
Harvey	251	0.0	5.6	8.0	66.9	16.7	2.8	0.0
Haskell	24	*	*	*	*	*	*	*
Hodgeman	12	*	*	*	*	*	*	*
Jackson	70	*	*	*	*	*	*	*
Jefferson	65	*	*	*	*	*	*	*
Jewell	14	*	*	*	*	*	*	*
Johnson	1,844	0.3	4.6	11.7	60.7	21.4	1.2	0.1
Kearny	21	*	*	*	*	*	*	*
Kingman	57	*	*	*	*	*	*	*
Kiowa	9	*	*	*	*	*	*	*
Labette	166	0.0	4.8	21.1	65.7	7.8	0.6	0.0
Lane	13	*	*	*	*	*	*	*
Leavenworth	356	0.0	5.9	19.4	61.5	11.0	2.2	0.0
Lincoln	22	*	*	*	*	*	*	*
Linn	54	*	*	*	*	*	*	*
Logan	13	*	*	*	*	*	*	*
Lyon	343	0.3	9.0	13.7	59.8	13.7	3.5	0.0
Marion	145	0.7	3.4	15.9	62.1	17.2	0.7	0.0
Marshall	46	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 4B**  
**2008 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Age Distribution by County**

<b>Comparison of Age Distribution</b> <sup>(2)</sup>								
<b>County</b>	<b>Number Included in Analysis</b>	<b>&lt; 15 Years %</b>	<b>15 - 17 Years %</b>	<b>18 - 19 Years %</b>	<b>20 - 29 Years %</b>	<b>30 - 39 Years %</b>	<b>&gt;= 40 Years %</b>	<b>Unknown %</b>
Mcpherson	63	*	*	*	*	*	*	*
Meade	49	*	*	*	*	*	*	*
Miami	136	0.0	7.4	11.0	67.6	14.0	0.0	0.0
Mitchell	43	*	*	*	*	*	*	*
Montgomery	330	0.0	7.3	18.2	63.0	10.6	0.6	0.3
Morris	32	*	*	*	*	*	*	*
Nemaha	37	*	*	*	*	*	*	*
Neosho	153	0.7	3.9	20.9	60.8	13.1	0.7	0.0
Ness	8	*	*	*	*	*	*	*
Norton	26	*	*	*	*	*	*	*
Osage	89	*	*	*	*	*	*	*
Osborne	19	*	*	*	*	*	*	*
Ottawa	22	*	*	*	*	*	*	*
Pawnee	43	*	*	*	*	*	*	*
Phillips	33	*	*	*	*	*	*	*
Pottawatom	40	*	*	*	*	*	*	*
Pratt	68	*	*	*	*	*	*	*
Rawlins	10	*	*	*	*	*	*	*
Reno	494	0.2	5.9	15.2	66.2	11.5	1.0	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 4B**  
**2008 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Age Distribution by County**

Comparison of Age Distribution <sup>(2)</sup>								
County	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Republic	33	*	*	*	*	*	*	*
Rice	70	*	*	*	*	*	*	*
Riley	826	0.0	1.5	13.0	73.8	11.1	0.6	0.0
Rooks	36	*	*	*	*	*	*	*
Rush	12	*	*	*	*	*	*	*
Russell	40	*	*	*	*	*	*	*
Saline	493	0.0	7.5	16.6	61.5	13.6	0.8	0.0
Scott	36	*	*	*	*	*	*	*
Sedgwick	4,149	0.3	6.7	14.3	61.4	15.9	1.4	0.0
Seward	496	0.4	6.7	15.1	57.3	19.2	1.4	0.0
Shawnee	1,351	0.2	6.7	13.8	61.7	16.4	1.2	0.0
Sheridan	7	*	*	*	*	*	*	*
Sherman	70	*	*	*	*	*	*	*
Smith	17	*	*	*	*	*	*	*
Stafford	40	*	*	*	*	*	*	*
Stanton	23	*	*	*	*	*	*	*
Stevens	66	*	*	*	*	*	*	*
Sumner	158	0.0	5.7	17.1	63.3	13.9	0.0	0.0
Thomas	50	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 4B**  
**2008 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Age Distribution by County**

<b>Comparison of Age Distribution</b> <sup>(2)</sup>								
<b>County</b>	<b>Number Included in Analysis</b>	<b>&lt; 15 Years %</b>	<b>15 - 17 Years %</b>	<b>18 - 19 Years %</b>	<b>20 - 29 Years %</b>	<b>30 - 39 Years %</b>	<b>&gt;= 40 Years %</b>	<b>Unknown %</b>
Trego	14	*	*	*	*	*	*	*
Wabaunsee	18	*	*	*	*	*	*	*
Wallace	1	*	*	*	*	*	*	*
Washington	25	*	*	*	*	*	*	*
Wichita	11	*	*	*	*	*	*	*
Wilson	81	*	*	*	*	*	*	*
Woodson	2	*	*	*	*	*	*	*
Wyandotte	2,112	0.7	8.6	13.4	58.0	18.1	1.2	0.0
<b>Kansas</b>	<b>20,703</b>	<b>0.3</b>	<b>6.5</b>	<b>14.6</b>	<b>61.6</b>	<b>15.9</b>	<b>1.2</b>	<b>0.0</b>
<b>Nation (Prior Year)</b>	<b>1,252,728</b>	<b>0.4</b>	<b>6.5</b>	<b>12.7</b>	<b>59.9</b>	<b>19.0</b>	<b>1.4</b>	<b>0.1</b>

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 5B**  
**2008 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Maternal Health Indicators by County**

<b>Comparison of Maternal Health Indicators</b> <sup>(2, 3)</sup>														
<b>County</b>	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	%	%	Number	%	%	Number	%	Number	%	Number	%	Number	%
	(Rank)	(Rank)		(Rank)	(Rank)		(Rank)		(Rank)		(Rank)		(Rank)	
Allen	112	9.8 (9)	49.1 (25)	108	21.3 (7)	49.1 (26)	20	* *	35	* *	97	* *	97	* *
Anderson	50	* *	* *	47	* *	* *	9	* *	8	* *	38	* *	38	* *
Atchison	116	13.8 (28)	44.0 (7)	106	27.4 (29)	43.4 (11)	20	* *	30	* *	90	* *	90	* *
Barber	39	* *	* *	33	* *	* *	2	* *	12	* *	28	* *	28	* *
Barton	274	10.6 (13)	45.6 (14)	260	27.3 (28)	41.5 (6)	27	* *	58	* *	219	4.1 (20)	219	7.3(20)
Bourbon	135	9.6 (8)	50.4 (30)	122	26.2 (26)	39.3 (2)	8	* *	31	* *	108	4.6 (23)	108	8.3(23)
Brown	110	6.4 (2)	52.7 (33)	99	* *	* *	19	* *	22	* *	95	* *	95	* *
Butler	283	12.7 (25)	41.3 (3)	269	21.6 (9)	46.8 (21)	32	* *	89	* *	208	2.4 (6)	208	9.6(26)
Chase	14	* *	* *	11	* *	* *	3	* *	2	* *	12	* *	12	* *
Chautauqua	32	* *	* *	27	* *	* *	2	* *	5	* *	25	* *	25	* *
Cherokee	218	12.8 (26)	46.3 (17)	196	22.4 (16)	43.9 (12)	24	* *	39	* *	168	4.2 (22)	168	5.4(7)
Clark	3	* *	* *	3	* *	* *	0	* *	2	* *	4	* *	4	* *
Clay	67	* *	* *	61	* *	* *	13	* *	13	* *	52	* *	52	* *
Cloud	72	* *	* *	65	* *	* *	5	* *	16	* *	55	* *	56	* *
Coffey	24	* *	* *	24	* *	* *	2	* *	7	* *	16	* *	16	* *
Comanche	6	* *	* *	5	* *	* *	1	* *	3	* *	3	* *	3	* *
Cowley	355	9.9 (10)	49.9 (29)	332	21.7 (11)	45.8 (18)	40	* *	58	* *	286	2.8 (10)	286	7.7(21)

- (1) Reporting period is January 1 through December 31.
  - (2) Excludes records with unknown data or errors.
  - (3) Rank compares one county's rate to other counties. Rank 1 = best rate.
  - (4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).
  - (5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).
  - (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- \* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 5B**  
**2008 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Maternal Health Indicators by County**

Comparison of Maternal Health Indicators <sup>(2, 3)</sup>														
County	Prepregnancy BMI <sup>(4)</sup>			Weight Gain <sup>(5)</sup>			Anemia (Low Hb/Hct) <sup>(6)</sup>				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Crawford	378	10.1 (11)	49.7 (28)	337	23.4 (21)	46.6 (20)	67	* *	72	* *	301	4.7 (24)	300	5.7(9)
Decatur	9	* *	* *	9	* *	* *	0	* *	3	* *	7	* *	7	* *
Dickinson	104	7.7 (3)	39.4 (1)	96	* *	* *	17	* *	24	* *	80	* *	80	* *
Doniphan	43	* *	* *	35	* *	* *	8	* *	6	* *	28	* *	28	* *
Douglas	459	13.1 (27)	41.8 (4)	409	20.3 (4)	53.1 (29)	67	* *	117	27.4 (5)	347	3.2 (15)	346	4.0(2)
Edwards	27	* *	* *	21	* *	* *	4	* *	7	* *	15	* *	15	* *
Elk	13	* *	* *	12	* *	* *	0	* *	2	* *	9	* *	9	* *
Ellis	142	10.6 (13)	46.5 (18)	127	22.8 (19)	42.5 (9)	33	* *	41	* *	101	3.0 (12)	101	6.9(19)
Ellsworth	28	* *	* *	28	* *	* *	3	* *	14	* *	20	* *	20	* *
Finney	682	14.8 (32)	40.6 (2)	640	28.3 (30)	40.0 (4)	99	* *	98	* *	570	6.7 (29)	570	3.5(1)
Ford	637	5.7 (1)	49.5 (27)	580	23.4 (21)	45.2 (16)	90	* *	78	* *	532	2.6 (8)	532	4.1(3)
Franklin	164	11.0 (17)	45.1 (11)	145	16.6 (1)	42.8 (10)	13	* *	32	* *	119	2.5 (7)	119	8.4(24)
Geary	336	11.0 (17)	44.3 (8)	283	21.2 (6)	48.1 (24)	55	* *	50	* *	256	5.1 (26)	256	9.8(27)
Gove	7	* *	* *	7	* *	* *	2	* *	0	* *	7	* *	7	* *
Graham	14	* *	* *	11	* *	* *	4	* *	1	* *	12	* *	12	* *
Grant	65	* *	* *	61	* *	* *	10	* *	20	* *	53	* *	53	* *
Gray	57	* *	* *	48	* *	* *	7	* *	19	* *	41	* *	41	* *

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one county's rate to other counties. Rank 1 = best rate.
- (4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).
- (5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).
- (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- \* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 5B**  
**2008 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Maternal Health Indicators by County**

Comparison of Maternal Health Indicators <sup>(2, 3)</sup>														
County	Prepregnancy BMI <sup>(4)</sup>			Weight Gain <sup>(5)</sup>			Anemia (Low Hb/Hct) <sup>(6)</sup>				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
		Number	% (Rank)		% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Greeley	10	* *	* *	10	* *	* *	2	* *	2	* *	10	* *	10	* *
Greenwood	44	* *	* *	44	* *	* *	3	* *	7	* *	40	* *	40	* *
Hamilton	18	* *	* *	14	* *	* *	2	* *	5	* *	14	* *	14	* *
Harper	51	* *	* *	44	* *	* *	5	* *	8	* *	38	* *	37	* *
Harvey	250	10.8 (15)	44.4 (9)	228	27.2 (27)	41.7 (8)	30	* *	46	* *	192	3.1 (14)	190	6.8(17)
Haskell	24	* *	* *	22	* *	* *	5	* *	8	* *	19	* *	19	* *
Hodgeman	12	* *	* *	12	* *	* *	2	* *	2	* *	9	* *	9	* *
Jackson	70	* *	* *	67	* *	* *	8	* *	12	* *	61	* *	61	* *
Jefferson	65	* *	* *	60	* *	* *	10	* *	18	* *	53	* *	53	* *
Jewell	14	* *	* *	14	* *	* *	1	* *	2	* *	13	* *	13	* *
Johnson	1,819	10.8 (15)	44.6 (10)	1,684	21.6 (9)	44.4 (14)	314	25.8 (2)	400	16.5 (1)	1,428	4.0 (19)	1,428	4.3(4)
Kearny	21	* *	* *	19	* *	* *	4	* *	7	* *	19	* *	19	* *
Kingman	57	* *	* *	51	* *	* *	7	* *	12	* *	41	* *	41	* *
Kiowa	9	* *	* *	7	* *	* *	2	* *	1	* *	8	* *	8	* *
Labette	165	9.1 (5)	50.9 (31)	150	22.7 (17)	53.3 (30)	26	* *	36	* *	132	3.0 (12)	132	11.4(29)
Lane	12	* *	* *	10	* *	* *	4	* *	1	* *	8	* *	8	* *
Leavenwort	352	10.5 (12)	48.3 (21)	323	21.7 (11)	49.8 (27)	55	* *	99	* *	253	1.6 (2)	254	5.1(5)

(1) Reporting period is January 1 through December 31.  
(2) Excludes records with unknown data or errors.  
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.  
(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).  
(5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).  
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.  
\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 5B**  
**2008 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Maternal Health Indicators by County**

Comparison of Maternal Health Indicators <sup>(2, 3)</sup>														
County	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight	Over- weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Lincoln	22	* *	* *	22	* *	* *	1	* *	5	* *	16	* *	16	* *
Linn	52	* *	* *	49	* *	* *	3	* *	10	* *	45	* *	45	* *
Logan	13	* *	* *	10	* *	* *	2	* *	3	* *	9	* *	9	* *
Lyon	341	11.4 (21)	52.5 (32)	312	22.8 (19)	52.2 (28)	48	* *	64	* *	266	2.3 (5)	266	9.8(27)
Marion	145	14.5 (30)	48.3 (21)	138	23.9 (24)	37.7 (1)	30	* *	30	* *	120	0.8 (1)	120	5.8(10)
Marshall	46	* *	* *	40	* *	* *	6	* *	6	* *	31	* *	31	* *
Mcperson	62	* *	* *	55	* *	* *	11	* *	13	* *	46	* *	46	* *
Meade	49	* *	* *	44	* *	* *	7	* *	12	* *	39	* *	39	* *
Miami	135	14.1 (29)	48.1 (20)	130	19.2 (2)	53.8 (31)	32	* *	42	* *	104	4.8 (25)	104	5.8(10)
Mitchell	43	* *	* *	43	* *	* *	8	* *	13	* *	38	* *	38	* *
Montgomery	328	16.8 (33)	45.4 (13)	298	30.2 (31)	39.6 (3)	52	* *	94	* *	258	1.9 (4)	258	5.8(10)
Morris	32	* *	* *	28	* *	* *	5	* *	6	* *	25	* *	25	* *
Nemaha	36	* *	* *	36	* *	* *	9	* *	16	* *	31	* *	31	* *
Neosho	151	14.6 (31)	48.3 (21)	142	19.7 (3)	40.1 (5)	23	* *	25	* *	117	1.7 (3)	117	6.0(14)
Ness	8	* *	* *	7	* *	* *	0	* *	5	* *	4	* *	4	* *
Norton	26	* *	* *	22	* *	* *	2	* *	7	* *	17	* *	17	* *
Osage	88	* *	* *	75	* *	* *	13	* *	22	* *	68	* *	68	* *

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one county's rate to other counties. Rank 1 = best rate.
- (4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).
- (5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).
- (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- \* Percentages are not calculated if <100 records are available for analysis after exclusions.





**Table 5B**  
**2008 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Maternal Health Indicators by County**

Comparison of Maternal Health Indicators <sup>(2, 3)</sup>														
County	Prepregnancy BMI <sup>(4)</sup>			Weight Gain <sup>(5)</sup>			Anemia (Low Hb/Hct) <sup>(6)</sup>				Gestational Diabetes During Pregnancy	Hypertension During Pregnancy		
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
							Number	% (Rank)	% (Rank)	Number			% (Rank)	Number
Osborne	19	* * * *	* * * *	18	* * * *	* * * *	1	* * *	3	* * *	17	* * *	17	* * *
Ottawa	22	* * * *	* * * *	20	* * * *	* * * *	3	* * *	7	* * *	16	* * *	16	* * *
Pawnee	43	* * * *	* * * *	42	* * * *	* * * *	4	* * *	11	* * *	36	* * *	36	* * *
Phillips	33	* * * *	* * * *	31	* * * *	* * * *	1	* * *	5	* * *	24	* * *	24	* * *
Pottawatom	40	* * * *	* * * *	35	* * * *	* * * *	7	* * *	17	* * *	31	* * *	31	* * *
Pratt	68	* * * *	* * * *	59	* * * *	* * * *	8	* * *	21	* * *	46	* * *	46	* * *
Rawlins	10	* * * *	* * * *	9	* * * *	* * * *	1	* * *	4	* * *	8	* * *	8	* * *
Reno	493	12.0 (24)	45.2 (12)	462	26.0 (25)	41.6 (7)	66	* * *	102	18.6 (2)	376	3.5 (16)	377	7.7(21)
Republic	33	* * * *	* * * *	33	* * * *	* * * *	5	* * *	5	* * *	30	* * *	30	* * *
Rice	70	* * * *	* * * *	60	* * * *	* * * *	9	* * *	15	* * *	53	* * *	53	* * *
Riley	820	9.4 (6)	42.4 (5)	701	21.3 (7)	48.8 (25)	106	27.4 (3)	175	34.9 (8)	566	3.7 (17)	566	5.8(10)
Rooks	36	* * * *	* * * *	35	* * * *	* * * *	8	* * *	6	* * *	30	* * *	30	* * *
Rush	12	* * * *	* * * *	11	* * * *	* * * *	2	* * *	2	* * *	10	* * *	10	* * *
Russell	40	* * * *	* * * *	37	* * * *	* * * *	5	* * *	5	* * *	31	* * *	31	* * *
Saline	490	11.4 (21)	47.6 (19)	458	21.0 (5)	44.3 (13)	109	25.7 (1)	126	25.4 (4)	413	4.1 (20)	413	6.8(17)
Scott	36	* * * *	* * * *	34	* * * *	* * * *	5	* * *	8	* * *	30	* * *	30	* * *
Sedgwick	4,121	11.2 (20)	45.8 (16)	3,907	22.7 (17)	45.6 (17)	822	33.8 (5)	1,156	25.2 (3)	3,347	3.7 (17)	3,346	6.3(15)

(1) Reporting period is January 1 through December 31.  
(2) Excludes records with unknown data or errors.  
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.  
(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).  
(5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).  
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.  
\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 5B**  
**2008 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Maternal Health Indicators by County**

<b>Comparison of Maternal Health Indicators</b> <sup>(2, 3)</sup>														
<b>County</b>	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Seward	493	8.7 (4)	45.6 (14)	450	22.0 (15)	44.4 (14)	82	* *	81	* *	393	5.3 (28)	391	6.6 (16)
Shawnee	1,337	11.0 (17)	49.4 (26)	1,217	21.9 (13)	47.7 (23)	139	35.3 (6)	233	28.8 (7)	1,063	5.2 (27)	1,066	5.4 (7)
Sheridan	7	* *	* *	5	* *	* *	3	* *	1	* *	5	* *	5	* *
Sherman	70	* *	* *	63	* *	* *	6	* *	17	* *	53	* *	53	* *
Smith	17	* *	* *	14	* *	* *	3	* *	8	* *	15	* *	15	* *
Stafford	39	* *	* *	33	* *	* *	3	* *	6	* *	27	* *	27	* *
Stanton	23	* *	* *	21	* *	* *	3	* *	6	* *	21	* *	21	* *
Stevens	66	* *	* *	55	* *	* *	8	* *	17	* *	44	* *	44	* *
Sumner	158	11.4 (21)	43.7 (6)	148	23.6 (23)	45.9 (19)	25	* *	43	* *	117	2.6 (8)	117	9.4 (25)
Thomas	48	* *	* *	41	* *	* *	5	* *	10	* *	37	* *	37	* *
Trego	14	* *	* *	13	* *	* *	0	* *	2	* *	12	* *	12	* *
Wabaunsee	18	* *	* *	18	* *	* *	0	* *	6	* *	15	* *	15	* *
Wallace	1	* *	* *	1	* *	* *	0	* *	1	* *	0	* *	0	* *
Washington	25	* *	* *	24	* *	* *	2	* *	5	* *	20	* *	19	* *
Wichita	11	* *	* *	11	* *	* *	1	* *	4	* *	10	* *	10	* *
Wilson	79	* *	* *	77	* *	* *	13	* *	14	* *	66	* *	66	* *
Woodson	2	* *	* *	2	* *	* *	0	* *	0	* *	2	* *	2	* *

- (1) Reporting period is January 1 through December 31.  
(2) Excludes records with unknown data or errors.  
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.  
(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).  
(5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).  
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.  
\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 5B**  
**2008 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Maternal Health Indicators by County**

<b>Comparison of Maternal Health Indicators</b> <sup>(2, 3)</sup>														
<b>County</b>	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	%	%	Number	%	%	Number	%	Number	%	Number	%	Number	%
	(Rank)	(Rank)		(Rank)	(Rank)		(Rank)	(Rank)	(Rank)	(Rank)	(Rank)	(Rank)	(Rank)	
Wyandotte	2,077	9.5 (7)	48.9 (24)	1,924	21.9 (13)	46.9 (22)	320	29.7 (4)	358	28.5 (6)	1,688	2.8 (10)	1,689	5.2(6)
<b>Kansas</b>	<b>20,516</b>	<b>10.9</b>	<b>46.6</b>	<b>18,896</b>	<b>22.9</b>	<b>45.6</b>	<b>3,213</b>	<b>29.3</b>	<b>4,533</b>	<b>24.0</b>	<b>16,226</b>	<b>3.7</b>	<b>16,224</b>	<b>6.2</b>
<b>Nation (Prior Year)</b>	<b>1,175,574</b>	<b>10.9</b>	<b>44.5</b>	<b>922,658</b>	<b>25.0</b>	<b>42.8</b>	<b>180,176</b>	<b>33.5</b>	<b>414,735</b>	<b>29.8</b>	<b>230,547</b>	<b>3.9</b>	<b>280,574</b>	<b>4.3</b>

(1) Reporting period is January 1 through December 31.  
(2) Excludes records with unknown data or errors.  
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.  
(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).  
(5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).  
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.  
\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 6B**  
**2008 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Maternal Behavioral Indicators by County**

<b>Comparison of Maternal Behavioral Indicators</b> <sup>(2, 3)</sup>														
<b>County</b>	Medical Care				WIC Enrollment			Multivitamin Consumption						
	1st Trimester		No Care		1st Trimester			Prenatal (4)		During Pregnancy (5)				
	Number	% (Rank)	%	% (Rank)	Number	%	% (Rank)	Number	%	% (Rank)				
Allen	102	95.1 (1)	0.0	(1)	96	*	*	69	*	*	64	*	*	
Anderson	48	*	*	*	*	40	*	*	38	*	*	29	*	*
Atchison	104	87.5 (6)	2.9	(14)	95	*	*	81	*	*	71	*	*	
Barber	33	*	*	*	*	32	*	*	24	*	*	21	*	*
Barton	254	81.1 (15)	1.2	(9)	217	56.7	(1)	157	19.7	(2)	132	63.6	(3)	
Bourbon	115	91.3 (2)	0.9	(7)	113	48.7	(6)	77	*	*	78	*	*	
Brown	96	*	*	*	*	95	*	*	64	*	*	83	*	*
Butler	254	70.5 (24)	13.4	(28)	193	48.2	(7)	154	12.3	(9)	117	37.6	(20)	
Chase	11	*	*	*	*	12	*	*	9	*	*	8	*	*
Chautauqua	27	*	*	*	*	28	*	*	16	*	*	17	*	*
Cherokee	177	76.3 (17)	9.6	(25)	167	50.3	(3)	115	9.6	(15)	108	48.1	(15)	
Clark	3	*	*	*	*	2	*	*	0	*	*	1	*	*
Clay	59	*	*	*	*	55	*	*	35	*	*	32	*	*
Cloud	64	*	*	*	*	58	*	*	44	*	*	36	*	*
Coffey	24	*	*	*	*	14	*	*	14	*	*	8	*	*
Comanche	5	*	*	*	*	3	*	*	3	*	*	2	*	*
Cowley	314	69.1 (26)	6.4	(22)	269	50.6	(2)	194	10.8	(14)	172	52.3	(12)	
Crawford	330	87.6 (5)	1.2	(9)	312	44.6	(10)	235	8.1	(16)	212	61.8	(4)	
Decatur	7	*	*	*	*	6	*	*	5	*	*	4	*	*
Dickinson	94	*	*	*	*	84	*	*	68	*	*	55	*	*

(1) Reporting period is January 1 through December 31.  
(2) Excludes records with unknown data or errors.  
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.  
(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.  
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.  
\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 6B**  
**2008 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Maternal Behavioral Indicators by County**

<b>Comparison of Maternal Behavioral Indicators</b> <sup>(2, 3)</sup>										
<b>County</b>	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Pregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	%	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Doniphan	36	* *	* *	* *	33	* *	27	* *	26	* *
Douglas	403	70.0 (25)	6.9 (23)		362	42.3 (14)	258	12.4 (7)	223	48.0 (16)
Edwards	18	* *	* *	* *	20	* *	19	* *	13	* *
Elk	11	* *	* *	* *	9	* *	5	* *	4	* *
Ellis	128	85.9 (8)	1.6 (11)		112	33.9 (26)	95	* *	84	* *
Ellsworth	28	* *	* *	* *	16	* *	16	* *	10	* *
Finney	578	66.1 (28)	19.0 (30)		551	37.4 (22)	413	5.8 (19)	357	29.7 (21)
Ford	549	72.7 (23)	14.2 (29)		538	49.1 (5)	347	5.5 (20)	336	49.4 (13)
Franklin	95	* *	* *	* *	129	41.1 (15)	91	* *	91	* *
Geary	274	73.0 (22)	0.7 (5)		284	45.8 (9)	195	15.9 (3)	183	47.0 (18)
Gove	7	* *	* *	* *	5	* *	3	* *	3	* *
Graham	12	* *	* *	* *	12	* *	4	* *	5	* *
Grant	54	* *	* *	* *	51	* *	36	* *	27	* *
Gray	49	* *	* *	* *	43	* *	32	* *	24	* *
Greeley	10	* *	* *	* *	7	* *	7	* *	7	* *
Greenwood	40	* *	* *	* *	36	* *	20	* *	21	* *
Hamilton	15	* *	* *	* *	14	* *	10	* *	11	* *
Harper	41	* *	* *	* *	43	* *	28	* *	29	* *
Harvey	212	73.6 (21)	12.7 (27)		202	48.0 (8)	138	11.6 (12)	128	61.7 (5)
Haskell	21	* *	* *	* *	21	* *	16	* *	15	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 6B**  
**2008 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Maternal Behavioral Indicators by County**

<b>Comparison of Maternal Behavioral Indicators</b> <sup>(2, 3)</sup>										
<b>County</b>	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Pregpregnancy (4)		During Pregpregnancy (5)	
	Number	% (Rank)	%	(Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Hodgeman	12	* *	* *	* *	8	* *	7	* *	4	* *
Jackson	42	* *	* *	* *	63	* *	39	* *	42	* *
Jefferson	52	* *	* *	* *	48	* *	36	* *	33	* *
Jewell	14	* *	* *	* *	12	* *	8	* *	10	* *
Johnson	1,525	62.8 (30)	3.3	(16)	1,406	34.1 (25)	1,022	11.5 (13)	891	60.6 (7)
Kearny	19	* *	* *	* *	17	* *	10	* *	9	* *
Kingman	49	* *	* *	* *	47	* *	39	* *	36	* *
Kiowa	8	* *	* *	* *	9	* *	6	* *	8	* *
Labette	128	85.9 (8)	2.3	(13)	132	40.2 (18)	92	* *	89	* *
Lane	9	* *	* *	* *	10	* *	7	* *	9	* *
Leavenwort	320	84.4 (10)	0.9	(7)	244	36.1 (24)	202	12.4 (7)	151	61.6 (6)
Lincoln	22	* *	* *	* *	15	* *	14	* *	9	* *
Linn	48	* *	* *	* *	43	* *	30	* *	31	* *
Logan	6	* *	* *	* *	10	* *	8	* *	7	* *
Lyon	312	81.4 (14)	0.6	(3)	265	44.2 (12)	193	14.0 (6)	169	58.0 (8)
Marion	138	88.4 (4)	0.7	(5)	114	37.7 (21)	74	* *	67	* *
Marshall	40	* *	* *	* *	34	* *	27	* *	21	* *
Mcpherson	56	* *	* *	* *	53	* *	33	* *	25	* *
Meade	44	* *	* *	* *	35	* *	26	* *	25	* *
Miami	129	79.1 (16)	0.0	(1)	94	* *	74	* *	62	* *

(1) Reporting period is January 1 through December 31.  
(2) Excludes records with unknown data or errors.  
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.  
(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.  
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.  
\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 6B**  
**2008 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Maternal Behavioral Indicators by County**

<b>Comparison of Maternal Behavioral Indicators</b> <sup>(2, 3)</sup>										
<b>County</b>	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	%	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Mitchell	43	* *	* *	* *	34	* *	24	* *	22	* *
Montgomery	254	86.2 (7)	6.3 (21)		257	44.0 (13)	186	14.5 (5)	160	52.5 (11)
Morris	26	* *	* *	* *	26	* *	14	* *	13	* *
Nemaha	36	* *	* *	* *	29	* *	16	* *	15	* *
Neosho	129	89.1 (3)	3.1 (15)		122	38.5 (20)	92	* *	77	* *
Ness	8	* *	* *	* *	3	* *	5	* *	1	* *
Norton	23	* *	* *	* *	15	* *	14	* *	8	* *
Osage	76	* *	* *	* *	65	* *	51	* *	37	* *
Osborne	18	* *	* *	* *	14	* *	6	* *	10	* *
Ottawa	19	* *	* *	* *	16	* *	11	* *	10	* *
Pawnee	41	* *	* *	* *	36	* *	28	* *	25	* *
Phillips	31	* *	* *	* *	25	* *	20	* *	21	* *
Pottawatom	28	* *	* *	* *	33	* *	19	* *	21	* *
Pratt	60	* *	* *	* *	46	* *	41	* *	29	* *
Rawlins	5	* *	* *	* *	9	* *	6	* *	7	* *
Reno	450	83.6 (11)	2.0 (12)		374	40.4 (17)	291	23.7 (1)	247	64.0 (2)
Republic	33	* *	* *	* *	26	* *	13	* *	14	* *
Rice	59	* *	* *	* *	59	* *	42	* *	41	* *
Riley	640	82.5 (13)	0.6 (3)		599	39.1 (19)	472	14.6 (4)	368	47.8 (17)
Rooks	35	* *	* *	* *	30	* *	17	* *	18	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 6B**  
**2008 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Maternal Behavioral Indicators by County**

<b>Comparison of Maternal Behavioral Indicators</b> <sup>(2, 3)</sup>										
<b>County</b>	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	%	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Rush	11	* *	* *	* *	11	* *	8	* *	7	* *
Russell	37	* *	* *	* *	30	* *	23	* *	21	* *
Saline	413	74.3 (18)	3.4	(17)	378	30.4 (27)	271	5.9 (18)	242	48.8 (14)
Scott	33	* *	* *	* *	28	* *	18	* *	18	* *
Sedgwick	3,873	74.0 (20)	3.8	(18)	3,211	28.5 (28)	2,306	11.9 (11)	1,978	42.7 (19)
Seward	427	66.0 (29)	11.5	(26)	398	37.4 (22)	267	6.0 (17)	238	64.3 (1)
Shawnee	1,162	74.2 (19)	4.7	(20)	1,069	49.6 (4)	785	5.0 (21)	739	57.8 (9)
Sheridan	5	* *	* *	* *	4	* *	2	* *	1	* *
Sherman	47	* *	* *	* *	58	* *	39	* *	38	* *
Smith	12	* *	* *	* *	13	* *	9	* *	10	* *
Stafford	33	* *	* *	* *	33	* *	25	* *	23	* *
Stanton	21	* *	* *	* *	21	* *	11	* *	12	* *
Stevens	55	* *	* *	* *	49	* *	38	* *	29	* *
Sumner	145	83.4 (12)	4.1	(19)	119	44.5 (11)	84	* *	68	* *
Thomas	36	* *	* *	* *	36	* *	23	* *	26	* *
Trego	13	* *	* *	* *	9	* *	6	* *	6	* *
Wabaunsee	16	* *	* *	* *	10	* *	10	* *	7	* *
Wallace	1	* *	* *	* *	0	* *	1	* *	0	* *
Washington	24	* *	* *	* *	20	* *	15	* *	14	* *
Wichita	11	* *	* *	* *	9	* *	6	* *	5	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.





**Table 6B**  
**2008 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Maternal Behavioral Indicators by County**

<b>Comparison of Maternal Behavioral Indicators</b> <sup>(2, 3)</sup>										
<b>County</b>	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Pregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	%	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Wilson	76	* *	* *	* *	71	* *	50	* *	47	* *
Woodson	2	* *	* *	* *	1	* *	0	* *	0	* *
Wyandotte	1,922	66.7 (27)	7.0	(24)	1,657	40.5 (16)	1,171	12.3 (9)	1,047	57.8 (9)
<b>Kansas</b>	<b>18,064</b>	<b>74.9</b>	<b>5.2</b>		<b>16,091</b>	<b>39.8</b>	<b>11,645</b>	<b>12.0</b>	<b>10,265</b>	<b>53.0</b>
<b>Nation (Prior Year)</b>	<b>612,462</b>	<b>79.4</b>	<b>4.7</b>		<b>1,034,046</b>	<b>31.3</b>	<b>98,170</b>	<b>15.7</b>	<b>238,509</b>	<b>75.9</b>

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 7B**  
**2008 Pregnancy Nutrition Surveillance** (1)  
**Kansas**  
**Comparison of Smoking Indicators by County**

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Allen	111	46.8 (28)	100	32.0 (27)	38	* *	96	* *	98	* *
Anderson	48	* *	43	* *	17	* *	39	* *	43	* *
Atchison	74	* *	38	* *	20	* *	67	* *	38	* *
Barber	39	* *	33	* *	14	* *	33	* *	33	* *
Barton	272	39.7 (16)	249	25.3 (19)	80	* *	218	10.1 (7)	252	6.7 (5)
Bourbon	128	43.8 (20)	102	25.5 (20)	34	* *	111	31.5 (24)	103	33.0 (24)
Brown	105	44.8 (25)	85	* *	26	* *	91	* *	85	* *
Butler	262	52.7 (30)	198	34.3 (28)	78	* *	191	14.7 (12)	196	7.7 (8)
Chase	14	* *	12	* *	7	* *	12	* *	12	* *
Chautauqua	30	* *	22	* *	6	* *	25	* *	21	* *
Cherokee	201	39.3 (14)	163	28.8 (24)	50	* *	160	30.0 (22)	169	29.6 (23)
Clark	1	* *	0	* *	0	* *	1	* *	0	* *
Clay	65	* *	57	* *	25	* *	53	* *	59	* *
Cloud	70	* *	61	* *	11	* *	57	* *	61	* *
Coffey	23	* *	23	* *	8	* *	14	* *	23	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 7B**  
**2008 Pregnancy Nutrition Surveillance** (1)  
**Kansas**  
**Comparison of Smoking Indicators by County**

<b>Comparison of Smoking Indicators</b> (2, 3)										
<b>County</b>	<b>Smoking</b>				<b>Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)</b>		<b>Smoking in Household</b>			
	<b>3 Months Prior to Pregnancy</b>		<b>Last 3 Months of Pregnancy</b>				<b>Prenatal</b>		<b>Postpartum</b>	
	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>
Comanche	5	* *	4	* *	0	* *	3	* *	4	* *
Cowley	339	44.5 (24)	292	28.8 (24)	100	27.0 (5)	270	21.1 (19)	295	21.0 (19)
Crawford	361	41.6 (19)	285	23.2 (15)	97	* *	310	40.0 (26)	286	38.8 (25)
Decatur	9	* *	7	* *	1	* *	6	* *	7	* *
Dickinson	100	45.0 (26)	87	* *	32	* *	84	* *	89	* *
Doniphan	31	* *	22	* *	9	* *	23	* *	24	* *
Douglas	447	44.1 (22)	381	20.7 (11)	118	44.9 (1)	358	10.1 (7)	382	7.3 (7)
Edwards	27	* *	20	* *	5	* *	20	* *	21	* *
Elk	13	* *	11	* *	2	* *	9	* *	11	* *
Ellis	140	29.3 (7)	121	9.1 (4)	34	* *	111	6.3 (2)	121	8.3 (9)
Ellsworth	27	* *	27	* *	10	* *	16	* *	28	* *
Finney	674	13.4 (3)	627	7.0 (3)	68	* *	556	17.3 (15)	626	15.7 (16)
Ford	612	7.7 (1)	533	3.8 (2)	29	* *	522	7.5 (5)	527	6.6 (4)
Franklin	163	44.2 (23)	147	25.2 (18)	47	* *	126	11.1 (9)	147	9.5 (11)
Geary	333	41.4 (18)	270	21.1 (13)	98	* *	284	39.8 (25)	273	38.8 (25)

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one county's rate to other counties. Rank 1 = best rate.
- (4) Among women who smoked 3 months prior to pregnancy.
- \* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 7B**  
**2008 Pregnancy Nutrition Surveillance** (1)  
**Kansas**  
**Comparison of Smoking Indicators by County**

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Gove	7	* *	7	* *	0	* *	5	* *	7	* *
Graham	14	* *	12	* *	7	* *	12	* *	12	* *
Grant	65	* *	62	* *	5	* *	50	* *	62	* *
Gray	55	* *	47	* *	3	* *	43	* *	47	* *
Greeley	10	* *	10	* *	3	* *	7	* *	10	* *
Greenwood	41	* *	33	* *	14	* *	36	* *	35	* *
Hamilton	18	* *	15	* *	2	* *	14	* *	15	* *
Harper	45	* *	25	* *	10	* *	41	* *	26	* *
Harvey	247	35.2 (11)	210	24.8 (17)	61	* *	203	5.9 (1)	211	6.2 (2)
Haskell	24	* *	23	* *	3	* *	21	* *	23	* *
Hodgeman	12	* *	9	* *	3	* *	8	* *	9	* *
Jackson	69	* *	64	* *	27	* *	62	* *	65	* *
Jefferson	16	* *	9	* *	1	* *	16	* *	11	* *
Jewell	14	* *	13	* *	3	* *	12	* *	13	* *
Johnson	1,801	24.6 (5)	1,597	11.5 (5)	277	40.8 (2)	1,400	18.9 (18)	1,606	19.6 (18)

(1) Reporting period is January 1 through December 31.  
(2) Excludes records with unknown data or errors.  
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.  
(4) Among women who smoked 3 months prior to pregnancy.  
\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 7B**  
**2008 Pregnancy Nutrition Surveillance** (1)  
**Kansas**  
**Comparison of Smoking Indicators by County**

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Kearny	21	* *	20	* *	2	* *	17	* *	20	* *
Kingman	50	* *	42	* *	9	* *	47	* *	48	* *
Kiowa	8	* *	4	* *	2	* *	8	* *	5	* *
Labette	161	33.5 (10)	139	20.9 (12)	33	* *	131	30.5 (23)	143	29.4 (22)
Lane	3	* *	0	* *	0	* *	4	* *	0	* *
Leavenworth	347	47.0 (29)	297	28.3 (23)	95	* *	243	46.5 (27)	301	40.5 (27)
Lincoln	20	* *	17	* *	6	* *	15	* *	17	* *
Linn	53	* *	47	* *	21	* *	43	* *	49	* *
Logan	13	* *	10	* *	4	* *	11	* *	10	* *
Lyon	338	30.2 (8)	304	16.1 (7)	63	* *	260	12.7 (10)	306	10.5 (12)
Marion	143	40.6 (17)	132	30.3 (26)	47	* *	114	7.0 (3)	133	4.5 (1)
Marshall	46	* *	40	* *	6	* *	34	* *	40	* *
Mcperson	63	* *	52	* *	24	* *	53	* *	53	* *
Meade	44	* *	37	* *	8	* *	35	* *	37	* *
Miami	114	54.4 (32)	98	* *	39	* *	85	* *	97	* *

(1) Reporting period is January 1 through December 31.  
(2) Excludes records with unknown data or errors.  
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.  
(4) Among women who smoked 3 months prior to pregnancy.  
\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 7B**  
**2008 Pregnancy Nutrition Surveillance** (1)  
**Kansas**  
**Comparison of Smoking Indicators by County**

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Mitchell	42	* *	41	* *	14	* *	34	* *	42	* *
Montgomery	318	39.6 (15)	273	28.2 (22)	84	* *	249	28.9 (21)	277	26.4 (21)
Morris	31	* *	27	* *	11	* *	26	* *	27	* *
Nemaha	37	* *	34	* *	17	* *	29	* *	34	* *
Neosho	144	36.8 (13)	109	17.4 (9)	29	* *	120	18.3 (16)	110	10.9 (13)
Ness	7	* *	7	* *	2	* *	3	* *	7	* *
Norton	26	* *	23	* *	6	* *	15	* *	23	* *
Osage	85	* *	76	* *	24	* *	64	* *	76	* *
Osborne	18	* *	18	* *	7	* *	13	* *	18	* *
Ottawa	18	* *	15	* *	4	* *	15	* *	15	* *
Pawnee	42	* *	41	* *	15	* *	36	* *	41	* *
Phillips	17	* *	3	* *	3	* *	17	* *	3	* *
Pottawatom	40	* *	33	* *	9	* *	33	* *	34	* *
Pratt	66	* *	58	* *	17	* *	46	* *	60	* *
Rawlins	10	* *	8	* *	1	* *	9	* *	8	* *

(1) Reporting period is January 1 through December 31.  
(2) Excludes records with unknown data or errors.  
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.  
(4) Among women who smoked 3 months prior to pregnancy.  
\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 7B**  
**2008 Pregnancy Nutrition Surveillance** (1)  
**Kansas**  
**Comparison of Smoking Indicators by County**

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Reno	478	45.6 (27)	417	24.7 (16)	151	31.1 (4)	372	18.3 (16)	419	17.4 (17)
Republic	32	* *	31	* *	13	* *	26	* *	32	* *
Rice	70	* *	58	* *	18	* *	59	* *	58	* *
Riley	783	28.9 (6)	578	17.8 (10)	109	22.9 (9)	592	15.0 (13)	648	14.4 (15)
Rooks	35	* *	26	* *	17	* *	30	* *	29	* *
Rush	12	* *	10	* *	4	* *	11	* *	10	* *
Russell	40	* *	36	* *	11	* *	30	* *	37	* *
Saline	433	43.9 (21)	357	27.5 (21)	130	24.6 (6)	347	8.6 (6)	360	6.4 (3)
Scott	34	* *	29	* *	7	* *	29	* *	29	* *
Sedgwick	4,069	33.4 (9)	3,782	16.4 (8)	933	35.4 (3)	3,197	7.1 (4)	3,775	6.9 (6)
Seward	482	8.9 (2)	430	3.3 (1)	32	* *	400	15.3 (14)	439	14.1 (14)
Shawnee	1,231	35.7 (12)	989	21.9 (14)	259	23.2 (8)	1,027	13.3 (11)	989	8.6 (10)
Sheridan	7	* *	5	* *	1	* *	4	* *	5	* *
Sherman	69	* *	58	* *	20	* *	57	* *	57	* *
Smith	14	* *	11	* *	3	* *	12	* *	12	* *

- (1) Reporting period is January 1 through December 31.  
(2) Excludes records with unknown data or errors.  
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.  
(4) Among women who smoked 3 months prior to pregnancy.  
\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 7B**  
**2008 Pregnancy Nutrition Surveillance** (1)  
**Kansas**  
**Comparison of Smoking Indicators by County**

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Stafford	39	* *	30	* *	9	* *	32	* *	31	* *
Stanton	22	* *	20	* *	2	* *	21	* *	20	* *
Stevens	66	* *	52	* *	10	* *	49	* *	52	* *
Sumner	150	54.0 (31)	135	34.8 (29)	57	* *	115	53.0 (28)	134	53.7 (28)
Thomas	47	* *	36	* *	10	* *	36	* *	36	* *
Trego	12	* *	10	* *	4	* *	11	* *	11	* *
Wabaunsee	18	* *	17	* *	5	* *	10	* *	17	* *
Wallace	0	* *	0	* *	0	* *	0	* *	0	* *
Washington	25	* *	24	* *	5	* *	20	* *	23	* *
Wichita	11	* *	9	* *	2	* *	9	* *	9	* *
Wilson	79	* *	73	* *	35	* *	70	* *	73	* *
Woodson	2	* *	2	* *	0	* *	1	* *	2	* *
Wyandotte	2,024	23.2 (4)	1,754	14.2 (6)	300	23.3 (7)	1,646	21.6 (20)	1,789	21.7 (20)
<b>Kansas</b>	<b>19,771</b>	<b>32.9</b>	<b>17,140</b>	<b>18.5</b>	<b>4,232</b>	<b>30.8</b>	<b>15,818</b>	<b>17.6</b>	<b>17,316</b>	<b>16.1</b>

(1) Reporting period is January 1 through December 31.  
(2) Excludes records with unknown data or errors.  
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.  
(4) Among women who smoked 3 months prior to pregnancy.  
\* Percentages are not calculated if <100 records are available for analysis after exclusions.





**Table 7B**  
**2008 Pregnancy Nutrition Surveillance** (1)  
**Kansas**  
**Comparison of Smoking Indicators by County**

<b>Comparison of Smoking Indicators</b> (2, 3)										
<b>County</b>	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
<b>Nation (Prior Year)</b>	<b>895,227</b>	<b>26.6</b>	<b>716,629</b>	<b>15.8</b>	<b>161,864</b>	<b>35.1</b>	<b>572,120</b>	<b>20.9</b>	<b>486,636</b>	<b>22.6</b>

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 8B**  
**2008 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Infant Health Indicators by County**

Comparison of Infant Health Indicators <sup>(2, 3)</sup>									
County	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)	High (5)		Number	%	Number	%	Number	%
	Number	% (Rank)	% (Rank)						
Allen	107	6.5 (17)	5.6 (10)	91	* *	85	* *	106	48.1 (30)
Anderson	46	* *	* *	37	* *	36	* *	43	* *
Atchison	101	5.9 (10)	5.9 (15)	84	* *	79	* *	105	53.3 (29)
Barber	31	* *	* *	26	* *	20	* *	32	* *
Barton	243	7.0 (21)	5.8 (14)	202	13.9 (25)	172	2.9 (14)	237	75.9 (6)
Bourbon	117	6.0 (13)	6.0 (18)	101	11.9 (19)	88	* *	116	64.7 (18)
Brown	86	* *	* *	88	* *	74	* *	90	* *
Butler	264	9.8 (31)	5.7 (12)	180	14.4 (26)	153	3.3 (19)	240	72.9 (11)
Chase	12	* *	* *	10	* *	10	* *	11	* *
Chautauqua	26	* *	* *	23	* *	20	* *	23	* *
Cherokee	194	7.7 (22)	5.7 (12)	147	12.9 (23)	128	3.1 (17)	182	61.0 (22)
Clark	4	* *	* *	2	* *	2	* *	3	* *
Clay	60	* *	* *	49	* *	41	* *	59	* *
Cloud	63	* *	* *	52	* *	42	* *	59	* *

(1) Reporting period is January 1 through December 31.  
(2) Excludes records with unknown data or errors.  
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.  
(4) <2500 g.  
(5) >4000 g.  
(6) <37 weeks gestation.  
(7) 37 or more weeks gestation and <2500 g.  
\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 8B**  
**2008 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Infant Health Indicators by County**

<b>Comparison of Infant Health Indicators</b> <sup>(2, 3)</sup>										
<b>County</b>	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)							
	Number	% (Rank)	%	(Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Coffey	23	* * * *	* *		14	* *	12	* *	24	* *
Comanche	5	* * * *	* *		2	* *	2	* *	5	* *
Cowley	320	6.9 (20)	7.2 (28)		245	11.8 (18)	215	5.1 (24)	311	55.3 (26)
Crawford	324	8.6 (26)	5.9 (15)		272	9.2 (7)	246	3.7 (20)	318	59.7 (23)
Decatur	7	* * * *	* *		6	* *	6	* *	6	* *
Dickinson	87	* * * *	* *		75	* *	70	* *	89	* *
Doniphan	37	* * * *	* *		27	* *	23	* *	34	* *
Douglas	407	5.4 (7)	7.1 (26)		313	12.1 (21)	273	2.6 (12)	389	75.6 (7)
Edwards	20	* * * *	* *		14	* *	9	* *	19	* *
Elk	12	* * * *	* *		8	* *	8	* *	11	* *
Ellis	118	8.5 (25)	0.8 (1)		96	* *	80	* *	115	81.7 (1)
Ellsworth	28	* * * *	* *		16	* *	14	* *	27	* *
Finney	618	5.7 (9)	6.5 (21)		514	10.1 (11)	457	2.0 (5)	341	78.6 (3)
Ford	562	4.8 (3)	5.9 (15)		483	9.7 (9)	432	1.4 (4)	440	74.3 (8)

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one county's rate to other counties. Rank 1 = best rate.
- (4) <2500 g.
- (5) >4000 g.
- (6) <37 weeks gestation.
- (7) 37 or more weeks gestation and <2500 g.
- \* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 8B**  
**2008 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Infant Health Indicators by County**

<b>Comparison of Infant Health Indicators</b> <sup>(2, 3)</sup>										
<b>County</b>	<b>Birthweight</b>				<b>Preterm</b> (6)		<b>Full Term LBW</b> (7)		<b>Ever Breastfed</b>	
	<b>Low</b> (4)		<b>High</b> (5)							
	<b>Number</b>	<b>% (Rank)</b>	<b>% (Rank)</b>		<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>
Franklin	145	6.2 (15)	5.5 (9)		110	10.9 (14)	98	* *	139	54.0 (27)
Geary	287	5.2 (5)	5.2 (6)		234	7.7 (3)	215	2.3 (8)	261	69.0 (14)
Gove	7	* *	* *		5	* *	4	* *	7	* *
Graham	11	* *	* *		10	* *	8	* *	12	* *
Grant	62	* *	* *		47	* *	41	* *	35	* *
Gray	37	* *	* *		34	* *	26	* *	44	* *
Greeley	10	* *	* *		7	* *	7	* *	10	* *
Greenwood	40	* *	* *		35	* *	29	* *	38	* *
Hamilton	13	* *	* *		11	* *	10	* *	5	* *
Harper	37	* *	* *		37	* *	28	* *	42	* *
Harvey	214	9.3 (28)	5.6 (10)		180	17.2 (27)	149	2.7 (13)	219	78.5 (4)
Haskell	23	* *	* *		20	* *	19	* *	12	* *
Hodgeman	12	* *	* *		8	* *	7	* *	10	* *
Jackson	66	* *	* *		60	* *	51	* *	62	* *

(1) Reporting period is January 1 through December 31.  
(2) Excludes records with unknown data or errors.  
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.  
(4) <2500 g.  
(5) >4000 g.  
(6) <37 weeks gestation.  
(7) 37 or more weeks gestation and <2500 g.  
\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 8B**  
**2008 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Infant Health Indicators by County**

Comparison of Infant Health Indicators <sup>(2, 3)</sup>										
County	Birthweight				Preterm <sup>(6)</sup>		Full Term LBW <sup>(7)</sup>		Ever Breastfed	
	Low <sup>(4)</sup>		High <sup>(5)</sup>		Number	% <sup>(Rank)</sup>	Number	% <sup>(Rank)</sup>	Number	% <sup>(Rank)</sup>
	Number	% <sup>(Rank)</sup>	% <sup>(Rank)</sup>							
Jefferson	59	* * * *	* * * *	44	* *	39	* *	57	* *	
Jewell	13	* * * *	* * * *	12	* *	8	* *	12	* *	
Johnson	1,568	6.7 (19)	7.0 (25)	1,270	8.8 (4)	1,100	2.5 (11)	1,531	78.8 (2)	
Kearny	19	* * * *	* * * *	16	* *	13	* *	9	* *	
Kingman	43	* * * *	* * * *	41	* *	33	* *	48	* *	
Kiowa	8	* * * *	* * * *	8	* *	7	* *	8	* *	
Labette	149	4.7 (2)	10.1 (31)	117	12.0 (20)	103	1.0 (2)	133	55.6 (25)	
Lane	10	* * * *	* * * *	7	* *	7	* *	10	* *	
Leavenworth	299	9.4 (29)	4.0 (2)	216	8.8 (4)	195	4.1 (22)	304	62.2 (20)	
Lincoln	22	* * * *	* * * *	15	* *	14	* *	20	* *	
Linn	46	* * * *	* * * *	39	* *	30	* *	45	* *	
Logan	10	* * * *	* * * *	8	* *	7	* *	9	* *	
Lyon	301	6.0 (13)	6.0 (18)	236	8.9 (6)	214	0.9 (1)	289	73.0 (10)	
Marion	133	3.8 (1)	5.3 (7)	106	4.7 (1)	100	2.0 (5)	132	74.2 (9)	

(1) Reporting period is January 1 through December 31.  
(2) Excludes records with unknown data or errors.  
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.  
(4) <2500 g.  
(5) >4000 g.  
(6) <37 weeks gestation.  
(7) 37 or more weeks gestation and <2500 g.  
\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 8B**  
**2008 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Infant Health Indicators by County**

<b>Comparison of Infant Health Indicators</b> <sup>(2, 3)</sup>										
<b>County</b>	<b>Birthweight</b>				<b>Preterm</b> (6)		<b>Full Term LBW</b> (7)		<b>Ever Breastfed</b>	
	<b>Low</b> (4)		<b>High</b> (5)							
	<b>Number</b>	<b>% (Rank)</b>	<b>% (Rank)</b>		<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>
Marshall	39	* * * *	* *		28	* *	24	* *	39	* *
Mcperson	54	* * * *	* *		46	* *	42	* *	52	* *
Meade	38	* * * *	* *		30	* *	25	* *	36	* *
Miami	126	5.6 (8)	6.3 (20)		87	* *	76	* *	98	* *
Mitchell	40	* * * *	* *		33	* *	30	* *	37	* *
Montgomery	288	9.4 (29)	4.2 (3)		225	13.3 (24)	193	4.1 (22)	270	62.2 (20)
Morris	29	* * * *	* *		23	* *	19	* *	28	* *
Nemaha	33	* * * *	* *		28	* *	21	* *	36	* *
Neosho	119	5.0 (4)	6.7 (23)		112	11.6 (15)	87	* *	142	53.5 (28)
Ness	8	* * * *	* *		3	* *	2	* *	8	* *
Norton	20	* * * *	* *		12	* *	10	* *	19	* *
Osage	83	* * * *	* *		64	* *	49	* *	74	* *
Osborne	17	* * * *	* *		13	* *	13	* *	16	* *
Ottawa	19	* * * *	* *		14	* *	13	* *	15	* *

(1) Reporting period is January 1 through December 31.  
(2) Excludes records with unknown data or errors.  
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.  
(4) <2500 g.  
(5) >4000 g.  
(6) <37 weeks gestation.  
(7) 37 or more weeks gestation and <2500 g.  
\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 8B**  
**2008 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Infant Health Indicators by County**

<b>Comparison of Infant Health Indicators</b> <sup>(2, 3)</sup>										
<b>County</b>	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)							
	Number	% (Rank)	%	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Pawnee	39	* * * *	* *	* *	35	* *	32	* *	38	* *
Phillips	29	* * * *	* *	* *	23	* *	21	* *	28	* *
Pottawatom	34	* * * *	* *	* *	27	* *	25	* *	33	* *
Pratt	57	* * * *	* *	* *	39	* *	38	* *	52	* *
Rawlins	9	* * * *	* *	* *	8	* *	8	* *	9	* *
Reno	438	5.9 (10)	7.5 (29)		338	10.4 (12)	299	2.3 (8)	418	58.9 (24)
Republic	32	* * * *	* *	* *	26	* *	17	* *	33	* *
Rice	59	* * * *	* *	* *	49	* *	41	* *	55	* *
Riley	699	5.9 (10)	6.6 (22)		484	9.7 (9)	436	2.1 (7)	679	78.2 (5)
Rooks	33	* * * *	* *	* *	29	* *	26	* *	32	* *
Rush	11	* * * *	* *	* *	10	* *	9	* *	11	* *
Russell	36	* * * *	* *	* *	27	* *	22	* *	33	* *
Saline	447	8.7 (27)	4.9 (4)		353	11.6 (15)	309	2.9 (14)	444	62.4 (19)
Scott	28	* * * *	* *	* *	26	* *	25	* *	27	* *

(1) Reporting period is January 1 through December 31.  
(2) Excludes records with unknown data or errors.  
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.  
(4) <2500 g.  
(5) >4000 g.  
(6) <37 weeks gestation.  
(7) 37 or more weeks gestation and <2500 g.  
\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 8B**  
**2008 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Infant Health Indicators by County**

Comparison of Infant Health Indicators <sup>(2, 3)</sup>									
County	Birthweight			Preterm <sup>(6)</sup>		Full Term LBW <sup>(7)</sup>		Ever Breastfed	
	Low <sup>(4)</sup>	High <sup>(5)</sup>		Number	%	Number	%	Number	%
	Number	% <sup>(Rank)</sup>	% <sup>(Rank)</sup>						
Sedgwick	3,833	8.1 (24)	7.1 (26)	2,993	9.5 (8)	2,691	3.2 (18)	3,383	67.8 (15)
Seward	424	5.2 (5)	5.0 (5)	356	10.4 (12)	315	1.3 (3)	432	71.1 (13)
Shawnee	1,171	6.6 (18)	5.4 (8)	963	11.7 (17)	839	2.3 (8)	1,158	67.5 (16)
Sheridan	6	*	*	3	*	2	*	6	*
Sherman	62	*	*	51	*	45	*	58	*
Smith	15	*	*	11	*	9	*	15	*
Stafford	30	*	*	27	*	23	*	28	*
Stanton	21	*	*	19	*	16	*	9	*
Stevens	32	*	*	38	*	21	*	55	*
Sumner	146	6.2 (15)	7.5 (29)	109	6.4 (2)	102	3.9 (21)	143	65.0 (17)
Thomas	44	*	*	30	*	28	*	40	*
Trego	11	*	*	9	*	6	*	11	*
Wabaunsee	17	*	*	10	*	8	*	16	*
Wallace	1	*	*	0	*	0	*	1	*

(1) Reporting period is January 1 through December 31.  
(2) Excludes records with unknown data or errors.  
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.  
(4) <2500 g.  
(5) >4000 g.  
(6) <37 weeks gestation.  
(7) 37 or more weeks gestation and <2500 g.  
\* Percentages are not calculated if <100 records are available for analysis after exclusions.





**Table 8B**  
**2008 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Infant Health Indicators by County**

<b>Comparison of Infant Health Indicators</b> <sup>(2, 3)</sup>										
<b>County</b>	<b>Birthweight</b>				<b>Preterm</b> (6)		<b>Full Term LBW</b> (7)		<b>Ever Breastfed</b>	
	<b>Low</b> (4)		<b>High</b> (5)							
	<b>Number</b>	<b>% (Rank)</b>	<b>% (Rank)</b>		<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>
Washington	23	* * * *	* *		20	* *	17	* *	23	* *
Wichita	11	* * * *	* *		9	* *	7	* *	11	* *
Wilson	76	* * * *	* *		67	* *	56	* *	62	* *
Woodson	2	* * * *	* *		1	* *	1	* *	2	* *
Wyandotte	1,855	8.0 (23)	6.8 (24)		1,485	12.1 (21)	1,287	2.9 (14)	1,785	71.8 (12)
<b>Kansas</b>	<b>18,210</b>	<b>7.2</b>	<b>6.5</b>		<b>14,503</b>	<b>10.7</b>	<b>12,744</b>	<b>2.8</b>	<b>16,938</b>	<b>69.4</b>
<b>Nation (Prior Year)</b>	<b>1,039,036</b>	<b>8.2</b>	<b>6.7</b>		<b>947,705</b>	<b>11.9</b>	<b>856,467</b>	<b>3.7</b>	<b>936,194</b>	<b>64.6</b>

(1) Reporting period is January 1 through December 31.  
(2) Excludes records with unknown data or errors.  
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.  
(4) <2500 g.  
(5) >4000 g.  
(6) <37 weeks gestation.  
(7) 37 or more weeks gestation and <2500 g.  
\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 3B**  
**2006 - 2008 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Racial and Ethnic Distribution by County**  
**Combined 3 Years**

<b>Comparison of Racial and Ethnic Distribution</b> <sup>(2)</sup>								
County	Number Included in Analysis	White, Not Hispanic  %	Black, Not Hispanic  %	Hispanic  %	American Indian/Alaska Native  %	Asian/Pacific Islander  %	Multiple Races  %	All Other/Unknown  %
Allen	300	92.0	4.3	2.0	0.7	0.7	0.3	0.0
Anderson	159	93.1	0.6	4.4	0.6	1.3	0.0	0.0
Atchison	396	87.1	10.1	1.0	0.0	0.5	1.3	0.0
Barber	103	95.1	2.9	0.0	0.0	1.9	0.0	0.0
Barton	845	73.4	1.4	24.9	0.1	0.0	0.2	0.0
Bourbon	371	92.5	3.5	2.7	0.0	0.5	0.8	0.0
Brown	288	77.4	2.4	3.1	14.9	0.0	2.1	0.0
Butler	802	91.4	2.0	4.5	0.4	0.1	1.6	0.0
Chase	41	*	*	*	*	*	*	*
Chautauqua	86	*	*	*	*	*	*	*
Cherokee	672	92.1	1.5	3.7	1.2	0.7	0.7	0.0
Cheyenne	5	*	*	*	*	*	*	*
Clark	13	*	*	*	*	*	*	*
Clay	206	96.6	0.5	1.5	1.0	0.5	0.0	0.0
Cloud	235	94.0	1.3	3.4	0.4	0.9	0.0	0.0
Coffey	75	*	*	*	*	*	*	*
Comanche	14	*	*	*	*	*	*	*
Cowley	1,008	80.6	1.9	14.2	1.0	1.9	0.5	0.0
Crawford	1,211	86.5	3.2	7.7	0.2	1.7	0.7	0.0
Decatur	31	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 3B**  
**2006 - 2008 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Racial and Ethnic Distribution by County**  
**Combined 3 Years**

<b>Comparison of Racial and Ethnic Distribution</b> <sup>(2)</sup>								
County	Number Included in Analysis	White, Not Hispanic  %	Black, Not Hispanic  %	Hispanic  %	American Indian/Alaska Native  %	Asian/Pacific Islander  %	Multiple Races  %	All Other/Unknown  %
Dickinson	340	93.2	0.6	4.4	1.2	0.3	0.3	0.0
Doniphan	123	86.2	4.1	7.3	1.6	0.0	0.8	0.0
Douglas	1,433	64.3	8.4	10.7	9.6	4.7	2.2	0.0
Edwards	76	*	*	*	*	*	*	*
Elk	56	*	*	*	*	*	*	*
Ellis	439	85.4	2.3	11.2	0.2	0.9	0.0	0.0
Ellsworth	88	*	*	*	*	*	*	*
Finney	2,082	34.0	1.1	62.8	0.0	2.0	0.1	0.0
Ford	1,872	23.2	0.8	75.1	0.2	0.7	0.1	0.0
Franklin	507	90.1	2.4	4.3	1.2	0.4	1.6	0.0
Geary	948	54.0	24.3	13.8	0.5	3.7	3.7	0.0
Gove	35	*	*	*	*	*	*	*
Graham	50	*	*	*	*	*	*	*
Grant	246	27.2	0.0	72.8	0.0	0.0	0.0	0.0
Gray	170	68.8	0.0	30.6	0.0	0.0	0.6	0.0
Greeley	23	*	*	*	*	*	*	*
Greenwood	130	93.8	0.8	3.8	0.0	0.8	0.8	0.0
Hamilton	55	*	*	*	*	*	*	*
Harper	154	81.8	0.0	16.2	0.6	0.0	1.3	0.0
Harvey	772	69.4	2.1	26.2	0.6	0.0	1.7	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 3B**  
**2006 - 2008 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Racial and Ethnic Distribution by County**  
**Combined 3 Years**

<b>Comparison of Racial and Ethnic Distribution</b> <sup>(2)</sup>								
County	Number Included in Analysis	White, Not Hispanic  %	Black, Not Hispanic  %	Hispanic  %	American Indian/Alaska Native  %	Asian/Pacific Islander  %	Multiple Races  %	All Other/Unknown  %
Haskell	65	*	*	*	*	*	*	*
Hodgeman	30	*	*	*	*	*	*	*
Jackson	242	79.3	0.0	5.4	14.5	0.0	0.8	0.0
Jefferson	199	95.5	0.0	1.5	2.0	0.5	0.5	0.0
Jewell	41	*	*	*	*	*	*	*
Johnson	5,464	46.7	12.6	36.2	0.5	3.1	1.0	0.0
Kearny	54	*	*	*	*	*	*	*
Kingman	164	93.3	0.0	4.3	1.8	0.6	0.0	0.0
Kiowa	40	*	*	*	*	*	*	*
Labette	528	88.3	6.4	3.8	0.6	0.2	0.8	0.0
Lane	39	*	*	*	*	*	*	*
Leavenworth	1,055	70.0	16.0	8.9	0.7	1.7	2.7	0.0
Lincoln	67	*	*	*	*	*	*	*
Linn	179	96.1	0.0	2.8	0.0	0.0	1.1	0.0
Logan	28	*	*	*	*	*	*	*
Lyon	1,091	54.5	5.1	38.7	0.1	1.1	0.5	0.0
Marion	435	88.7	1.4	8.0	1.4	0.2	0.2	0.0
Marshall	149	92.6	0.7	6.0	0.7	0.0	0.0	0.0
Mcperson	172	93.0	0.6	2.9	2.3	0.0	1.2	0.0
Meade	123	60.2	0.0	39.8	0.0	0.0	0.0	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 3B**  
**2006 - 2008 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Racial and Ethnic Distribution by County**  
**Combined 3 Years**

<b>Comparison of Racial and Ethnic Distribution</b> <sup>(2)</sup>								
County	Number Included in Analysis	White, Not Hispanic  %	Black, Not Hispanic  %	Hispanic  %	American Indian/Alaska Native  %	Asian/Pacific Islander  %	Multiple Races  %	All Other/Unknown  %
Miami	389	93.1	2.3	3.1	0.0	0.5	1.0	0.0
Mitchell	105	95.2	0.0	4.8	0.0	0.0	0.0	0.0
Montgomery	1,001	77.3	10.3	8.2	2.6	0.9	0.7	0.0
Morris	87	*	*	*	*	*	*	*
Nemaha	107	92.5	0.0	5.6	0.0	0.9	0.9	0.0
Neosho	452	91.4	2.4	4.9	0.4	0.7	0.2	0.0
Ness	23	*	*	*	*	*	*	*
Norton	81	*	*	*	*	*	*	*
Osage	241	97.9	0.0	1.2	0.4	0.0	0.4	0.0
Osborne	55	*	*	*	*	*	*	*
Ottawa	76	*	*	*	*	*	*	*
Pawnee	148	83.1	2.0	13.5	0.0	0.7	0.7	0.0
Phillips	103	93.2	1.0	4.9	0.0	1.0	0.0	0.0
Pottawatom	129	92.2	0.0	7.0	0.0	0.0	0.8	0.0
Pratt	234	86.3	0.0	13.2	0.0	0.4	0.0	0.0
Rawlins	23	*	*	*	*	*	*	*
Reno	1,446	83.2	2.9	11.8	0.6	0.9	0.6	0.0
Republic	91	*	*	*	*	*	*	*
Rice	211	79.6	1.9	18.5	0.0	0.0	0.0	0.0
Riley	2,681	75.4	11.0	8.5	0.6	4.3	0.2	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 3B**  
**2006 - 2008 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Racial and Ethnic Distribution by County**  
**Combined 3 Years**

<b>Comparison of Racial and Ethnic Distribution</b> <sup>(2)</sup>								
<b>County</b>	<b>Number Included in Analysis</b>	<b>White, Not Hispanic</b>	<b>Black, Not Hispanic</b>	<b>Hispanic</b>	<b>American Indian/Alaska Native</b>	<b>Asian/Pacific Islander</b>	<b>Multiple Races</b>	<b>All Other/Unknown</b>
		%	%	%	%	%	%	%
Rooks	119	95.0	0.8	1.7	0.8	0.8	0.8	0.0
Rush	47	*	*	*	*	*	*	*
Russell	118	93.2	4.2	1.7	0.8	0.0	0.0	0.0
Saline	1,351	73.1	4.3	19.5	0.3	1.8	1.1	0.0
Scott	142	66.2	0.7	33.1	0.0	0.0	0.0	0.0
Sedgwick	12,379	47.5	17.8	28.2	1.6	3.6	1.3	0.0
Seward	1,442	18.8	1.5	76.8	1.7	0.8	0.3	0.0
Shawnee	4,278	55.3	17.0	23.3	2.2	0.8	1.5	0.0
Sheridan	29	*	*	*	*	*	*	*
Sherman	232	77.6	0.0	20.7	0.4	0.4	0.9	0.0
Smith	62	*	*	*	*	*	*	*
Stafford	121	71.1	0.0	28.1	0.0	0.8	0.0	0.0
Stanton	58	*	*	*	*	*	*	*
Stevens	205	45.9	0.0	53.2	0.0	1.0	0.0	0.0
Sumner	427	94.4	0.5	4.2	0.5	0.2	0.2	0.0
Thomas	158	86.1	1.3	9.5	0.6	1.3	1.3	0.0
Trego	48	*	*	*	*	*	*	*
Wabaunsee	55	*	*	*	*	*	*	*
Wallace	8	*	*	*	*	*	*	*
Washington	81	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 3B**  
**2006 - 2008 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Racial and Ethnic Distribution by County**  
**Combined 3 Years**

<b>Comparison of Racial and Ethnic Distribution</b> <sup>(2)</sup>								
<b>County</b>	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Wichita	57	*	*	*	*	*	*	*
Wilson	294	95.9	0.3	2.7	0.0	0.3	0.7	0.0
Woodson	28	*	*	*	*	*	*	*
Wyandotte	6,483	25.8	31.6	39.8	0.2	2.3	0.3	0.0
<b>Kansas</b>	<b>62,530</b>	<b>58.3</b>	<b>11.4</b>	<b>26.2</b>	<b>1.2</b>	<b>2.0</b>	<b>0.9</b>	<b>0.0</b>

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 4B**  
**2006 - 2008 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Age Distribution by County**  
**Combined 3 Years**

<b>Comparison of Age Distribution</b> <sup>(2)</sup>								
<b>County</b>	<b>Number Included in Analysis</b>	<b>&lt; 15 Years %</b>	<b>15 - 17 Years %</b>	<b>18 - 19 Years %</b>	<b>20 - 29 Years %</b>	<b>30 - 39 Years %</b>	<b>&gt;= 40 Years %</b>	<b>Unknown %</b>
Allen	300	0.7	9.0	15.7	63.7	10.0	0.3	0.7
Anderson	159	0.0	8.2	12.6	66.7	11.9	0.6	0.0
Atchison	396	0.0	7.6	18.7	61.1	11.6	0.8	0.3
Barber	103	0.0	4.9	16.5	66.0	12.6	0.0	0.0
Barton	845	0.1	6.3	17.6	61.7	14.0	0.4	0.0
Bourbon	371	0.3	5.9	18.9	65.5	8.9	0.5	0.0
Brown	288	0.0	6.9	17.4	64.6	9.7	1.0	0.3
Butler	802	0.1	6.4	16.5	62.0	14.2	0.9	0.0
Chase	41	*	*	*	*	*	*	*
Chautauqua	86	*	*	*	*	*	*	*
Cherokee	672	0.1	7.9	18.0	60.1	12.9	0.6	0.3
Cheyenne	5	*	*	*	*	*	*	*
Clark	13	*	*	*	*	*	*	*
Clay	206	0.0	4.4	12.6	67.0	15.0	1.0	0.0
Cloud	235	0.4	7.2	15.3	62.1	13.2	1.7	0.0
Coffey	75	*	*	*	*	*	*	*
Comanche	14	*	*	*	*	*	*	*
Cowley	1,008	0.5	8.3	17.3	59.4	14.0	0.5	0.0
Crawford	1,211	0.2	7.0	15.9	63.3	12.6	0.7	0.2

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.





**Table 4B**  
**2006 - 2008 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Age Distribution by County**  
**Combined 3 Years**

Comparison of Age Distribution <sup>(2)</sup>								
County	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Decatur	31	*	*	*	*	*	*	*
Dickinson	340	0.3	6.5	13.5	63.2	16.2	0.3	0.0
Doniphan	123	0.0	4.9	17.1	65.0	13.0	0.0	0.0
Douglas	1,433	0.1	4.0	12.9	64.2	17.3	1.4	0.0
Edwards	76	*	*	*	*	*	*	*
Elk	56	*	*	*	*	*	*	*
Ellis	439	0.2	5.2	12.5	68.8	11.2	1.8	0.2
Ellsworth	88	*	*	*	*	*	*	*
Finney	2,082	0.5	10.3	13.6	54.9	19.3	1.2	0.1
Ford	1,872	0.4	6.9	12.0	58.1	21.4	1.0	0.2
Franklin	507	1.0	7.7	16.8	61.5	12.4	0.4	0.2
Geary	948	0.4	5.8	16.2	66.2	10.8	0.5	0.0
Gove	35	*	*	*	*	*	*	*
Graham	50	*	*	*	*	*	*	*
Grant	246	0.4	6.5	14.2	56.9	19.9	2.0	0.0
Gray	170	0.0	5.9	9.4	51.8	30.0	2.4	0.6
Greeley	23	*	*	*	*	*	*	*
Greenwood	130	0.0	6.9	16.9	63.1	13.1	0.0	0.0
Hamilton	55	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 4B**  
**2006 - 2008 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Age Distribution by County**  
**Combined 3 Years**

<b>Comparison of Age Distribution</b> <sup>(2)</sup>								
<b>County</b>	<b>Number Included in Analysis</b>	<b>&lt; 15 Years %</b>	<b>15 - 17 Years %</b>	<b>18 - 19 Years %</b>	<b>20 - 29 Years %</b>	<b>30 - 39 Years %</b>	<b>&gt;= 40 Years %</b>	<b>Unknown %</b>
Harper	154	0.0	7.1	11.7	67.5	13.6	0.0	0.0
Harvey	772	0.1	6.0	11.4	65.7	14.9	1.9	0.0
Haskell	65	*	*	*	*	*	*	*
Hodgeman	30	*	*	*	*	*	*	*
Jackson	242	0.0	7.0	13.2	69.8	9.9	0.0	0.0
Jefferson	199	0.0	10.6	21.6	56.3	10.6	1.0	0.0
Jewell	41	*	*	*	*	*	*	*
Johnson	5,464	0.2	4.3	11.7	62.5	20.1	1.2	0.1
Kearny	54	*	*	*	*	*	*	*
Kingman	164	0.0	4.9	13.4	64.0	17.7	0.0	0.0
Kiowa	40	*	*	*	*	*	*	*
Labette	528	0.0	6.8	17.4	65.7	9.5	0.6	0.0
Lane	39	*	*	*	*	*	*	*
Leavenworth	1,055	0.3	5.8	18.3	63.9	10.6	1.1	0.0
Lincoln	67	*	*	*	*	*	*	*
Linn	179	0.0	6.1	16.8	63.7	11.7	1.7	0.0
Logan	28	*	*	*	*	*	*	*
Lyon	1,091	0.5	6.6	13.3	61.9	15.8	2.0	0.0
Marion	435	0.2	5.3	13.6	63.9	16.1	0.9	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 4B**  
**2006 - 2008 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Age Distribution by County**  
**Combined 3 Years**

<b>Comparison of Age Distribution</b> <sup>(2)</sup>								
<b>County</b>	<b>Number Included in Analysis</b>	<b>&lt; 15 Years %</b>	<b>15 - 17 Years %</b>	<b>18 - 19 Years %</b>	<b>20 - 29 Years %</b>	<b>30 - 39 Years %</b>	<b>&gt;= 40 Years %</b>	<b>Unknown %</b>
Marshall	149	0.0	3.4	8.1	68.5	17.4	2.7	0.0
Mcperson	172	0.0	5.2	10.5	69.8	14.5	0.0	0.0
Meade	123	0.0	5.7	11.4	64.2	17.9	0.8	0.0
Miami	389	0.0	7.7	15.4	65.8	10.5	0.5	0.0
Mitchell	105	0.0	4.8	19.0	54.3	21.9	0.0	0.0
Montgomery	1,001	0.7	7.7	17.2	63.1	10.8	0.4	0.1
Morris	87	*	*	*	*	*	*	*
Nemaha	107	0.0	5.6	15.9	61.7	15.9	0.9	0.0
Neosho	452	0.7	5.8	20.8	61.3	11.1	0.4	0.0
Ness	23	*	*	*	*	*	*	*
Norton	81	*	*	*	*	*	*	*
Osage	241	0.4	3.3	17.8	63.9	14.1	0.0	0.4
Osborne	55	*	*	*	*	*	*	*
Ottawa	76	*	*	*	*	*	*	*
Pawnee	148	0.7	4.1	16.9	66.9	10.1	1.4	0.0
Phillips	103	1.0	1.9	15.5	60.2	21.4	0.0	0.0
Pottawatom	129	0.0	7.0	7.8	54.3	25.6	4.7	0.8
Pratt	234	0.4	8.5	16.2	64.1	10.3	0.4	0.0
Rawlins	23	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 4B**  
**2006 - 2008 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Age Distribution by County**  
**Combined 3 Years**

<b>Comparison of Age Distribution</b> <sup>(2)</sup>								
<b>County</b>	<b>Number Included in Analysis</b>	<b>&lt; 15 Years %</b>	<b>15 - 17 Years %</b>	<b>18 - 19 Years %</b>	<b>20 - 29 Years %</b>	<b>30 - 39 Years %</b>	<b>&gt;= 40 Years %</b>	<b>Unknown %</b>
Reno	1,446	0.3	4.8	15.5	66.7	12.0	0.6	0.1
Republic	91	*	*	*	*	*	*	*
Rice	211	0.5	7.1	14.7	57.8	18.0	1.9	0.0
Riley	2,681	0.1	2.0	12.2	74.2	11.0	0.4	0.0
Rooks	119	0.0	4.2	13.4	63.0	16.8	2.5	0.0
Rush	47	*	*	*	*	*	*	*
Russell	118	0.0	7.6	15.3	68.6	8.5	0.0	0.0
Saline	1,351	0.2	8.0	16.7	61.4	13.0	0.7	0.1
Scott	142	0.7	7.0	6.3	71.8	12.0	1.4	0.7
Sedgwick	12,379	0.3	6.8	13.9	61.5	16.0	1.3	0.1
Seward	1,442	0.5	7.6	13.5	56.6	20.3	1.5	0.1
Shawnee	4,278	0.5	6.5	14.1	63.2	14.7	0.8	0.1
Sheridan	29	*	*	*	*	*	*	*
Sherman	232	0.0	5.2	20.7	58.2	13.4	2.6	0.0
Smith	62	*	*	*	*	*	*	*
Stafford	121	0.0	4.1	11.6	56.2	26.4	1.7	0.0
Stanton	58	*	*	*	*	*	*	*
Stevens	205	0.5	5.9	15.6	60.5	16.6	1.0	0.0
Sumner	427	0.2	6.1	16.9	63.5	13.1	0.2	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 4B**  
**2006 - 2008 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Age Distribution by County**  
**Combined 3 Years**

<b>Comparison of Age Distribution</b> <sup>(2)</sup>								
<b>County</b>	<b>Number Included in Analysis</b>	<b>&lt; 15 Years %</b>	<b>15 - 17 Years %</b>	<b>18 - 19 Years %</b>	<b>20 - 29 Years %</b>	<b>30 - 39 Years %</b>	<b>&gt;= 40 Years %</b>	<b>Unknown %</b>
Thomas	158	0.0	4.4	12.0	66.5	15.8	0.6	0.6
Trego	48	*	*	*	*	*	*	*
Wabaunsee	55	*	*	*	*	*	*	*
Wallace	8	*	*	*	*	*	*	*
Washington	81	*	*	*	*	*	*	*
Wichita	57	*	*	*	*	*	*	*
Wilson	294	0.0	11.6	12.9	63.3	12.2	0.0	0.0
Woodson	28	*	*	*	*	*	*	*
Wyandotte	6,483	0.7	8.2	14.3	58.5	17.1	1.2	0.1
<b>Kansas</b>	<b>62,530</b>	<b>0.3</b>	<b>6.4</b>	<b>14.3</b>	<b>62.2</b>	<b>15.6</b>	<b>1.1</b>	<b>0.1</b>

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 5B**  
**2006 - 2008 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Maternal Health Indicators by County**  
**Combined 3 Years**

Comparison of Maternal Health Indicators <sup>(2, 3)</sup>														
County	Prepregnancy BMI <sup>(4)</sup>			Weight Gain <sup>(5)</sup>			Anemia (Low Hb/Hct) <sup>(6)</sup>				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Allen	297	11.8 (42)	48.5 (50)	270	21.9 (28)	47.0 (37)	49	* *	86	* *	271	4.1 (58)	271	7.4(59)
Anderson	151	13.9 (53)	41.1 (3)	142	24.6 (47)	43.0 (13)	22	* *	28	* *	144	0.7 (7)	144	6.9(57)
Atchison	383	13.3 (50)	47.5 (45)	283	26.9 (54)	43.1 (14)	66	* *	66	* *	345	2.3 (46)	345	7.0(58)
Barber	100	7.0 (5)	56.0 (61)	90	* *	* *	11	* *	22	* *	89	* *	89	* *
Barton	818	10.0 (21)	45.5 (28)	746	25.2 (49)	43.6 (16)	96	* *	145	22.8(11)	731	2.1 (42)	731	3.7(41)
Bourbon	366	11.5 (39)	50.8 (57)	330	23.3 (38)	42.7 (11)	47	* *	73	* *	333	2.4 (47)	333	3.3(37)
Brown	276	8.3 (10)	46.0 (36)	228	18.0 (5)	56.6 (59)	51	* *	47	* *	252	0.0 (1)	252	4.8(51)
Butler	772	13.5 (52)	45.5 (28)	716	20.5 (17)	48.7 (48)	90	* *	212	28.8(23)	686	1.2 (19)	686	5.2(55)
Chase	37	* *	* *	30	* *	* *	5	* *	5	* *	37	* *	37	* *
Chautauqua	85	* *	* *	64	* *	* *	4	* *	10	* *	70	* *	70	* *
Cherokee	624	10.6 (27)	45.5 (28)	503	19.3 (10)	49.5 (52)	72	* *	146	24.0(14)	571	1.8 (36)	571	1.9(11)
Cheyenne	5	* *	* *	5	* *	* *	0	* *	1	* *	5	* *	5	* *
Clark	11	* *	* *	5	* *	* *	0	* *	3	* *	9	* *	9	* *
Clay	198	8.1 (9)	49.5 (53)	181	12.7 (1)	52.5 (57)	25	* *	43	* *	181	3.9 (56)	181	2.2(18)
Cloud	228	13.2 (49)	43.0 (10)	202	24.3 (43)	49.0 (49)	27	* *	49	* *	199	1.5 (26)	200	2.5(23)
Coffey	75	* *	* *	73	* *	* *	6	* *	21	* *	66	* *	66	* *
Comanche	14	* *	* *	12	* *	* *	2	* *	5	* *	10	* *	10	* *

(1) Reporting period is January 1 through December 31.  
(2) Excludes records with unknown data or errors.  
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.  
(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).  
(5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).  
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.  
\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 5B**  
**2006 - 2008 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Maternal Health Indicators by County**  
**Combined 3 Years**

Comparison of Maternal Health Indicators <sup>(2, 3)</sup>														
County	Prepregnancy BMI <sup>(4)</sup>			Weight Gain <sup>(5)</sup>			Anemia (Low Hb/Hct) <sup>(6)</sup>				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
							Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)
Cowley	976	11.8 (42)	45.0 (22)	885	23.2 (37)	44.3 (22)	138	31.2 (11)	124	21.8 (9)	875	1.4 (24)	875	3.2(33)
Crawford	1,144	13.1 (48)	44.6 (19)	963	20.8 (20)	48.5 (46)	163	14.1 (1)	189	16.9 (4)	1,031	1.8 (36)	1,030	2.4(20)
Decatur	31	* *	* *	27	* *	* *	6	* *	5	* *	27	* *	27	* *
Dickinson	325	10.2 (23)	41.8 (6)	293	24.6 (47)	46.4 (33)	44	* *	71	* *	296	2.4 (47)	296	3.0(30)
Doniphan	118	5.1 (1)	56.8 (63)	90	* *	* *	25	* *	24	* *	98	* *	98	* *
Douglas	1,378	11.7 (41)	43.1 (12)	1,157	18.8 (8)	51.4 (56)	192	31.8 (12)	351	29.3(24)	1,168	1.7 (32)	1,167	2.0(15)
Edwards	74	* *	* *	61	* *	* *	9	* *	17	* *	59	* *	59	* *
Elk	53	* *	* *	47	* *	* *	2	* *	8	* *	50	* *	50	* *
Ellis	421	14.0 (55)	37.8 (2)	381	19.2 (9)	43.8 (18)	89	* *	103	13.6 (1)	370	2.2 (44)	370	4.6(47)
Ellsworth	85	* *	* *	76	* *	* *	9	* *	22	* *	76	* *	76	* *
Finney	2,025	11.6 (40)	44.7 (20)	1,863	25.6 (51)	39.9 (7)	257	29.2 (9)	265	23.8(13)	1,860	2.6 (51)	1,860	1.4(4)
Ford	1,824	7.0 (5)	45.3 (25)	1,652	24.3 (43)	43.2 (15)	258	22.5 (3)	187	19.8 (7)	1,661	1.2 (19)	1,661	1.9(11)
Franklin	484	11.4 (37)	46.3 (40)	423	17.5 (3)	48.2 (43)	48	* *	115	17.4 (5)	425	0.7 (7)	425	4.7(49)
Geary	916	10.9 (31)	45.4 (26)	771	20.4 (16)	48.5 (46)	150	38.0 (16)	137	28.5(21)	775	2.5 (50)	775	4.6(47)
Gove	33	* *	* *	29	* *	* *	6	* *	8	* *	32	* *	32	* *
Graham	50	* *	* *	42	* *	* *	6	* *	9	* *	46	* *	46	* *
Grant	242	9.1 (14)	53.7 (60)	233	20.2 (12)	40.8 (9)	41	* *	67	* *	226	1.3 (21)	226	0.4(2)

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one county's rate to other counties. Rank 1 = best rate.
- (4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).
- (5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).
- (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- \* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 5B**  
**2006 - 2008 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Maternal Health Indicators by County**  
**Combined 3 Years**

<b>Comparison of Maternal Health Indicators</b> <sup>(2, 3)</sup>														
<b>County</b>	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Gray	164	5.5 (3)	45.7 (33)	144	31.3 (56)	34.7 (2)	22	* *	42	* *	143	0.7 (7)	143	4.9(52)
Greeley	23	* *	* *	22	* *	* *	3	* *	6	* *	22	* *	22	* *
Greenwood	127	7.9 (8)	52.0 (58)	105	21.9 (28)	46.7 (35)	15	* *	27	* *	118	0.8 (10)	118	3.4(38)
Hamilton	55	* *	* *	46	* *	* *	9	* *	19	* *	47	* *	47	* *
Harper	146	17.8 (64)	42.5 (8)	118	33.1 (59)	39.0 (6)	17	* *	17	* *	130	1.5 (26)	129	4.7(49)
Harvey	755	10.7 (28)	47.5 (45)	687	22.6 (35)	46.7 (35)	108	28.7 (8)	126	26.2(17)	678	2.1 (42)	676	2.4(20)
Haskell	64	* *	* *	58	* *	* *	14	* *	25	* *	59	* *	59	* *
Hodgeman	30	* *	* *	29	* *	* *	6	* *	7	* *	26	* *	26	* *
Jackson	234	10.7 (28)	46.6 (41)	202	23.8 (42)	44.1 (21)	37	* *	48	* *	221	0.5 (4)	221	1.4(4)
Jefferson	191	14.7 (57)	48.2 (48)	161	18.0 (5)	48.4 (45)	26	* *	51	* *	173	2.9 (53)	173	1.2(3)
Jewell	39	* *	* *	39	* *	* *	4	* *	6	* *	40	* *	40	* *
Johnson	5,232	10.5 (26)	44.1 (16)	4,534	20.8 (20)	45.8 (27)	896	23.8 (5)	1,423	16.3 (3)	4,686	3.9 (56)	4,686	2.2(18)
Kearny	53	* *	* *	50	* *	* *	15	* *	19	* *	52	* *	52	* *
Kingman	158	13.9 (53)	44.3 (17)	120	22.5 (34)	47.5 (38)	21	* *	40	* *	136	4.4 (59)	136	5.1(53)
Kiowa	38	* *	* *	29	* *	* *	3	* *	5	* *	34	* *	34	* *
Labette	517	10.1 (22)	48.4 (49)	431	23.0 (36)	48.3 (44)	86	* *	114	18.4 (6)	471	1.3 (21)	471	4.2(45)
Lane	37	* *	* *	30	* *	* *	9	* *	4	* *	31	* *	31	* *

- (1) Reporting period is January 1 through December 31.
  - (2) Excludes records with unknown data or errors.
  - (3) Rank compares one county's rate to other counties. Rank 1 = best rate.
  - (4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).
  - (5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).
  - (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- \* Percentages are not calculated if <100 records are available for analysis after exclusions.





**Table 5B**  
**2006 - 2008 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Maternal Health Indicators by County**  
**Combined 3 Years**

Comparison of Maternal Health Indicators <sup>(2, 3)</sup>														
County	Prepregnancy BMI <sup>(4)</sup>			Weight Gain <sup>(5)</sup>			Anemia (Low Hb/Hct) <sup>(6)</sup>				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Leavenworth	1,006	11.0 (32)	46.0 (36)	884	20.9 (22)	50.1 (55)	161	41.0 (17)	210	22.4 (10)	867	0.9 (14)	868	2.0 (15)
Lincoln	65	* *	* *	54	* *	* *	9	* *	12	* *	57	* *	57	* *
Linn	173	11.0 (32)	49.1 (51)	160	28.8 (55)	43.8 (18)	23	* *	33	* *	162	0.6 (6)	162	3.1 (31)
Logan	27	* *	* *	22	* *	* *	5	* *	6	* *	22	* *	22	* *
Lyon	1,058	8.8 (13)	50.1 (56)	970	23.5 (40)	47.7 (40)	157	17.2 (2)	183	14.2 (2)	958	1.5 (26)	958	5.2 (55)
Marion	418	13.4 (51)	47.1 (44)	379	23.7 (41)	42.2 (10)	69	* *	83	* *	380	1.1 (16)	380	3.4 (38)
Marshall	144	10.4 (25)	44.4 (18)	132	20.5 (17)	34.8 (3)	16	* *	29	* *	128	0.8 (10)	128	3.9 (43)
Mcperson	166	9.6 (17)	45.2 (23)	124	26.6 (52)	42.7 (11)	27	* *	28	* *	136	0.0 (1)	136	1.5 (6)
Meade	122	6.6 (4)	50.0 (55)	106	24.5 (46)	49.1 (51)	18	* *	21	* *	106	1.9 (40)	106	1.9 (11)
Miami	368	14.7 (57)	45.4 (26)	343	20.1 (11)	49.6 (53)	78	* *	89	* *	334	1.8 (36)	334	1.8 (10)
Mitchell	103	9.7 (18)	45.6 (32)	94	* *	* *	18	* *	32	* *	91	* *	91	* *
Montgomery	992	15.4 (63)	42.4 (7)	827	26.6 (52)	43.7 (17)	169	33.1 (14)	260	26.9 (19)	875	0.8 (10)	875	3.2 (33)
Morris	84	* *	* *	68	* *	* *	11	* *	15	* *	75	* *	75	* *
Nemaha	104	8.7 (12)	34.6 (1)	98	* *	* *	25	* *	33	* *	97	* *	97	* *
Neosho	439	15.3 (62)	46.2 (39)	401	17.5 (3)	46.4 (33)	77	* *	68	* *	400	1.3 (21)	400	2.0 (15)
Ness	22	* *	* *	19	* *	* *	4	* *	11	* *	17	* *	17	* *
Norton	80	* *	* *	61	* *	* *	7	* *	16	* *	69	* *	69	* *

(1) Reporting period is January 1 through December 31.  
(2) Excludes records with unknown data or errors.  
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.  
(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).  
(5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).  
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.  
\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 5B**  
**2006 - 2008 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Maternal Health Indicators by County**  
**Combined 3 Years**

Comparison of Maternal Health Indicators <sup>(2, 3)</sup>														
County	Prepregnancy BMI <sup>(4)</sup>			Weight Gain <sup>(5)</sup>			Anemia (Low Hb/Hct) <sup>(6)</sup>				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Osage	235	11.1 (34)	45.5 (28)	189	23.3 (38)	47.6 (39)	28	* *	48	* *	209	2.4 (47)	209	2.9(28)
Osborne	54	* *	* *	49	* *	* *	8	* *	7	* *	51	* *	51	* *
Ottawa	75	* *	* *	66	* *	* *	9	* *	20	* *	65	* *	65	* *
Pawnee	145	15.2 (61)	53.1 (59)	128	32.0 (57)	40.6 (8)	11	* *	22	* *	128	0.8 (10)	128	3.1(31)
Phillips	98	* *	* *	82	* *	* *	5	* *	13	* *	86	* *	86	* *
Pottawatom	128	7.0 (5)	56.3 (62)	115	24.3 (43)	44.3 (22)	15	* *	53	* *	116	1.7 (32)	116	1.7(9)
Pratt	223	9.9 (19)	43.0 (10)	193	20.2 (12)	46.1 (29)	29	* *	56	* *	191	0.5 (4)	191	1.6(7)
Rawlins	23	* *	* *	21	* *	* *	3	* *	10	* *	21	* *	21	* *
Reno	1,419	11.8 (42)	46.7 (42)	1,282	21.8 (27)	46.3 (31)	213	28.6 (7)	346	22.8(11)	1,243	1.7 (32)	1,244	2.4(20)
Republic	87	* *	* *	81	* *	* *	12	* *	12	* *	83	* *	83	* *
Rice	204	9.3 (16)	49.5 (53)	181	21.5 (26)	44.8 (24)	23	* *	34	* *	184	1.1 (16)	184	1.6(7)
Riley	2,553	10.2 (23)	41.3 (4)	2,074	18.4 (7)	49.0 (49)	362	23.5 (4)	433	28.6(22)	2,120	1.6 (31)	2,120	2.5(23)
Rooks	116	12.1 (47)	43.1 (12)	102	25.5 (50)	38.2 (4)	17	* *	25	* *	110	0.9 (14)	110	3.6(40)
Rush	46	* *	* *	41	* *	* *	8	* *	7	* *	41	* *	41	* *
Russell	116	14.7 (57)	41.4 (5)	102	32.4 (58)	30.4 (1)	18	* *	34	* *	103	1.9 (40)	103	1.9(11)
Saline	1,328	11.4 (37)	44.8 (21)	1,216	21.2 (23)	45.4 (26)	248	23.8 (5)	297	20.9 (8)	1,201	1.5 (26)	1,201	3.2(33)
Scott	140	14.3 (56)	42.9 (9)	123	17.1 (2)	55.3 (58)	19	* *	31	* *	127	0.0 (1)	127	0.0(1)

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one county's rate to other counties. Rank 1 = best rate.
- (4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).
- (5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).
- (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- \* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 5B**  
**2006 - 2008 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Maternal Health Indicators by County**  
**Combined 3 Years**

Comparison of Maternal Health Indicators <sup>(2, 3)</sup>														
County	Prepregnancy BMI <sup>(4)</sup>			Weight Gain <sup>(5)</sup>			Anemia (Low Hb/Hct) <sup>(6)</sup>				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Sedgwick	11,992	11.2 (35)	45.2 (23)	11,054	20.6 (19)	46.3 (31)	2,467	32.4 (13)	3,230	25.4 (16)	10,966	1.8 (36)	10,965	2.9 (28)
Seward	1,393	9.1 (14)	45.9 (35)	1,269	20.3 (15)	45.3 (25)	242	29.3 (10)	203	24.6 (15)	1,271	2.2 (44)	1,269	3.2 (33)
Shawnee	4,153	10.7 (28)	45.8 (34)	3,671	21.3 (24)	48.0 (42)	425	41.4 (18)	711	28.4 (20)	3,718	2.9 (53)	3,721	2.6 (26)
Sheridan	28	* *	* *	22	* *	* *	3	* *	5	* *	24	* *	24	* *
Sherman	217	12.0 (45)	47.9 (47)	185	22.2 (32)	45.9 (28)	16	* *	33	* *	195	1.5 (26)	195	5.1 (53)
Smith	61	* *	* *	42	* *	* *	11	* *	22	* *	54	* *	54	* *
Stafford	118	8.5 (11)	56.8 (63)	98	* *	* *	15	* *	19	* *	99	* *	99	* *
Stanton	58	* *	* *	54	* *	* *	7	* *	21	* *	54	* *	54	* *
Stevens	193	5.2 (2)	49.2 (52)	173	20.2 (12)	49.7 (54)	29	* *	48	* *	178	3.4 (55)	178	4.5 (46)
Sumner	415	12.0 (45)	43.1 (12)	285	21.4 (25)	47.7 (40)	87	* *	93	* *	357	1.7 (32)	357	3.9 (43)
Thomas	154	14.9 (60)	46.1 (38)	140	22.1 (30)	38.6 (5)	20	* *	28	* *	141	2.8 (52)	141	2.8 (27)
Trego	44	* *	* *	37	* *	* *	5	* *	9	* *	45	* *	45	* *
Wabaunsee	54	* *	* *	48	* *	* *	5	* *	19	* *	50	* *	50	* *
Wallace	7	* *	* *	7	* *	* *	1	* *	1	* *	7	* *	7	* *
Washington	76	* *	* *	69	* *	* *	3	* *	15	* *	71	* *	70	* *
Wichita	57	* *	* *	54	* *	* *	7	* *	10	* *	53	* *	53	* *
Wilson	284	11.3 (36)	44.0 (15)	232	22.4 (33)	44.0 (20)	48	* *	45	* *	267	1.1 (16)	267	3.7 (41)

(1) Reporting period is January 1 through December 31.  
(2) Excludes records with unknown data or errors.  
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.  
(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).  
(5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).  
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.  
\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 5B**  
**2006 - 2008 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Maternal Health Indicators by County**  
**Combined 3 Years**

<b>Comparison of Maternal Health Indicators</b> <sup>(2, 3)</sup>														
<b>County</b>	<b>Prepregnancy BMI</b> (4)			<b>Weight Gain</b> (5)			<b>Anemia</b> (Low Hb/Hct) (6)				<b>Gestational Diabetes</b> During Pregnancy		<b>Hypertension</b> During Pregnancy	
	<b>Under-weight</b>		<b>Over-weight</b>	<b>&lt; Ideal</b>		<b>&gt; Ideal</b>	<b>3rd Trimester</b>		<b>Postpartum</b>					
	<b>Number</b>	<b>% (Rank)</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>
Woodson	26	* *	* *	22	* *	* *	4	* *	3	* *	25	* *	25	* *
Wyandotte	6,225	9.9 (19)	46.8 (43)	5,748	22.1 (30)	46.1 (29)	1,103	36.0 (15)	1,132	26.4 (18)	5,747	1.4 (24)	5,748	2.5 (23)
<b>Kansas</b>	<b>60,398</b>	<b>10.9</b>	<b>45.5</b>	<b>53,675</b>	<b>21.7</b>	<b>46.1</b>	<b>9,707</b>	<b>30.3</b>	<b>12,867</b>	<b>23.4</b>	<b>54,316</b>	<b>1.9</b>	<b>54,314</b>	<b>2.9</b>

(1) Reporting period is January 1 through December 31.  
(2) Excludes records with unknown data or errors.  
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.  
(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).  
(5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).  
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.  
\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 6B**  
**2006 - 2008 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Maternal Behavioral Indicators by County**  
**Combined 3 Years**

<b>Comparison of Maternal Behavioral Indicators</b> <sup>(2, 3)</sup>										
<b>County</b>	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Pregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	%	(Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Allen	272	90.1 (5)	1.1 (16)		253	41.5 (39)	69	* *	151	84.1 (18)
Anderson	154	87.7 (14)	1.3 (18)		121	49.6 (10)	38	* *	66	* *
Atchison	352	90.1 (5)	1.7 (26)		329	37.1 (48)	81	* *	158	74.7 (37)
Barber	94	* *	* *		81	* *	24	* *	44	* *
Barton	747	80.3 (35)	0.9 (11)		671	57.8 (4)	157	19.7 (2)	367	86.9 (3)
Bourbon	332	93.7 (1)	0.9 (11)		298	45.0 (24)	77	* *	156	78.8 (31)
Brown	245	82.9 (29)	6.9 (56)		231	45.5 (23)	64	* *	165	81.2 (24)
Butler	715	76.8 (44)	6.7 (55)		557	44.2 (29)	154	12.3 (9)	317	77.0 (35)
Chase	36	* *	* *		30	* *	9	* *	18	* *
Chautauqua	70	* *	* *		74	* *	16	* *	41	* *
Cherokee	513	80.3 (35)	4.9 (47)		470	47.4 (18)	115	9.6 (15)	277	79.8 (28)
Cheyenne	5	* *	* *		4	* *	0	* *	2	* *
Clark	8	* *	* *		11	* *	0	* *	8	* *
Clay	187	82.9 (29)	3.7 (40)		154	48.1 (15)	35	* *	55	* *
Cloud	206	85.9 (18)	1.9 (27)		181	54.7 (5)	44	* *	41	* *
Coffey	74	* *	* *		52	* *	14	* *	28	* *
Comanche	12	* *	* *		10	* *	3	* *	6	* *
Cowley	882	73.1 (57)	3.6 (38)		797	44.5 (26)	194	10.8 (14)	416	80.3 (27)
Crawford	1,036	89.0 (7)	0.8 (8)		947	46.4 (21)	235	8.1 (16)	571	85.8 (11)
Decatur	27	* *	* *		28	* *	5	* *	18	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 6B**  
**2006 - 2008 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Maternal Behavioral Indicators by County**  
**Combined 3 Years**

<b>Comparison of Maternal Behavioral Indicators</b> <sup>(2, 3)</sup>										
<b>County</b>	Medical Care			WIC Enrollment		Multivitamin Consumption				
	1st Trimester		No Care	1st Trimester		Pregnancy (4)		During Pregnancy (5)		
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	
Dickinson	304	83.6 (26)	3.6 (38)	252	48.4 (13)	68	* *	136	81.6 (22)	
Doniphan	105	80.0 (38)	4.8 (46)	98	* *	27	* *	66	* *	
Douglas	1,199	74.1 (54)	5.8 (53)	1,122	44.0 (30)	258	12.4 (7)	609	81.0 (26)	
Edwards	62	* *	* *	62	* *	19	* *	29	* *	
Elk	51	* *	* *	39	* *	5	* *	17	* *	
Ellis	396	86.1 (17)	1.0 (14)	349	37.0 (49)	95	* *	215	85.6 (13)	
Ellsworth	83	* *	* *	61	* *	16	* *	30	* *	
Finney	1,836	70.6 (58)	9.3 (59)	1,682	41.7 (37)	413	5.8 (19)	904	72.2 (38)	
Ford	1,654	75.0 (52)	9.4 (60)	1,562	47.4 (18)	347	5.5 (20)	774	78.0 (32)	
Franklin	386	81.3 (33)	1.6 (24)	346	39.9 (44)	91	* *	161	86.3 (7)	
Geary	752	73.7 (55)	0.4 (3)	769	45.6 (22)	195	15.9 (3)	431	77.5 (34)	
Gove	32	* *	* *	24	* *	3	* *	14	* *	
Graham	46	* *	* *	41	* *	4	* *	21	* *	
Grant	226	85.0 (22)	4.9 (47)	195	36.9 (51)	36	* *	109	86.2 (9)	
Gray	149	83.2 (28)	2.0 (28)	137	51.1 (9)	32	* *	69	* *	
Greeley	22	* *	* *	16	* *	7	* *	10	* *	
Greenwood	105	73.3 (56)	18.1 (62)	108	51.9 (7)	20	* *	56	* *	
Hamilton	45	* *	* *	41	* *	10	* *	21	* *	
Harper	127	76.4 (48)	7.1 (57)	123	49.6 (10)	28	* *	63	* *	
Harvey	665	74.3 (53)	9.8 (61)	633	47.9 (16)	138	11.6 (12)	312	84.3 (17)	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 6B**  
**2006 - 2008 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Maternal Behavioral Indicators by County**  
**Combined 3 Years**

<b>Comparison of Maternal Behavioral Indicators</b> <sup>(2, 3)</sup>										
<b>County</b>	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Pregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	%	(Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Haskell	59	* *	* *	* *	57	* *	16	* *	29	* *
Hodgeman	28	* *	* *	* *	20	* *	7	* *	10	* *
Jackson	188	84.6 (24)	3.7 (40)		188	42.6 (34)	39	* *	101	90.1 (1)
Jefferson	164	87.8 (12)	1.2 (17)		138	31.9 (57)	36	* *	87	* *
Jewell	40	* *	* *	* *	29	* *	8	* *	17	* *
Johnson	4,618	66.4 (62)	2.1 (30)		4,197	32.0 (56)	1,022	11.5 (13)	2,340	85.0 (15)
Kearny	51	* *	* *	* *	45	* *	10	* *	20	* *
Kingman	140	79.3 (42)	7.1 (57)		128	44.5 (26)	39	* *	69	* *
Kiowa	33	* *	* *	* *	36	* *	6	* *	23	* *
Labette	444	87.8 (12)	1.4 (19)		443	43.8 (31)	92	* *	262	86.6 (5)
Lane	32	* *	* *	* *	31	* *	7	* *	17	* *
Leavenworth	923	85.5 (19)	0.7 (4)		738	39.0 (46)	202	12.4 (7)	429	86.5 (6)
Lincoln	61	* *	* *	* *	47	* *	14	* *	23	* *
Linn	162	79.6 (40)	3.7 (40)		146	43.8 (31)	30	* *	82	* *
Logan	18	* *	* *	* *	24	* *	8	* *	17	* *
Lyon	992	81.7 (32)	0.8 (8)		823	43.3 (33)	193	14.0 (6)	458	84.5 (16)
Marion	395	88.4 (10)	1.0 (14)		327	44.6 (25)	74	* *	190	86.3 (7)
Marshall	136	80.1 (37)	1.5 (20)		106	58.5 (3)	27	* *	60	* *
Mcperson	146	91.1 (3)	0.7 (4)		131	40.5 (42)	33	* *	50	* *
Meade	109	75.2 (51)	5.5 (51)		101	46.5 (20)	26	* *	49	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 6B**  
**2006 - 2008 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Maternal Behavioral Indicators by County**  
**Combined 3 Years**

Comparison of Maternal Behavioral Indicators <sup>(2, 3)</sup>										
County	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Pregnancy <sup>(4)</sup>		During Pregnancy <sup>(5)</sup>	
	Number	% (Rank)	% (Rank)		Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Miami	353	76.8 (44)	0.3 (2)		263	35.4 (53)	74	* *	138	76.8 (36)
Mitchell	96	* *	* *		77	* *	24	* *	43	* *
Montgomery	822	86.4 (15)	2.9 (34)		831	42.1 (35)	186	14.5 (5)	477	84.1 (18)
Morris	73	* *	* *		65	* *	14	* *	30	* *
Nemaha	102	88.2 (11)	3.9 (43)		80	* *	16	* *	35	* *
Neosho	397	90.4 (4)	2.0 (28)		369	44.4 (28)	92	* *	181	81.2 (24)
Ness	21	* *	* *		16	* *	5	* *	7	* *
Norton	74	* *	* *		59	* *	14	* *	32	* *
Osage	205	91.2 (2)	1.5 (20)		183	51.9 (7)	51	* *	61	* *
Osborne	50	* *	* *		41	* *	6	* *	23	* *
Ottawa	68	* *	* *		56	* *	11	* *	29	* *
Pawnee	131	84.0 (25)	1.5 (20)		126	63.5 (2)	28	* *	62	* *
Phillips	92	* *	* *		74	* *	20	* *	58	* *
Pottawatom	104	80.8 (34)	0.0 (1)		100	35.0 (54)	19	* *	52	* *
Pratt	202	70.3 (59)	6.4 (54)		178	47.8 (17)	41	* *	104	85.6 (13)
Rawlins	16	* *	* *		19	* *	6	* *	12	* *
Reno	1,303	85.1 (20)	1.5 (20)		1,123	42.0 (36)	291	23.7 (1)	678	86.9 (3)
Republic	86	* *	* *		73	* *	13	* *	18	* *
Rice	188	76.6 (47)	1.6 (24)		171	40.9 (40)	42	* *	61	* *
Riley	1,836	82.5 (31)	0.7 (4)		1,966	40.1 (43)	472	14.6 (4)	1,160	83.4 (20)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.





**Table 6B**  
**2006 - 2008 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Maternal Behavioral Indicators by County**  
**Combined 3 Years**

<b>Comparison of Maternal Behavioral Indicators</b> <sup>(2, 3)</sup>										
<b>County</b>	<b>Medical Care</b>			<b>WIC Enrollment</b>			<b>Multivitamin Consumption</b>			
	<b>1st Trimester</b>		<b>No Care</b>	<b>1st Trimester</b>		<b>Prepregnancy</b> (4)		<b>During Pregnancy</b> (5)		
	<b>Number</b>	<b>% (Rank)</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	
Rooks	112	84.8 (23)	2.7 (33)	97	* *	17	* *	59	* *	
Rush	41	* *	* *	41	* *	8	* *	22	* *	
Russell	109	89.0 (7)	0.9 (11)	95	* *	23	* *	57	* *	
Saline	1,190	76.1 (49)	3.2 (36)	1,050	34.4 (55)	271	5.9 (18)	612	79.7 (29)	
Scott	128	79.7 (39)	5.5 (51)	113	39.8 (45)	18	* *	66	* *	
Sedgwick	11,318	76.7 (46)	2.6 (32)	9,638	28.8 (58)	2,306	11.9 (11)	5,099	77.8 (33)	
Seward	1,287	68.6 (61)	5.1 (50)	1,151	37.4 (47)	267	6.0 (17)	596	85.7 (12)	
Shawnee	3,733	75.7 (50)	3.0 (35)	3,399	48.2 (14)	785	5.0 (21)	1,691	81.5 (23)	
Sheridan	23	* *	* *	21	* *	2	* *	14	* *	
Sherman	174	85.1 (20)	4.0 (44)	176	54.5 (6)	39	* *	106	88.7 (2)	
Smith	39	* *	* *	51	* *	9	* *	31	* *	
Stafford	103	79.6 (40)	4.9 (47)	101	64.4 (1)	25	* *	61	* *	
Stanton	54	* *	* *	48	* *	11	* *	25	* *	
Stevens	188	77.1 (43)	3.2 (36)	146	37.0 (49)	38	* *	52	* *	
Sumner	348	86.2 (16)	2.3 (31)	320	40.6 (41)	84	* *	167	86.2 (9)	
Thomas	132	83.3 (27)	0.8 (8)	119	49.6 (10)	23	* *	87	* *	
Trego	46	* *	* *	37	* *	6	* *	17	* *	
Wabaunsee	51	* *	* *	33	* *	10	* *	19	* *	
Wallace	8	* *	* *	5	* *	1	* *	4	* *	
Washington	75	* *	* *	63	* *	15	* *	35	* *	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 6B**  
**2006 - 2008 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Maternal Behavioral Indicators by County**  
**Combined 3 Years**

<b>Comparison of Maternal Behavioral Indicators</b> <sup>(2, 3)</sup>										
<b>County</b>	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Pregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	%	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Wichita	54	* *	* *	* *	46	* *	6	* *	30	* *
Wilson	271	88.6 (9)	0.7 (4)	(4)	252	41.7 (37)	50	* *	139	79.1 (30)
Woodson	24	* *	* *	* *	21	* *	0	* *	11	* *
Wyandotte	5,883	69.3 (60)	4.2 (45)	(45)	5,137	35.7 (52)	1,171	12.3 (9)	2,640	83.3 (21)
<b>Kansas</b>	<b>54,561</b>	<b>76.9</b>	<b>3.2</b>	<b>(45)</b>	<b>48,744</b>	<b>39.3</b>	<b>11,645</b>	<b>12.0</b>	<b>26,186</b>	<b>81.6</b>

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 7B**  
**2006 - 2008 Pregnancy Nutrition Surveillance** (1)  
**Kansas**  
**Comparison of Smoking Indicators by County**  
**Combined 3 Years**

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Allen	294	45.2 (40)	263	32.7 (48)	96	* *	247	36.0 (48)	261	32.6 (48)
Anderson	153	57.5 (61)	140	33.6 (49)	59	* *	119	33.6 (46)	141	20.6 (32)
Atchison	321	44.9 (37)	235	21.3 (18)	74	* *	290	32.8 (43)	239	28.5 (45)
Barber	102	43.1 (33)	91	* *	33	* *	81	* *	92	* *
Barton	826	39.8 (23)	745	24.4 (25)	229	29.7 (20)	662	18.4 (17)	747	7.4 (9)
Bourbon	359	47.6 (50)	322	28.0 (39)	111	26.1 (26)	289	28.4 (38)	323	30.7 (47)
Brown	270	42.2 (31)	218	25.7 (29)	65	* *	221	41.6 (51)	220	38.6 (52)
Butler	762	53.8 (56)	646	33.6 (49)	248	30.6 (19)	539	19.5 (19)	645	9.5 (12)
Chase	39	* *	35	* *	18	* *	30	* *	35	* *
Chautauqua	84	* *	61	* *	18	* *	70	* *	61	* *
Cherokee	623	41.7 (30)	519	25.6 (28)	133	21.8 (28)	450	32.9 (45)	537	27.6 (44)
Cheyenne	5	* *	5	* *	0	* *	4	* *	5	* *
Clark	8	* *	5	* *	0	* *	9	* *	5	* *
Clay	202	52.0 (55)	179	30.2 (42)	74	* *	149	30.2 (40)	179	25.1 (41)
Cloud	229	41.0 (27)	195	21.5 (21)	60	* *	179	6.7 (1)	194	2.6 (1)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 7B**  
**2006 - 2008 Pregnancy Nutrition Surveillance** (1)  
**Kansas**  
**Comparison of Smoking Indicators by County**  
**Combined 3 Years**

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Coffey	74	* *	72	* *	24	* *	51	* *	73	* *
Comanche	12	* *	11	* *	4	* *	10	* *	11	* *
Cowley	976	47.1 (47)	857	31.3 (44)	325	25.5 (27)	780	21.5 (25)	863	17.5 (26)
Crawford	1,156	44.8 (36)	963	26.5 (30)	337	33.5 (14)	924	38.6 (49)	963	34.1 (50)
Decatur	31	* *	27	* *	8	* *	28	* *	27	* *
Dickinson	328	47.0 (45)	293	27.6 (37)	107	26.2 (25)	248	12.5 (7)	297	7.1 (6)
Doniphan	99	* *	68	* *	29	* *	79	* *	70	* *
Douglas	1,390	44.0 (35)	1,151	20.7 (17)	362	44.2 (4)	1,096	12.6 (9)	1,157	7.1 (6)
Edwards	76	* *	64	* *	21	* *	61	* *	65	* *
Elk	56	* *	52	* *	13	* *	37	* *	52	* *
Ellis	431	37.8 (19)	377	14.6 (10)	122	53.3 (1)	344	12.5 (7)	375	8.8 (11)
Ellsworth	85	* *	77	* *	30	* *	61	* *	79	* *
Finney	2,041	15.7 (5)	1,866	7.1 (4)	227	48.9 (2)	1,676	19.7 (20)	1,871	17.7 (27)
Ford	1,808	10.7 (2)	1,519	4.9 (2)	119	35.3 (10)	1,544	9.2 (3)	1,528	7.3 (8)
Franklin	499	46.7 (43)	434	27.2 (35)	128	35.2 (11)	335	20.9 (24)	447	14.1 (18)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 7B**  
**2006 - 2008 Pregnancy Nutrition Surveillance** (1)  
**Kansas**  
**Comparison of Smoking Indicators by County**  
**Combined 3 Years**

<b>Comparison of Smoking Indicators</b> (2, 3)										
<b>County</b>	<b>Smoking</b>				<b>Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)</b>		<b>Smoking in Household</b>			
	<b>3 Months Prior to Pregnancy</b>		<b>Last 3 Months of Pregnancy</b>				<b>Prenatal</b>		<b>Postpartum</b>	
	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>
Geary	914	43.5 (34)	752	21.3 (18)	268	38.8 (5)	756	39.9 (50)	755	41.1 (53)
Gove	34	* *	31	* *	9	* *	23	* *	31	* *
Graham	49	* *	44	* *	21	* *	40	* *	44	* *
Grant	243	13.2 (3)	235	5.5 (3)	22	* *	194	17.5 (15)	235	14.0 (17)
Gray	167	13.8 (4)	146	8.9 (5)	16	* *	137	15.3 (12)	148	12.2 (15)
Greeley	23	* *	22	* *	4	* *	16	* *	22	* *
Greenwood	121	47.1 (47)	73	* *	30	* *	107	33.6 (46)	107	22.4 (37)
Hamilton	54	* *	47	* *	3	* *	40	* *	47	* *
Harper	141	40.4 (24)	105	21.9 (22)	32	* *	122	27.9 (34)	109	15.6 (21)
Harvey	759	35.2 (17)	660	23.5 (23)	188	27.1 (23)	621	6.8 (2)	663	4.8 (4)
Haskell	63	* *	61	* *	8	* *	54	* *	61	* *
Hodgeman	28	* *	24	* *	5	* *	20	* *	24	* *
Jackson	230	47.0 (45)	207	26.6 (32)	74	* *	183	27.9 (34)	212	19.3 (29)
Jefferson	127	54.3 (57)	87	* *	33	* *	102	44.1 (54)	96	* *
Jewell	41	* *	39	* *	8	* *	30	* *	39	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 7B**  
**2006 - 2008 Pregnancy Nutrition Surveillance** (1)  
**Kansas**  
**Comparison of Smoking Indicators by County**  
**Combined 3 Years**

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Johnson	5,311	26.2 (9)	4,684	11.1 (6)	837	47.2 (3)	4,065	20.2 (21)	4,697	18.7 (28)
Kearny	53	* *	52	* *	9	* *	45	* *	52	* *
Kingman	139	32.4 (13)	121	12.4 (8)	19	* *	125	32.8 (43)	134	32.8 (49)
Kiowa	37	* *	28	* *	6	* *	33	* *	29	* *
Labette	513	40.9 (26)	447	24.6 (26)	134	20.9 (30)	437	30.2 (40)	467	26.1 (43)
Lane	29	* *	21	* *	3	* *	22	* *	22	* *
Leavenworth	1,028	48.7 (53)	867	27.8 (38)	283	28.6 (22)	738	42.4 (52)	878	42.6 (54)
Lincoln	62	* *	53	* *	21	* *	45	* *	53	* *
Linn	177	59.3 (63)	159	44.0 (55)	77	* *	144	20.8 (23)	161	14.9 (19)
Logan	27	* *	22	* *	7	* *	24	* *	22	* *
Lyon	1,070	30.7 (12)	974	15.7 (11)	210	36.7 (8)	808	16.5 (13)	977	13.4 (16)
Marion	427	41.0 (27)	387	26.6 (32)	123	26.8 (24)	320	10.6 (4)	390	4.6 (3)
Marshall	148	33.8 (15)	134	20.1 (15)	33	* *	105	11.4 (5)	135	5.2 (5)
Mcperson	170	58.8 (62)	135	43.0 (54)	61	* *	131	54.2 (56)	140	47.9 (56)
Meade	115	33.0 (14)	98	* *	23	* *	98	* *	99	* *

(1) Reporting period is January 1 through December 31.  
(2) Excludes records with unknown data or errors.  
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.  
(4) Among women who smoked 3 months prior to pregnancy.  
\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 7B**  
**2006 - 2008 Pregnancy Nutrition Surveillance** (1)  
**Kansas**  
**Comparison of Smoking Indicators by County**  
**Combined 3 Years**

<b>Comparison of Smoking Indicators</b> (2, 3)										
<b>County</b>	<b>Smoking</b>				<b>Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)</b>		<b>Smoking in Household</b>			
	<b>3 Months Prior to Pregnancy</b>		<b>Last 3 Months of Pregnancy</b>				<b>Prenatal</b>		<b>Postpartum</b>	
	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>
Miami	344	54.9 (58)	282	32.3 (46)	106	35.8 (9)	245	27.8 (33)	280	20.0 (31)
Mitchell	104	39.4 (21)	94	* *	31	* *	77	* *	95	* *
Montgomery	961	40.7 (25)	829	26.5 (30)	263	21.3 (29)	813	31.1 (42)	851	25.4 (42)
Morris	81	* *	70	* *	27	* *	63	* *	72	* *
Nemaha	106	47.2 (49)	99	* *	36	* *	80	* *	98	* *
Neosho	441	45.1 (39)	382	25.1 (27)	136	33.8 (13)	366	27.9 (34)	383	21.4 (34)
Ness	22	* *	20	* *	6	* *	16	* *	20	* *
Norton	80	* *	71	* *	24	* *	59	* *	70	* *
Osage	231	46.8 (44)	200	30.5 (43)	70	* *	180	28.3 (37)	200	25.0 (40)
Osborne	53	* *	50	* *	15	* *	40	* *	50	* *
Ottawa	72	* *	63	* *	17	* *	55	* *	63	* *
Pawnee	146	48.6 (52)	128	29.7 (41)	47	* *	124	24.2 (29)	128	22.7 (38)
Phillips	83	* *	58	* *	22	* *	64	* *	58	* *
Pottawatom	125	30.4 (11)	112	20.5 (16)	29	* *	96	* *	113	3.5 (2)
Pratt	223	38.1 (20)	126	32.5 (47)	51	* *	153	26.8 (32)	158	23.4 (39)

(1) Reporting period is January 1 through December 31.  
(2) Excludes records with unknown data or errors.  
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.  
(4) Among women who smoked 3 months prior to pregnancy.  
\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 7B**  
**2006 - 2008 Pregnancy Nutrition Surveillance** (1)  
**Kansas**  
**Comparison of Smoking Indicators by County**  
**Combined 3 Years**

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Rawlins	22	* *	20	* *	5	* *	19	* *	20	* *
Reno	1,421	46.1 (42)	1,261	23.9 (24)	443	34.5 (12)	1,109	22.5 (28)	1,267	17.4 (25)
Republic	89	* *	83	* *	30	* *	72	* *	84	* *
Rice	208	48.1 (51)	181	34.3 (52)	64	* *	173	30.1 (39)	182	22.0 (35)
Riley	2,509	29.9 (10)	1,949	16.2 (12)	388	38.7 (6)	1,905	18.6 (18)	2,102	17.2 (24)
Rooks	117	55.6 (59)	99	* *	47	* *	95	* *	105	30.5 (46)
Rush	45	* *	40	* *	19	* *	41	* *	40	* *
Russell	117	45.3 (41)	105	36.2 (53)	41	* *	95	* *	108	44.4 (55)
Saline	1,239	44.9 (37)	1,020	27.0 (34)	365	30.7 (18)	991	14.8 (10)	1,023	7.9 (10)
Scott	134	24.6 (8)	116	17.2 (14)	25	* *	111	26.1 (30)	118	22.0 (35)
Sedgwick	12,115	33.8 (15)	10,995	16.9 (13)	2,710	37.3 (7)	9,460	12.4 (6)	11,020	10.3 (14)
Seward	1,404	9.2 (1)	1,267	3.8 (1)	89	* *	1,137	17.6 (16)	1,273	16.2 (22)
Shawnee	4,027	36.9 (18)	3,345	21.4 (20)	883	29.6 (21)	3,266	17.4 (14)	3,340	10.2 (13)
Sheridan	25	* *	18	* *	3	* *	20	* *	18	* *
Sherman	221	43.0 (32)	187	28.3 (40)	60	* *	167	20.4 (22)	186	15.1 (20)

(1) Reporting period is January 1 through December 31.  
(2) Excludes records with unknown data or errors.  
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.  
(4) Among women who smoked 3 months prior to pregnancy.  
\* Percentages are not calculated if <100 records are available for analysis after exclusions.





**Table 7B**  
**2006 - 2008 Pregnancy Nutrition Surveillance** (1)  
**Kansas**  
**Comparison of Smoking Indicators by County**  
**Combined 3 Years**

<b>Comparison of Smoking Indicators</b> (2, 3)										
<b>County</b>	<b>Smoking</b>				<b>Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)</b>	<b>Smoking in Household</b>				
	<b>3 Months Prior to Pregnancy</b>		<b>Last 3 Months of Pregnancy</b>			<b>Prenatal</b>		<b>Postpartum</b>		
	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>
Smith	56	* *	40	* *	13	* *	49	* *	42	* *
Stafford	119	41.2 (29)	97	* *	34	* *	100	15.0 (11)	99	* *
Stanton	57	* *	52	* *	3	* *	47	* *	53	* *
Stevens	203	22.7 (6)	180	11.1 (6)	29	* *	145	22.1 (26)	181	19.3 (29)
Sumner	388	51.8 (54)	297	31.3 (44)	116	31.0 (16)	306	47.7 (55)	299	49.5 (57)
Thomas	152	39.5 (22)	135	27.4 (36)	37	* *	117	26.5 (31)	136	16.2 (22)
Trego	46	* *	35	* *	14	* *	37	* *	37	* *
Wabaunsee	54	* *	51	* *	14	* *	32	* *	51	* *
Wallace	6	* *	6	* *	3	* *	5	* *	6	* *
Washington	81	* *	74	* *	18	* *	63	* *	73	* *
Wichita	57	* *	52	* *	7	* *	47	* *	52	* *
Wilson	289	57.1 (60)	261	33.7 (51)	129	31.0 (16)	246	42.7 (53)	265	38.5 (51)
Woodson	27	* *	21	* *	8	* *	21	* *	22	* *
Wyandotte	6,269	24.2 (7)	5,632	13.3 (9)	1,006	32.4 (15)	5,036	22.4 (27)	5,688	21.0 (33)
<b>Kansas</b>	<b>60,208</b>	<b>34.1</b>	<b>52,408</b>	<b>18.6</b>	<b>13,124</b>	<b>34.0</b>	<b>47,541</b>	<b>20.5</b>	<b>52,940</b>	<b>17.1</b>

(1) Reporting period is January 1 through December 31.  
(2) Excludes records with unknown data or errors.  
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.  
(4) Among women who smoked 3 months prior to pregnancy.  
\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 8B**  
**2006 - 2008 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Infant Health Indicators by County**  
**Combined 3 Years**

<b>Comparison of Infant Health Indicators</b> <sup>(2, 3)</sup>									
<b>County</b>	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Allen	273	7.3 (25)	4.4 (10)	232	12.9 (39)	200	3.5 (30)	275	50.5 (57)
Anderson	149	6.7 (20)	4.7 (14)	115	11.3 (25)	102	5.9 (45)	144	50.7 (56)
Atchison	342	7.6 (29)	5.6 (21)	294	9.5 (13)	261	3.1 (23)	337	49.9 (58)
Barber	89	* * *	* *	72	* *	57	* *	89	* *
Barton	735	7.6 (29)	4.8 (15)	596	12.2 (31)	519	3.5 (30)	710	72.3 (22)
Bourbon	335	6.9 (21)	6.0 (27)	275	12.4 (32)	236	3.4 (27)	332	58.1 (46)
Brown	167	7.2 (24)	7.2 (42)	203	10.3 (17)	140	2.9 (19)	214	53.3 (54)
Butler	721	9.2 (47)	5.8 (24)	504	14.9 (51)	426	3.5 (30)	692	73.4 (19)
Chase	35	* * *	* *	27	* *	25	* *	35	* *
Chautauqua	63	* * *	* *	59	* *	45	* *	63	* *
Cherokee	586	7.7 (31)	7.3 (43)	400	13.0 (41)	347	2.3 (11)	561	57.9 (47)
Cheyenne	5	* * *	* *	4	* *	4	* *	5	* *
Clark	7	* * *	* *	7	* *	5	* *	8	* *
Clay	186	7.5 (26)	3.8 (6)	136	14.0 (48)	117	3.4 (27)	182	63.2 (37)

(1) Reporting period is January 1 through December 31.  
(2) Excludes records with unknown data or errors.  
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.  
(4) <2500 g.  
(5) >4000 g.  
(6) <37 weeks gestation.  
(7) 37 or more weeks gestation and <2500 g.  
\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 8B**  
**2006 - 2008 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Infant Health Indicators by County**  
**Combined 3 Years**

Comparison of Infant Health Indicators <sup>(2, 3)</sup>										
County	Birthweight				Preterm <sup>(6)</sup>		Full Term LBW <sup>(7)</sup>		Ever Breastfed	
	Low <sup>(4)</sup>		High <sup>(5)</sup>							
	Number	% <sup>(Rank)</sup>	% <sup>(Rank)</sup>		Number	% <sup>(Rank)</sup>	Number	% <sup>(Rank)</sup>	Number	% <sup>(Rank)</sup>
Cloud	200	8.0 (37)	3.5 (3)		154	16.2 (53)	128	3.1 (23)	193	67.4 (30)
Coffey	72	* *	* *		51	* *	47	* *	73	* *
Comanche	12	* *	* *		8	* *	7	* *	12	* *
Cowley	894	7.7 (31)	7.5 (44)		710	11.0 (23)	622	4.5 (40)	869	55.9 (51)
Crawford	1,002	9.2 (47)	4.9 (16)		803	12.6 (35)	695	4.3 (38)	1,009	60.0 (44)
Decatur	27	* *	* *		26	* *	26	* *	26	* *
Dickinson	297	6.1 (10)	7.7 (47)		223	8.5 (4)	201	2.0 (4)	294	73.8 (18)
Doniphan	89	* *	* *		82	* *	57	* *	101	46.5 (59)
Douglas	1,200	6.5 (16)	6.9 (36)		918	11.3 (25)	802	2.6 (17)	1,145	77.6 (6)
Edwards	61	* *	* *		51	* *	42	* *	62	* *
Elk	48	* *	* *		36	* *	28	* *	50	* *
Ellis	380	5.8 (6)	4.5 (11)		306	8.5 (4)	274	2.2 (8)	375	77.9 (5)
Ellsworth	79	* *	* *		56	* *	47	* *	78	* *
Finney	1,867	6.2 (11)	6.3 (30)		1,535	10.7 (20)	1,357	2.4 (15)	1,416	74.4 (17)

(1) Reporting period is January 1 through December 31.  
(2) Excludes records with unknown data or errors.  
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.  
(4) <2500 g.  
(5) >4000 g.  
(6) <37 weeks gestation.  
(7) 37 or more weeks gestation and <2500 g.  
\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 8B**  
**2006 - 2008 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Infant Health Indicators by County**  
**Combined 3 Years**

<b>Comparison of Infant Health Indicators</b> <sup>(2, 3)</sup>										
<b>County</b>	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)							
	Number	% (Rank)	%	(Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Ford	1,629	5.5 (4)	5.8	(24)	1,398	10.3 (17)	1,232	2.2 (8)	1,423	72.7 (20)
Franklin	435	6.9 (21)	5.3	(19)	287	13.2 (42)	246	2.8 (18)	419	53.9 (53)
Geary	797	6.6 (19)	7.0	(39)	628	8.3 (3)	571	2.3 (11)	734	68.5 (29)
Gove	18	* * *	* *	*	21	* *	9	* *	30	* *
Graham	40	* * *	* *	*	34	* *	30	* *	43	* *
Grant	228	9.2 (47)	7.5	(44)	186	11.3 (25)	161	3.7 (34)	183	75.4 (11)
Gray	121	5.8 (6)	6.6	(32)	116	12.1 (30)	90	* *	136	83.8 (1)
Greeley	15	* * *	* *	*	15	* *	11	* *	21	* *
Greenwood	115	8.7 (42)	3.5	(3)	99	* *	83	* *	109	62.4 (39)
Hamilton	44	* * *	* *	*	34	* *	32	* *	32	* *
Harper	92	* * *	* *	*	106	11.3 (25)	67	* *	129	76.0 (10)
Harvey	592	9.1 (46)	5.9	(26)	572	13.6 (45)	419	3.6 (33)	681	78.3 (4)
Haskell	62	* * *	* *	*	54	* *	51	* *	44	* *
Hodgeman	29	* * *	* *	*	19	* *	18	* *	25	* *

(1) Reporting period is January 1 through December 31.  
(2) Excludes records with unknown data or errors.  
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.  
(4) <2500 g.  
(5) >4000 g.  
(6) <37 weeks gestation.  
(7) 37 or more weeks gestation and <2500 g.  
\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 8B**  
**2006 - 2008 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Infant Health Indicators by County**  
**Combined 3 Years**

<b>Comparison of Infant Health Indicators</b> <sup>(2, 3)</sup>									
<b>County</b>	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Jackson	195	8.2 (39)	8.2 (49)	173	14.5 (50)	131	2.3 (11)	214	76.2 (9)
Jefferson	169	6.5 (16)	8.3 (50)	121	13.2 (42)	104	3.8 (36)	172	57.0 (50)
Jewell	40	* * *	* *	29	* *	25	* *	36	* *
Johnson	4,646	6.4 (13)	7.7 (47)	3,675	9.0 (9)	3,233	2.5 (16)	4,540	79.4 (3)
Kearny	51	* * *	* *	44	* *	36	* *	36	* *
Kingman	93	* * *	* *	107	15.9 (52)	67	* *	126	75.4 (11)
Kiowa	31	* * *	* *	30	* *	25	* *	32	* *
Labette	474	7.8 (33)	7.0 (39)	400	10.8 (21)	355	3.7 (34)	456	54.6 (52)
Lane	30	* * *	* *	25	* *	20	* *	29	* *
Leavenworth	865	8.3 (41)	4.5 (11)	625	9.3 (10)	553	2.9 (19)	893	61.4 (42)
Lincoln	54	* * *	* *	43	* *	34	* *	59	* *
Linn	160	6.9 (21)	5.0 (17)	132	13.6 (45)	113	4.4 (39)	156	60.3 (43)
Logan	22	* * *	* *	20	* *	19	* *	21	* *
Lyon	983	6.2 (11)	6.7 (33)	734	9.9 (16)	660	2.1 (5)	941	72.5 (21)

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one county's rate to other counties. Rank 1 = best rate.
- (4) <2500 g.
- (5) >4000 g.
- (6) <37 weeks gestation.
- (7) 37 or more weeks gestation and <2500 g.
- \* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 8B**  
**2006 - 2008 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Infant Health Indicators by County**  
**Combined 3 Years**

<b>Comparison of Infant Health Indicators</b> <sup>(2, 3)</sup>										
<b>County</b>	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed		
	Low (4)		High (5)							
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	
Marion	388	6.4 (13)	6.7 (33)	288	8.7 (8)	261	2.3 (11)	383	77.3 (7)	
Marshall	133	7.5 (26)	3.0 (1)	93	* *	81	* *	131	82.4 (2)	
Mcperson	129	11.6 (53)	3.9 (7)	105	10.5 (19)	87	* *	140	57.9 (47)	
Meade	99	* *	* *	88	* *	71	* *	94	* *	
Miami	352	6.5 (16)	5.7 (22)	233	12.4 (32)	202	4.5 (40)	305	67.2 (31)	
Mitchell	88	* *	* *	66	* *	61	* *	85	* *	
Montgomery	890	8.2 (39)	3.3 (2)	739	12.4 (32)	641	3.4 (27)	854	57.7 (49)	
Morris	66	* *	* *	57	* *	40	* *	70	* *	
Nemaha	81	* *	* *	75	* *	49	* *	101	65.3 (35)	
Neosho	336	8.0 (37)	7.1 (41)	341	13.8 (47)	240	3.3 (26)	419	51.6 (55)	
Ness	21	* *	* *	14	* *	13	* *	21	* *	
Norton	72	* *	* *	53	* *	44	* *	68	* *	
Osage	199	5.5 (4)	6.0 (27)	170	12.9 (39)	135	1.5 (2)	200	59.0 (45)	
Osborne	39	* *	* *	38	* *	25	* *	50	* *	

(1) Reporting period is January 1 through December 31.  
(2) Excludes records with unknown data or errors.  
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.  
(4) <2500 g.  
(5) >4000 g.  
(6) <37 weeks gestation.  
(7) 37 or more weeks gestation and <2500 g.  
\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 8B**  
**2006 - 2008 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Infant Health Indicators by County**  
**Combined 3 Years**

<b>Comparison of Infant Health Indicators</b> <sup>(2, 3)</sup>										
<b>County</b>	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)							
	Number	% (Rank)	%	(Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Ottawa	56	* * * *	*	*	49	* *	37	* *	61	* *
Pawnee	125	8.8 (44)	4.0	(9)	111	14.4 (49)	95	* *	122	68.9 (28)
Phillips	88	* * * *	*	*	63	* *	53	* *	87	* *
Pottawatom	115	8.7 (42)	6.1	(29)	90	* *	83	* *	113	75.2 (14)
Pratt	195	9.2 (47)	8.7	(53)	149	9.4 (12)	131	4.6 (42)	180	71.1 (24)
Rawlins	21	* * * *	*	*	17	* *	15	* *	20	* *
Reno	1,271	6.4 (13)	8.6	(51)	994	11.0 (23)	872	2.1 (5)	1,239	63.5 (36)
Republic	58	* * * *	*	*	66	* *	38	* *	83	* *
Rice	186	7.5 (26)	6.5	(31)	151	12.6 (35)	131	4.6 (42)	180	65.6 (33)
Riley	2,217	5.4 (3)	8.6	(51)	1,547	9.7 (15)	1,390	2.1 (5)	2,152	77.2 (8)
Rooks	101	5.9 (8)	5.0	(17)	90	* *	77	* *	104	70.2 (25)
Rush	40	* * * *	*	*	35	* *	32	* *	39	* *
Russell	103	9.7 (52)	3.9	(7)	86	* *	73	* *	98	* *
Saline	1,211	9.4 (51)	5.5	(20)	949	11.9 (29)	830	4.0 (37)	1,206	61.8 (41)

(1) Reporting period is January 1 through December 31.  
(2) Excludes records with unknown data or errors.  
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.  
(4) <2500 g.  
(5) >4000 g.  
(6) <37 weeks gestation.  
(7) 37 or more weeks gestation and <2500 g.  
\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 8B**  
**2006 - 2008 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Infant Health Indicators by County**  
**Combined 3 Years**

<b>Comparison of Infant Health Indicators</b> <sup>(2, 3)</sup>										
<b>County</b>	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)							
	Number	% (Rank)	%	(Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Scott	98	* * * *	* *		101	6.9 (2)	81	* *	117	70.1 (27)
Sedgwick	11,211	7.9 (35)	6.8 (35)		8,765	9.3 (10)	7,879	3.1 (23)	10,274	65.8 (32)
Seward	1,253	5.3 (2)	7.5 (44)		1,040	9.6 (14)	914	1.9 (3)	1,283	71.9 (23)
Shawnee	3,610	7.8 (33)	5.7 (22)		3,003	12.8 (37)	2,517	2.9 (19)	3,664	65.4 (34)
Sheridan	24	* * * *	* *		17	* *	15	* *	23	* *
Sherman	200	15.0 (54)	3.5 (3)		149	13.4 (44)	129	10.1 (46)	192	74.5 (16)
Smith	53	* * * *	* *		43	* *	37	* *	53	* *
Stafford	94	* * * *	* *		83	* *	75	* *	89	* *
Stanton	52	* * * *	* *		45	* *	40	* *	35	* *
Stevens	99	* * * *	* *		129	8.5 (4)	64	* *	179	75.4 (11)
Sumner	334	6.0 (9)	6.9 (36)		280	8.6 (7)	224	2.2 (8)	350	62.9 (38)
Thomas	146	3.4 (1)	8.9 (54)		109	6.4 (1)	102	1.0 (1)	140	75.0 (15)
Trego	35	* * * *	* *		34	* *	24	* *	41	* *
Wabaunsee	51	* * * *	* *		31	* *	28	* *	47	* *

(1) Reporting period is January 1 through December 31.  
(2) Excludes records with unknown data or errors.  
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.  
(4) <2500 g.  
(5) >4000 g.  
(6) <37 weeks gestation.  
(7) 37 or more weeks gestation and <2500 g.  
\* Percentages are not calculated if <100 records are available for analysis after exclusions.





**Table 8B**  
**2006 - 2008 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Kansas**  
**Comparison of Infant Health Indicators by County**  
**Combined 3 Years**

<b>Comparison of Infant Health Indicators</b> <sup>(2, 3)</sup>									
<b>County</b>	<b>Birthweight</b>			<b>Preterm</b> (6)		<b>Full Term LBW</b> (7)		<b>Ever Breastfed</b>	
	<b>Low</b> (4)		<b>High</b> (5)						
	<b>Number</b>	<b>% (Rank)</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>
Wallace	8	* * * *	*	5	* *	5	* *	8	* *
Washington	45	* * * *	*	58	* *	37	* *	72	* *
Wichita	37	* * * *	*	43	* *	22	* *	53	* *
Wilson	267	9.0 (45)	4.5 (11)	234	12.8 (37)	201	5.0 (44)	244	61.9 (40)
Woodson	23	* * * *	*	18	* *	14	* *	22	* *
Wyandotte	5,745	7.9 (35)	6.9 (36)	4,637	10.9 (22)	4,080	3.0 (22)	5,600	70.2 (25)
<b>Kansas</b>	<b>54,016</b>	<b>7.3</b>	<b>6.6</b>	<b>43,191</b>	<b>10.7</b>	<b>37,507</b>	<b>2.9</b>	<b>52,161</b>	<b>68.6</b>

(1) Reporting period is January 1 through December 31.  
(2) Excludes records with unknown data or errors.  
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.  
(4) <2500 g.  
(5) >4000 g.  
(6) <37 weeks gestation.  
(7) 37 or more weeks gestation and <2500 g.  
\* Percentages are not calculated if <100 records are available for analysis after exclusions.

