

Table 3B
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Racial and Ethnic Distribution by County

Comparison of Racial and Ethnic Distribution (2)								
County	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Allen	91	*	*	*	*	*	*	*
Anderson	50	*	*	*	*	*	*	*
Atchison	155	88.4	7.1	1.3	1.9	0.0	1.3	0.0
Barber	46	*	*	*	*	*	*	*
Barton	257	65.0	1.6	31.1	0.4	0.8	1.2	0.0
Bourbon	117	92.3	1.7	3.4	0.9	0.0	1.7	0.0
Brown	99	*	*	*	*	*	*	*
Butler	282	88.3	1.1	7.1	0.7	0.7	2.1	0.0
Chase	16	*	*	*	*	*	*	*
Chautauqua	32	*	*	*	*	*	*	*
Cherokee	212	88.7	0.9	3.8	1.9	3.8	0.9	0.0
Cheyenne	3	*	*	*	*	*	*	*
Clark	4	*	*	*	*	*	*	*
Clay	59	*	*	*	*	*	*	*
Cloud	89	*	*	*	*	*	*	*
Coffey	25	*	*	*	*	*	*	*
Comanche	11	*	*	*	*	*	*	*
Cowley	363	73.0	2.8	20.9	0.8	1.4	1.1	0.0
Crawford	401	84.3	2.7	10.0	0.0	1.5	1.5	0.0
Decatur	10	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3B
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Racial and Ethnic Distribution by County

Comparison of Racial and Ethnic Distribution (2)								
County	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Dickinson	134	91.8	0.0	5.2	1.5	0.7	0.7	0.0
Doniphan	37	*	*	*	*	*	*	*
Douglas	507	63.1	10.5	11.6	8.1	5.3	1.4	0.0
Edwards	25	*	*	*	*	*	*	*
Elk	20	*	*	*	*	*	*	*
Ellis	180	84.4	0.0	13.9	1.1	0.6	0.0	0.0
Ellsworth	33	*	*	*	*	*	*	*
Finney	718	30.8	0.6	65.3	0.1	2.9	0.3	0.0
Ford	686	21.6	1.0	76.7	0.1	0.6	0.0	0.0
Franklin	187	90.4	0.5	7.0	0.5	0.0	1.6	0.0
Geary	393	49.9	24.9	15.5	0.8	3.6	5.3	0.0
Gove	12	*	*	*	*	*	*	*
Graham	17	*	*	*	*	*	*	*
Grant	71	*	*	*	*	*	*	*
Gray	45	*	*	*	*	*	*	*
Greeley	15	*	*	*	*	*	*	*
Greenwood	49	*	*	*	*	*	*	*
Hamilton	31	*	*	*	*	*	*	*
Harper	55	*	*	*	*	*	*	*
Harvey	284	71.5	2.5	22.2	1.8	0.7	1.4	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3B
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Racial and Ethnic Distribution by County

Comparison of Racial and Ethnic Distribution (2)								
County	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Haskell	20	*	*	*	*	*	*	*
Hodgeman	16	*	*	*	*	*	*	*
Jackson	75	*	*	*	*	*	*	*
Jefferson	68	*	*	*	*	*	*	*
Jewell	9	*	*	*	*	*	*	*
Johnson	1,916	45.5	13.3	35.8	0.6	3.4	1.4	0.0
Kearny	19	*	*	*	*	*	*	*
Kingman	59	*	*	*	*	*	*	*
Kiowa	16	*	*	*	*	*	*	*
Labette	194	86.6	7.7	1.5	2.1	0.5	1.5	0.0
Lane	13	*	*	*	*	*	*	*
Leavenworth	359	71.6	13.1	10.6	0.3	0.8	3.6	0.0
Lincoln	21	*	*	*	*	*	*	*
Linn	66	*	*	*	*	*	*	*
Logan	15	*	*	*	*	*	*	*
Lyon	323	54.8	4.3	36.5	0.6	2.5	1.2	0.0
Marion	144	89.6	1.4	6.9	0.7	0.0	1.4	0.0
Marshall	47	*	*	*	*	*	*	*
Mcpherson	48	*	*	*	*	*	*	*
Meade	44	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3B
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Racial and Ethnic Distribution by County

Comparison of Racial and Ethnic Distribution (2)								
County	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Miami	146	92.5	3.4	2.1	0.0	0.7	1.4	0.0
Mitchell	54	*	*	*	*	*	*	*
Montgomery	344	76.7	7.8	10.5	3.8	0.3	0.9	0.0
Morris	23	*	*	*	*	*	*	*
Nemaha	43	*	*	*	*	*	*	*
Neosho	146	91.1	3.4	4.1	0.0	0.0	1.4	0.0
Ness	5	*	*	*	*	*	*	*
Norton	32	*	*	*	*	*	*	*
Osage	95	*	*	*	*	*	*	*
Osborne	14	*	*	*	*	*	*	*
Ottawa	28	*	*	*	*	*	*	*
Pawnee	62	*	*	*	*	*	*	*
Phillips	36	*	*	*	*	*	*	*
Pottawatom	38	*	*	*	*	*	*	*
Pratt	85	*	*	*	*	*	*	*
Rawlins	12	*	*	*	*	*	*	*
Reno	499	80.4	2.6	15.4	0.2	0.4	1.0	0.0
Republic	43	*	*	*	*	*	*	*
Rice	75	*	*	*	*	*	*	*
Riley	953	74.0	10.7	11.4	0.3	3.0	0.5	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3B
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Racial and Ethnic Distribution by County

Comparison of Racial and Ethnic Distribution (2)								
County	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaska Native	Asian/Pacific Islander	Multiple Races	All Other/Unknown
		%	%	%	%	%	%	%
Rooks	50	*	*	*	*	*	*	*
Rush	9	*	*	*	*	*	*	*
Russell	42	*	*	*	*	*	*	*
Saline	531	68.4	4.9	21.3	0.4	2.6	2.4	0.0
Scott	51	*	*	*	*	*	*	*
Sedgwick	4,721	44.6	18.0	29.8	1.8	4.0	1.8	0.0
Seward	532	15.0	1.7	82.1	0.6	0.4	0.2	0.0
Shawnee	1,510	56.0	15.7	24.2	2.1	0.7	1.3	0.0
Sheridan	10	*	*	*	*	*	*	*
Sherman	60	*	*	*	*	*	*	*
Smith	20	*	*	*	*	*	*	*
Stafford	43	*	*	*	*	*	*	*
Stanton	14	*	*	*	*	*	*	*
Stevens	89	*	*	*	*	*	*	*
Sumner	157	88.5	1.3	4.5	1.9	0.0	3.8	0.0
Thomas	60	*	*	*	*	*	*	*
Trego	18	*	*	*	*	*	*	*
Wabaunsee	25	*	*	*	*	*	*	*
Washington	27	*	*	*	*	*	*	*
Wichita	18	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3B
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Racial and Ethnic Distribution by County

Comparison of Racial and Ethnic Distribution (2)								
County	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaska Native	Asian/Pacific Islander	Multiple Races	All Other/Unknown
		%	%	%	%	%	%	%
Wilson	101	95.0	1.0	3.0	1.0	0.0	0.0	0.0
Woodson	8	*	*	*	*	*	*	*
Wyandotte	2,286	26.2	29.9	40.3	0.3	2.8	0.4	0.0
Unknown Counties	3	*	*	*	*	*	*	*
Kansas	22,411	56.4	11.3	27.6	1.3	2.2	1.3	0.0
Nation (Prior Year)	1,307,887	42.0	24.2	28.1	1.0	3.1	1.1	0.5

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4B
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Age Distribution by County

Comparison of Age Distribution (2)								
County	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Allen	91	*	*	*	*	*	*	*
Anderson	50	*	*	*	*	*	*	*
Atchison	155	0.0	7.1	16.8	62.6	13.5	0.0	0.0
Barber	46	*	*	*	*	*	*	*
Barton	257	0.0	8.6	14.4	61.9	14.0	1.2	0.0
Bourbon	117	0.0	7.7	18.8	59.0	13.7	0.9	0.0
Brown	99	*	*	*	*	*	*	*
Butler	282	0.0	7.8	19.9	59.9	11.7	0.4	0.4
Chase	16	*	*	*	*	*	*	*
Chautauqua	32	*	*	*	*	*	*	*
Cherokee	212	0.0	8.0	19.8	59.9	10.8	0.9	0.5
Cheyenne	3	*	*	*	*	*	*	*
Clark	4	*	*	*	*	*	*	*
Clay	59	*	*	*	*	*	*	*
Cloud	89	*	*	*	*	*	*	*
Coffey	25	*	*	*	*	*	*	*
Comanche	11	*	*	*	*	*	*	*
Cowley	363	0.8	8.0	16.5	61.2	12.7	0.8	0.0
Crawford	401	0.2	7.0	16.5	62.8	12.7	0.5	0.2

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4B
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Age Distribution by County

Comparison of Age Distribution (2)								
County	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Decatur	10	*	*	*	*	*	*	*
Dickinson	134	0.0	9.7	13.4	61.2	14.9	0.7	0.0
Doniphan	37	*	*	*	*	*	*	*
Douglas	507	0.0	3.9	11.4	63.1	20.1	1.4	0.0
Edwards	25	*	*	*	*	*	*	*
Elk	20	*	*	*	*	*	*	*
Ellis	180	0.0	3.3	10.6	73.9	10.6	1.7	0.0
Ellsworth	33	*	*	*	*	*	*	*
Finney	718	0.6	9.1	15.7	54.0	18.8	1.8	0.0
Ford	686	0.4	7.9	14.0	53.9	21.7	1.9	0.1
Franklin	187	0.0	5.3	21.9	56.7	14.4	1.6	0.0
Geary	393	0.3	4.8	14.5	67.9	12.2	0.3	0.0
Gove	12	*	*	*	*	*	*	*
Graham	17	*	*	*	*	*	*	*
Grant	71	*	*	*	*	*	*	*
Gray	45	*	*	*	*	*	*	*
Greeley	15	*	*	*	*	*	*	*
Greenwood	49	*	*	*	*	*	*	*
Hamilton	31	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4B
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Age Distribution by County

Comparison of Age Distribution (2)								
County	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Harper	55	*	*	*	*	*	*	*
Harvey	284	0.7	7.4	18.3	59.9	13.4	0.4	0.0
Haskell	20	*	*	*	*	*	*	*
Hodgeman	16	*	*	*	*	*	*	*
Jackson	75	*	*	*	*	*	*	*
Jefferson	68	*	*	*	*	*	*	*
Jewell	9	*	*	*	*	*	*	*
Johnson	1,916	0.2	3.8	10.9	59.5	24.0	1.5	0.1
Kearny	19	*	*	*	*	*	*	*
Kingman	59	*	*	*	*	*	*	*
Kiowa	16	*	*	*	*	*	*	*
Labette	194	0.0	10.3	19.1	59.3	10.3	1.0	0.0
Lane	13	*	*	*	*	*	*	*
Leavenworth	359	0.0	6.7	17.3	62.7	12.8	0.6	0.0
Lincoln	21	*	*	*	*	*	*	*
Linn	66	*	*	*	*	*	*	*
Logan	15	*	*	*	*	*	*	*
Lyon	323	0.6	3.7	13.3	66.3	14.6	1.5	0.0
Marion	144	0.0	4.9	9.7	68.8	16.7	0.0	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4B
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Age Distribution by County

Comparison of Age Distribution (2)								
County	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Marshall	47	*	*	*	*	*	*	*
Mcpherson	48	*	*	*	*	*	*	*
Meade	44	*	*	*	*	*	*	*
Miami	146	0.7	4.8	17.1	63.0	13.7	0.7	0.0
Mitchell	54	*	*	*	*	*	*	*
Montgomery	344	0.3	2.0	16.6	68.0	12.8	0.0	0.3
Morris	23	*	*	*	*	*	*	*
Nemaha	43	*	*	*	*	*	*	*
Neosho	146	0.0	9.6	19.2	59.6	11.0	0.7	0.0
Ness	5	*	*	*	*	*	*	*
Norton	32	*	*	*	*	*	*	*
Osage	95	*	*	*	*	*	*	*
Osborne	14	*	*	*	*	*	*	*
Ottawa	28	*	*	*	*	*	*	*
Pawnee	62	*	*	*	*	*	*	*
Phillips	36	*	*	*	*	*	*	*
Pottawatom	38	*	*	*	*	*	*	*
Pratt	85	*	*	*	*	*	*	*
Rawlins	12	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4B
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Age Distribution by County

Comparison of Age Distribution (2)								
County	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Reno	499	0.0	7.0	17.0	62.5	13.0	0.4	0.0
Republic	43	*	*	*	*	*	*	*
Rice	75	*	*	*	*	*	*	*
Riley	953	0.0	1.9	12.4	74.4	11.0	0.3	0.0
Rooks	50	*	*	*	*	*	*	*
Rush	9	*	*	*	*	*	*	*
Russell	42	*	*	*	*	*	*	*
Saline	531	0.2	7.2	14.5	64.6	12.4	1.1	0.0
Scott	51	*	*	*	*	*	*	*
Sedgwick	4,721	0.3	6.9	13.6	60.6	17.2	1.2	0.0
Seward	532	0.0	7.7	14.3	54.7	21.8	1.5	0.0
Shawnee	1,510	0.3	6.8	15.9	60.3	15.5	1.3	0.0
Sheridan	10	*	*	*	*	*	*	*
Sherman	60	*	*	*	*	*	*	*
Smith	20	*	*	*	*	*	*	*
Stafford	43	*	*	*	*	*	*	*
Stanton	14	*	*	*	*	*	*	*
Stevens	89	*	*	*	*	*	*	*
Sumner	157	0.0	5.7	19.7	60.5	12.7	1.3	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4B
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Age Distribution by County

Comparison of Age Distribution (2)								
County	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Thomas	60	*	*	*	*	*	*	*
Trego	18	*	*	*	*	*	*	*
Wabaunsee	25	*	*	*	*	*	*	*
Washington	27	*	*	*	*	*	*	*
Wichita	18	*	*	*	*	*	*	*
Wilson	101	0.0	6.9	19.8	61.4	11.9	0.0	0.0
Woodson	8	*	*	*	*	*	*	*
Wyandotte	2,286	0.4	7.6	13.7	57.9	18.9	1.4	0.0
Unknown Counties	3	*	*	*	*	*	*	*
Kansas	22,411	0.3	6.3	14.6	61.1	16.6	1.1	0.1
Nation (Prior Year)	1,307,887	0.3	6.3	12.6	59.7	19.4	1.4	0.3

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5B
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Maternal Health Indicators by County

Comparison of Maternal Health Indicators (2, 3)														
County	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Allen	89	* *	* *	80	* *	* *	19	* *	24	* *	82	* *	82	* *
Anderson	49	* *	* *	43	* *	* *	8	* *	6	* *	45	* *	45	* *
Atchison	147	6.8 (25)	59.9 (31)	133	15.0 (3)	56.4 (27)	23	* *	29	* *	134	7.5 (24)	133	15.8(29)
Barber	44	* *	* *	33	* *	* *	7	* *	10	* *	37	* *	37	* *
Barton	252	4.8 (17)	55.6 (20)	213	15.5 (4)	52.6 (21)	28	* *	40	* *	217	3.7 (4)	215	13.5(24)
Bourbon	114	3.5 (6)	58.8 (29)	99	* *	* *	17	* *	21	* *	101	5.9 (14)	101	11.9(20)
Brown	97	* *	* *	87	* *	* *	8	* *	26	* *	87	* *	87	* *
Butler	263	6.8 (25)	51.7 (7)	234	17.9 (9)	49.1 (11)	41	* *	55	* *	255	8.6 (27)	255	12.9(22)
Chase	15	* *	* *	13	* *	* *	1	* *	0	* *	14	* *	14	* *
Chautauqua	32	* *	* *	32	* *	* *	4	* *	4	* *	32	* *	32	* *
Cherokee	192	7.8 (30)	54.7 (17)	169	13.0 (1)	58.0 (30)	24	* *	45	* *	184	4.9 (7)	185	13.5(24)
Cheyenne	3	* *	* *	3	* *	* *	0	* *	1	* *	3	* *	3	* *
Clark	4	* *	* *	3	* *	* *	1	* *	0	* *	3	* *	3	* *
Clay	59	* *	* *	51	* *	* *	4	* *	16	* *	53	* *	53	* *
Cloud	89	* *	* *	80	* *	* *	13	* *	27	* *	78	* *	81	* *
Coffey	23	* *	* *	19	* *	* *	1	* *	9	* *	20	* *	20	* *
Comanche	11	* *	* *	11	* *	* *	4	* *	2	* *	11	* *	11	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5B
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Maternal Health Indicators by County

Comparison of Maternal Health Indicators (2, 3)														
County	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Cowley	356	4.8 (17)	52.8 (10)	313	21.4 (23)	53.4 (23)	52	* * *	45	* * *	316	5.7 (12)	317	12.0(21)
Crawford	389	7.2 (28)	53.2 (12)	336	19.3 (13)	56.3 (26)	54	* * *	54	* * *	348	7.2 (21)	348	13.8(26)
Decatur	10	* * *	* * *	9	* * *	* * *	1	* * *	1	* * *	9	* * *	9	* * *
Dickinson	128	4.7 (15)	57.0 (25)	114	14.9 (2)	57.0 (29)	13	* * *	33	* * *	121	6.6 (18)	117	11.1(16)
Doniphan	35	* * *	* * *	31	* * *	* * *	5	* * *	9	* * *	33	* * *	33	* * *
Douglas	491	3.7 (7)	48.1 (2)	421	16.9 (7)	55.1 (24)	70	* * *	137	21.2 (5)	432	6.7 (19)	432	9.0(7)
Edwards	25	* * *	* * *	25	* * *	* * *	3	* * *	4	* * *	25	* * *	25	* * *
Elk	19	* * *	* * *	18	* * *	* * *	1	* * *	4	* * *	19	* * *	19	* * *
Ellis	166	3.0 (5)	51.8 (8)	140	19.3 (13)	47.9 (9)	42	* * *	43	* * *	157	1.9 (1)	157	5.7(1)
Ellsworth	33	* * *	* * *	32	* * *	* * *	4	* * *	5	* * *	31	* * *	31	* * *
Finney	701	7.6 (29)	50.6 (5)	630	23.3 (26)	46.5 (8)	99	* * *	104	15.4 (2)	645	7.4 (23)	643	7.2(4)
Ford	677	2.5 (4)	55.5 (19)	604	22.8 (25)	45.5 (5)	80	* * *	61	* * *	601	6.3 (17)	600	9.8(9)
Franklin	180	2.2 (3)	55.6 (20)	160	18.1 (10)	56.9 (28)	19	* * *	56	* * *	128	4.7 (6)	128	10.9(15)
Geary	378	5.3 (21)	51.1 (6)	301	15.9 (5)	52.5 (20)	55	* * *	83	* * *	314	3.5 (3)	314	10.5(13)
Gove	9	* * *	* * *	8	* * *	* * *	0	* * *	4	* * *	11	* * *	11	* * *
Graham	17	* * *	* * *	14	* * *	* * *	2	* * *	2	* * *	14	* * *	14	* * *
Grant	69	* * *	* * *	63	* * *	* * *	7	* * *	22	* * *	66	* * *	64	* * *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5B
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Maternal Health Indicators by County

Comparison of Maternal Health Indicators (2, 3)														
County	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Gray	45	* *	* *	40	* *	* *	2	* *	7	* *	40	* *	40	* *
Greeley	13	* *	* *	12	* *	* *	1	* *	2	* *	14	* *	14	* *
Greenwood	47	* *	* *	43	* *	* *	5	* *	5	* *	44	* *	45	* *
Hamilton	28	* *	* *	25	* *	* *	3	* *	10	* *	27	* *	27	* *
Harper	53	* *	* *	44	* *	* *	7	* *	9	* *	43	* *	43	* *
Harvey	277	4.7 (15)	53.1 (11)	241	26.6 (30)	45.2 (4)	40	* *	37	* *	246	6.1 (15)	246	11.8(19)
Haskell	20	* *	* *	17	* *	* *	3	* *	7	* *	17	* *	17	* *
Hodgeman	15	* *	* *	15	* *	* *	3	* *	4	* *	16	* *	16	* *
Jackson	70	* *	* *	62	* *	* *	15	* *	19	* *	64	* *	64	* *
Jefferson	67	* *	* *	61	* *	* *	11	* *	20	* *	62	* *	62	* *
Jewell	9	* *	* *	9	* *	* *	1	* *	1	* *	9	* *	9	* *
Johnson	1,812	4.5 (13)	54.8 (18)	1,609	19.4 (15)	51.0 (14)	307	24.1 (2)	592	17.9 (4)	1,686	8.3 (26)	1,683	7.3(5)
Kearny	19	* *	* *	17	* *	* *	2	* *	4	* *	17	* *	17	* *
Kingman	59	* *	* *	51	* *	* *	7	* *	6	* *	51	* *	51	* *
Kiowa	15	* *	* *	12	* *	* *	1	* *	4	* *	13	* *	13	* *
Labette	190	7.9 (31)	52.6 (9)	178	25.8 (29)	42.7 (2)	31	* *	43	* *	179	5.6 (10)	179	11.2(17)
Lane	13	* *	* *	12	* *	* *	0	* *	2	* *	12	* *	12	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5B
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Maternal Health Indicators by County

Comparison of Maternal Health Indicators (2, 3)														
County	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Leavenworth	347	4.6 (14)	57.3 (26)	297	19.2 (12)	51.5 (16)	61	* *	75	* *	303	8.9 (29)	302	9.9(10)
Lincoln	19	* *	* *	17	* *	* *	4	* *	7	* *	19	* *	19	* *
Linn	66	* *	* *	62	* *	* *	10	* *	10	* *	61	* *	61	* *
Logan	15	* *	* *	13	* *	* *	2	* *	2	* *	13	* *	13	* *
Lyon	306	2.0 (2)	63.1 (32)	267	23.6 (27)	43.8 (3)	45	* *	51	* *	282	7.1 (20)	282	14.2(27)
Marion	142	5.6 (22)	50.0 (4)	126	19.8 (19)	46.0 (6)	18	* *	35	* *	128	5.5 (9)	129	10.1(11)
Marshall	44	* *	* *	38	* *	* *	5	* *	14	* *	42	* *	42	* *
Mcpherson	47	* *	* *	41	* *	* *	5	* *	10	* *	42	* *	42	* *
Meade	44	* *	* *	37	* *	* *	4	* *	6	* *	37	* *	37	* *
Miami	137	6.6 (24)	57.7 (27)	120	18.3 (11)	49.2 (12)	22	* *	44	* *	131	9.2 (31)	131	13.0(23)
Mitchell	52	* *	* *	49	* *	* *	4	* *	12	* *	50	* *	50	* *
Montgomery	337	6.2 (23)	59.1 (30)	294	21.1 (22)	48.0 (10)	61	* *	87	* *	298	5.7 (12)	298	10.1(11)
Morris	22	* *	* *	20	* *	* *	3	* *	8	* *	21	* *	21	* *
Nemaha	37	* *	* *	34	* *	* *	8	* *	10	* *	40	* *	40	* *
Neosho	140	8.6 (32)	49.3 (3)	128	21.9 (24)	52.3 (19)	22	* *	19	* *	133	9.0 (30)	133	14.3(28)
Ness	4	* *	* *	4	* *	* *	0	* *	3	* *	5	* *	5	* *
Norton	32	* *	* *	29	* *	* *	4	* *	7	* *	29	* *	29	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5B
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Maternal Health Indicators by County

Comparison of Maternal Health Indicators (2, 3)															
County	Prepregnancy BMI (4)				Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight		Over-weight		< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)		Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Osage	94	* *	* *		82	* *	* *		9	* *		18	* *	86	* *
Osborne	14	* *	* *		10	* *	* *		2	* *		5	* *	10	* *
Ottawa	26	* *	* *		23	* *	* *		3	* *		4	* *	25	* *
Pawnee	60	* *	* *		46	* *	* *		5	* *		12	* *	52	* *
Phillips	34	* *	* *		30	* *	* *		2	* *		9	* *	32	* *
Pottawatom	37	* *	* *		34	* *	* *		1	* *		18	* *	35	* *
Pratt	81	* *	* *		67	* *	* *		10	* *		17	* *	71	* *
Rawlins	12	* *	* *		11	* *	* *		1	* *		3	* *	11	* *
Reno	489	4.9 (19)	55.8 (23)		426	19.7 (18)	50.9 (13)		54	* *		64	* *	438	2.5 (2)
Republic	41	* *	* *		32	* *	* *		5	* *		11	* *	34	* *
Rice	72	* *	* *		63	* *	* *		7	* *		18	* *	68	* *
Riley	898	5.0 (20)	48.0 (1)		716	16.2 (6)	55.9 (25)		160	20.6 (1)		208	17.8 (3)	765	4.1 (5)
Rooks	50	* *	* *		45	* *	* *		5	* *		7	* *	45	* *
Rush	9	* *	* *		9	* *	* *		1	* *		3	* *	9	* *
Russell	42	* *	* *		37	* *	* *		6	* *		9	* *	37	* *
Saline	516	4.1 (8)	53.9 (14)		473	16.9 (7)	53.1 (22)		89	* *		105	14.3 (1)	482	5.6 (10)
Scott	50	* *	* *		45	* *	* *		8	* *		6	* *	45	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5B
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Maternal Health Indicators by County

Comparison of Maternal Health Indicators (2, 3)														
County	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Sedgwick	4,574	4.4 (11)	53.5 (13)	4,150	20.1 (20)	51.6 (17)	914	31.4 (3)	988	25.3 (7)	4,278	6.1 (15)	4,281	11.2(17)
Seward	518	1.9 (1)	58.7 (28)	468	24.1 (28)	46.2 (7)	93	* *	64	* *	478	8.8 (28)	482	6.4(3)
Shawnee	1,461	4.4 (11)	53.9 (14)	1,273	19.6 (17)	52.1 (18)	152	43.4 (5)	236	30.9 (8)	1,303	7.3 (22)	1,305	7.7(6)
Sheridan	10	* *	* *	10	* *	* *	2	* *	2	* *	10	* *	10	* *
Sherman	59	* *	* *	51	* *	* *	5	* *	5	* *	52	* *	52	* *
Smith	20	* *	* *	17	* *	* *	4	* *	4	* *	16	* *	16	* *
Stafford	43	* *	* *	38	* *	* *	4	* *	5	* *	38	* *	38	* *
Stanton	14	* *	* *	14	* *	* *	1	* *	7	* *	14	* *	14	* *
Stevens	86	* *	* *	76	* *	* *	14	* *	26	* *	79	* *	79	* *
Sumner	147	4.1 (8)	54.4 (16)	134	20.1 (20)	41.0 (1)	20	* *	37	* *	144	7.6 (25)	144	16.7(31)
Thomas	57	* *	* *	52	* *	* *	11	* *	11	* *	54	* *	54	* *
Trego	16	* *	* *	14	* *	* *	3	* *	3	* *	16	* *	16	* *
Wabaunsee	24	* *	* *	24	* *	* *	3	* *	10	* *	24	* *	24	* *
Washington	25	* *	* *	21	* *	* *	1	* *	8	* *	23	* *	23	* *
Wichita	18	* *	* *	15	* *	* *	3	* *	2	* *	15	* *	15	* *
Wilson	101	6.9 (27)	56.4 (24)	93	* *	* *	17	* *	14	* *	93	* *	93	* *
Woodson	8	* *	* *	8	* *	* *	1	* *	4	* *	8	* *	8	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5B
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Maternal Health Indicators by County

Comparison of Maternal Health Indicators (2, 3)														
County	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Wyandotte	2,228	4.2 (10)	55.7 (22)	1,973	19.5 (16)	51.2 (15)	351	31.9 (4)	393	22.4 (6)	2,004	5.2 (8)	2,005	9.5 (8)
Unknown Counties	3	* *	* *	3	* *	* *	0	* *	1	* *	3	* *	3	* *
Kansas	21,646	4.6	54.3	19,129	19.9	51.1	3,394	29.5	4,491	22.7	19,697	6.4	19,699	10.3
Nation (Prior Year)	1,224,766	4.7	52.1	963,624	21.4	48.2	188,530	33.8	416,710	29.6	346,704	4.4	395,960	5.2

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6B
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Maternal Behavioral Indicators by County

Comparison of Maternal Behavioral Indicators (2, 3)										
County	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	% (Rank)		Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Allen	77	* *	* *		91	* *	91	* *	83	* *
Anderson	36	* *	* *		50	* *	50	* *	45	* *
Atchison	136	89.0 (1)	2.2 (10)		155	31.0 (27)	155	9.7 (27)	127	66.9 (6)
Barber	37	* *	* *		45	* *	46	* *	40	* *
Barton	214	85.0 (9)	1.4 (6)		257	51.0 (1)	257	19.8 (5)	215	49.8 (27)
Bourbon	97	* *	* *		117	47.0 (3)	117	12.8 (24)	104	59.6 (17)
Brown	86	* *	* *		98	* *	99	* *	85	* *
Butler	241	64.7 (27)	13.3 (26)		282	38.3 (14)	282	16.7 (10)	207	31.9 (31)
Chase	14	* *	* *		16	* *	16	* *	12	* *
Chautauqua	32	* *	* *		31	* *	32	* *	30	* *
Cherokee	143	79.0 (15)	5.6 (20)		210	28.6 (31)	212	15.1 (16)	151	55.6 (21)
Cheyenne	1	* *	* *		3	* *	3	* *	2	* *
Clark	3	* *	* *		4	* *	4	* *	3	* *
Clay	53	* *	* *		59	* *	59	* *	56	* *
Cloud	81	* *	* *		88	* *	89	* *	77	* *
Coffey	20	* *	* *		25	* *	25	* *	17	* *
Comanche	11	* *	* *		11	* *	11	* *	9	* *
Cowley	283	66.1 (25)	6.0 (22)		358	42.2 (9)	363	12.9 (23)	307	58.0 (19)
Crawford	329	83.3 (13)	3.6 (13)		401	42.1 (10)	401	13.2 (21)	350	59.7 (16)
Decatur	4	* *	* *		10	* *	10	* *	7	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6B
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Maternal Behavioral Indicators by County

Comparison of Maternal Behavioral Indicators (2, 3)											
County	Medical Care				WIC Enrollment			Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester			Prepregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	% (Rank)		Number	% (Rank)		Number	% (Rank)	Number	% (Rank)
Dickinson	122	84.4 (10)	0.8 (4)		133	46.6 (5)		134	19.4 (7)	102	74.5 (2)
Doniphan	33	* *	* *		37	* *		37	* *	25	* *
Douglas	428	75.7 (18)	4.7 (16)		507	33.1 (24)		507	20.5 (4)	397	63.0 (12)
Edwards	23	* *	* *		24	* *		25	* *	21	* *
Elk	18	* *	* *		20	* *		20	* *	16	* *
Ellis	154	84.4 (10)	2.6 (11)		179	30.2 (30)		180	26.7 (1)	139	74.8 (1)
Ellsworth	32	* *	* *		33	* *		33	* *	27	* *
Finney	581	58.9 (29)	25.6 (29)		712	34.0 (21)		718	13.0 (22)	593	35.1 (30)
Ford	584	70.0 (22)	16.4 (28)		682	43.0 (7)		686	7.4 (29)	611	52.7 (25)
Franklin	32	* *	* *		184	42.9 (8)		187	4.3 (32)	152	53.3 (24)
Geary	285	67.0 (23)	1.4 (6)		392	33.2 (23)		393	18.6 (8)	321	60.7 (13)
Gove	10	* *	* *		12	* *		12	* *	8	* *
Graham	14	* *	* *		17	* *		17	* *	14	* *
Grant	60	* *	* *		70	* *		71	* *	56	* *
Gray	36	* *	* *		44	* *		45	* *	39	* *
Greeley	14	* *	* *		15	* *		15	* *	10	* *
Greenwood	42	* *	* *		49	* *		49	* *	40	* *
Hamilton	27	* *	* *		31	* *		31	* *	24	* *
Harper	44	* *	* *		53	* *		55	* *	45	* *
Harvey	225	74.7 (20)	14.7 (27)		281	47.0 (3)		284	16.2 (12)	238	68.9 (4)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6B
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Maternal Behavioral Indicators by County

Comparison of Maternal Behavioral Indicators (2, 3)										
County	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	% (Rank)		Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Haskell	15	* *	* *		20	* *	20	* *	17	* *
Hodgeman	16	* *	* *		16	* *	16	* *	13	* *
Jackson	38	* *	* *		74	* *	75	* *	62	* *
Jefferson	60	* *	* *		68	* *	68	* *	54	* *
Jewell	9	* *	* *		9	* *	9	* *	9	* *
Johnson	1,518	66.5 (24)	3.8 (14)		1,905	30.6 (29)	1,916	14.0 (20)	1,527	64.2 (9)
Kearny	15	* *	* *		19	* *	19	* *	15	* *
Kingman	49	* *	* *		58	* *	59	* *	53	* *
Kiowa	13	* *	* *		16	* *	16	* *	12	* *
Labette	151	86.8 (5)	5.3 (19)		193	39.4 (12)	194	16.0 (13)	166	55.4 (22)
Lane	12	* *	* *		13	* *	13	* *	10	* *
Leavenworth	304	88.5 (3)	0.0 (1)		357	34.7 (20)	359	22.6 (3)	275	72.4 (3)
Lincoln	19	* *	* *		21	* *	21	* *	16	* *
Linn	62	* *	* *		66	* *	66	* *	57	* *
Logan	2	* *	* *		15	* *	15	* *	12	* *
Lyon	279	86.7 (6)	0.7 (3)		322	38.8 (13)	323	15.5 (15)	265	63.4 (11)
Marion	126	88.1 (4)	0.8 (4)		144	38.2 (15)	144	22.9 (2)	121	65.3 (7)
Marshall	40	* *	* *		47	* *	47	* *	31	* *
Mcperson	42	* *	* *		48	* *	48	* *	40	* *
Meade	37	* *	* *		43	* *	44	* *	36	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6B
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Maternal Behavioral Indicators by County

Comparison of Maternal Behavioral Indicators (2, 3)										
County	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	% (Rank)		Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Miami	129	85.3 (8)	1.6 (8)		146	30.8 (28)	146	14.4 (17)	107	54.2 (23)
Mitchell	48	* *	* *		53	* *	54	* *	44	* *
Montgomery	241	82.6 (14)	5.0 (18)		342	33.3 (22)	344	14.2 (19)	292	64.4 (8)
Morris	18	* *	* *		23	* *	23	* *	19	* *
Nemaha	40	* *	* *		43	* *	43	* *	39	* *
Neosho	124	88.7 (2)	6.5 (23)		146	37.7 (16)	146	8.9 (28)	127	68.5 (5)
Ness	5	* *	* *		5	* *	5	* *	3	* *
Norton	29	* *	* *		32	* *	32	* *	29	* *
Osage	73	* *	* *		90	* *	95	* *	89	* *
Osborne	10	* *	* *		14	* *	14	* *	12	* *
Ottawa	25	* *	* *		28	* *	28	* *	24	* *
Pawnee	50	* *	* *		61	* *	62	* *	50	* *
Phillips	32	* *	* *		36	* *	36	* *	32	* *
Pottawatom	34	* *	* *		38	* *	38	* *	30	* *
Pratt	70	* *	* *		84	* *	85	* *	63	* *
Rawlins	0	* *	* *		12	* *	12	* *	11	* *
Reno	424	84.0 (12)	2.1 (9)		495	40.0 (11)	499	19.8 (5)	414	59.4 (18)
Republic	34	* *	* *		43	* *	43	* *	35	* *
Rice	67	* *	* *		73	* *	75	* *	61	* *
Riley	712	85.5 (7)	0.6 (2)		952	32.8 (26)	953	16.7 (10)	763	60.0 (15)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6B
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Maternal Behavioral Indicators by County

Comparison of Maternal Behavioral Indicators (2, 3)										
County	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Rooks	44	* *	* *	* *	50	* *	50	* *	40	* *
Rush	9	* *	* *	* *	8	* *	9	* *	7	* *
Russell	37	* *	* *	* *	42	* *	42	* *	34	* *
Saline	419	74.0 (21)	5.7 (21)		529	32.9 (25)	531	4.9 (31)	431	51.5 (26)
Scott	44	* *	* *	* *	49	* *	51	* *	45	* *
Sedgwick	4,225	75.4 (19)	4.1 (15)		4,705	28.0 (32)	4,721	14.4 (17)	3,847	47.1 (29)
Seward	420	64.0 (28)	10.2 (25)		528	34.8 (19)	532	10.2 (26)	438	63.5 (10)
Shawnee	1,167	78.5 (17)	3.2 (12)		1,502	47.2 (2)	1,510	7.4 (29)	1,264	49.5 (28)
Sheridan	10	* *	* *	* *	10	* *	10	* *	8	* *
Sherman	16	* *	* *	* *	60	* *	60	* *	52	* *
Smith	15	* *	* *	* *	19	* *	20	* *	18	* *
Stafford	38	* *	* *	* *	43	* *	43	* *	38	* *
Stanton	14	* *	* *	* *	14	* *	14	* *	7	* *
Stevens	79	* *	* *	* *	88	* *	89	* *	72	* *
Sumner	143	79.0 (15)	4.9 (17)		157	35.7 (18)	157	15.9 (14)	121	60.3 (14)
Thomas	20	* *	* *	* *	60	* *	60	* *	49	* *
Trego	15	* *	* *	* *	18	* *	18	* *	15	* *
Wabaunsee	21	* *	* *	* *	25	* *	25	* *	20	* *
Washington	23	* *	* *	* *	27	* *	27	* *	20	* *
Wichita	15	* *	* *	* *	18	* *	18	* *	15	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6B
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Maternal Behavioral Indicators by County

Comparison of Maternal Behavioral Indicators (2, 3)														
County	Medical Care				WIC Enrollment			Multivitamin Consumption						
	1st Trimester		No Care		1st Trimester			Prepregnancy (4)		During Pregnancy (5)				
	Number	%	%	%	Number	%	%	Number	%	Number	%			
	(Rank)	(Rank)	(Rank)	(Rank)	(Rank)	(Rank)	(Rank)	(Rank)	(Rank)	(Rank)	(Rank)			
Wilson	92	* *	* *	* *	101	46.5	(6)	101	16.8	(9)	88	* *		
Woodson	8	* *	* *	* *	8	* *	*	8	* *	*	6	* *		
Wyandotte	1,981	65.6	(26)	8.0	(24)	2,267	35.9	(17)	2,286	10.4	(25)	1,910	56.9	(20)
Unknown Counties	3	* *	* *	* *	3	* *	*	3	* *	*	1	* *		
Kansas	18,492	75.4	5.7	5.7	22,287	35.7	35.7	22,411	14.6	14.6	18,386	55.7		
Nation (Prior Year)	580,485	80.1	3.1	3.1	1,187,387	31.7	31.7	266,567	19.3	19.3	269,548	80.6		

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7B
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by County

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Allen	86	* *	68	* *	22	* *	80	* *	68	* *
Anderson	48	* *	37	* *	18	* *	44	* *	38	* *
Atchison	109	48.6 (29)	81	* *	27	* *	89	* *	84	* *
Barber	46	* *	37	* *	18	* *	40	* *	37	* *
Barton	255	36.9 (11)	215	23.3 (13)	68	* *	215	14.0 (12)	214	4.7 (1)
Bourbon	110	52.7 (30)	82	* *	34	* *	97	* *	82	* *
Brown	95	* *	70	* *	25	* *	82	* *	70	* *
Butler	275	44.0 (24)	234	26.5 (20)	77	* *	204	12.7 (10)	236	7.6 (8)
Chase	16	* *	14	* *	6	* *	12	* *	14	* *
Chautauqua	23	* *	13	* *	5	* *	22	* *	13	* *
Cherokee	204	37.7 (15)	182	24.7 (18)	53	* *	148	33.1 (25)	183	32.8 (25)
Cheyenne	3	* *	3	* *	0	* *	2	* *	3	* *
Clark	0	* *	1	* *	0	* *	2	* *	3	* *
Clay	59	* *	53	* *	25	* *	56	* *	53	* *
Cloud	89	* *	79	* *	29	* *	77	* *	81	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7B
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by County

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Coffey	22	* *	20	* *	8	* *	16	* *	20	* *
Comanche	11	* *	11	* *	5	* *	9	* *	11	* *
Cowley	347	41.8 (19)	249	28.5 (24)	86	* *	297	22.2 (20)	249	22.1 (21)
Crawford	364	41.2 (18)	282	23.8 (16)	93	* *	328	40.2 (28)	283	42.4 (27)
Decatur	10	* *	9	* *	2	* *	7	* *	9	* *
Dickinson	133	43.6 (23)	120	29.2 (25)	39	* *	101	10.9 (7)	120	5.8 (6)
Doniphan	34	* *	26	* *	7	* *	24	* *	27	* *
Douglas	497	44.3 (25)	422	21.6 (12)	150	41.3 (2)	392	10.2 (4)	425	5.6 (5)
Edwards	25	* *	24	* *	5	* *	20	* *	24	* *
Elk	19	* *	16	* *	5	* *	15	* *	15	* *
Ellis	179	36.9 (11)	151	13.2 (6)	38	* *	139	7.2 (1)	151	5.3 (3)
Ellsworth	33	* *	30	* *	7	* *	27	* *	30	* *
Finney	706	11.9 (3)	618	5.8 (3)	54	* *	588	17.2 (16)	619	16.0 (17)
Ford	673	11.1 (2)	568	5.5 (2)	44	* *	601	10.5 (6)	570	10.5 (12)
Franklin	186	47.8 (28)	165	30.9 (26)	61	* *	151	13.9 (11)	166	8.4 (10)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7B
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by County

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Geary	386	35.0 (9)	307	19.2 (10)	82	* *	320	37.5 (26)	307	29.6 (23)
Gove	11	* *	11	* *	1	* *	7	* *	11	* *
Graham	17	* *	13	* *	6	* *	14	* *	14	* *
Grant	71	* *	67	* *	1	* *	56	* *	67	* *
Gray	45	* *	37	* *	1	* *	38	* *	38	* *
Greeley	15	* *	14	* *	3	* *	10	* *	14	* *
Greenwood	38	* *	14	* *	6	* *	35	* *	31	* *
Hamilton	30	* *	27	* *	1	* *	24	* *	27	* *
Harper	47	* *	33	* *	13	* *	35	* *	28	* *
Harvey	275	36.0 (10)	234	19.7 (11)	70	* *	231	9.5 (3)	236	5.1 (2)
Haskell	20	* *	17	* *	3	* *	17	* *	17	* *
Hodgeman	16	* *	16	* *	7	* *	12	* *	15	* *
Jackson	75	* *	63	* *	20	* *	61	* *	63	* *
Jefferson	17	* *	1	* *	0	* *	19	* *	2	* *
Jewell	9	* *	9	* *	6	* *	9	* *	9	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7B
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by County

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Johnson	1,888	23.2 (5)	1,665	9.8 (4)	266	42.9 (1)	1,517	18.4 (19)	1,666	17.5 (19)
Kearny	19	* *	17	* *	3	* *	15	* *	17	* *
Kingman	56	* *	43	* *	17	* *	52	* *	45	* *
Kiowa	15	* *	10	* *	1	* *	11	* *	11	* *
Labette	189	37.0 (13)	169	26.6 (21)	52	* *	165	30.9 (22)	170	28.8 (22)
Lane	3	* *	1	* *	0	* *	3	* *	1	* *
Leavenworth	351	45.9 (27)	294	25.2 (19)	95	* *	274	40.1 (27)	295	41.0 (26)
Lincoln	21	* *	15	* *	5	* *	16	* *	15	* *
Linn	66	* *	60	* *	26	* *	57	* *	60	* *
Logan	15	* *	13	* *	3	* *	12	* *	13	* *
Lyon	319	27.9 (7)	279	15.4 (8)	59	* *	265	12.1 (8)	278	8.6 (11)
Marion	143	38.5 (16)	129	23.3 (13)	43	* *	120	9.2 (2)	129	5.4 (4)
Marshall	47	* *	41	* *	8	* *	31	* *	41	* *
Mcperson	45	* *	37	* *	8	* *	38	* *	38	* *
Meade	38	* *	32	* *	9	* *	33	* *	33	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7B
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by County

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)		Number	% (Rank)	Number	% (Rank)	
Miami	138	42.0 (20)	116	26.7 (23)	38	* *	102	16.7 (15)	116	14.7 (15)
Mitchell	53	* *	48	* *	14	* *	44	* *	49	* *
Montgomery	337	40.7 (17)	275	30.9 (26)	87	* *	285	32.3 (23)	276	32.2 (24)
Morris	23	* *	18	* *	4	* *	17	* *	18	* *
Nemaha	40	* *	33	* *	15	* *	34	* *	33	* *
Neosho	112	45.5 (26)	54	* *	21	* *	100	33.0 (24)	58	* *
Ness	5	* *	4	* *	1	* *	3	* *	4	* *
Norton	31	* *	26	* *	12	* *	28	* *	28	* *
Osage	91	* *	74	* *	29	* *	83	* *	75	* *
Osborne	14	* *	10	* *	5	* *	12	* *	10	* *
Ottawa	27	* *	19	* *	9	* *	24	* *	21	* *
Pawnee	60	* *	46	* *	13	* *	49	* *	46	* *
Phillips	30	* *	24	* *	3	* *	19	* *	24	* *
Pottawatom	38	* *	32	* *	6	* *	30	* *	34	* *
Pratt	83	* *	69	* *	25	* *	62	* *	68	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7B
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by County

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Rawlins	12	* *	11	* *	4	* *	11	* *	11	* *
Reno	490	43.3 (22)	415	23.6 (15)	143	37.8 (3)	407	18.2 (18)	419	16.0 (17)
Republic	42	* *	31	* *	10	* *	33	* *	32	* *
Rice	75	* *	67	* *	23	* *	61	* *	68	* *
Riley	937	27.5 (6)	705	14.8 (7)	132	28.0 (5)	759	14.9 (14)	722	12.5 (14)
Rooks	49	* *	42	* *	19	* *	40	* *	43	* *
Rush	9	* *	9	* *	5	* *	7	* *	9	* *
Russell	42	* *	36	* *	16	* *	34	* *	36	* *
Saline	503	43.1 (21)	406	26.6 (21)	135	24.4 (7)	410	17.8 (17)	408	15.4 (16)
Scott	51	* *	44	* *	9	* *	44	* *	46	* *
Sedgwick	4,680	31.0 (8)	4,250	16.2 (9)	1,000	35.3 (4)	3,827	10.3 (5)	4,253	8.3 (9)
Seward	514	5.8 (1)	457	3.1 (1)	17	* *	435	14.7 (13)	463	11.7 (13)
Shawnee	1,411	37.1 (14)	1,103	23.8 (16)	315	27.9 (6)	1,191	12.3 (9)	1,113	5.8 (6)
Sheridan	9	* *	10	* *	1	* *	8	* *	10	* *
Sherman	59	* *	52	* *	10	* *	50	* *	52	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7B
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by County

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Smith	19	* *	14	* *	4	* *	17	* *	15	* *
Stafford	41	* *	36	* *	15	* *	35	* *	36	* *
Stanton	13	* *	12	* *	1	* *	7	* *	12	* *
Stevens	87	* *	77	* *	13	* *	70	* *	79	* *
Sumner	153	52.9 (31)	135	32.6 (28)	55	* *	117	46.2 (29)	138	49.3 (28)
Thomas	59	* *	53	* *	15	* *	48	* *	53	* *
Trego	16	* *	12	* *	5	* *	14	* *	12	* *
Wabaunsee	25	* *	24	* *	7	* *	20	* *	24	* *
Washington	27	* *	22	* *	5	* *	20	* *	22	* *
Wichita	18	* *	12	* *	2	* *	15	* *	12	* *
Wilson	99	* *	88	* *	40	* *	87	* *	88	* *
Woodson	8	* *	8	* *	2	* *	6	* *	8	* *
Wyandotte	2,255	19.8 (4)	2,000	12.1 (5)	299	23.7 (8)	1,897	22.5 (21)	2,004	19.4 (20)
Unknown Counties	3	* *	3	* *	1	* *	1	* *	3	* *
Kansas	21,732	31.5	18,530	17.8	4,411	31.7	17,945	18.8	18,640	15.9

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7B
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Smoking Indicators by County

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Nation (Prior Year)	999,199	23.1	856,953	14.2	146,933	35.5	655,095	18.9	578,852	19.9

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8B
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Infant Health Indicators by County

Comparison of Infant Health Indicators (2, 3)										
County	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)		Number	%	Number	%	Number	%
	Number	% (Rank)	% (Rank)	Number						
Allen	81	* * * *	* *	82	* *	71	* *	82	* *	
Anderson	33	* * * *	* *	43	* *	30	* *	43	* *	
Atchison	136	6.6 (14)	5.9 (9)	139	6.5 (4)	127	3.1 (6)	139	54.7 (29)	
Barber	35	* * * *	* *	34	* *	29	* *	35	* *	
Barton	204	4.4 (1)	2.9 (1)	206	9.2 (15)	185	3.2 (7)	204	72.1 (8)	
Bourbon	98	* * * *	* *	99	* *	88	* *	99	* *	
Brown	80	* * * *	* *	88	* *	76	* *	88	* *	
Butler	246	5.7 (9)	7.3 (17)	250	5.2 (3)	233	3.4 (9)	248	70.2 (13)	
Chase	13	* * * *	* *	13	* *	12	* *	13	* *	
Chautauqua	30	* * * *	* *	31	* *	27	* *	32	* *	
Cherokee	185	5.4 (6)	8.1 (21)	183	4.9 (1)	174	2.9 (4)	183	59.0 (27)	
Cheyenne	2	* * * *	* *	2	* *	2	* *	2	* *	
Clark	3	* * * *	* *	3	* *	2	* *	3	* *	
Clay	52	* * * *	* *	52	* *	46	* *	52	* *	

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8B
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Infant Health Indicators by County

Comparison of Infant Health Indicators (2, 3)										
County	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)		Number	%	Number	%	Number	%
	Number	% (Rank)	% (Rank)	(Rank)						
Cloud	78	* * * *	* * * *	77	* *	69	* *	78	* *	
Coffey	19	* * * *	* * * *	19	* *	18	* *	19	* *	
Comanche	11	* * * *	* * * *	11	* *	9	* *	11	* *	
Cowley	306	9.2 (26)	7.5 (18)	309	11.7 (28)	266	5.6 (25)	310	63.5 (22)	
Crawford	330	9.4 (27)	5.2 (6)	342	11.1 (26)	292	5.1 (19)	340	55.6 (28)	
Decatur	8	* * * *	* * * *	8	* *	7	* *	8	* *	
Dickinson	116	6.0 (11)	10.3 (28)	115	9.6 (17)	104	2.9 (4)	116	71.6 (11)	
Doniphan	31	* * * *	* * * *	32	* *	30	* *	32	* *	
Douglas	422	6.4 (13)	5.2 (6)	430	7.7 (8)	390	3.8 (12)	427	77.8 (4)	
Edwards	23	* * * *	* * * *	23	* *	19	* *	23	* *	
Elk	19	* * * *	* * * *	19	* *	17	* *	19	* *	
Ellis	136	4.4 (1)	10.3 (28)	150	7.3 (7)	124	1.6 (2)	150	72.0 (10)	
Ellsworth	29	* * * *	* * * *	29	* *	26	* *	29	* *	
Finney	611	7.2 (16)	6.7 (15)	625	10.1 (25)	544	4.0 (15)	621	77.9 (3)	

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8B
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Infant Health Indicators by County

Comparison of Infant Health Indicators (2, 3)										
County	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)		Number	%	Number	%	Number	%
	Number	% (Rank)	% (Rank)	Number						
Ford	568	4.6 (3)	6.0 (11)	588	10.0 (24)	510	1.6 (2)	587	72.1 (8)	
Franklin	164	5.5 (7)	4.9 (4)	163	9.2 (15)	146	3.4 (9)	165	68.5 (19)	
Geary	314	8.9 (23)	5.1 (5)	313	9.9 (21)	282	5.3 (23)	311	71.4 (12)	
Gove	11	* * *	* *	11	* *	10	* *	9	* *	
Graham	12	* * *	* *	13	* *	12	* *	13	* *	
Grant	65	* * *	* *	66	* *	59	* *	67	* *	
Gray	36	* * *	* *	38	* *	32	* *	37	* *	
Greeley	13	* * *	* *	13	* *	12	* *	13	* *	
Greenwood	40	* * *	* *	40	* *	38	* *	40	* *	
Hamilton	26	* * *	* *	26	* *	23	* *	26	* *	
Harper	43	* * *	* *	43	* *	37	* *	45	* *	
Harvey	234	6.0 (11)	8.5 (25)	234	8.5 (10)	211	3.8 (12)	235	80.0 (1)	
Haskell	15	* * *	* *	16	* *	14	* *	16	* *	
Hodgeman	15	* * *	* *	15	* *	13	* *	15	* *	

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8B
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Infant Health Indicators by County

Comparison of Infant Health Indicators (2, 3)										
County	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)		Number	%	Number	%	Number	%
	Number	% (Rank)	% (Rank)	Number						
Jackson	65	* * * *	* * * *	65	* *	57	* *	65	* *	
Jefferson	61	* * * *	* * * *	62	* *	49	* *	61	* *	
Jewell	9	* * * *	* * * *	9	* *	8	* *	9	* *	
Johnson	1,561	6.6 (14)	7.7 (19)	1,662	7.0 (6)	1,444	3.9 (14)	1,663	79.9 (2)	
Kearny	17	* * * *	* * * *	17	* *	15	* *	17	* *	
Kingman	34	* * * *	* * * *	47	* *	32	* *	48	* *	
Kiowa	13	* * * *	* * * *	13	* *	10	* *	13	* *	
Labette	167	4.8 (4)	6.0 (11)	168	8.9 (13)	153	3.3 (8)	166	51.2 (30)	
Lane	10	* * * *	* * * *	11	* *	9	* *	11	* *	
Leavenworth	298	9.1 (25)	8.4 (22)	303	8.6 (12)	272	6.3 (27)	304	62.2 (24)	
Lincoln	17	* * * *	* * * *	17	* *	17	* *	16	* *	
Linn	59	* * * *	* * * *	60	* *	57	* *	60	* *	
Logan	12	* * * *	* * * *	12	* *	11	* *	12	* *	
Lyon	273	8.4 (20)	5.9 (9)	272	9.6 (17)	246	5.7 (26)	269	72.9 (6)	

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8B
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Infant Health Indicators by County

Comparison of Infant Health Indicators (2, 3)											
County	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed		
	Low (4)		High (5)		Number	%	Number	%	Number	%	
	Number	% (Rank)	% (Rank)	(Rank)							(Rank)
Marion	122	4.9 (5)	6.6 (14)		123	6.5 (4)		114	0.9 (1)	123	69.1 (16)
Marshall	32	*	*	*	34	*	*	31	*	33	*
Mcpherson	40	*	*	*	41	*	*	36	*	41	*
Meade	30	*	*	*	33	*	*	27	*	33	*
Miami	128	10.2 (28)	8.6 (26)		130	8.5 (10)		118	6.8 (28)	128	68.8 (18)
Mitchell	36	*	*	*	47	*	*	34	*	47	*
Montgomery	293	7.8 (17)	3.4 (2)		292	9.9 (21)		262	4.2 (17)	293	61.1 (26)
Morris	15	*	*	*	18	*	*	15	*	17	*
Nemaha	26	*	*	*	40	*	*	24	*	39	*
Neosho	70	*	*	*	133	12.8 (30)		61	*	132	62.1 (25)
Ness	4	*	*	*	4	*	*	3	*	4	*
Norton	28	*	*	*	28	*	*	22	*	27	*
Osage	85	*	*	*	82	*	*	67	*	87	*
Osborne	9	*	*	*	10	*	*	9	*	9	*

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8B
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Infant Health Indicators by County

Comparison of Infant Health Indicators (2, 3)														
County	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed						
	Low (4)	High (5)		Number	%	Number	%	Number	%					
	Number	% (Rank)	% (Rank)							% (Rank)	% (Rank)			
Ottawa	25	*	*	*	*	25	*	*	24	*	*	25	*	*
Pawnee	48	*	*	*	*	47	*	*	44	*	*	48	*	*
Phillips	32	*	*	*	*	32	*	*	25	*	*	32	*	*
Pottawatom	34	*	*	*	*	34	*	*	30	*	*	34	*	*
Pratt	66	*	*	*	*	65	*	*	61	*	*	66	*	*
Rawlins	11	*	*	*	*	11	*	*	10	*	*	11	*	*
Reno	428	8.2 (18)	3.5 (3)			427	9.6 (17)			384	5.2 (22)	431	66.6 (20)	
Republic	32	*	*	*	*	33	*	*	32	*	*	33	*	*
Rice	65	*	*	*	*	63	*	*	54	*	*	65	*	*
Riley	756	5.6 (8)	7.9 (20)			759	4.9 (1)			718	3.5 (11)	759	76.5 (5)	
Rooks	45	*	*	*	*	45	*	*	44	*	*	45	*	*
Rush	9	*	*	*	*	8	*	*	6	*	*	9	*	*
Russell	35	*	*	*	*	35	*	*	32	*	*	35	*	*
Saline	466	8.8 (22)	8.4 (22)			478	9.8 (20)			418	5.3 (23)	476	63.2 (23)	

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8B
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Infant Health Indicators by County

Comparison of Infant Health Indicators (2, 3)										
County	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)		Number	%	Number	%	Number	%
	Number	% (Rank)	% (Rank)	Number						
Scott	45	* * * *	* * * *	43	* *	40	* *	45	* *	
Sedgwick	4,182	9.0 (24)	6.8 (16)	4,198	8.2 (9)	3,828	5.1 (19)	4,190	68.9 (17)	
Seward	442	5.9 (10)	8.4 (22)	466	9.9 (21)	394	4.1 (16)	464	64.0 (21)	
Shawnee	1,273	8.6 (21)	5.5 (8)	1,287	11.5 (27)	1,119	5.1 (19)	1,285	72.3 (7)	
Sheridan	10	* * * *	* * * *	10	* *	8	* *	10	* *	
Sherman	52	* * * *	* * * *	52	* *	51	* *	52	* *	
Smith	15	* * * *	* * * *	15	* *	12	* *	16	* *	
Stafford	35	* * * *	* * * *	37	* *	31	* *	34	* *	
Stanton	12	* * * *	* * * *	14	* *	12	* *	14	* *	
Stevens	46	* * * *	* * * *	75	* *	42	* *	76	* *	
Sumner	135	11.1 (29)	8.9 (27)	140	12.1 (29)	118	6.8 (28)	140	70.0 (15)	
Thomas	54	* * * *	* * * *	54	* *	47	* *	54	* *	
Trego	16	* * * *	* * * *	16	* *	16	* *	16	* *	
Wabaunsee	24	* * * *	* * * *	24	* *	24	* *	24	* *	

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8B
2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Infant Health Indicators by County

Comparison of Infant Health Indicators (2, 3)														
County	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed					
	Low (4)		High (5)		Number	%	Number	%	Number	%				
	Number	% (Rank)	% (Rank)	Number							% (Rank)	Number	% (Rank)	
Washington	21	*	*	*	*	22	*	*	16	*	*	22	*	*
Wichita	15	*	*	*	*	15	*	*	12	*	*	15	*	*
Wilson	93	*	*	*	*	94	*	*	81	*	*	93	*	*
Woodson	8	*	*	*	*	8	*	*	7	*	*	8	*	*
Wyandotte	1,925	8.3 (19)	6.4 (13)			1,966	8.9 (13)		1,742	4.8 (18)		1,970	70.2 (13)	
Unknown Counties	3	*	*	*	*	3	*	*	3	*	*	3	*	*
Kansas	18,865	7.7	6.5			19,327	8.8		17,104	4.4		19,312	70.3	
Nation (Prior Year)	1,085,134	8.2	6.6			1,025,278	11.5		890,981	3.6		952,980	65.4	

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3B
2007 - 2009 Pregnancy Nutrition Surveillance
Kansas
Comparison of Racial and Ethnic Distribution by County
Combined 3 Years

(1)

Comparison of Racial and Ethnic Distribution (2)								
County	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Allen	298	92.3	3.4	1.3	1.3	0.7	1.0	0.0
Anderson	155	92.9	0.6	4.5	0.6	1.3	0.0	0.0
Atchison	413	88.1	8.2	1.5	0.7	0.0	1.5	0.0
Barber	115	95.7	2.6	0.9	0.0	0.9	0.0	0.0
Barton	815	69.6	1.5	28.0	0.2	0.2	0.5	0.0
Bourbon	390	93.3	2.8	2.1	0.3	0.3	1.3	0.0
Brown	293	76.1	2.7	4.1	14.7	0.0	2.4	0.0
Butler	815	90.8	1.3	5.4	0.4	0.4	1.7	0.0
Chase	40	*	*	*	*	*	*	*
Chautauqua	89	*	*	*	*	*	*	*
Cherokee	626	90.7	1.3	3.8	1.8	1.6	0.8	0.0
Cheyenne	6	*	*	*	*	*	*	*
Clark	10	*	*	*	*	*	*	*
Clay	185	95.7	0.5	1.6	1.1	0.5	0.5	0.0
Cloud	240	93.8	1.3	2.9	0.8	0.8	0.4	0.0
Coffey	69	*	*	*	*	*	*	*
Comanche	24	*	*	*	*	*	*	*
Cowley	1,020	76.5	2.4	17.8	1.0	1.6	0.8	0.0
Crawford	1,157	84.1	3.5	9.7	0.1	1.5	1.2	0.0
Decatur	30	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3B
2007 - 2009 Pregnancy Nutrition Surveillance
Kansas
Comparison of Racial and Ethnic Distribution by County
Combined 3 Years

(1)

Comparison of Racial and Ethnic Distribution (2)								
County	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Dickinson	348	92.8	0.0	5.2	1.4	0.3	0.3	0.0
Doniphan	122	86.1	4.1	7.4	0.8	0.0	1.6	0.0
Douglas	1,442	63.9	9.1	10.9	9.2	5.2	1.8	0.0
Edwards	69	*	*	*	*	*	*	*
Elk	48	*	*	*	*	*	*	*
Ellis	471	84.7	1.5	13.0	0.4	0.4	0.0	0.0
Ellsworth	94	*	*	*	*	*	*	*
Finney	2,024	30.0	0.9	66.2	0.0	2.6	0.2	0.0
Ford	1,930	22.8	0.9	75.4	0.2	0.7	0.1	0.0
Franklin	509	90.6	1.8	4.7	1.2	0.4	1.4	0.0
Geary	1,056	53.1	24.4	15.2	0.6	2.8	3.9	0.0
Gove	31	*	*	*	*	*	*	*
Graham	51	*	*	*	*	*	*	*
Grant	210	19.5	0.0	80.0	0.0	0.0	0.5	0.0
Gray	157	67.5	0.0	31.8	0.0	0.0	0.6	0.0
Greeley	31	*	*	*	*	*	*	*
Greenwood	140	92.9	0.7	5.0	0.0	0.7	0.7	0.0
Hamilton	74	*	*	*	*	*	*	*
Harper	147	81.0	0.0	16.3	1.4	0.0	1.4	0.0
Harvey	790	69.9	2.5	24.7	1.1	0.3	1.5	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3B
2007 - 2009 Pregnancy Nutrition Surveillance
Kansas
Comparison of Racial and Ethnic Distribution by County
Combined 3 Years

(1)

Comparison of Racial and Ethnic Distribution (2)								
County	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Haskell	63	*	*	*	*	*	*	*
Hodgeman	39	*	*	*	*	*	*	*
Jackson	220	76.4	0.0	5.9	15.5	0.0	2.3	0.0
Jefferson	193	96.4	0.5	1.0	2.1	0.0	0.0	0.0
Jewell	39	*	*	*	*	*	*	*
Johnson	5,514	46.4	12.8	35.9	0.5	3.1	1.2	0.0
Kearny	60	*	*	*	*	*	*	*
Kingman	173	91.9	0.0	5.2	2.3	0.6	0.0	0.0
Kiowa	33	*	*	*	*	*	*	*
Labette	536	87.7	7.3	2.6	1.3	0.4	0.7	0.0
Lane	38	*	*	*	*	*	*	*
Leavenworth	1,045	69.4	15.6	9.8	0.7	1.4	3.2	0.0
Lincoln	60	*	*	*	*	*	*	*
Linn	186	96.2	0.0	2.7	0.0	0.0	1.1	0.0
Logan	36	*	*	*	*	*	*	*
Lyon	1,027	54.2	5.0	38.6	0.2	1.4	0.7	0.0
Marion	406	89.2	1.5	7.6	1.2	0.0	0.5	0.0
Marshall	136	92.6	0.0	5.9	0.7	0.7	0.0	0.0
Mcpherson	171	91.8	0.0	3.5	2.9	0.0	1.8	0.0
Meade	137	62.0	0.0	37.2	0.0	0.0	0.7	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3B
2007 - 2009 Pregnancy Nutrition Surveillance
Kansas
Comparison of Racial and Ethnic Distribution by County
Combined 3 Years

(1)

Comparison of Racial and Ethnic Distribution (2)								
County	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Miami	409	92.7	2.4	2.9	0.0	0.7	1.2	0.0
Mitchell	129	96.1	0.0	3.9	0.0	0.0	0.0	0.0
Montgomery	1,010	77.1	9.4	8.8	3.1	0.8	0.8	0.0
Morris	81	*	*	*	*	*	*	*
Nemaha	116	89.7	0.0	7.8	0.0	0.9	1.7	0.0
Neosho	438	90.6	2.5	5.3	0.5	0.5	0.7	0.0
Ness	18	*	*	*	*	*	*	*
Norton	85	*	*	*	*	*	*	*
Osage	264	96.6	0.0	1.9	0.8	0.0	0.8	0.0
Osborne	53	*	*	*	*	*	*	*
Ottawa	76	*	*	*	*	*	*	*
Pawnee	164	81.7	2.4	14.6	0.0	0.0	1.2	0.0
Phillips	107	93.5	0.0	5.6	0.0	0.9	0.0	0.0
Pottawatom	130	90.8	0.0	6.9	0.8	0.0	1.5	0.0
Pratt	227	87.2	0.0	12.3	0.0	0.4	0.0	0.0
Rawlins	31	*	*	*	*	*	*	*
Reno	1,434	81.8	2.9	13.2	0.5	0.8	0.9	0.0
Republic	103	91.3	1.9	4.9	0.0	0.0	1.9	0.0
Rice	209	77.0	1.4	21.5	0.0	0.0	0.0	0.0
Riley	2,639	73.4	11.4	10.3	0.6	3.9	0.3	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3B
2007 - 2009 Pregnancy Nutrition Surveillance
Kansas
Comparison of Racial and Ethnic Distribution by County
Combined 3 Years

(1)

Comparison of Racial and Ethnic Distribution (2)								
County	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Rooks	130	95.4	0.8	0.8	1.5	1.5	0.0	0.0
Rush	35	*	*	*	*	*	*	*
Russell	126	92.1	4.8	1.6	1.6	0.0	0.0	0.0
Saline	1,437	71.7	4.4	19.7	0.4	2.3	1.5	0.0
Scott	143	59.4	0.7	39.9	0.0	0.0	0.0	0.0
Sedgwick	12,755	45.6	18.4	29.1	1.7	3.8	1.4	0.0
Seward	1,478	17.4	1.4	78.3	1.8	0.7	0.3	0.0
Shawnee	4,303	55.5	15.8	24.4	2.2	0.7	1.3	0.0
Sheridan	27	*	*	*	*	*	*	*
Sherman	190	75.3	0.0	22.6	0.5	0.5	1.1	0.0
Smith	61	*	*	*	*	*	*	*
Stafford	118	66.9	0.0	32.2	0.0	0.8	0.0	0.0
Stanton	56	*	*	*	*	*	*	*
Stevens	250	45.2	0.0	53.6	0.8	0.4	0.0	0.0
Sumner	443	91.2	0.9	5.0	1.1	0.2	1.6	0.0
Thomas	160	86.9	1.9	10.0	0.0	0.0	1.3	0.0
Trego	50	*	*	*	*	*	*	*
Wabaunsee	62	*	*	*	*	*	*	*
Wallace	4	*	*	*	*	*	*	*
Washington	83	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3B
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Racial and Ethnic Distribution by County
Combined 3 Years

Comparison of Racial and Ethnic Distribution (2)								
County	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Wichita	46	*	*	*	*	*	*	*
Wilson	281	96.4	0.7	2.1	0.4	0.0	0.4	0.0
Woodson	25	*	*	*	*	*	*	*
Wyandotte	6,467	26.0	31.1	39.9	0.2	2.5	0.3	0.0
Unknown Counties	3	*	*	*	*	*	*	*
Kansas	63,102	57.2	11.4	27.0	1.3	2.1	1.0	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4B
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Age Distribution by County
Combined 3 Years

Comparison of Age Distribution (2)								
County	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Allen	298	0.3	8.1	15.4	63.8	11.4	0.3	0.7
Anderson	155	0.6	8.4	12.3	67.1	11.0	0.6	0.0
Atchison	413	0.0	6.5	17.4	61.5	13.8	0.7	0.0
Barber	115	0.0	4.3	22.6	60.0	13.0	0.0	0.0
Barton	815	0.1	6.6	17.8	59.9	15.0	0.6	0.0
Bourbon	390	0.0	6.9	19.7	61.8	11.0	0.5	0.0
Brown	293	0.0	6.5	15.4	64.8	12.3	1.0	0.0
Butler	815	0.1	7.6	17.2	61.7	12.8	0.5	0.1
Chase	40	*	*	*	*	*	*	*
Chautauqua	89	*	*	*	*	*	*	*
Cherokee	626	0.2	8.6	18.7	58.9	12.5	0.8	0.3
Cheyenne	6	*	*	*	*	*	*	*
Clark	10	*	*	*	*	*	*	*
Clay	185	0.0	7.6	17.8	60.0	14.1	0.5	0.0
Cloud	240	0.0	5.4	17.5	65.4	10.4	1.3	0.0
Coffey	69	*	*	*	*	*	*	*
Comanche	24	*	*	*	*	*	*	*
Cowley	1,020	0.8	8.1	16.4	60.9	13.4	0.4	0.0
Crawford	1,157	0.2	7.2	16.2	63.1	12.5	0.6	0.2

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4B
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Age Distribution by County
Combined 3 Years

Comparison of Age Distribution (2)								
County	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Decatur	30	*	*	*	*	*	*	*
Dickinson	348	0.3	7.2	14.7	62.4	15.2	0.3	0.0
Doniphan	122	0.8	4.1	14.8	63.9	16.4	0.0	0.0
Douglas	1,442	0.1	3.8	13.2	63.9	17.7	1.4	0.0
Edwards	69	*	*	*	*	*	*	*
Elk	48	*	*	*	*	*	*	*
Ellis	471	0.0	3.8	11.7	70.9	11.5	1.9	0.2
Ellsworth	94	*	*	*	*	*	*	*
Finney	2,024	0.4	9.9	14.8	54.2	19.2	1.4	0.1
Ford	1,930	0.4	7.6	12.5	56.1	22.1	1.3	0.1
Franklin	509	0.8	6.3	19.8	58.2	14.1	0.6	0.2
Geary	1,056	0.5	4.9	15.4	67.2	11.6	0.4	0.0
Gove	31	*	*	*	*	*	*	*
Graham	51	*	*	*	*	*	*	*
Grant	210	0.0	4.8	15.7	60.0	17.6	1.9	0.0
Gray	157	0.0	5.7	10.8	53.5	28.0	1.9	0.0
Greeley	31	*	*	*	*	*	*	*
Greenwood	140	0.0	5.7	17.1	62.1	15.0	0.0	0.0
Hamilton	74	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4B
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Age Distribution by County
Combined 3 Years

Comparison of Age Distribution (2)								
County	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Harper	147	0.0	7.5	12.2	70.1	10.2	0.0	0.0
Harvey	790	0.3	6.5	13.4	63.3	15.3	1.3	0.0
Haskell	63	*	*	*	*	*	*	*
Hodgeman	39	*	*	*	*	*	*	*
Jackson	220	0.0	7.3	15.9	66.8	10.0	0.0	0.0
Jefferson	193	0.0	8.3	17.6	60.1	13.5	0.5	0.0
Jewell	39	*	*	*	*	*	*	*
Johnson	5,514	0.2	4.4	11.6	60.8	21.7	1.3	0.1
Kearny	60	*	*	*	*	*	*	*
Kingman	173	0.0	5.8	14.5	64.2	15.0	0.6	0.0
Kiowa	33	*	*	*	*	*	*	*
Labette	536	0.0	7.5	19.4	62.1	10.1	0.9	0.0
Lane	38	*	*	*	*	*	*	*
Leavenworth	1,045	0.1	5.9	18.4	63.3	11.3	1.1	0.0
Lincoln	60	*	*	*	*	*	*	*
Linn	186	0.0	5.4	17.2	62.9	13.4	1.1	0.0
Logan	36	*	*	*	*	*	*	*
Lyon	1,027	0.4	6.2	13.7	62.7	14.7	2.2	0.0
Marion	406	0.2	4.9	13.1	65.8	15.5	0.5	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4B
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Age Distribution by County
Combined 3 Years

Comparison of Age Distribution (2)								
County	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Marshall	136	0.0	2.2	12.5	66.9	14.7	3.7	0.0
Mcpherson	171	0.0	5.3	9.9	69.0	15.8	0.0	0.0
Meade	137	0.0	7.3	10.9	60.6	19.7	1.5	0.0
Miami	409	0.2	7.3	14.4	65.5	12.0	0.5	0.0
Mitchell	129	0.0	2.3	16.3	61.2	20.2	0.0	0.0
Montgomery	1,010	0.3	5.7	17.1	65.2	11.1	0.3	0.2
Morris	81	*	*	*	*	*	*	*
Nemaha	116	0.0	5.2	12.9	62.9	15.5	2.6	0.9
Neosho	438	0.2	7.1	19.6	61.4	11.2	0.5	0.0
Ness	18	*	*	*	*	*	*	*
Norton	85	*	*	*	*	*	*	*
Osage	264	0.0	5.7	18.2	65.5	9.8	0.0	0.8
Osborne	53	*	*	*	*	*	*	*
Ottawa	76	*	*	*	*	*	*	*
Pawnee	164	1.2	6.7	14.6	68.3	7.9	1.2	0.0
Phillips	107	0.0	2.8	15.0	63.6	18.7	0.0	0.0
Pottawatom	130	0.0	8.5	7.7	56.9	24.6	2.3	0.0
Pratt	227	0.4	9.3	17.6	58.1	13.7	0.9	0.0
Rawlins	31	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4B
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Age Distribution by County
Combined 3 Years

Comparison of Age Distribution (2)								
County	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Reno	1,434	0.1	5.6	16.5	65.8	11.4	0.6	0.0
Republic	103	0.0	3.9	22.3	55.3	16.5	1.9	0.0
Rice	209	0.5	6.2	13.4	58.9	20.1	1.0	0.0
Riley	2,639	0.0	1.9	12.1	74.6	11.0	0.4	0.0
Rooks	130	0.0	3.8	13.8	63.8	16.9	1.5	0.0
Rush	35	*	*	*	*	*	*	*
Russell	126	0.0	6.3	16.7	65.1	11.9	0.0	0.0
Saline	1,437	0.2	7.7	15.1	63.1	12.9	1.0	0.1
Scott	143	0.7	5.6	10.5	66.4	15.4	1.4	0.0
Sedgwick	12,755	0.3	6.9	13.9	60.8	16.7	1.3	0.0
Seward	1,478	0.3	7.7	13.9	55.8	20.6	1.6	0.0
Shawnee	4,303	0.4	6.9	14.7	61.6	15.3	1.1	0.0
Sheridan	27	*	*	*	*	*	*	*
Sherman	190	0.0	4.7	18.4	57.4	16.3	3.2	0.0
Smith	61	*	*	*	*	*	*	*
Stafford	118	0.0	4.2	12.7	59.3	21.2	2.5	0.0
Stanton	56	*	*	*	*	*	*	*
Stevens	250	0.0	6.8	14.8	59.2	18.8	0.4	0.0
Sumner	443	0.0	5.0	18.7	62.8	12.9	0.7	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4B
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Age Distribution by County
Combined 3 Years

Comparison of Age Distribution (2)								
County	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Thomas	160	0.0	7.5	15.6	65.0	11.9	0.0	0.0
Trego	50	*	*	*	*	*	*	*
Wabaunsee	62	*	*	*	*	*	*	*
Wallace	4	*	*	*	*	*	*	*
Washington	83	*	*	*	*	*	*	*
Wichita	46	*	*	*	*	*	*	*
Wilson	281	0.0	8.9	16.0	62.6	12.5	0.0	0.0
Woodson	25	*	*	*	*	*	*	*
Wyandotte	6,467	0.6	8.2	13.9	58.0	18.0	1.3	0.1
Unknown Counties	3	*	*	*	*	*	*	*
Kansas	63,102	0.3	6.5	14.5	61.6	16.1	1.1	0.1

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5B
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Maternal Health Indicators by County
Combined 3 Years

Comparison of Maternal Health Indicators (2, 3)														
County	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Allen	296	5.1 (42)	54.4 (39)	273	18.3 (20)	56.4 (59)	51	* *	92	* *	267	6.7 (60)	267	12.4(61)
Anderson	154	4.5 (29)	50.6 (13)	143	22.4 (48)	47.6 (12)	26	* *	27	* *	137	2.2 (6)	137	10.9(59)
Atchison	400	5.8 (47)	56.5 (50)	349	19.5 (33)	53.9 (45)	61	* *	87	* *	354	4.0 (38)	353	12.2(60)
Barber	113	3.5 (15)	57.5 (55)	94	* *	* *	15	* *	27	* *	93	* *	93	* *
Barton	803	4.5 (29)	52.6 (24)	724	20.9 (41)	48.8 (16)	97	* *	156	25.0 (17)	692	3.2 (21)	690	7.7(44)
Bourbon	384	3.1 (11)	56.8 (53)	343	19.5 (33)	52.8 (38)	47	* *	78	* *	339	3.2 (21)	339	6.5(24)
Brown	290	3.1 (11)	55.2 (44)	256	16.8 (6)	56.3 (58)	45	* *	65	* *	257	1.6 (4)	257	9.7(57)
Butler	791	6.3 (53)	50.2 (11)	728	18.7 (25)	52.5 (37)	103	38.8 (18)	216	28.7 (24)	692	4.2 (42)	692	8.8(53)
Chase	38	* *	* *	33	* *	* *	4	* *	3	* *	35	* *	35	* *
Chautauqua	89	* *	* *	79	* *	* *	7	* *	12	* *	79	* *	79	* *
Cherokee	604	5.0 (41)	51.8 (18)	526	16.2 (3)	54.8 (50)	74	* *	139	27.3 (23)	525	3.0 (18)	526	6.7(31)
Cheyenne	6	* *	* *	6	* *	* *	0	* *	1	* *	6	* *	6	* *
Clark	8	* *	* *	6	* *	* *	1	* *	2	* *	7	* *	7	* *
Clay	183	3.8 (19)	56.3 (48)	163	13.5 (1)	61.3 (62)	21	* *	47	* *	158	5.7 (58)	158	7.0(33)
Cloud	240	4.6 (31)	51.3 (16)	214	21.5 (45)	53.3 (42)	28	* *	62	* *	202	4.0 (38)	206	8.7(52)
Coffey	67	* *	* *	62	* *	* *	6	* *	25	* *	56	* *	56	* *
Comanche	24	* *	* *	22	* *	* *	6	* *	7	* *	20	* *	20	* *

- (1) Reporting period is January 1 through December 31.
 - (2) Excludes records with unknown data or errors.
 - (3) Rank compares one county's rate to other counties. Rank 1 = best rate.
 - (4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
 - (5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
 - (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5B
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Maternal Health Indicators by County
Combined 3 Years

Comparison of Maternal Health Indicators (2, 3)														
County	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Cowley	1,009	5.1 (42)	52.9 (26)	915	20.8 (39)	52.0 (32)	145	32.4 (16)	129	23.3 (11)	877	3.3 (24)	878	7.4(42)
Crawford	1,129	5.9 (48)	53.3 (29)	978	18.8 (27)	55.8 (56)	176	17.0 (2)	190	18.4 (7)	977	4.4 (46)	976	7.3(40)
Decatur	30	* *	* *	28	* *	* *	4	* *	5	* *	27	* *	27	* *
Dickinson	340	4.7 (34)	52.1 (20)	312	20.5 (38)	53.8 (44)	45	* *	84	* *	304	4.9 (49)	300	7.3(40)
Doniphan	119	2.5 (6)	63.9 (65)	97	* *	* *	25	* *	24	* *	98	* *	98	* *
Douglas	1,421	4.4 (25)	49.3 (7)	1,246	17.3 (10)	56.1 (57)	202	31.2 (14)	403	24.8 (13)	1,199	3.9 (35)	1,198	5.2(14)
Edwards	69	* *	* *	59	* *	* *	8	* *	16	* *	55	* *	55	* *
Elk	47	* *	* *	44	* *	* *	1	* *	9	* *	42	* *	42	* *
Ellis	453	4.4 (25)	49.0 (5)	397	18.6 (23)	49.4 (19)	106	23.6 (4)	126	16.7 (4)	394	2.5 (11)	394	5.6(18)
Ellsworth	94	* *	* *	91	* *	* *	9	* *	23	* *	84	* *	84	* *
Finney	1,987	6.7 (56)	50.1 (10)	1,839	23.3 (54)	45.7 (7)	270	28.1 (10)	289	24.9 (16)	1,805	5.0 (51)	1,803	4.0(6)
Ford	1,906	2.4 (5)	54.7 (43)	1,727	21.1 (43)	48.2 (14)	252	21.8 (3)	200	16.0 (2)	1,690	3.3 (24)	1,689	5.1(11)
Franklin	499	3.2 (13)	54.3 (38)	437	16.7 (5)	55.4 (55)	49	* *	133	16.5 (3)	381	2.4 (9)	381	8.4(49)
Geary	1,031	4.6 (31)	51.6 (17)	860	17.6 (12)	54.3 (49)	165	38.8 (18)	193	29.5 (26)	854	3.4 (28)	854	7.8(45)
Gove	28	* *	* *	26	* *	* *	3	* *	10	* *	29	* *	29	* *
Graham	51	* *	* *	44	* *	* *	8	* *	8	* *	46	* *	46	* *
Grant	205	3.9 (20)	66.3 (66)	194	19.6 (35)	53.1 (41)	27	* *	60	* *	189	3.2 (21)	187	3.2(2)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5B
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Maternal Health Indicators by County
Combined 3 Years

Comparison of Maternal Health Indicators (2, 3)														
County	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Gray	157	2.5 (6)	57.3 (54)	135	24.4 (56)	49.6 (21)	17	* *	42	* *	129	2.3 (7)	129	7.0(33)
Greeley	29	* *	* *	27	* *	* *	4	* *	6	* *	29	* *	29	* *
Greenwood	137	2.9 (8)	60.6 (61)	128	22.7 (50)	52.3 (34)	16	* *	25	* *	126	4.0 (38)	127	9.4(55)
Hamilton	71	* *	* *	60	* *	* *	9	* *	25	* *	63	* *	63	* *
Harper	144	7.6 (61)	49.3 (7)	123	22.8 (51)	47.2 (11)	16	* *	22	* *	118	3.4 (28)	117	6.0(22)
Harvey	779	4.2 (22)	54.0 (35)	707	22.8 (51)	48.8 (16)	114	30.7 (12)	133	25.6 (20)	679	3.5 (30)	677	6.6(28)
Haskell	61	* *	* *	53	* *	* *	12	* *	22	* *	54	* *	54	* *
Hodgeman	38	* *	* *	38	* *	* *	7	* *	10	* *	36	* *	36	* *
Jackson	214	5.1 (42)	48.6 (4)	193	17.6 (12)	54.9 (51)	34	* *	46	* *	192	2.1 (5)	192	6.8(32)
Jefferson	192	7.3 (59)	55.2 (44)	172	16.9 (8)	55.2 (53)	29	* *	58	* *	169	4.7 (48)	169	6.5(24)
Jewell	39	* *	* *	39	* *	* *	5	* *	5	* *	38	* *	38	* *
Johnson	5,359	4.2 (22)	53.9 (34)	4,828	18.2 (18)	51.3 (28)	920	24.9 (6)	1,632	17.7 (6)	4,702	5.3 (56)	4,699	4.6(8)
Kearny	60	* *	* *	56	* *	* *	15	* *	19	* *	56	* *	56	* *
Kingman	173	4.6 (31)	45.7 (2)	148	17.6 (12)	54.1 (47)	19	* *	37	* *	140	5.0 (51)	140	7.1(39)
Kiowa	31	* *	* *	26	* *	* *	4	* *	6	* *	28	* *	28	* *
Labette	530	6.8 (57)	55.8 (47)	487	23.4 (55)	51.1 (27)	90	* *	125	23.2 (10)	478	3.3 (24)	478	8.4(49)
Lane	35	* *	* *	30	* *	* *	7	* *	4	* *	30	* *	30	* *

- (1) Reporting period is January 1 through December 31.
 - (2) Excludes records with unknown data or errors.
 - (3) Rank compares one county's rate to other counties. Rank 1 = best rate.
 - (4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
 - (5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
 - (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5B
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Maternal Health Indicators by County
Combined 3 Years

Comparison of Maternal Health Indicators (2, 3)														
County	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy	Hypertension During Pregnancy		
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)			Number	% (Rank)
Leavenworth	1,019	4.9 (38)	53.5 (30)	905	18.6 (23)	55.0 (52)	171	29.2 (11)	246	19.5 (9)	857	3.7 (33)	857	5.5(17)
Lincoln	57	* *	* *	54	* *	* *	9	* *	16	* *	51	* *	51	* *
Linn	184	5.4 (46)	58.2 (56)	171	29.8 (62)	43.9 (3)	24	* *	30	* *	168	3.0 (18)	168	4.2(7)
Logan	36	* *	* *	29	* *	* *	6	* *	7	* *	28	* *	28	* *
Lyon	1,004	3.5 (15)	59.2 (58)	913	20.8 (39)	51.0 (26)	143	13.3 (1)	194	17.0 (5)	887	3.3 (24)	887	9.6(56)
Marion	399	7.5 (60)	51.9 (19)	367	19.1 (28)	46.6 (10)	62	* *	92	* *	357	2.5 (11)	358	7.0(33)
Marshall	133	9.0 (66)	42.9 (1)	119	18.5 (22)	44.5 (4)	14	* *	33	* *	114	0.9 (3)	114	7.0(33)
Mcperson	167	7.2 (58)	50.3 (12)	140	22.9 (53)	47.9 (13)	23	* *	35	* *	137	0.7 (2)	137	5.1(11)
Meade	136	3.7 (17)	59.6 (59)	116	21.6 (46)	50.9 (25)	18	* *	23	* *	115	5.2 (54)	115	5.2(14)
Miami	395	8.1 (63)	54.2 (36)	362	18.2 (18)	53.3 (42)	78	* *	116	25.0 (17)	352	5.1 (53)	352	6.5(24)
Mitchell	127	3.9 (20)	58.3 (57)	122	28.7 (61)	45.9 (8)	18	* *	37	* *	118	5.9 (59)	118	8.5(51)
Montgomery	999	6.3 (53)	53.6 (32)	882	22.0 (47)	49.5 (20)	169	30.8 (13)	278	23.7 (12)	866	2.8 (13)	866	6.6(28)
Morris	79	* *	* *	72	* *	* *	12	* *	19	* *	71	* *	71	* *
Nemaha	109	1.8 (2)	53.2 (28)	104	19.2 (29)	51.9 (30)	25	* *	37	* *	106	2.8 (13)	106	2.8(1)
Neosho	424	8.5 (65)	50.7 (14)	400	17.5 (11)	50.3 (23)	72	* *	71	* *	387	3.9 (35)	387	7.0(33)
Ness	17	* *	* *	15	* *	* *	1	* *	10	* *	13	* *	13	* *
Norton	85	* *	* *	75	* *	* *	10	* *	19	* *	73	* *	73	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5B
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Maternal Health Indicators by County
Combined 3 Years

Comparison of Maternal Health Indicators (2, 3)														
County	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Osage	262	6.1 (50)	53.1 (27)	229	17.9 (16)	52.8 (38)	29	* *	59	* *	228	3.9 (35)	228	5.7(20)
Osborne	53	* *	* *	47	* *	* *	6	* *	11	* *	47	* *	47	* *
Ottawa	74	* *	* *	67	* *	* *	9	* *	17	* *	66	* *	66	* *
Pawnee	162	8.0 (62)	55.6 (46)	141	27.0 (59)	46.1 (9)	11	* *	26	* *	142	2.8 (13)	142	4.9(9)
Phillips	105	2.9 (8)	61.0 (62)	93	* *	* *	6	* *	21	* *	90	* *	90	* *
Pottawatom	128	4.7 (34)	56.3 (48)	120	22.5 (49)	45.0 (5)	12	* *	61	* *	117	4.3 (44)	117	3.4(4)
Pratt	222	3.2 (13)	52.3 (21)	188	17.0 (9)	56.4 (59)	29	* *	61	* *	181	0.6 (1)	181	6.1(23)
Rawlins	30	* *	* *	28	* *	* *	4	* *	11	* *	28	* *	28	* *
Reno	1,419	4.9 (38)	54.2 (36)	1,280	20.1 (36)	51.9 (30)	180	26.7 (7)	314	24.8 (13)	1,216	2.3 (7)	1,216	8.3(48)
Republic	101	5.9 (48)	52.5 (23)	88	* *	* *	15	* *	19	* *	88	* *	88	* *
Rice	205	2.0 (3)	56.6 (51)	181	21.0 (42)	50.8 (24)	24	* *	45	* *	181	5.5 (57)	180	5.6(18)
Riley	2,565	4.4 (25)	49.2 (6)	2,119	16.5 (4)	54.2 (48)	386	24.4 (5)	566	26.0 (21)	2,054	2.9 (16)	2,053	6.5(24)
Rooks	129	6.2 (52)	49.6 (9)	119	26.1 (57)	42.9 (2)	17	* *	27	* *	118	4.2 (42)	118	7.6(43)
Rush	35	* *	* *	32	* *	* *	5	* *	8	* *	31	* *	31	* *
Russell	125	6.4 (55)	48.0 (3)	114	28.1 (60)	39.5 (1)	17	* *	32	* *	110	3.6 (32)	110	3.6(5)
Saline	1,415	4.7 (34)	53.6 (32)	1,307	18.4 (21)	52.3 (34)	264	27.7 (8)	340	19.4 (8)	1,277	3.5 (30)	1,281	5.0(10)
Scott	141	4.3 (24)	54.6 (42)	125	17.6 (12)	55.2 (53)	19	* *	27	* *	126	2.4 (9)	126	3.2(2)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5B
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Maternal Health Indicators by County
Combined 3 Years

Comparison of Maternal Health Indicators (2, 3)														
County	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Sedgwick	12,538	4.9 (38)	52.4 (22)	11,593	19.4 (32)	51.8 (29)	2,553	32.3 (15)	3,421	25.4 (19)	11,241	3.8 (34)	11,243	7.0(33)
Seward	1,450	2.9 (8)	54.4 (39)	1,328	21.4 (44)	50.1 (22)	249	27.7 (8)	218	24.8 (13)	1,294	5.2 (54)	1,296	5.3(16)
Shawnee	4,220	4.7 (34)	52.8 (25)	3,763	19.2 (29)	52.1 (33)	448	41.1 (20)	764	28.9 (25)	3,688	4.9 (49)	3,693	5.1(11)
Sheridan	26	* *	* *	23	* *	* *	5	* *	3	* *	24	* *	24	* *
Sherman	187	3.7 (17)	61.5 (63)	164	20.1 (36)	59.1 (61)	14	* *	27	* *	157	4.5 (47)	157	8.9(54)
Smith	61	* *	* *	51	* *	* *	11	* *	20	* *	52	* *	52	* *
Stafford	117	0.9 (1)	63.2 (64)	102	26.5 (58)	45.1 (6)	13	* *	15	* *	97	* *	97	* *
Stanton	56	* *	* *	54	* *	* *	7	* *	21	* *	54	* *	54	* *
Stevens	247	2.0 (3)	60.3 (60)	220	16.8 (6)	49.1 (18)	38	* *	63	* *	213	8.9 (61)	213	6.6(28)
Sumner	433	5.3 (45)	51.0 (15)	385	19.2 (29)	48.3 (15)	77	* *	113	12.4 (1)	378	4.0 (38)	378	9.8(58)
Thomas	155	8.4 (64)	53.5 (30)	141	15.6 (2)	53.9 (45)	23	* *	37	* *	139	2.9 (16)	139	7.9(47)
Trego	47	* *	* *	41	* *	* *	6	* *	9	* *	45	* *	45	* *
Wabaunsee	61	* *	* *	58	* *	* *	5	* *	25	* *	57	* *	57	* *
Wallace	4	* *	* *	4	* *	* *	1	* *	1	* *	3	* *	3	* *
Washington	81	* *	* *	71	* *	* *	4	* *	19	* *	70	* *	69	* *
Wichita	46	* *	* *	42	* *	* *	7	* *	10	* *	41	* *	41	* *
Wilson	279	6.1 (50)	56.6 (51)	265	18.1 (17)	52.8 (38)	46	* *	50	* *	255	4.3 (44)	255	7.8(45)

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one county's rate to other counties. Rank 1 = best rate.
- (4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
- (5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
- (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5B
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Maternal Health Indicators by County
Combined 3 Years

Comparison of Maternal Health Indicators (2, 3)														
County	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Woodson	24	* *	* *	23	* *	* *	4	* *	5	* *	24	* *	24	* *
Wyandotte	6,336	4.4 (25)	54.4 (39)	5,782	18.7 (25)	52.3 (34)	1,060	34.3 (17)	1,196	26.3 (22)	5,623	3.1 (20)	5,625	5.7 (20)
Unknown Counties	3	* *	* *	3	* *	* *	0	* *	1	* *	3	* *	3	* *
Kansas	61,869	4.7	53.2	55,802	19.5	51.6	9,864	30.1	14,091	23.5	54,295	3.8	54,295	6.4

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6B
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Maternal Behavioral Indicators by County
Combined 3 Years

Comparison of Maternal Behavioral Indicators (2, 3)										
County	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	% (Rank)		Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Allen	265	91.7 (2)	1.9 (15)		261	42.5 (29)	160	13.1 (24)	162	54.9 (38)
Anderson	138	89.1 (9)	2.2 (20)		133	45.1 (19)	88	* *	86	* *
Atchison	367	89.6 (8)	2.5 (23)		366	35.8 (56)	236	7.6 (36)	221	62.9 (24)
Barber	98	* *	* *		100	45.0 (21)	70	* *	67	* *
Barton	714	80.0 (31)	1.1 (9)		698	54.6 (4)	414	19.8 (5)	398	60.8 (28)
Bourbon	335	91.9 (1)	1.5 (12)		330	46.4 (17)	194	10.3 (31)	199	62.3 (26)
Brown	251	79.3 (35)	10.4 (56)		257	42.4 (30)	163	11.7 (27)	197	69.0 (8)
Butler	716	70.8 (55)	10.9 (57)		638	43.9 (23)	436	15.1 (17)	364	41.2 (43)
Chase	34	* *	* *		34	* *	25	* *	21	* *
Chautauqua	79	* *	* *		80	* *	48	* *	55	* *
Cherokee	463	77.3 (40)	6.7 (48)		515	39.6 (41)	327	13.1 (24)	296	58.4 (35)
Cheyenne	4	* *	* *		5	* *	3	* *	2	* *
Clark	6	* *	* *		8	* *	4	* *	5	* *
Clay	164	80.5 (30)	6.1 (45)		159	47.8 (14)	94	* *	92	* *
Cloud	214	83.2 (22)	2.8 (25)		205	49.8 (9)	133	16.5 (12)	113	59.3 (32)
Coffey	64	* *	* *		47	* *	39	* *	27	* *
Comanche	22	* *	* *		20	* *	14	* *	14	* *
Cowley	866	68.4 (57)	5.5 (43)		870	44.9 (22)	557	12.2 (26)	535	60.6 (30)
Crawford	975	86.9 (11)	2.1 (18)		1,016	43.8 (24)	636	11.3 (28)	653	66.0 (16)
Decatur	22	* *	* *		27	* *	15	* *	17	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6B
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Maternal Behavioral Indicators by County
Combined 3 Years

Comparison of Maternal Behavioral Indicators (2, 3)										
County	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	% (Rank)		Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Dickinson	318	82.1 (25)	3.8 (34)		294	48.6 (12)	202	18.8 (8)	173	70.5 (6)
Doniphan	106	78.3 (38)	5.7 (44)		103	39.8 (40)	64	* *	72	* *
Douglas	1,243	72.3 (52)	6.9 (51)		1,242	39.3 (42)	765	17.8 (9)	719	63.4 (22)
Edwards	56	* *	* *		59	* *	44	* *	35	* *
Elk	42	* *	* *		40	* *	25	* *	20	* *
Ellis	418	85.4 (16)	1.9 (15)		406	33.3 (60)	275	25.8 (1)	262	74.8 (2)
Ellsworth	92	* *	* *		73	* *	49	* *	40	* *
Finney	1,721	64.4 (60)	18.3 (61)		1,741	37.7 (49)	1,131	10.3 (31)	1,052	39.5 (44)
Ford	1,668	72.1 (53)	14.0 (60)		1,712	45.7 (18)	1,033	6.8 (37)	1,074	57.3 (36)
Franklin	254	79.5 (33)	3.1 (28)		414	41.8 (35)	278	3.2 (42)	292	68.2 (9)
Geary	822	70.7 (56)	0.7 (4)		923	40.4 (39)	588	17.7 (10)	569	60.8 (28)
Gove	28	* *	* *		22	* *	15	* *	12	* *
Graham	46	* *	* *		44	* *	21	* *	23	* *
Grant	183	77.0 (42)	9.8 (55)		166	36.1 (55)	107	11.2 (29)	94	* *
Gray	133	79.7 (32)	2.3 (21)		133	43.6 (25)	77	* *	75	* *
Greeley	29	* *	* *		27	* *	22	* *	17	* *
Greenwood	118	61.9 (62)	29.7 (62)		124	54.8 (3)	69	* *	73	* *
Hamilton	61	* *	* *		62	* *	41	* *	40	* *
Harper	120	72.5 (50)	12.5 (58)		129	48.8 (11)	83	* *	85	* *
Harvey	659	72.1 (53)	13.8 (59)		688	47.2 (16)	422	14.7 (19)	417	70.5 (6)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6B
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Maternal Behavioral Indicators by County
Combined 3 Years

Comparison of Maternal Behavioral Indicators (2, 3)										
County	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	% (Rank)		Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Haskell	52	* *	* *		57	* *	36	* *	35	* *
Hodgeman	38	* *	* *		29	* *	23	* *	19	* *
Jackson	139	81.3 (29)	5.0 (41)		188	42.6 (28)	114	8.8 (33)	115	80.9 (1)
Jefferson	162	90.7 (4)	1.9 (15)		153	39.2 (43)	104	6.7 (38)	105	54.3 (39)
Jewell	38	* *	* *		31	* *	17	* *	21	* *
Johnson	4,545	64.0 (61)	3.0 (27)		4,617	31.6 (62)	2,938	13.2 (23)	2,727	67.1 (10)
Kearny	54	* *	* *		53	* *	29	* *	27	* *
Kingman	142	81.7 (27)	4.9 (40)		147	49.7 (10)	98	* *	102	71.6 (5)
Kiowa	28	* *	* *		32	* *	22	* *	23	* *
Labette	434	85.5 (15)	3.2 (29)		471	42.0 (33)	286	15.4 (15)	300	63.7 (21)
Lane	31	* *	* *		34	* *	20	* *	21	* *
Leavenworth	915	85.4 (16)	0.7 (4)		823	36.5 (53)	561	18.9 (6)	485	72.4 (4)
Lincoln	55	* *	* *		46	* *	35	* *	31	* *
Linn	170	79.4 (34)	6.5 (46)		164	45.1 (19)	96	* *	109	66.1 (15)
Logan	14	* *	* *		32	* *	23	* *	23	* *
Lyon	927	81.6 (28)	1.1 (9)		841	41.5 (36)	516	14.9 (18)	476	64.7 (19)
Marion	372	86.6 (12)	1.3 (11)		334	38.6 (47)	218	20.2 (4)	206	67.0 (11)
Marshall	120	72.5 (50)	3.3 (30)		109	42.2 (32)	74	* *	62	* *
Mcperson	147	90.5 (5)	0.7 (4)		138	42.0 (33)	81	* *	72	* *
Meade	119	75.6 (45)	6.7 (48)		115	52.2 (6)	70	* *	72	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6B
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Maternal Behavioral Indicators by County
Combined 3 Years

Comparison of Maternal Behavioral Indicators (2, 3)										
County	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Pregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	% (Rank)		Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Miami	371	77.9 (39)	0.8 (7)		323	32.5 (61)	220	15.5 (14)	195	58.5 (33)
Mitchell	121	91.7 (2)	0.0 (1)		110	34.5 (58)	78	* *	72	* *
Montgomery	774	84.0 (19)	4.7 (38)		869	38.7 (46)	530	14.3 (21)	540	66.7 (14)
Morris	69	* *	* *		67	* *	37	* *	36	* *
Nemaha	111	86.5 (13)	5.4 (42)		96	* *	59	* *	59	* *
Neosho	377	88.9 (10)	4.0 (37)		382	42.4 (30)	238	6.7 (38)	223	66.8 (13)
Ness	17	* *	* *		13	* *	10	* *	4	* *
Norton	79	* *	* *		68	* *	46	* *	41	* *
Osage	214	89.7 (7)	2.3 (21)		216	51.4 (7)	146	21.2 (3)	136	41.9 (42)
Osborne	48	* *	* *		43	* *	20	* *	27	* *
Ottawa	69	* *	* *		63	* *	39	* *	37	* *
Pawnee	144	81.9 (26)	2.8 (25)		148	57.4 (2)	90	* *	87	* *
Phillips	96	* *	* *		88	* *	56	* *	71	* *
Pottawatom	109	77.1 (41)	0.0 (1)		105	39.0 (44)	57	* *	58	* *
Pratt	194	78.9 (36)	6.7 (48)		188	47.3 (15)	126	7.9 (35)	116	64.7 (19)
Rawlins	12	* *	* *		27	* *	18	* *	19	* *
Reno	1,267	83.4 (21)	2.1 (18)		1,195	41.2 (37)	790	21.3 (2)	753	65.9 (18)
Republic	91	* *	* *		92	* *	56	* *	51	* *
Rice	184	76.1 (43)	1.6 (14)		181	40.9 (38)	117	14.5 (20)	104	43.3 (41)
Riley	1,938	83.6 (20)	0.6 (3)		2,178	36.3 (54)	1,425	16.0 (13)	1,358	63.4 (22)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6B
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Maternal Behavioral Indicators by County
Combined 3 Years

Comparison of Maternal Behavioral Indicators (2, 3)										
County	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	%	(Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Rooks	119	83.2 (22)	3.4	(31)	113	43.4 (26)	67	* *	68	* *
Rush	32	* *	* *		32	* *	17	* *	18	* *
Russell	116	90.5 (5)	0.9	(8)	105	51.4 (7)	65	* *	65	* *
Saline	1,209	74.0 (49)	4.8	(39)	1,205	33.9 (59)	802	5.2 (41)	775	57.0 (37)
Scott	127	78.7 (37)	8.7	(54)	122	36.9 (52)	69	* *	81	* *
Sedgwick	11,643	74.7 (47)	3.6	(33)	10,832	28.2 (63)	7,027	13.6 (22)	6,444	50.8 (40)
Seward	1,260	66.4 (59)	8.4	(52)	1,279	37.1 (50)	799	8.8 (33)	740	66.9 (12)
Shawnee	3,602	75.4 (46)	3.8	(34)	3,706	48.5 (13)	2,295	6.6 (40)	2,290	58.5 (33)
Sheridan	23	* *	* *		20	* *	12	* *	12	* *
Sherman	106	84.9 (18)	8.5	(53)	168	52.4 (5)	99	* *	108	74.1 (3)
Smith	42	* *	* *		52	* *	29	* *	37	* *
Stafford	103	75.7 (44)	3.9	(36)	105	62.9 (1)	68	* *	72	* *
Stanton	54	* *	* *		47	* *	25	* *	22	* *
Stevens	224	74.6 (48)	2.7	(24)	210	35.2 (57)	127	18.9 (6)	115	60.0 (31)
Sumner	398	82.7 (24)	3.5	(32)	362	38.1 (48)	241	16.6 (11)	209	66.0 (16)
Thomas	98	* *	* *		129	38.8 (45)	83	* *	92	* *
Trego	45	* *	* *		41	* *	24	* *	28	* *
Wabaunsee	55	* *	* *		46	* *	35	* *	31	* *
Wallace	4	* *	* *		2	* *	1	* *	2	* *
Washington	74	* *	* *		74	* *	42	* *	47	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6B
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Maternal Behavioral Indicators by County
Combined 3 Years

Comparison of Maternal Behavioral Indicators (2, 3)										
County	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Pregnancy (4)		During Pregnancy (5)	
	Number	%	%	%	Number	%	Number	%	Number	%
	(Rank)	(Rank)	(Rank)	(Rank)	(Rank)	(Rank)	(Rank)	(Rank)	(Rank)	(Rank)
Wichita	42	* *	* *	* *	40	* *	24	* *	24	* *
Wilson	260	85.8 (14)	1.5 (12)	1.5 (12)	254	42.9 (27)	151	15.2 (16)	156	62.8 (25)
Woodson	23	* *	* *	* *	20	* *	8	* *	9	* *
Wyandotte	5,808	66.5 (58)	6.6 (47)	6.6 (47)	5,566	37.0 (51)	3,457	11.0 (30)	3,286	61.5 (27)
Unknown Counties	3	* *	* *	* *	3	* *	3	* *	1	* *
Kansas	54,171	75.0	4.9	4.9	53,570	38.0	34,056	13.7	32,418	60.0

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7B
2007 - 2009 Pregnancy Nutrition Surveillance
Kansas
Comparison of Smoking Indicators by County
Combined 3 Years

(1)

Page:
Run Date:

67
7 / 9 / 10

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Allen	291	44.3 (38)	249	28.5 (41)	82	* *	250	37.2 (48)	247	32.8 (47)
Anderson	148	56.8 (62)	126	32.5 (50)	57	* *	127	27.6 (37)	128	22.7 (37)
Atchison	298	47.0 (44)	195	28.7 (42)	65	* *	264	28.0 (38)	202	24.8 (39)
Barber	115	48.7 (51)	96	* *	41	* *	96	* *	97	* *
Barton	803	37.6 (20)	714	23.7 (23)	222	29.3 (17)	657	11.1 (10)	716	5.6 (5)
Bourbon	373	50.1 (54)	311	28.9 (43)	113	26.5 (24)	307	34.5 (47)	312	34.0 (49)
Brown	274	42.3 (32)	208	25.5 (27)	66	* *	234	44.4 (53)	210	44.8 (56)
Butler	771	50.5 (55)	639	31.0 (49)	235	30.6 (13)	558	14.3 (17)	641	8.6 (13)
Chase	38	* *	34	* *	16	* *	30	* *	34	* *
Chautauqua	78	* *	52	* *	15	* *	68	* *	52	* *
Cherokee	587	39.2 (23)	489	26.8 (32)	138	20.3 (27)	448	31.7 (44)	503	30.2 (46)
Cheyenne	6	* *	6	* *	0	* *	4	* *	6	* *
Clark	2	* *	1	* *	0	* *	5	* *	3	* *
Clay	181	54.7 (59)	157	32.5 (50)	74	* *	154	22.1 (29)	159	15.7 (23)
Cloud	235	38.3 (21)	200	21.0 (19)	60	* *	193	6.7 (2)	202	2.0 (1)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7B
2007 - 2009 Pregnancy Nutrition Surveillance
Kansas
Comparison of Smoking Indicators by County
Combined 3 Years

(1)

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Coffey	65	* *	62	* *	21	* *	38	* *	63	* *
Comanche	22	* *	21	* *	8	* *	18	* *	21	* *
Cowley	981	44.5 (39)	802	30.3 (47)	297	26.6 (23)	810	19.4 (24)	808	18.8 (29)
Crawford	1,087	43.3 (35)	866	25.5 (27)	304	34.9 (8)	937	40.0 (51)	868	39.3 (52)
Decatur	30	* *	27	* *	7	* *	24	* *	27	* *
Dickinson	340	44.7 (40)	305	27.9 (39)	104	30.8 (12)	261	9.2 (7)	308	6.5 (6)
Doniphan	98	* *	62	* *	23	* *	76	* *	65	* *
Douglas	1,407	43.7 (36)	1,195	20.6 (18)	392	42.9 (3)	1,121	10.1 (9)	1,200	6.5 (6)
Edwards	69	* *	59	* *	15	* *	54	* *	60	* *
Elk	47	* *	41	* *	12	* *	35	* *	40	* *
Ellis	463	37.1 (19)	396	14.1 (9)	117	48.7 (1)	366	8.5 (4)	394	7.4 (9)
Ellsworth	91	* *	85	* *	29	* *	67	* *	87	* *
Finney	1,979	13.2 (5)	1,796	6.6 (4)	179	40.8 (4)	1,625	17.4 (21)	1,797	16.0 (24)
Ford	1,855	10.0 (4)	1,531	4.9 (3)	105	28.6 (18)	1,618	8.9 (5)	1,532	8.3 (11)
Franklin	504	46.0 (42)	444	27.3 (36)	136	31.6 (11)	378	14.8 (18)	446	10.5 (17)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7B
2007 - 2009 Pregnancy Nutrition Surveillance
Kansas
Comparison of Smoking Indicators by County
Combined 3 Years

(1)

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Geary	1,032	39.2 (23)	842	20.4 (17)	265	36.6 (5)	850	39.8 (50)	844	37.0 (51)
Gove	30	* *	28	* *	3	* *	18	* *	28	* *
Graham	50	* *	43	* *	22	* *	41	* *	44	* *
Grant	207	7.7 (2)	199	4.0 (2)	9	* *	152	12.5 (12)	199	12.6 (20)
Gray	154	9.7 (3)	129	7.0 (5)	7	* *	127	11.8 (11)	132	9.8 (15)
Greeley	31	* *	29	* *	6	* *	22	* *	29	* *
Greenwood	124	54.8 (60)	72	* *	31	* *	110	31.8 (45)	104	26.0 (41)
Hamilton	72	* *	63	* *	3	* *	55	* *	63	* *
Harper	129	48.8 (52)	85	* *	32	* *	109	22.0 (28)	84	* *
Harvey	773	35.1 (16)	658	22.5 (22)	189	30.2 (14)	638	5.8 (1)	662	4.1 (2)
Haskell	61	* *	56	* *	7	* *	53	* *	56	* *
Hodgeman	37	* *	34	* *	11	* *	25	* *	33	* *
Jackson	212	48.1 (50)	184	29.3 (45)	67	* *	176	19.9 (25)	187	18.2 (27)
Jefferson	70	* *	28	* *	9	* *	69	* *	32	* *
Jewell	39	* *	37	* *	11	* *	32	* *	37	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7B
2007 - 2009 Pregnancy Nutrition Surveillance
Kansas
Comparison of Smoking Indicators by County
Combined 3 Years

(1)

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Johnson	5,398	24.9 (8)	4,746	10.7 (6)	812	44.5 (2)	4,213	18.4 (23)	4,761	18.3 (28)
Kearny	60	* *	57	* *	8	* *	49	* *	57	* *
Kingman	146	40.4 (27)	118	19.5 (16)	29	* *	139	39.6 (49)	133	36.1 (50)
Kiowa	30	* *	21	* *	4	* *	25	* *	23	* *
Labette	521	38.6 (22)	455	25.1 (26)	133	15.8 (29)	443	28.0 (38)	467	27.0 (43)
Lane	18	* *	9	* *	1	* *	17	* *	10	* *
Leavenworth	1,010	47.6 (46)	853	27.1 (35)	281	29.9 (15)	742	42.7 (52)	862	40.5 (53)
Lincoln	56	* *	43	* *	16	* *	42	* *	43	* *
Linn	185	56.2 (61)	166	40.4 (57)	76	* *	155	14.2 (16)	168	10.7 (19)
Logan	36	* *	29	* *	9	* *	30	* *	29	* *
Lyon	1,011	29.1 (12)	909	15.3 (10)	188	35.6 (6)	778	13.2 (14)	911	10.6 (18)
Marion	400	40.3 (26)	367	26.2 (30)	120	26.7 (22)	309	7.1 (3)	370	4.1 (2)
Marshall	135	25.9 (10)	119	19.3 (15)	22	* *	93	* *	120	5.0 (4)
Mcperson	166	50.6 (56)	132	32.6 (52)	46	* *	128	52.3 (56)	137	49.6 (57)
Meade	125	30.4 (14)	104	18.3 (14)	23	* *	106	22.6 (34)	106	19.8 (33)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7B
2007 - 2009 Pregnancy Nutrition Surveillance
Kansas
Comparison of Smoking Indicators by County
Combined 3 Years

(1)

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Miami	359	47.9 (48)	287	30.3 (47)	99	* *	265	21.5 (27)	286	16.8 (26)
Mitchell	127	39.4 (25)	119	26.9 (33)	40	* *	101	8.9 (5)	121	9.1 (14)
Montgomery	975	41.0 (30)	815	29.2 (44)	258	16.7 (28)	804	29.9 (43)	827	28.1 (44)
Morris	78	* *	68	* *	24	* *	61	* *	68	* *
Nemaha	113	47.8 (47)	101	26.7 (31)	41	* *	88	* *	100	42.0 (55)
Neosho	393	43.0 (33)	292	25.7 (29)	102	29.4 (16)	335	24.2 (35)	297	21.5 (35)
Ness	17	* *	15	* *	5	* *	11	* *	15	* *
Norton	84	* *	73	* *	30	* *	64	* *	74	* *
Osage	252	46.0 (42)	210	29.5 (46)	73	* *	205	28.3 (40)	211	26.1 (42)
Osborne	52	* *	47	* *	18	* *	40	* *	47	* *
Ottawa	71	* *	58	* *	20	* *	58	* *	60	* *
Pawnee	161	47.2 (45)	138	27.5 (37)	49	* *	136	19.9 (25)	138	18.8 (29)
Phillips	81	* *	54	* *	18	* *	62	* *	54	* *
Pottawatom	129	30.2 (13)	113	21.2 (20)	29	* *	97	* *	116	6.9 (8)
Pratt	223	41.3 (31)	167	28.1 (40)	60	* *	165	32.7 (46)	187	28.9 (45)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7B
2007 - 2009 Pregnancy Nutrition Surveillance
Kansas
Comparison of Smoking Indicators by County
Combined 3 Years

(1)

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Rawlins	30	* *	27	* *	6	* *	26	* *	27	* *
Reno	1,401	45.9 (41)	1,214	24.9 (25)	424	34.4 (9)	1,107	17.5 (22)	1,222	16.4 (25)
Republic	100	43.0 (33)	85	* *	29	* *	82	* *	87	* *
Rice	206	49.0 (53)	181	33.7 (54)	68	* *	170	22.4 (31)	183	19.7 (32)
Riley	2,550	27.7 (11)	1,887	16.2 (12)	353	28.0 (20)	1,976	15.6 (19)	2,036	14.3 (21)
Rooks	127	59.1 (63)	107	38.3 (56)	53	* *	102	25.5 (36)	111	33.3 (48)
Rush	34	* *	31	* *	14	* *	31	* *	31	* *
Russell	125	48.0 (49)	112	42.9 (58)	45	* *	98	* *	114	50.9 (59)
Saline	1,303	44.2 (37)	1,029	26.9 (33)	348	27.3 (21)	1,040	13.2 (14)	1,034	10.3 (16)
Scott	136	25.7 (9)	115	15.7 (11)	25	* *	119	29.4 (42)	119	22.7 (37)
Sedgwick	12,530	32.2 (15)	11,365	16.4 (13)	2,701	35.4 (7)	9,927	9.7 (8)	11,365	8.3 (11)
Seward	1,428	7.5 (1)	1,278	3.5 (1)	69	* *	1,188	15.7 (20)	1,292	14.4 (22)
Shawnee	3,990	36.4 (17)	3,164	22.3 (21)	856	26.2 (26)	3,331	13.0 (13)	3,177	7.6 (10)
Sheridan	22	* *	19	* *	2	* *	18	* *	19	* *
Sherman	183	40.4 (27)	155	27.7 (38)	54	* *	156	22.4 (31)	154	22.1 (36)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7B
2007 - 2009 Pregnancy Nutrition Surveillance
Kansas
Comparison of Smoking Indicators by County
Combined 3 Years

(1)

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Smith	54	* *	36	* *	11	* *	49	* *	39	* *
Stafford	114	40.4 (27)	93	* *	33	* *	96	* *	95	* *
Stanton	54	* *	50	* *	3	* *	41	* *	51	* *
Stevens	247	21.9 (7)	214	11.2 (7)	36	* *	192	22.4 (31)	217	18.9 (31)
Sumner	418	53.6 (58)	357	33.9 (55)	147	28.6 (18)	316	49.1 (55)	362	50.0 (58)
Thomas	156	36.5 (18)	134	24.6 (24)	33	* *	117	29.1 (41)	135	25.9 (40)
Trego	46	* *	33	* *	13	* *	39	* *	35	* *
Wabaunsee	61	* *	58	* *	17	* *	41	* *	58	* *
Wallace	2	* *	2	* *	1	* *	2	* *	2	* *
Washington	83	* *	72	* *	17	* *	67	* *	71	* *
Wichita	46	* *	37	* *	5	* *	37	* *	37	* *
Wilson	276	53.3 (57)	247	33.6 (53)	113	32.7 (10)	239	45.6 (54)	250	41.6 (54)
Woodson	24	* *	21	* *	6	* *	18	* *	22	* *
Wyandotte	6,305	21.6 (6)	5,597	12.7 (8)	906	26.3 (25)	5,183	22.2 (30)	5,641	21.0 (34)
Unknown Counties	3	* *	3	* *	1	* *	1	* *	3	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7B
2007 - 2009 Pregnancy Nutrition Surveillance
Kansas
Comparison of Smoking Indicators by County
Combined 3 Years

(1)

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Kansas	60,725	32.6	52,064	18.3	12,628	31.8	48,870	18.4	52,549	16.3

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8B
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Infant Health Indicators by County
Combined 3 Years

Comparison of Infant Health Indicators (2, 3)										
County	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)		Number	%	Number	%	Number	%
	Number	% (Rank)	% (Rank)	Number						
Allen	274	9.5 (59)	4.0 (6)	239	12.6 (48)	207	5.3 (44)	273	52.4 (59)	
Anderson	132	9.1 (52)	5.3 (19)	122	9.0 (17)	101	7.9 (50)	137	53.3 (58)	
Atchison	361	6.9 (26)	5.5 (20)	328	8.5 (12)	296	3.4 (22)	364	52.2 (60)	
Barber	93	* * * *		81	* *	65	* *	92	* *	
Barton	693	6.5 (20)	4.6 (10)	607	12.2 (45)	529	3.0 (12)	675	73.5 (14)	
Bourbon	343	7.6 (33)	5.2 (18)	293	11.6 (42)	257	3.9 (33)	339	58.7 (53)	
Brown	223	7.2 (29)	7.2 (45)	231	7.8 (8)	195	3.6 (27)	235	50.6 (62)	
Butler	727	8.1 (41)	5.6 (21)	575	10.4 (33)	509	3.7 (28)	700	72.1 (20)	
Chase	33	* * * *		29	* *	27	* *	32	* *	
Chautauqua	74	* * * *		72	* *	63	* *	73	* *	
Cherokee	549	5.8 (12)	7.7 (53)	446	9.9 (24)	402	2.2 (6)	525	59.6 (51)	
Cheyenne	5	* * * *		4	* *	4	* *	5	* *	
Clark	7	* * * *		5	* *	4	* *	6	* *	
Clay	165	7.3 (30)	3.6 (4)	140	13.6 (50)	121	3.3 (20)	162	62.3 (44)	

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8B
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Infant Health Indicators by County
Combined 3 Years

Comparison of Infant Health Indicators (2, 3)										
County	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)							
	Number	% (Rank)	% (Rank)		Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Cloud	206	8.3 (44)	2.4 (1)		177	15.3 (55)	149	4.0 (35)	199	66.8 (36)
Coffey	61	* *	* *		41	* *	38	* *	62	* *
Comanche	22	* *	* *		18	* *	15	* *	22	* *
Cowley	894	7.8 (36)	7.3 (46)		771	10.9 (39)	675	4.9 (42)	875	58.3 (54)
Crawford	964	9.3 (55)	4.9 (12)		870	12.1 (44)	752	4.5 (40)	958	59.0 (52)
Decatur	26	* *	* *		25	* *	24	* *	25	* *
Dickinson	301	6.6 (24)	10.0 (61)		260	8.1 (10)	236	3.0 (12)	301	73.4 (15)
Doniphan	104	5.8 (12)	5.8 (26)		87	* *	78	* *	102	52.0 (61)
Douglas	1,234	6.1 (14)	6.6 (32)		1,060	10.6 (36)	933	3.0 (12)	1,188	77.2 (8)
Edwards	55	* *	* *		50	* *	40	* *	56	* *
Elk	45	* *	* *		37	* *	31	* *	43	* *
Ellis	386	5.7 (10)	5.7 (23)		350	8.3 (11)	300	1.7 (3)	395	74.9 (12)
Ellsworth	89	* *	* *		69	* *	61	* *	86	* *
Finney	1,799	6.4 (17)	6.6 (32)		1,587	10.8 (38)	1,388	2.8 (9)	1,347	74.8 (13)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8B
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Infant Health Indicators by County
Combined 3 Years

Comparison of Infant Health Indicators (2, 3)										
County	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)							
	Number	% (Rank)	% (Rank)		Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Ford	1,658	5.1 (5)	5.7 (23)		1,519	10.4 (33)	1,335	1.9 (4)	1,467	72.1 (20)
Franklin	438	6.2 (16)	5.0 (14)		350	10.3 (30)	311	3.2 (18)	425	60.7 (48)
Gearly	881	7.4 (32)	6.6 (32)		755	8.7 (13)	686	3.5 (24)	826	69.5 (29)
Gove	28	* * *	* *		20	* *	18	* *	26	* *
Graham	39	* * *	* *		36	* *	31	* *	43	* *
Grant	196	8.7 (49)	7.7 (53)		158	9.5 (20)	140	4.3 (38)	148	70.3 (27)
Gray	114	6.1 (14)	7.9 (57)		111	11.7 (43)	89	* *	124	87.1 (1)
Greeley	28	* * *	* *		24	* *	23	* *	27	* *
Greenwood	121	6.6 (24)	3.3 (2)		109	10.1 (26)	97	* *	116	60.3 (50)
Hamilton	59	* * *	* *		51	* *	46	* *	47	* *
Harper	107	6.5 (20)	7.5 (47)		109	10.1 (26)	85	* *	121	76.9 (9)
Harvey	668	7.3 (30)	7.0 (42)		604	12.3 (46)	518	3.3 (20)	680	78.8 (4)
Haskell	56	* * *	* *		51	* *	47	* *	39	* *
Hodgeman	38	* * *	* *		28	* *	25	* *	36	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8B
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Infant Health Indicators by County
Combined 3 Years

Comparison of Infant Health Indicators (2, 3)										
County	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)		Number	%	Number	%	Number	%
	Number	% (Rank)	% (Rank)	Number						
Jackson	196	9.2 (53)	9.7 (60)		167	14.4 (52)	143	5.6 (46)	190	78.4 (5)
Jefferson	172	8.1 (41)	7.6 (51)		138	16.7 (56)	114	5.3 (44)	170	61.8 (46)
Jewell	38	* * * *			31	* *	26	* *	34	* *
Johnson	4,628	6.5 (20)	7.5 (47)		4,066	8.0 (9)	3,551	3.0 (12)	4,612	79.2 (3)
Kearny	56	* * * *			50	* *	43	* *	40	* *
Kingman	106	4.7 (1)	7.5 (47)		120	9.2 (18)	85	* *	134	80.6 (2)
Kiowa	28	* * * *			27	* *	23	* *	27	* *
Labette	479	6.5 (20)	7.5 (47)		421	10.5 (35)	377	3.7 (28)	456	54.6 (57)
Lane	28	* * * *			27	* *	22	* *	29	* *
Leavenworth	862	9.4 (57)	5.0 (14)		713	9.4 (19)	631	4.8 (41)	892	63.2 (43)
Lincoln	54	* * * *			41	* *	39	* *	51	* *
Linn	166	8.4 (47)	4.8 (11)		150	10.0 (25)	133	6.8 (49)	163	68.7 (32)
Logan	27	* * * *			25	* *	23	* *	26	* *
Lyon	908	6.9 (26)	6.5 (30)		739	10.1 (26)	663	3.0 (12)	869	71.6 (25)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8B
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Infant Health Indicators by County
Combined 3 Years

Comparison of Infant Health Indicators (2, 3)										
County	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)							
	Number	% (Rank)	% (Rank)		Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Marion	361	5.0 (4)	6.9 (38)		297	7.7 (7)	271	1.1 (2)	358	72.1 (20)
Marshall	110	5.5 (8)	6.4 (29)		88	* *	80	* *	110	77.3 (7)
Mcpherson	138	9.4 (57)	5.1 (17)		113	10.6 (36)	99	* *	140	56.4 (55)
Meade	105	5.7 (10)	6.7 (35)		95	* *	77	* *	106	76.4 (11)
Miami	370	7.6 (33)	5.9 (27)		290	9.7 (22)	260	5.8 (47)	322	72.7 (17)
Mitchell	104	8.7 (49)	6.7 (35)		100	7.0 (4)	83	* *	111	65.8 (38)
Montgomery	886	7.9 (38)	3.5 (3)		760	11.4 (41)	670	3.4 (22)	853	60.5 (49)
Morris	69	* *	* *		58	* *	50	* *	69	* *
Nemaha	89	* *	* *		91	* *	63	* *	109	64.2 (39)
Neosho	302	7.9 (38)	6.3 (28)		355	12.7 (49)	231	3.5 (24)	407	55.5 (56)
Ness	16	* *	* *		11	* *	9	* *	16	* *
Norton	75	* *	* *		61	* *	47	* *	72	* *
Osage	235	6.4 (17)	4.3 (8)		201	14.9 (54)	167	3.0 (12)	228	61.0 (47)
Osborne	46	* *	* *		38	* *	34	* *	45	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8B
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Infant Health Indicators by County
Combined 3 Years

Comparison of Infant Health Indicators (2, 3)									
County	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)	High (5)		Number	%	Number	%	Number	%
	Number	% (Rank)	% (Rank)						
Ottawa	66	* * * *		57	* *	52	* *	62	* *
Pawnee	137	8.8 (51)	4.4 (9)	127	8.7 (13)	116	2.6 (7)	134	72.4 (19)
Phillips	94	* * * *		77	* *	62	* *	92	* *
Pottawatom	116	7.8 (36)	6.9 (38)	94	* *	85	* *	113	72.6 (18)
Pratt	187	6.4 (17)	9.1 (58)	152	5.9 (3)	143	3.5 (24)	171	68.4 (34)
Rawlins	28	* * * *		24	* *	22	* *	27	* *
Reno	1,247	7.1 (28)	7.1 (43)	1,048	10.3 (30)	930	3.2 (18)	1,218	63.3 (42)
Republic	85	* * * *		79	* *	65	* *	89	* *
Rice	181	8.3 (44)	7.7 (53)	157	14.6 (53)	133	3.8 (32)	174	71.8 (24)
Riley	2,166	5.6 (9)	7.8 (56)	1,734	7.0 (4)	1,604	2.6 (7)	2,109	76.9 (9)
Rooks	121	8.3 (44)	5.0 (14)	106	4.7 (2)	101	5.0 (43)	118	72.0 (23)
Rush	31	* * * *		28	* *	24	* *	29	* *
Russell	109	9.2 (53)	3.7 (5)	93	* *	81	* *	103	64.1 (40)
Saline	1,283	9.3 (55)	6.5 (30)	1,095	11.0 (40)	957	4.4 (39)	1,285	62.3 (44)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8B
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Infant Health Indicators by County
Combined 3 Years

Comparison of Infant Health Indicators (2, 3)											
County	Birthweight				Preterm (6)			Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)								
	Number	% (Rank)	% (Rank)		Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	
Scott	118	5.1 (5)	4.2 (7)		108	3.7 (1)	102	2.9 (11)	121	69.4 (30)	
Sedgwick	11,520	8.4 (47)	6.9 (38)		9,833	8.8 (15)	8,896	4.0 (35)	10,628	66.7 (37)	
Seward	1,270	5.2 (7)	7.1 (43)		1,148	9.7 (22)	998	2.8 (9)	1,298	69.4 (30)	
Shawnee	3,609	7.7 (35)	5.6 (21)		3,255	12.3 (46)	2,755	3.7 (28)	3,667	67.9 (35)	
Sheridan	24	* * *	* *		18	* *	15	* *	24	* *	
Sherman	163	11.0 (61)	4.9 (12)		145	7.6 (6)	134	6.7 (48)	155	69.7 (28)	
Smith	51	* * *	* *		43	* *	35	* *	52	* *	
Stafford	94	* * *	* *		90	* *	76	* *	89	* *	
Stanton	50	* * *	* *		46	* *	40	* *	35	* *	
Stevens	143	4.9 (3)	9.1 (58)		181	8.8 (15)	104	1.0 (1)	217	77.4 (6)	
Sumner	393	8.1 (41)	7.6 (51)		324	10.2 (29)	286	4.2 (37)	375	68.5 (33)	
Thomas	145	4.8 (2)	6.9 (38)		115	9.6 (21)	104	1.9 (4)	139	73.4 (15)	
Trego	44	* * *	* *		38	* *	35	* *	43	* *	
Wabaunsee	58	* * *	* *		44	* *	41	* *	55	* *	

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8B
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Kansas
Comparison of Infant Health Indicators by County
Combined 3 Years

Comparison of Infant Health Indicators (2, 3)										
County	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)							
	Number	% (Rank)	% (Rank)		Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Wallace	4	* * * *	* *		2	* *	2	* *	4	* *
Washington	62	* * * *	* *		65	* *	50	* *	71	* *
Wichita	42	* * * *	* *		36	* *	27	* *	42	* *
Wilson	261	9.6 (60)	5.7 (23)		240	14.2 (51)	205	3.9 (33)	235	63.8 (41)
Woodson	23	* * * *	* *		19	* *	15	* *	22	* *
Wyandotte	5,635	8.0 (40)	6.8 (37)		4,952	10.3 (30)	4,358	3.7 (28)	5,555	70.7 (26)
Unknown Counties	3	* * * *	* *		3	* *	3	* *	3	* *
Kansas	54,513	7.4	6.6		47,348	10.0	41,621	3.5	52,595	69.2

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

